



# Characteristics of two tsunamis generated by successive $M_w$ 7.4 and $M_w$ 8.1 earthquakes in Kermadec Islands on March 4, 2021

Yuchen Wang<sup>1,2</sup>, Mohammad Heidarzadeh<sup>3</sup>, Kenji Satake<sup>1</sup>, Gui Hu<sup>4</sup>

<sup>1</sup>Earthquake Research Institute, The University of Tokyo, Tokyo, 113-0032, Japan

5 <sup>2</sup>Now at Japan Agency for Marine–Earth Science and Technology, Kanazawa, Yokohama, 236-0001, Japan

<sup>3</sup>Department of Civil and Environmental Engineering, Brunel University London, Uxbridge, UB8 3PH, UK

<sup>4</sup>Guangdong Provincial Key Laboratory of Geodynamics and Geohazards, School of Earth Sciences and Engineering, Sun Yat-Sen University, Guangzhou, 510275, China

Correspondence to: Yuchen Wang (ywang@jamstec.go.jp)

10 **Abstract.** On March 4, 2021, two tsunamigenic earthquakes ( $M_w$  7.4 and  $M_w$  8.1) occurred successively within 2 h in Kermadec Islands. We examined sea level records at tide gauges located at ~100 km to ~ 2,000 km from the epicenters, conducted Fourier and Wavelet analyses as well as numerical modelling of both tsunamis. Fourier analyses indicated that the energy of the first tsunami is mainly distributed in the period range of 5–17 min, whereas it is 8–28 min for the second tsunami. Wavelet plots showed that the oscillation of the first tsunami continued even after the arrival of the second tsunami.  
15 As the epicenters of two earthquakes are close (~ 55 km), we reconstructed the source spectrum of the second tsunami by using the first tsunami as the empirical Green's function. The main spectral peaks are 25.6 min, 16.0 min, and 9.8 min. The results are similar to those calculated using tsunami/background ratio method and also consistent with source models.

## 1 Introduction

The Kermadec Islands are an island arc in the southwestern Pacific Ocean, formed at the convergent boundary where  
20 the Pacific Plate subducts under the Indo-Australian Plate (Figure 1) (Billen et al., 2003). The Kermadec Trench, which accommodates westward subduction of the Pacific Plate beneath the active Kermadec volcanic arc, is identified as the key region for concern regarding seismic/tsunami hazards in New Zealand (e.g., Power et al., 2012). The historical records of earthquakes in Kermadec Islands are short but there are three great earthquakes over the 20th century: the 1917 ( $M_w$  8.6; Morgan, 1918), the 1976 ( $M_w$  8.0; Wyss et al., 1984), and the 1986 earthquakes ( $M_w$  7.9; Lundgren et al., 1989).

25 On March 4, 2021, two earthquakes occurred successively in Kermadec Islands. The first event ( $M_w$  7.4; foreshock) occurred at 17:41:23 (UTC) whose epicenter was at 29.677 S, 177.840 W, south of Raoul Island in the Kermadec Islands region with a depth of 43.0 km (United States Geological Survey (USGS);  
<https://earthquake.usgs.gov/earthquakes/eventpage/us7000dfk3/executive>) (Figure 1). The earthquake generated a small tsunami. The second earthquake ( $M_w$  8.1; mainshock) occurred at 19:28:33 (UTC), approximately two hours after the  
30 foreshock. The epicenter was located at 29.723 S, 177.279 W with a depth of 28.9 km (USGS). The epicenters of these two



35 successive tsunamigenic earthquakes are very close to each other (~55 km; Figure 1) and their focal mechanisms are similar; both are thrust earthquakes. The National Emergency Management Agency, New Zealand, issued a tsunami warning for coastal areas of the North Island after the  $M_w$  8.1 earthquake. The tsunami propagated across the Pacific Ocean and reached South America. No casualties were reported. The situation of these two consequent tsunamigenic earthquakes resembles the earthquake events also in Kermadec Islands on January 14, 1976, where two great earthquakes ( $M_w$  7.8 and  $M_w$  8.0) occurred approximately within one hour (Power et al., 2012). The mainshock of the 1976 events ( $M_w$  8.0) generated a moderate tsunami recorded by tide gauges.

40 The occurrence of two successive earthquakes provides us with a rare opportunity to study their source characteristics by different methods. The spectra of tsunami waveforms at tide gauges contain the effects of source, propagation path, and local topography (Rabinovich, 1997; Heidarzadeh et al., 2016; Cortés et al., 2017). For a single event, it is a common practice to reconstruct tsunami source spectrum from tide gauge records by calculating the ratio of tsunami spectra to background spectra (Rabinovich, 1997). This is based on the assumption that the effects of transmission path on the characteristics of tsunami spectra is small compared to local topographic effects when the tsunami source is not too far away from the observation station (Miller, 1972; Rabinovich, 1997). This assumption can be readily made for two successive tsunami events in Kermadec Islands. If the sources of two earthquakes are close to each other, we can adopt the method of empirical Green's function (EGF) to reconstruct tsunami source spectrum (Heidarzadeh et al., 2016). The smaller tsunami event is adopted as the EGF for the larger one. The spectral deconvolution separates the effects of propagation path and local topography around the tide gauge and gives the source spectrum of the larger tsunami. Heidarzadeh et al. (2016) successfully applied the EGF method to the 2015 ( $M_w$  7.0) and 2013 ( $M_w$  8.0) earthquakes in the Solomon Islands.

50 Here, we studied the characteristics of tsunamis generated by the two successive Kermadec Islands earthquakes and calculated the source spectrum of the tsunami generated by the mainshock (hereafter, the second tsunami). Fourier analysis was applied to the sea level records at 15 tide gauges. Wavelet analysis was adopted to examine the temporal changes of the dominant spectral peaks. Finally, we used two different methods, tsunami/background ratio method and EGF method, to reconstruct tsunami source spectra. We compared two alternative methods of determining tsunami source spectra and also compared their results with USGS source models. This is a unique and rare incident and thus the data and analyses would greatly help to further understand tsunami generation and propagation.




Figure 1: Bathymetry map of the southwestern Pacific region and the epicenters of two successive earthquakes in Kermadec Islands on March 4, 2021. Pink and red beachballs represent the focal mechanisms (according to the Global Centroid Moment



60 **Tensor Project; <https://www.globalcmt.org/CMTsearch.html> of the  $M_w$  7.4 and  $M_w$  8.1 earthquakes, respectively. Green squares indicate tide gauges. The travel time of the second tsunami is marked by grey dashed contours with an interval of 0.5 h.**

## 2 Data and Method

### 2.1 Tsunami Data

We collected sea level records of 15 tide gauges from the Intergovernmental Oceanographic Commission  
65 (<http://www.ioc-sealevelmonitoring.org/list.php>). These stations are located at distances between ~100 km and ~ 2,000 km from the epicenters (Figure 1). The sampling rate is 60 s. First, we conducted quality control of these data, removing spikes and filling short gaps by linear interpolation. Then, we applied a second-order high-pass filter with the corner frequency of 0.00014 Hz (7,200 s) to remove the tidal components (Figure 2) (Heidarzadeh and Satake, 2013). The time series between 17:00:00 (UTC), March 4 and 05:00:00 (UTC), March 5 was selected for analysis. However, as the Raoul Island Fishing  
70 Rock (RIFR) tide gauge (Figure 1) lost its data communication during the mainshock, we only used its records before 17:41:00 (UTC) (Figure 2).

### 2.2 Spectral Analyses


Two types of spectral analyses were conducted to tide gauge records: Fourier analysis and wavelet (frequency-time) analysis. For Fourier analysis, a Hanning window with 40% overlaps was applied following the method of Heidarzadeh and  
75 Mulia (2021). The tsunami generated by the foreshock (hereafter, the first tsunami) was clearly recorded at eight tide gauges: North Cape (NC), Great Barrier Island (GBI), East Cape (EC), Suva Viti Levu (SVL), Lenakel, Quinne, Kingston Jetty Norfolk (KJN), and RIFR, but it is difficult to accurately distinguish the arrival times at these stations. We used the 2 h segment before the arrival of the second tsunami as the input for spectral analyses. For RIFR, we utilized the 2 h segment before the time it lost connection. The second tsunami generated by the mainshock was clearly recorded at all the tide gauges  
80 except for RIFR. We selected a 2 h time segment after the tsunami arrival as the input for Fourier analysis. Additionally, background spectra were calculated using the 2 h records before the arrival time of the first tsunami at the same stations. We ensured that there were no storms or other atmospheric events at the time period of the background signals, so the background spectra could exclusively reflect the frequency response of local topography. Tidal components were removed by applying a high-pass filter in a similar way to preparation of the tsunami records.

85 Wavelet analysis was adopted to study the temporal changes of the dominant spectral peaks and the superposition of two tsunamis. We applied the wavelet package provided by Torrence and Compo (1998), which uses the Morlet wavelet mother function. We only conducted wavelet analysis to seven tide gauges that clearly recorded both tsunamis. The waveform segments for wavelet analyses are the same as those used for Fourier analyses.



### 2.3 Earthquake Slip Models and Tsunami Numerical Simulation

90 We used numerical simulation to study the propagation paths of the two tsunamis. For the initial condition, the finite  
fault models provided by the USGS were adopted (Slip model of the first event at:  
<https://earthquake.usgs.gov/earthquakes/eventpage/us7000dfk3/finite-fault>; Slip model of the second event at:  
<https://earthquake.usgs.gov/earthquakes/eventpage/us7000dflf/finite-fault>). The source model of the first ( $M_w$  7.4)  
95 earthquake has a rectangular dimension of 80 km (length)  $\times$  80 km (width). The strike and dip angles are  $196^\circ$  and  $32^\circ$ ,  
respectively. The spatial intervals of sub-faults are 4 km  $\times$  4 km. The source model of the second ( $M_w$  8.1) earthquake has a  
rectangular dimension of 240 km (length)  $\times$  190 km (width) with strike and dip angles of  $210^\circ$  and  $16^\circ$ , respectively. The  
spatial intervals of sub-faults are 10 km  $\times$  10 km. We used Okada's dislocation model (Okada, 1985) to compute the  
seafloor deformation and used it as the initial condition for tsunami propagation modeling. The bathymetric grids were  
obtained from the General Bathymetric Chart of the Oceans and resampled to 0.9 arc-min. The *JAGURS* numerical package  
100 (Baba et al., 2015) was adopted to simulate the tsunami propagation based on linear long-wave equations. The time step for  
numerical simulations is 1.0 s. We simulated two tsunamis separately up to 12 h after the earthquake origin time. Tsunami  
travel time (TTT) calculation was performed for the second tsunami by the TTT software of Geoware (2011).



105 **Figure 2:** Sea level change recorded by tide gauges showing two tsunamis. The pink and red vertical lines show the origin times of the  $M_w$  7.4 and  $M_w$  8.1 earthquakes, respectively. The arrival times of the first and second tsunamis are marked by pink and red arrows, respectively.

### 3 Tsunami Waveform Analysis

110 Two tsunamis arrive successively at 15 tide gauges (Figure 2). The first tsunami arrived at the RIFR station 11 min after the foreshock. The first peak and the maximum amplitudes are 0.21 m and 0.34 m, respectively. It reached EC at 117 min after the earthquake with a maximum amplitude of 0.09 m. It is noteworthy that the Lenakel tide gauge, located in the direction parallel to the short axis of the earthquake source fault, recorded a maximum amplitude of 0.31 m. It is the largest value among all stations except for the RIFR. The first tsunami was also recorded by KJN with a maximum amplitude of 0.21 m.



The arrival times of the second tsunami are generally consistent with the results of the TTT calculations (Figure 1; 115 dashed contours) with clear large amplitude signals. It arrived at EC at 20:59 (UTC); 91 min after the mainshock with an amplitude of 0.09 m. Yet the maximum amplitude at this station (0.17 m) appeared in a later time (approximately one hour after the first arrival) (Figure 2). The largest waves were also observed in later time in other stations such as NC, GBI, Nukualofa, Owenga, Ouinne, and Hienghene (Figure 2). Such late arrival of largest tsunami wave was reported for other tsunamis in the past such as for the 2004 Indian Ocean tsunami (Rabinovich and Thomson, 2007) and the 2020 Aegean Sea 120 tsunami (Heidarzadeh et al., 2021). The largest tsunami amplitude (0.64 m) was recorded by KJN, which is also located in the direction parallel to the short axis of the fault. Besides, the second tsunami has a longer wavelength than the first tsunami, leading to longer tsunami periods. At stations without clear signal of the first tsunami (e.g., Lautoka), the tsunami waveforms are dominant by the long-period components which are generated by the second tsunami. However, the stations that recorded both events show the superposition of two successive tsunamis. For example, the waveform of the second tsunami at EC is 125 mixed up with the short-period components in the few hours after its arrival (21:00–24:00), likely caused by the oscillation of the first tsunami (Figure 2). After 24:00, the tsunami waveforms mainly present long-period components, possibly due to the dissipation of the short-period waves of the first tsunami. Similar patterns were also be observed at Ouinne (Figure 2).

#### 4 Spectral Analyses


Figures 3 presents the results of Fourier analyses for two tsunamis and background signals. The gap between the 130 tsunami and background spectra is attributed to the tsunami energy. The periods of the tsunami spectral peaks generally contain the effects of tsunami source, propagation path, and local topography. According to Figure 3, the second tsunami has larger energy (spectral power) than the first tsunami due to the larger magnitude of the second earthquake. The background spectra are smoother than tsunami spectra at most stations. At most stations, the peak periods of the first tsunami are distributed in the range of 5–17 min, whereas the dominant periods range for the second tsunami is approximately 8–28 min. 135 In other words, the second tsunami has generated longer-period waves, which is natural as the source dimension of the second earthquake (240 km × 190 km) is much larger than that of the first earthquake (80 km × 80 km).

Wavelet analyses reveal the variations of dominant tsunami peak periods over time (Figure 4). At EC, Lenakel, and KJN, the arrivals of two successive tsunamis can be clearly identified on the wavelet plots. The second tsunami has larger energy levels and longer-period waves than the first tsunami (Figure 4), which is consistent with the results of Fourier 140 analyses. At other stations such as GBI, the arrival time of the first tsunami is not clear. At most tide gauges, there are two oscillation patterns visible on the wavelet plots. One oscillation pattern has dominant periods in the range of 5–17 min. The other one has dominant periods over 15 min and up to approximately 30 min, which occurs after the first pattern. These two period bands on the wavelet plots reflect the first and the second tsunamis. For example, the wavelet plot of EC reveals a persisting wave in the period range of 5–11 min. The oscillation begins at approximately 2 h after the  $M_w$  7.4 earthquake,



145 which corresponds to the arrival time of the first tsunami. Another oscillation with longer dominant periods begins at  
approximately 2 h after the  $M_w$  8.1 earthquake, which is consistent with the arrival time of the second tsunami. The  
oscillation lasts for more than 5 h. It is noteworthy that after the arrival of the second tsunami, the oscillation pattern with  
short dominant periods still exists and becomes even stronger sometimes, indicating the contribution of both tsunamis at EC.  
Two oscillation patterns simultaneously exist for several hours and almost simultaneously diminish. The wavelet plot of KJN  
150 shows similar patterns to that of EC. Although there are also two oscillation patterns on the wavelet plot of Lenakel, the  
long-period oscillation only lasts for less than 5 h. To the contrary, at GBI, the oscillation pattern with long dominant periods  
lasts for approximately 3 h, whereas the oscillation pattern with short dominant periods lasts for more than 6 h. At Ouinne, a  
persisting wave in the period band of 20–30 min is visible with high energy. Another oscillation pattern (6–12 min)  
diminishes earlier, which explains the reason why the tsunami waveforms at Ouinne has less short-period components in the  
155 later phases (Figure 2). Nevertheless, different from other stations, we cannot find a long-lasting wave at SVL. There is only  
one oscillation period band (20–40 min) and it diminishes rapidly.





**Figure 3:** Fourier analyses for tsunamis generated by two successive earthquakes ( $M_w$  7.4 and  $M_w$  8.1) in Kermadec Islands. Pink and red curves represent the spectra of the first tsunami and the second tsunami, respectively. The background spectra (black curves) are also plotted for comparison.

160




Figure 4: Wavelet (frequency-time) analyses for tsunamis generated by two successive earthquakes ( $M_w$  7.4 and  $M_w$  8.1) in Kermadec Islands. The colormap shows levels of spectral energy at different times and periods. For guidance, we marked the



165 **dominant periods of two tsunamis by pink ( $M_w$  7.4) and red ( $M_w$  8.1) rectangles at East Cape. The pink and red vertical lines show the origin times of the  $M_w$  7.4 and  $M_w$  8.1 earthquakes, respectively.**

## 5 Reconstructing the Tsunami Source Spectrum

A tsunami source spectrum reveals the earthquake source characteristics without the effects of tsunami propagation path or local topography. In our study, the epicenters of two earthquakes are close to each other (~55 km; Figure 1); the earthquakes are of similar mechanism (both thrust events), and the propagation paths are similar (Figures 5a and 5b). Hence, it enables us to reconstruct tsunami source spectrum using EGF method which assumes that the smaller event acts as the EGF for the larger event (Heidarzadeh et al., 2016). We computed the spectral ratio of the second tsunami to the first tsunami at seven tide gauges (Figure 5c; gray curves), and then calculated their normalized average values (Figure 5c; blue curve) through adjusting the peak energy of all spectra to that of the largest one. The source spectrum shows that the energy of the second tsunami is mainly distributed in the period range of 8–30 min, with spectral peaks at 25.6 min, 16.0 min, and 9.8 min. The period range is generally consistent with the results of Fourier analyses of each station for the second event (Figure 3). Figure 5d shows the results of the method proposed by Rabinovich (1997) which is based on dividing the tsunami spectra to that of the background to construct tsunami source spectrum. We computed the spectral ratio of the second tsunami to the background signals at all tide gauges except RIFR and calculated their normalized average. The period range of main energy (7–28 min) contains spectral peaks at 25.6 min, 16.0 min and 8.5 min, which are close to the spectral peaks calculated by EGF method.

Theoretically, the tsunami source period is related to earthquake rupture length and water depth (Rabinovich, 1997; Heidarzadeh and Satake, 2013; Wang et al., 2021). It can be estimated as:


$$T_n = \frac{2L}{n\sqrt{gh}} \quad n = 1, 2, 3, \dots \quad (1)$$

where  $L$  is the typical size of tsunami source area (length or width),  $g$  is the gravity acceleration, and  $h$  is the average water depth in source area. According to the USGS model, the size of the  $M_w$  8.1 earthquake is 240 km (length of the fault)  $\times$  190 km (width of the fault). The average water depth in the source area is ~ 5,000 m. Hence, the first three source periods of the short axis (width) are 28.6 min, 14.3 min, and 9.5 min. The first three source periods of the long axis (length) are 36.1 min, 18.1 min, and 12.0 min. The source periods of the short axis match well with the peaks of tsunami source spectrum, which confirms the validity of the model geometry. We note that the periods of tsunami waves are mainly influenced by the short axis (width) (Heidarzadeh and Satake, 2013).

In addition, the results of two methods (i.e., EGF and tsunami/background spectral ratio) show similarities in shapes and peak periods of tsunami source spectrum (Figures 5c and 5d). It is noted that the EGF method has the capability to remove both propagation-path effects and local bathymetric effects from the tide gauge records whereas the



195 tsunami/background spectral ratio method would remove mainly local bathymetric effects. Hence, our results may imply that the effects of propagation path are negligibly small for this case, where the tide gauges are located at distances between ~100 km and ~ 2,000 km from the source. Both of the methods are able to reveal the source characteristics merely based on tsunami observations rather than seismological data.



200 **Figure 5:** (a, b) Maximum simulated amplitudes for two tsunamis during the entire simulation time. The source models used for numerical simulation are from the USGS. (c) Spectral ratio of two tsunamis by dividing the spectral energy of the second tsunami



to that of the first tsunami (EGF method). Blue curve is the normalized average of tsunami spectral energy at different tide gauges. (d) Spectral ratio of the second tsunami spectrum to the background signal spectrum. Green curve is the normalized average of different tide gauges.

## 205 6 Conclusion

We studied the characteristics of tsunamis generated by two earthquakes ( $M_w$  7.4 and  $M_w$  8.1) that occurred in Kermadec Islands successively on March 4, 2021, within ~55 km from each other and within approximately 2 h interval. We used the sea level records of 15 tide gauges. The spectra of Fourier analyses show that the dominant period bands of the first tsunami and the second tsunami are 5–17 min and 8–28 min, respectively. Two oscillation patterns with different period  
210 ranges are visible on the wavelet plot at most stations which belong to the first and the second tsunamis. We observed that after the arrival of the second tsunami, the oscillation in the period range of the first tsunami still exists. We calculated the tsunami source spectrum of the larger event (i.e., the second tsunami) by two approaches: empirical Green's function (EGF) method and tsunami/background ratio method. Using the first tsunami as the EGF, spectral deconvolution indicated that energy of the second tsunami is mainly distributed in the period range of 8–30 min, with spectral peaks at 25.6 min, 16.0 min,  
215 and 9.8 min. The method of tsunami/background ratio yielded similar results to the EGF method. The source characteristics were obtained merely based on tsunami data and thus these two methods could be complementary to seismological approaches in source analysis.

### Data availability

The datasets were derived from sources in the public domain. The tide gauge data used in this research were provided  
220 by the Sea Level Station Monitoring Facility of the Intergovernmental Oceanographic Commission (<http://www.ioc-sealevelmonitoring.org/list.php>). We used bathymetric and topographic data of the General Bathymetric Chart of the Ocean ([https://www.gebco.net/data\\_and\\_products/gridded\\_bathymetry\\_data/](https://www.gebco.net/data_and_products/gridded_bathymetry_data/)). We used earthquake focal mechanism catalog of the Global Centroid Moment Tensor Project (<https://www.globalcmt.org/CMTsearch.html>) and earthquake source model of the United States Geological Survey (<https://earthquake.usgs.gov/earthquakes/eventpage/us7000dflf/executive>). We used the  
225 GMT software for drafting some of the figures (Wessel and Smith, 1998).

### Author contribution

YW is responsible for methodology, data curation, software, and writing - original draft preparation. MH is responsible for methodology, software, and writing- reviewing and editing. KS is responsible for methodology, supervision, and writing- reviewing and editing. GH is responsible for data curation and software.



## 230 **Competing Interests**

The authors declare that they have no conflict of interest.

## **Acknowledgements**

The authors declare that they have no conflict of interest. We thank Dr. Takane Hori and Dr. Kentaro Imai from Japan Agency for Marine-Earth Science Technology for their valuable suggestions. MH is funded by the Royal Society (the United Kingdom), grant number CHL\R1\180173. YW is funded by Japan Society for the Promotion of Science, grant number 19J20293.

## **References**

- Baba, T., Takahashi, N. Kaneda, Y. Ando, K. Matsuoka, D., and Kato, T.: Parallel implementation of dispersive tsunami wave modeling with a nesting algorithm for the 2011 Tohoku Tsunami, *Pure Appl. Geophys.*, 172, 3433–3472, doi:10.1007/s00024-015-1049-2, 2015.
- 240 Billen, M. I., Gurnis, M., and Simons, M. Multiscale dynamics of the Tonga-Kermadec subduction zone, *Geophys. J. Int.*, 153(2), 359–388. doi:10.1046/j.1365-246X.2003.01915.x, 2003.
- Cortés, P., Catalán, P. A., Aránguiz, R., and Bellotti, G. Tsunami and shelf resonance on the northern Chile coast, *J. Geophys. Res. Oceans*, 122, 7364–7379. doi:10.1002/2017JC012922, 2017.
- 245 Geoware, The Tsunami Travel Times (TTT) Package, <http://www.geoware-online.com/tsunami.html>, 2011.
- Heidarzadeh, M., and Mulia, I. E., Ultra-long period and small-amplitude tsunami generated following the July 2020 Alaska Mw7.8 tsunamigenic earthquake, *Ocean Eng.*, 234, 109243. doi:10.1016/j.oceaneng.2021.109243, 2021.
- Heidarzadeh, M., Pranantyo, I. R., Okuwaki, R., Dogan, G. G., and Yalciner, A.C., Long tsunami oscillations following the 30 October 2020 Mw 7.0 Aegean Sea earthquake: Observations and modelling, *Pure Appl. Geophys.*, 178, 1531–
- 250 1548. doi:10.1007/s00024-021-02761-8, 2021.
- Heidarzadeh, M., and Satake, K., Waveform and spectral analyses of the 2011 Japan tsunami records on tide gauge and DART stations across the Pacific Ocean, *Pure Appl. Geophys.*, 170, 1275–1293, doi:10.1007/s00024-012-0558-5, 2013.
- Heidarzadeh, M., and Satake, K., New insights into the source of the Makran tsunami of 27 November 1945 from tsunami waveforms and coastal deformation data, *Pure Appl. Geophys.*, 172, 621–640, doi:10.1007/s00024-014-0948-y, 2015.
- 255 Heidarzadeh, M., Harada, T., Satake, K., Ishibe, T., and Gusman, A. R., Comparative study of two tsunamigenic earthquakes in the Solomon Islands: 2015 Mw 7.0 normal-fault and 2013 Santa Cruz Mw 8.0 megathrust earthquakes, *Geophys. Res. Lett.*, 43, 4340–4349. doi:10.1002/2016GL068601, 2016.



- 260 Lundgren, P. R., Okal, E. A., and Wiens, D. A., Rupture characteristics of the 1982 Tonga and 1986 Kermadec earthquakes, *J. Geophys. Res. Solid Earth*, 94, 15521–15539, doi:10.1029/JB094iB11p15521, 1989.
- Morgan, P. G., The New Zealand earthquake of August 6, 1917, *Bull. Seismol. Soc. Am.*, 8(4), 127–128, 1918.
- Okada, Y., Surface deformation due to shear and tensile faults in a half-space, *Bull. Seismol. Soc. Am.*, 75(4), 1135–1154, 1985.
- 265 Power, W., Wallace, L., Wang, X., and Reynier, M., Tsunami Hazard Posed to New Zealand by the Kermadec and Southern New Hebrides Subduction Margins: An Assessment Based on Plate Boundary Kinematics, Interseismic Coupling, and Historical Seismicity, *Pure Appl. Geophys.*, 169, 1–36, doi:10.1007/s00024-011-0299-x, 2012.
- Rabinovich, A. B., Spectral analysis of tsunami waves: Separation of source and topography effects, *J. Geophys. Res. Oceans*, 102, 12663–12676. doi:10.1029/97JC00479, 1997.
- 270 Rabinovich, A. B., and Thomson, R. E., The 26 December 2004 Sumatra tsunami: Analysis of tide gauge data from the world ocean Part 1. Indian Ocean and South Africa, *Pure Appl. Geophys.*, 164, 261–308, doi:10.1007/978-3-7643-8364-0\_2, 2007.
- Torrence, C., and Compo, G., A practical guide to wavelet analysis, *Bull. Am. Meteorol. Soc.*, 79, 61–78, doi:10.1175/1520-0477(1998)079<0061:APGTWA>2.0.CO;2, 1998.
- 275 Wang, Y., Zamora, N., Quiroz, M., Satake, K., and Cienfuegos, R., Tsunami resonance characterization in Japan due to trans-Pacific sources: Response on the bay and continental shelf, *J. Geophys. Res. Oceans*, 126, e2020JC017037, doi:10.1029/2020JC017037, 2021.
- Wessel, P., and Smith, W. H. F., New, improved version of generic mapping tools released, *Eos Transactions of AGU*, 79(47), 579. doi:10.1029/98EO00426, 1998.
- 280 Wyss, M., Habermann, R. E., and Griesser, J.-C., Seismic quiescence and asperities in the Tonga-Kermadec Arc, *J. Geophys. Res. Solid Earth*, 89, 9293–9304, doi:10.1029/JB089iB11p09293, 1984.