1 Invited perspectives: Landslide populations - can they be predicted?

2

3 Fausto Guzzetti^(1,2)

4 (1) Dipartimento della Protezione Civile, Presidenza del Consiglio dei Ministri, via Vitorchiano 2,

5 00189 Roma

6 (2) Istituto di Ricerca per la Protezione Idrogeologica, Consiglio Nazionale delle Ricerche, via
7 della Madonna Alta 126, 06129 Perugia

8

9 Landslides are different from other natural hazards. Unlike volcanoes, they do not threaten human civilization (Papale and Marzocchi, 2019). Unlike tsunamis, they do not affect simultaneously 10 11 several thousands of kilometres of coastline – although a submarine landslide in Norway has caused 12 a tsunami to hit Scotland (Dawson et al., 1988). Unlike floods and earthquakes, they do not cause hundreds of thousands of casualties in a single event – although a landslide has killed thousands in 13 14 Peru (Evans et al., 2009) and debris flows tens of thousands in Colombia (Wieczorek et al., 2001). 15 But the human toll of landslides is high (Froude and Petley, 2018), and their economic and societal 16 consequences are largely undetermined. Compared to other hazards, landslides are subtle, often go 17 unnoticed, and their consequences are underestimated. 18 As with other hazards, the design and implementation of effective risk reduction strategies depend 19 on the ability to predict (forecast, project, anticipate) landslides. I have argued that "our ability to

20 predict landslides and their consequences measures our ability to understand the underlying [...]

21 processes that control or condition landslides, as well as their spatial and temporal occurrence"

22 (Guzzetti, 2021). This assumes that landslide prediction is possible; something that has not been

23 demonstrated (or disproved), theoretically. Yet, there is nothing in the literature that prevents

24 landslide prediction; provided that one clarifies the meaning of "prediction" (Guzzetti, 2021), that

25 the prediction is scientifically based (Guzzetti, 2015), and that we understand the limits of the

26 prediction (Wolpert, 2001). Efforts are needed to determine the limits of landslide predictions, for

all landslide types (Hungr et al., 2014) and at all geographic and temporal scales (Figure 1).

28 Here, I outline what I consider to be the main problems that need to be addressed in order to 29 advance our ability to predict landslide hazards and risk. The field is vast, and I limit my perspective 30 to populations of landslides - that is, the hazards and risk posed by many landslides caused by one 31 triggering event, or by multiple events in a short period. In this context, predicting landslide hazard 32 means anticipating where, when, how frequently, how many, and how large populations of 33 landslides are expected (Guzzetti et al., 2005; Lombardo et al., 2020; Guzzetti, 2021). Predicting 34 landslide risk is about anticipating the consequences of landslide populations to different 35 vulnerable elements (Alexander, 2005; Glade et al., 2005; Galli and Guzzetti, 2007; Salvati et al., 2018). 36

37 Landslides tend to occur where they have previously occurred (Temme et al., 2020). Therefore, 38 one way to assess *where* they are expected is to map past and new landslides. The technology is 39 mature for regional and even global landslide detection and mapping services based on the 40 automatic or semi-automatic processing of aerial and satellite imagery; optical, SAR and LiDAR 41 data (Guzzetti et al., 2012; Mondini et al., 2021). An alternative – and complementary – way is through susceptibility modelling; an approach for which there is no shortage of data-driven 42 43 methods, but rather of suitable environmental and landslide data (Reichenbach et al., 2018). The 44 increasing availability of satellite imagery, some of which repeated over time and free of charge 45 (Aschbacher, 2017), opens unprecedented opportunities to prepare event and multi-temporal 46 inventory maps covering very large areas, which are essential to build space-time prediction models 47 (Lombardo et al., 2020), to investigate the legacy of old landslides on new ones (Samia et al., 2017; 48 Temme et al., 2020), to obtain accurate thematic data for susceptibility modelling (Reichenbach et 49 al., 2018), and to validate geographical landslide early warning systems (Piciullo et al., 2018; 50 Guzzetti et al., 2020). However, the literature reveals a systematic lack of standards for 51 constructing, validating, and ranking the quality of landslide maps and prediction models (Guzzetti 52 et al., 2012; Mondini et al., 2021; Reichenbach et al., 2018). This reduces the credibility of the 53 maps and models. A gap that urgently needs to be bridged (Guzzetti, 2021).

54 Predicting *when* or *how frequently* landslides will occur can be done for short and for long periods.

55 For short periods – from hours to weeks – the prediction is obtained through process-based models,

- 56 rainfall thresholds, or their combination. Process-based models rely upon the understanding of the
- 57 physical laws controlling the slope instability conditions of a landscape forced by a transient trigger

e.g., a rainfall, snow melt, seismic, or volcanic event (Bogaard and Greco, 2016, 2018). The major
limitation of physically-based models is the scarcity of relevant data, which are hard to obtain for
very large areas. New approaches to obtain relevant, spatially-distributed data are needed, as well
as novel models able to extrapolate what is learned in sample areas to vast territories (Bellugi et
al., 2011; Alvioli and Baum, 2016; Alvioli et al., 2018; Mirus et al., 2020).

63 Thresholds are empirical or statistical models that link physical quantities (e.g., cumulative rainfall, rainfall duration) to the occurrence - or lack of occurrence - of known landslides. Reviews of the 64 65 literature (Guzzetti et al., 2008; Segoni et al.; 2018) have highlighted conceptual problems with the 66 definition and use of rainfall thresholds for operational landslide forecasting and early warning 67 (Piciullo et al., 2018; Guzzetti et al., 2020), including the lack of standards for defining the 68 thresholds and their associated uncertainty (Melillo et al., 2018), and for the validation of the 69 threshold models (Piciullo et al., 2017, 2018; Guzzetti et al., 2020). The community needs shared 70 criteria and algorithms coded into open-source software for the objective definition of rainfall 71 events, of the rainfall conditions that can result in landslides, of rainfall thresholds (Melillo et al., 72 2015, 2018), and for the validation of the threshold models (Piciullo et al., 2017). This will not 73 only provide reliable and comparable thresholds, allowing for regional and global studies (Guzzetti 74 et al., 2008; Segoni et al.; 2018), but also increase the credibility of early warning systems based 75 on rainfall threshold models (Guzzetti et al., 2020).

76 The projection of landslide frequency for long periods – decades to millennia – is much more 77 difficult and uncertain, as it depends on climatic and environmental characteristics that are poorly 78 know and difficult to measure and model (Crozier, 2010; Gariano and Guzzetti, 2016), as well as 79 on the inherent incompleteness of the historical landslide records (Rossi et al., 2010). The literature 80 on the analysis of historical landslide records remains scarce, but the number of studies projecting 81 the future occurrence of landslides is increasing (Gariano et al., 2017; Peres and Cancelliere, 2018; 82 Schlögl and Matulla, 2018; Patton et al., 2019; Schlögel et al., 2020; Gariano and Guzzetti, 2021). 83 In this field, studies will be relevant if they compare analyses and validation methods in different

84 areas. This requires the exchange of data and information.

85 Predicting how many and how large landslides are expected means anticipating the size (e.g., area,

86 volume, length, width, depth) and number of landslides in an area – with size and number correlated

87 in a population of landslides. This information is obtained by constructing and modelling probability distributions of landslide sizes obtained typically from landslide event inventory maps 88 89 (Stark and Hovius, 2011; Malamud et al., 2004). The literature on the topic is limited, and with 90 differences in the way the distributions are modelled. This hampers comparisons from different 91 areas. Although models have been proposed to explain the probability size distributions (Katz and 92 Aharanov, 2006; Stark and Guzzetti, 2009; Klar et al., 2011; Bellugi et al., 2021), further efforts 93 are needed to explain the observed distributions of landslide sizes, and to evaluate their variability 94 and uncertainty.

By combining probabilistic information on *where*, *when* or *how frequently*, and *how many* or *how large* landslides are, one can evaluate landslide hazards for different landslide types. However, the existing models are crude, they work under assumptions that are difficult to prove (Guzzetti et al., 2005), and the possibility to export them in different areas is limited, or untested. Novel efforts are needed to prepare reliable landslide hazard models (Lombardo et al., 2020; Guzzetti, 2021).

100 Assessing landslide hazard is important but, for social applications what is needed is the estimation 101 of the landslide consequences, which means assessing the vulnerability to landslides of various 102 elements at risk (Alexander 1999; Galli and Guzzetti 2007), and evaluating landslide risk (Cruden 103 and Fell, 1997; Glade et al., 2005; Porter and Morgenstern, 2013), including risk to the population 104 (Petley, 2012; Froude and Petley, 2018; Salvati et al., 2018; Rossi et al., 2019). Here, the main 105 limitation is the difficulty to obtain data on landslide vulnerability, and reliable records of landslide 106 events and their consequences (Petley, 2012; Froude and Petley, 2018; Salvati et al. 2018). Where 107 the information is available, comprehensive landslide risk models can be constructed, and validated 108 (Rossi et al., 2019). It is important that efforts are made to collect reliable records of landslides and 109 their consequences, and that the records are shared to test different risk models.

Of the various factors governing landslide hazard the most uncertain and the one requiring more urgent efforts is the time prediction (*when, how frequently*), followed by the prediction of the size and number of expected failures. For both, multi-temporal inventories and landslide catalogues are essential to build innovative predictive models. To construct the records, systematic efforts are needed for landslide detection and mapping (Mondini et al., 2021). For susceptibility (*where*), the challenge is to prepare reliable regional, continental, or global assessments (Stanley and Kirschbaum, 2017; Broeckx et al., 2018; Wilde et al., 2018; Mirus et al., 2020). Critical are also novel modelling frameworks combining the hazard factors (Lombardo et al., 2020). But the goal is to reduce risk (Glade et al., 2005). For that, vulnerability studies, improved early warning capabilities, quantification of the benefits of prevention, and better risk communication strategies

120 are crucial (Guzzetti, 2018). Much work is needed on these largely unexplored subjects.

121 Ultimately, I note that in medicine – a field of science conceptually close to the field of landslide

- 122 hazard assessment and risk mitigation (Guzzetti, 2015) the paradigm of "convergence research"
- 123 is emerging (Sharp and Hockfield, 2017), where "convergence comes as a result of the sharing of

124 methods and ideas ... It is the integration of insights and approaches from historically distinct

scientific and technological disciplines" (Sharp et al., 2016). The community of landslide scientists

126 should embrace the paradigm of "converge research", exploiting the vast amount of data,

127 measurements, and observations that are available and will be collected, expanding the making and

128 use of predictions, assessing the economic and social costs of landslides, designing sustainable

129 mitigation and adaptation strategies, and addressing the ethical issues posed by natural hazards,

130 including landslides (Bohle, 2019). I am persuaded that this will contribute to advancing

131 knowledge and building a safer society (Guzzetti, 2018).

132 **References**

- Alexander, E.D.: Vulnerability to landslides. In: Glade, T., Anderson, M.G., Crozier, M.J. (eds.)
 Landslide risk assessment. John Wiley, pp. 175–198,
 https://doi.org/10.1002/9780470012659.ch5, 2005.
- Alvioli, M. and Baum, R.L.: Parallelization of the TRIGRS model for rainfall-induced landslides
 using the message passing interface. Env. Model. Soft. 81, 122–135.
 https://doi.org/10.1016/j.envsoft.2016.04.002, 2016.
- Alvioli, M., Melillo, M., Guzzetti, F., Rossi, M., Palazzi, E., von Hardenberg, J., Brunetti, M.T.
 and Peruccacci, S.: Implications of climate change on landslide hazard in Central Italy. Sci Tot
 Env 630, 1528-1543, doi:10.1016/j.scitotenv.2018.02.315, 2018.
- Aschbacher, J.: ESA's Earth Observation Strategy and Copernicus. In: Onoda, M., Young, O.R.
 (eds.) Satellite Earth Observations and Their Impact on Society and Policy. Springer
 Singapore, Singapore, Ch. 5, pp. 81-86, https://doi.org/10.1007/978-981-10-3713-9 5, 2017.
- Bellugi, D., Dietrich, W.E., Stock, J.D., McKean, J.A., Kazian, B. and Hargrove, P.: Spatially
 Explicit Shallow Landslide Susceptibility Mapping Over Large Areas. It. J. Eng. Geol.
 Environ 399–407, https://doi.org/10.4408/IJEGE.2011-03.B-04, 2011.
- Bellugi, D.G., Milledge, D.G., Cuffey, K.M., Dietrich, W.E. and Larsen, L.G.: Controls on the size
 distributions of shallow landslides. Proc. Nat. Ac. Sci. 118, e2021855118,


- 150 https://doi.org/10.1073/pnas.2021855118, 2021.
- Broeckx, J., Vanmaercke, M., Duchateau, R. and Poesen, J.: A data-based landslide susceptibility
 map of Africa. Earth-Science Reviews 185, 102–121.
 https://doi.org/10.1016/j.earscirev.2018.05.002, 2018.
- Bogaard, T. and Greco, R.: Invited perspectives: Hydrological perspectives on precipitation
 intensity-duration thresholds for landslide initiation: proposing hydro-meteorological
 thresholds. Nat. Haz. Earth Sys. Sci. 18, 31–39, https://doi.org/10.5194/nhess-18-31-2018,
 2018.
- Bogaard, T.A. and Greco, R.: Landslide hydrology: from hydrology to pore pressure. Wiley
 Interdisciplinary Reviews: Water 3, 3, 439–459, https://doi.org/10.1002/wat2.1126, 2016.
- Bohle, M. (ed.): Exploring Geoethics: Ethical Implications, Societal Contexts, and Professional
 Obligations of the Geosciences. Springer Int. Pub., Cham, https://doi.org/10.1007/978-3-03012010-8, 2019.
- 163 Crozier, M.J.: Deciphering the effect of climate change on landslide activity: A review.
 164 Geomorphology 124, 260–267, https://doi.org/10.1016/j.geomorph.2010.04.009, 2010.
- 165 Cruden, D.M. and Fell, R.: Landslide Risk Assessment. CRC Press, 1997.
- Dawson, A.G., Long, D. and Smith, D.E.: The Storegga slides: evidence from eastern Scotland for
 a possible tsunami. Marine Geol. 82, 271–276, https://doi.org/10.1016/0025-3227(88)901466, 1988.
- Evans, S.G., Bishop, N.F., Fidel Smoll, L., Valderrama Murillo, P., Delaney, K.B., and OliverSmith, A.: A re-examination of the mechanism and human impact of catastrophic mass flows
 originating on Nevado Huascarán, Cordillera Blanca, Peru in 1962 and 1970. Eng. Geol. 108,
 96–118, https://doi.org/10.1016/j.enggeo.2009.06.020, 2009.
- Froude, M.J. and Petley, D.N.: Global fatal landslide occurrence from 2004 to 2016. Nat. Haz.
 Earth Sys. Sci. 18, 2161–2181, https://doi.org/10.5194/nhess-18-2161-2018, 2018.
- Galli, M. and Guzzetti, F.: Landslide vulnerability criteria: a case study from Umbria, Central Italy.
 Env. Manag. 40, 649–665, https://doi.org/10.1007/s00267-006-0325-4, 2007.
- Gariano, S.L. and Guzzetti, F.: Landslides in a changing climate. Earth-Science Reviews 162, 227–
 252, https://doi.org/10.1016/j.earscirev.2016.08.011, 2016.
- Gariano, S.L. and Guzzetti, F.: Mass-Movements and Climate Change, in: Reference Module in
 Earth Sys. Env. Sci., Elsevier, p. B9780128182345000000, https://doi.org/10.1016/B978-012-818234-5.00043-2, 2021.
- Gariano, S.L., Rianna, G., Petrucci, O. and Guzzetti, F. (2017) Assessing future changes in the
 occurrence of rainfall-induced landslides at a regional scale. Sci. Tot. Env. 596-597, 417–426,
 https://doi.org/10.1016/j.scitotenv.2017.03.103
- 185 Glade, T., Anderson, M.G. and Crozier, M.J. (eds.): Landslide hazard and risk. John Wiley Sons,
 186 ISBN-13:978-0471486633, 2005.
- 187Guzzetti, F.: Forecasting natural hazards, performance of scientists, ethics, and the need for188transparency.Toxicol.Env.Chem.98,1043–1059,189https://doi.org/10.1080/02772248.2015.1030664, 2015.
- 190 Guzzetti, F.: Rischi naturali: l'urgenza di una scienza nuova, in: Caporale, C., Maffei, L., Marchis,
 191 V., de Martin, J.C. (eds.) Europa: Le sfide della scienza, Europa. Istituto della Enciclopedia

- 192 Italiana, Roma, pp. 127–133, 2018, (in Italian).
- Guzzetti, F.: On the Prediction of Landslides and Their Consequences, in: Sassa, K., Mikoš, M.,
 Sassa, S., Bobrowsky, P.T., Takara, K. and Dang, K. (eds.), Understanding and Reducing
 Landslide Disaster Risk: Volume 1 Sendai Landslide Partnerships and Kyoto Landslide
 Commitment. Springer International Publishing, Cham, pp. 3–32, https://doi.org/10.1007/9783-030-60196-6 1, 2021.
- Guzzetti, F., Gariano, S.L., Peruccacci, S., Brunetti, M.T., Marchesini, I., Rossi, M. and Melillo,
 M.: Geographical landslide early warning systems. Earth-Science Reviews 200, 102973,
 https://doi.org/10.1016/j.earscirev.2019.102973, 2020.
- Guzzetti, F., Mondini, A.C., Cardinali, M., Fiorucci, F., Santangelo, M. and Chang, K.-T.:
 Landslide inventory maps: New tools for an old problem. Earth-Science Reviews 112, 42–66, https://doi.org/10.1016/j.earscirev.2012.02.001, 2012.
- Guzzetti, F., Peruccacci, S., Rossi, M. and Stark, C.P.: The rainfall intensity-duration control of
 shallow landslides and debris flows: an update. Landslides 5, 3–17,
 https://doi.org/10.1007/s10346-007-0112-1(2008), 2008.
- Guzzetti, F., Reichenbach, P., Cardinali, M., Galli, M. and Ardizzone, F.: Probabilistic landslide
 hazard assessment at the basin scale. Geomorphology 72, 272–299,
 https://doi.org/10.1016/j.geomorph.2005.06.002, 2005.
- Haque, U., da Silva, P.F., Devoli, G., Pilz, J., Zhao, B., Khaloua, A., Wilopo, W., Andersen, P.,
 Lu, P., Lee, J., Yamamoto, T., Keellings, D., Wu, J.-H. and Glass, G.E.: The human cost of
 global warming: Deadly landslides and their triggers (1995–2014). Sci. Tot. Env. 682, 673–
 684, https://doi.org/10.1016/j.scitotenv.2019.03.415, 2019.
- Hungr, O., Leroueil, S. and Picarelli, L.: The Varnes classification of landslide types, an update.
 Landslides 11, 167–194, https://doi.org/10.1007/s10346-013-0436-y, 2014.
- Katz, O. and Aharonov, E.: Landslides in vibrating sand box: What controls types of slope failure
 and frequency magnitude relations? Earth Plan. Sci. Let. 247, 280–294,
 https://doi.org/10.1007/978-1-4020-4370-3, 2006.
- Klar, A., Aharonov, E., Kalderon-Asael, B. and Katz, O.: Analytical and observational relations
 between landslide volume and surface area. J. Geophy. Res. 116, F02001,
 https://doi.org/10.1029/2009JF001604, 2011.
- Lombardo, L., Opitz, T., Ardizzone, F., Guzzetti, F. and Huser, R.: Space-time landslide predictive
 modelling. Earth-Science Reviews 209, 103318,
 https://doi.org/10.1016/j.earscirev.2020.103318, 2020.
- Malamud, B.D., Turcotte, D.L., Guzzetti, F. and Reichenbach, P.: Landslide inventories and their
 statistical properties. Earth Surf. Proc. Land. 29, 687–711, https://doi.org/10.1002/esp.1064,
 2004.
- Melillo, M., Brunetti, M.T., Peruccacci, S., Gariano, S.L. and Guzzetti, F.: An algorithm for the
 objective reconstruction of rainfall events responsible for landslides. Landslides 12, 311–320.
 https://doi.org/10.1007/s10346-014-0471-3, 2015.
- Melillo, M., Brunetti, M.T., Peruccacci, S., Gariano, S.L., Roccati, A. and Guzzetti, F.: A tool for
 the automatic calculation of rainfall thresholds for landslide occurrence. Env. Mod. & Soft.
 105, 230, 242, https://doi.org/10.1016/j.epu/oft.2018.03.024, 2018
- 233 105, 230–243, https://doi.org/10.1016/j.envsoft.2018.03.024, 2018.

- Mirus, B.B., Jones, E.S., Baum, R.L., Godt, J.W., Slaughter, S., Crawford, M.M., Lancaster, J.,
 Stanley, T., Kirschbaum, D.B., Burns, W.J., Schmitt, R.G., Lindsey, K.O. and McCoy, K.M.:
 Landslides across the USA: occurrence, susceptibility, and data limitations. Landslides.
 https://doi.org/10.1007/s10346-020-01424-4, 2020.
- Mondini, A.C., Guzzetti, F., Chang, K.-T., Monserrat, O., Martha, T.R. and Manconi, A.: Landslide
 failures detection and mapping using Synthetic Aperture Radar: Past, present and future. EarthScience Reviews 216, 103574, https://doi.org/10.1016/j.earscirev.2021.103574, 2021.
- Papale, P. and Marzocchi, W.: Volcanic threats to global society. Science 363, 1275–1276,
 https://doi.org/10.1126/science.aaw7201, 2019.
- Patton, A.I., Rathburn, S.L. and Capps, D.M.: Landslide response to climate change in permafrost
 regions. Geomorphology 340, 116–128, https://doi.org/10.1016/j.geomorph.2019.04.029,
 2019.
- Peres, D.J. and Cancelliere, A.: Modeling impacts of climate change on return period of landslide
 triggering. J. Hydrology 567, 420–434, https://doi.org/10.1016/j.jhydrol.2018.10.036, 2018.
- Petley, D.: Global patterns of loss of life from landslides. Geology 40(10), 927–930, https://doi.org/10.1130/g33217.1, 2012.
- Piciullo, L., Calvello, M. and Cepeda, J.M.: Territorial early warning systems for rainfall-induced
 landslides. Earth-Science Reviews 179, 228–247,
 https://doi.org/10.1016/j.earscirev.2018.02.013, 2018.
- Piciullo, L., Gariano, S.L., Melillo, M., Brunetti, M.T., Peruccacci, S., Guzzetti, F. and Calvello,
 M.: Definition and performance of a threshold-based regional early warning model for rainfall induced landslides. Landslides 14, 995–1008, https://doi.org/10.1007/s10346-016-0750-2,
 2017.
- Porter, M. and Morgenstern, N.: Landslide risk evaluation Canadian technical guidelines and
 best practices related to landslides: a national initiative for loss reduction. Geological Survey
 of Canada, Open File 7312:21, 2013.
- Reichenbach, P., Rossi, M., Malamud, B.D., Mihir, M. and Guzzetti, F.: A review of statisticallybased landslide susceptibility models. Earth-Science Reviews 180, 60–91,
 https://doi.org/10.1016/j.earscirev.2018.03.001, 2018.
- Rossi, M., Guzzetti, F., Salvati, P., Donnini, M., Napolitano, E. and Bianchi, C.: A predictive model
 of societal landslide risk in Italy. Earth-Science Reviews 196, 102849,
 https://doi.org/10.1016/j.earscirev.2019.04.021, 2019.
- Rossi, M., Witt, A., Guzzetti, F., Malamud, B.D. and Peruccacci, S.: Analysis of historical landslide
 time series in the Emilia-Romagna region, northern Italy. Earth Surf. Proc. Land. 35, 1123–
 1137, https://doi.org/10.1002/esp.1858, 2010.
- Salvati, P., Petrucci, O., Rossi, M., Bianchi, C., Pasqua, A.A. and Guzzetti, F.: Gender, age and
 circumstances analysis of flood and landslide fatalities in Italy. Sci. Tot. Env. 610–611, 867–
 879, https://doi.org/10.1016/j.scitotenv.2017.08.064, 2018.
- Samia, J., Temme, A.J.A.M., Bregt, A., Wallinga, J., Guzzetti, F., Ardizzone, F. and Rossi, M.: Do
 landslides follow landslides? Insights in path dependency from a multi-temporal landslide
 inventory. Landslides 14, 547–558, https://doi.org/10.1007/s10346-016-0739-x, 2017.
- 275 Schlögel, R., Kofler, C., Gariano, S.L., Campenhout, J.V. and Plummer, S.: Changes in climate

- patterns and their association to natural hazard distribution in South Tyrol (Eastern Italian
 Alps). Scientific Reports 10, 5022, https://doi.org/10.1038/s41598-020-61615-w, 2020.
- Schlögl, M. and Matulla, C.: Potential future exposure of European land transport infrastructure to
 rainfall-induced landslides throughout the 21st century. Nat. Haz. Earth Sys. Sci. 18, 1121–
 1132, https://doi.org/10.5194/nhess-18-1121-2018, 2018.
- Segoni, S., Piciullo, L. and Gariano, S.L.: A review of the recent literature on rainfall thresholds
 for landslide occurrence. Landslides 15, 1483–1501, https://doi.org/10.1007/s10346-0180966-4, 2018.
- Sharp, P. and Hockfield, S.: Convergence: The future of health. Science 355, 589.1–589,
 https://doi.org/10.1126/science.aam8563, 2017.
- Sharp, P., Jacks, T. and Hockfield, S.: Convergence: the future of health.
 ConvergenceRevolution.net, Cambridge, Massachusetts, 2016.
- Stanley, T and Kirschbaum, D: A heuristic approach to global landslide susceptibility mapping.
 Natural Hazards 87, 145–164. https://doi.org/10.1007/s11069-017-2757-y, 2017.
- Stark, C.P. and Guzzetti, F.: Landslide rupture and the probability distribution of mobilized debris
 volumes. J. Geophy. Res. 114, F00A02, https://doi.org/10.1029/2008JF001008, 2009.
- Stark, C.P. and Hovius, N.: The characterization of landslide size distributions. Geophy. Res. Let.
 28, 1091–1094, https://doi.org/10.1029/2000gl008527, 2001.
- Temme, A., Guzzetti, F., Samia, J. and Mirus, B.B.: The future of landslides' past—a framework
 for assessing consecutive landsliding systems. Landslides 17, 1519–1528,
 https://doi.org/10.1007/s10346-020-01405-7, 2020.
- 297 Wieczorek, G.F., Larsen, M.C., Eaton, L.S., Morgan, B.A. and Blair, J.L.: Debris-flow and 298 flooding hazards associated with the December 1999 storm in coastal Venezuela and strategies 299 for mitigation. U.S. Geological Survey Open File Report 01-0144, 300 https://pubs.usgs.gov/of/2001/ofr-01-0144/, 2001.
- Wilde, M., Günther, A., Reichenbach, P., Malet, J.-P. and Hervás, J.: Pan-European landslide
 susceptibility mapping: ELSUS Version 2. J. Maps 14, 97–104.
 https://doi.org/10.1080/17445647.2018.1432511, 2018.
- Wolpert, D.H.: Computational capabilities of physical systems. Physical Review E 65, 016128–1,
 https://doi.org/10.1103/PhysRevE.65.016128, 2001.

306



307

Figure 1. Space (lower x-axis) – time (y-axes) chart showing main geomorphological and geomechanical landslide domains, and typical length-scale of main meteorological and geophysical
triggers and drivers of populations of landslides. Coloured polygons show approximate subdomains for typical landslide hazards and risk mapping and modelling efforts. Modified after
Guzzetti (2021).

313