

1 **Review Article: A European Perspective on Wind and storm damage: From**
2 **Meteorologythe meteorological background to index-based approaches to assess**
3 **Impacts**

4
5 Daniel Glikzman^{1,2}, Paul Averbeck³, Nico Becker^{4,5}, Barry Gardiner^{6,7}, Valeri Goldberg¹, Jens
6 Grieger⁴, Dörthe Handorf⁸, Karsten Haustein^{9,#}, Alexia Karwat¹⁰, Florian Knutzen⁹, Hilke S.
7 Lentink¹¹, Rike Lorenz⁴, Deborah Niermann¹², Joaquim G. Pinto¹¹, Ronald Queck¹, Astrid
8 Ziemann¹, and Christian L. E. Franzke^{13,14}

9
10 ¹ Technische Universität Dresden, Faculty of Environmental Sciences, Institute for
11 Hydrology and Meteorology, Chair of Meteorology, Piennner Str. 23, 01737 Tharandt
12 Germany

13 ² Chair of Computational Landscape Ecology, Institute of Geography, Technische Universität
14 Dresden, Helmholtzstr. 10, 01069, Dresden, Germany

15 ³ iES Landau, Institute for Environmental Sciences, University of Koblenz-Landau, Fortstraße
16 7, 76829 Landau, Germany

17 ⁴ Freie Universität Berlin, Institut für Meteorologie, Berlin, Germany

18 ⁵ Hans-Ertel-Centre for Weather Research, Berlin, Germany

19 ⁶ Faculty of Environment and Natural Resources, Albert-Ludwigs University, Freiburg,
20 Germany.

21 ⁷ Institut Européen de la Forêt Cultivée, Cestas, France.

22 ⁸ Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Research
23 Department Potsdam, Telegrafenberg A45-ND-14473 Potsdam, Germany

24 ⁹ Climate Service Center Germany (GERICS), Helmholtz-Zentrum hereon, Fischertwiete 1,
25 20095 Hamburg, Germany

26 ¹⁰ Universität Hamburg, Meteorological Institute, Grindelberg 5, 20144 Hamburg, Germany

27 ¹¹ Institute of Meteorology and Climate Research, Department of Tropospheric Research
28 (IMK-TRO), Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany

29 ¹² Deutscher Wetterdienst, Frankfurter Straße 135, 63067 Offenbach, Germany

30 ¹³ Center for Climate Physics, Institute for Basic Science, Busan, Republic of Korea

31 ¹⁴ Pusan National University, Busan, Republic of Korea

32
33 [#] Now at Institute for Meteorology, University of Leipzig, Leipzig, Germany

34
35 Corresponding author: christian.franzke@pusan.ac.kr

36
37 **Abstract**

38 **Wind and windstorms cause severe damage to natural and human-made environments.**
39 **Thus, wind-related risk assessment is vital for the preparation and mitigation of calamities.**
40 **However, the cascade of events leading to damage depends on many factors that are**
41 **environment-specific and the available methods to address wind-related damage often require**
42 **sophisticated analysis and specialization. Fortunately, simple indices and thresholds are as**
43 **effective as complex mechanistic models for many applications. Nonetheless, the multitude of**
44 **indices and thresholds available requires a careful selection process according to the target**
45 **environmentsector. Here, we first provide a basic background on wind and storm formation**
46 **and characteristics, followed by a comprehensive collection of both indices and thresholds**
47 **that can be used to predict the occurrence and magnitude of wind and storm damage. We**

48 focused on five key environmentsectors: forests, urban areas, transport, agriculture, and wind-
49 based energy production. For each environmentsector we described indices and thresholds
50 relating to physical properties such as topography and land cover but also to economic
51 aspects (e.g. disruptions in transportation or energy production). In the face of increased
52 climatic variability, the promotion of more effective analysis of wind and storm damage could
53 reduce the impact on society and the environment.

54 1. General introduction

55 Wind is a common feature of our day-to-day weather just like air temperature and precipitation.
56 Wind is per definition a sustained air movement in the atmosphere, which can range from still
57 conditions to extraordinary values, from very local to global spatial scales, and has a wide
58 range of temporal scales from seconds to decades. Winds can have both a beneficial and
59 detrimental effect on society, infrastructure, and agriculture. On one hand, storms, which have
60 very strong winds, can lead to considerable damage in infrastructure and forestry, e.g. storm
61 Kyrill in 2007 (Fink et al., 2009), contribute to widespread forest fires, e.g. Australia 2020 (van
62 Oldenborgh et al., 2021), or enhance evaporation, thus, drying out the soil (e.g., Bittelli et al.,
63 2008). We view damage as a disadvantageous change in the quantities, quality, or function of
64 an object. On the other hand, moderately strong winds can have positive effects on wind
65 energy production and cause a stronger mixing in the boundary layer (cancelling detrimental
66 thermal inversions to agriculture) or – in the case of nightly slope winds - alleviate summer
67 heat conditions in valleys and cities (e.g. Ganbat et al., 2015).

68

69 The damage associated with strong winds is primarily due to short-term wind gusts, and
70 leadleads to a substantial increase in wind speed (e.g. Brasseur, 2001). Wind gusts are
71 sudden increases in windspeed, which last typically less than 20 seconds, while strong
72 winds refer to sustained wind speed over longer time periods. Strong wind gusts often lead
73 to uprooting or breaking of trees, damaging of damage to crops in fields (Gardiner et al. 2016),
74 lifting of roofs, and damaging critical infrastructure like bridges and roads (e.g., Klawa and
75 Ulbrich, 2003; Mitchell-Wallace et al. 2017). In coastal areas, strong winds and wind gusts
76 may lead to storm surges and coastal flooding (e.g. Flather, 2001). The exact impacts of
77 strong winds depend also on other factors besides wind speed thresholds. For example,
78 damage to forests dependdepends on many other factors like precipitation and topography
79 (Gardiner, 2021). Thus, to predict damage or identify areas at risk of wind or storm damage,
80 indices are a vital tool in assessing the likelihood and magnitude of damage for an
81 environment. in a given sector or environment. For example, Merz et al. (2020) explore in
82 their review the current state of knowledge on skillful forecasts of impacts for many hazards,
83 for which indices are very useful. With storm damage we refer to damage, mainly to
84 properties and forests, caused by severe wind storms, while wind damage is more general
85 and includes all adverse effects of wind, including storm damage. We define risk as the
86 likelihood here that wind causes some damage, and their consequences and risk can be
87 quantified as the function of hazard probability, exposure and vulnerability (e.g. Kelman
88 2003; Hoeppe 2016; Franzke 2017).

89

90 For wind indices and wind impact models different wind parameters are in use. These are
91 often derived from modeled data like reanalysis datasets. While these model parameters are
92 strongly related to observed wind parameters, they are not the same and their definitions

93 cannot be used interchangeably. Since observational data is rare and it is more common to
94 work with modeled data the following parameter definitions focus on parameters derived
95 from models. It is often assumed that the maximum daily or hourly gust speed [m/s] at 10m
96 height relates strongest to damage. The WMO defines a wind gust as the maximum of the
97 wind averaged over 3 second intervals which is in most cases shorter than the model time
98 step. Thus, many models rely on parametrization for gust speed. For example, the ECMWF
99 Integrated Forecasting System deduces the magnitude of a gust within each time step from
100 the time-step-averaged surface stress, surface friction, wind shear and stability. Other
101 common parameters in use are daily or hourly mean or maximum wind speeds at 10m
102 height which express the mean or maximum values of all model time steps in an hour or a
103 day. The parametrized gust speed as well as mean wind speeds in a model grid cell can
104 deviate widely from local observations.

105

106 Indices can be used to predict damage caused directly by wind, or to quantify how the wind
107 modulates the damage caused by another process such as fire or drought. Furthermore, the
108 choice of indices depends also on land use as it influences the interaction between land
109 surfaces and the wind; tree species and forest structures can have considerable influence on
110 the damage probability (e.g. Gardiner, 2021). The understanding of wind, storm dynamics,
111 and the ability to predict the damage they cause, requires an interdisciplinary approach.
112 However, much of the relevant literature is in specialized journals. Here, we aim to bring
113 these different disciplines together to provide an interdisciplinary synthesis of the topic. To
114 bridge the gap between the different communities we, within the ClimXtreme consortium, we
115 created a work group and invited specialists from outside the consortium to broaden our
116 research expertise. During regular joint meetings we identified the following sectors: forests,
117 urban areas, transport, agriculture, and energy as the most relevant terrestrial environments
118 that could be impacted by wind and storm damage. We focused on literature resources
119 stemming mainly from Europe, but in cases of relevance and to further expand the scope of
120 the review we also incorporated examples from other regions.

121 We provide a basic background on wind and storm formation and intra-seasonal variability in
122 section 2. Section 3 focuses on the ~~interaction~~ interactions between wind and surface
123 structures which are prone to wind-damage. Section 4 focusses on wind- and storm-related
124 indices and thresholds. ~~Indices and thresholds are typically easy to use and are efficient in terms~~
125 ~~of the time required to understand and use them, as compared with complex mechanistic modelling~~
126 ~~approaches.~~ In particular, we cover the following ~~environments~~ sectors: forests, urban areas,
127 transport, agriculture, and energy. Additionally, we discuss compound indices and thresholds
128 used by national weather services. Finally, in section 5 we provide an outlook and discuss
129 open research questions. Due to the location of the authors, we provide mainly a European
130 perspective on this topic, but believe our synthesis is more widely applicable.

131

132 **2. Wind and storm formation – mechanisms and concepts**

133 **2.1. The general circulation and wind generation**

134 The general circulation of the atmosphere is driven by the differential heating of the Earth
135 (Held 2019); the equatorial regions receive more solar radiation than higher latitudes, while in
136 the polar regions the atmosphere is losing heat into space. This differential heating ~~in~~ of the
137 Earths' surface causes pressure differences in the atmosphere. As a result, a pressure
138 gradient force acts on the air masses, leading to a movement from high to low pressure centers

139 to alleviate this pressure difference. Since the atmosphere moves toward an equilibrium, it
140 causes a meridional heat transport towards the poles through the atmosphere and ocean,
141 which takes place mainly through the movements of circulation systems and storms (Bjerknes
142 1922; Schultz et al. 2019; Ma et al. 2021).

143

144 Mid-latitude weather systems include both cyclones and anticyclones, but strong wind
145 situations are primarily associated with intense cyclones. The main paths that weather
146 systems and storms take, are called storm tracks (Hoskins and Valdez 1990; Blender et al.
147 1997; Chang et al. 2002; Ulbrich et al. 2009). Storm tracks form over the major ocean basins
148 of the Northern and Southern hemispheres and are closely related to atmospheric jet-streams,
149 which are areas of maximum upper-level wind ~~speeds~~ speed and determine the areas that are
150 prone to storms as discussed below in section 2.4 ~~below~~. These regimes set the propensity
151 with which weather systems take a more poleward or equatorward path on intra-seasonal time
152 scales, thus, offering potential predictability.

153

154 In its most basic form, atmospheric jet-streams (e.g. Feldstein and Franzke 2017) are a product
155 of the pressure gradient force, induced by the above-mentioned latitudinal air temperature
156 gradients, and the Coriolis force. For large-scale movements in the atmosphere, the wind is
157 diverted to the right (left) in the ~~Northern (Southern) Hemisphere~~ northern (southern) hemisphere
158 due to the Coriolis force. The resulting winds in the free atmosphere, above the boundary
159 layer, ~~blows~~ blow parallel to lines of equal pressure, in a balance between the pressure gradient
160 and the Coriolis force; also called geostrophic wind. The strength of the dominant westerly
161 winds over Western Europe is determined by the pressure difference between the subpolar
162 and subtropical regions over the eastern North Atlantic. The stronger the pressure difference,
163 the stronger the mid-latitude westerlies.

164

165 ~~In the boundary layer, the pressure gradient and Coriolis force are not in balance, because the surface~~
166 ~~characteristics, local conditions, vertical stability, and other effects play crucial roles in modifying the~~
167 ~~winds. Under the influence of the surface friction, the air movements are not parallel to the lines~~
168 ~~of equal pressure but have a tangential component from high to low pressure centers. On the~~
169 ~~regional to local scale, wind systems like the land sea breeze, and mountain valley wind~~
170 ~~systems develop due to differential heating conditions within comparatively small distances,~~
171 ~~which vary between day and nighttime.~~

172

173 Under hypothetical unperturbed conditions, the bands of maximum wind speed ~~—called jet-~~
174 ~~streams—~~ sit at 30° and 60° latitude in either hemisphere at upper levels of the troposphere,
175 due to surface friction. However, differential diabatic heating over land and the ocean, or
176 orographic surface features, such as mountains, do perturb the jet-stream in multiple ways.
177 As a result, in the extra-tropics of the northern hemisphere the jet-stream is commonly split
178 into a subtropical and mid-litudinal branch. While the former is mainly driven by angular
179 momentum transport by the thermally direct Hadley circulation (Held and Hou 1980), the latter
180 is primarily driven by the eddy momentum flux convergence provided by short waves that form

181 in regions of enhanced baroclinicity (Held 1975). Accordingly, the mid-latitude jet-stream is
182 referred to as an eddy-driven or polar jet-stream due to its proximity to polar latitudes.

183

184 In the atmosphere unstable conditions are needed for weather systems to form (Holton and
185 Hakim 2012). So-called baroclinically unstable conditions occur where we find strong
186 horizontal and vertical air temperature gradients. ~~These areas are conducive to the generation of~~
187 ~~positive vorticity, which is linked to upper-level divergence, which in turn causes the formation of~~
188 ~~short wave troughs. Vorticity is a variable which measures the local rotation of the atmosphere~~
189 ~~(Holton and Hakim 2012).~~ For example, the North Atlantic is an ideal source region for
190 baroclinically unstable conditions as very cold polar air is advected over moderately warm
191 ocean waters, leading to excessive air temperature gradients and, thus, pressure gradients,
192 which – under the influence of the Coriolis force – generate ~~positive vorticity and~~ enhanced
193 baroclinicity.

194

195 In the boundary layer, the pressure gradient and Coriolis forces are not in balance, because
196 the surface characteristics, local conditions, vertical stability, and other effects play crucial
197 roles in modifying the winds. Under the influence of surface friction, the air movements are not
198 parallel to the lines of equal pressure but have a tangential component from high to low
199 pressure centers. On the regional to local scale, wind systems like the land-sea-breeze, and
200 mountain-valley wind systems develop due to differential heating conditions within
201 comparatively small distances, which vary between day- and nighttime.

202

203 **2.2. How do cyclones form?**

204 While anti-cyclones are primarily associated with low wind conditions in ~~its~~their center and
205 strong winds are only found around its edges (i.e. co-located with another pressure system),
206 cyclones feature typically strong pressure gradients and are thus associated with strong winds
207 and wind gusts. Many extra-tropical cyclones develop under the influence of the mid-latitude
208 jet-stream, its associated baroclinicity and upper-air flow divergence. Other cyclones develop
209 as secondary cyclones in the trailing cold fronts of pre-existing systems and are more
210 influenced by lower-level processes such as latent heat release (Parker, 1998; Dacre and
211 Gray, 2009). Another large group of cyclones develop by the interaction of atmospheric waves
212 with ~~orography~~topography (McGinley, 1982; Radinovic, 1986). Focusing on the North Atlantic
213 sector for a European perspective, baroclinically driven (primary) cyclones develop typically
214 over the North Atlantic (~~e.g.,~~ Dacre and Gray, 2009), secondary cyclones develop further
215 downstream often close to the eastern North Atlantic (~~e.g.,~~ Priestley et al., 2020a), and the
216 orographically driven cyclones dominate in the Mediterranean basin (~~e.g.,~~ Trigo et al., 1999).

217

218 The most common conceptual models to describe extra-tropical cyclone development are the
219 Norwegian and the Shapiro-Keyser models (Bjerknes, 1922;₁ Schultz et al. 2019;₁ Dacre
220 2020). According to the Norwegian model, a stationary front forms between cold and warm air,
221 initiating strong vertical wind shear within the troposphere. A front is a density discontinuity
222 and, hence, separates cold and warm air masses. Typically triggered by an upper-level trough,
223 a cyclone begins to grow along this front where it develops a warm and a cold front. As the

224 cyclone deepens, both fronts become better defined and a warm sector develops. When the
225 cold front catches up to the warm front, the so-called occlusion process starts. At this stage,
226 the cyclone reaches its most intense period (Bjerknes 1922), followed by cyclone decay. In
227 the Shapiro-Keyser model, the initial development is similar, but the cold front does not
228 overtake the warm front, but rather builds a T-bone structure (see Fig. 16-24 of Schultz et al.
229 (Schultz et al., 1999)) instead of a narrowing warm sector during occlusion as in the classical model
230 (Shapiro and Keyser 1990).

231
232 Windstorms produce winds which are strong enough to cause damage; they typically have
233 windspeeds in excess of 15m/s (Wallace and Hobbs 2006). In order to quantify the impact of
234 ~~storms~~windstorms, it is important to know the parts of a storm where the strongest wind
235 ~~speeds~~speed typically ~~occur~~occurs. There are three zones where strong winds can occur: the
236 warm jet, the cold jet, and the sting jet (Clark and Gray, 2018). Hewson and Neu (2015; see
237 their Fig. 1) have developed a conceptual windstorm model to describe how strong winds may
238 develop associated with the passage of a cyclone during different stages of its development.
239 In most cases, the strongest winds are often associated with the passage of the cold jet at the
240 cold front. However, Shapiro-Keyser cyclones may rarely on occasion feature sting jets, which,
241 if reaching they reach the surface, may lead to even more damaging wind ~~speeds~~speed (Clark
242 and Gray, 2018).

243
244 The potentially most damaging events affecting Europe are commonly assigned to slow
245 movers, rapid developers, or serial storms (Mailier et al. 2006). Slow mover cyclones lead to
246 large accumulations of precipitation in the same area, often triggering severe floodings (Grams
247 et al., 2014). Rapid developers are fast deepening cyclones, often fulfilling the conditions for
248 a “bomb” (Gyakum and Danielson, 2000). When occurring close to Europe, many of these are
249 secondary cyclones. Finally, serial storms (also known as cyclone families) indicate that
250 multiple and related cyclones affect the same area within a comparatively short period of time,
251 leading, potentially, to severe cumulative losses (Mailier et al., 2006; Pinto et al., 2014). In
252 these clustering periods, the passage of storms occurs more frequently than may be expected
253 from a random process (e.g. if they would occur independently from each other (Vitolo et al.,
254 2009; Franzke, 2013; Blender et al., 2015). Two physical reasons are given in the literature
255 (Economou et al., 2015; Dacre and Gray 2020): i) the steering through the large-scale flow,
256 typically characterized by an intensified, quasi-stationary jet-stream extending towards Europe
257 and ii) the occurrence of secondary cyclogenesis.

259 **2.3. Storm's spatial Spatial characteristics of storms**

260 To analyse cyclones and storms, objective identification and tracking methods are needed
261 (Ulbrich et al., 2009; Neu et al., 2013). This leads to a Lagrangian perspective where certain
262 properties during the life cycle of the cyclone can be defined, by e.g. radius, propagation
263 speed, and spatial wind distribution. Various objective methods for the identification and
264 tracking of extra-tropical cyclones have been used to investigate their characteristics (Neu et
265 al., 2013; ~~Pinto et al., 2005; Ulbrich et al., 2009; Zappa et al., 2013;~~ Priestley et al., 2020a).

266
267 In the North Atlantic-European region, cyclone track densities show maximum values over the
268 western North Atlantic with a second maximum over the Mediterranean (Ulbrich et al., 2009;
269 Pinto et al., 2005). North Atlantic cyclone activity shows a tilt towards the northern North


270 Atlantic. While this can be found in different reanalysis products, [CMIP5 Coupled Model](#)
271 [Intercomparison Project phase 5 \(CMIP5\)](#) simulations are characterised by a bias of the
272 maximum and tilt in the North Atlantic, leading to more zonally oriented storm tracks (Zappa
273 et al. 2013). While many cyclones can be identified in the extra-tropics, only a subset of strong
274 cyclones lead to a high wind ~~speeds~~ [speed](#). See section 4.1 for related storm indices.
275

276 **2.4. Large-scale ~~Circulation Characteristics~~ [circulation characteristics](#) and their impact on** 277 **wind**


278 Storms and their related wind gusts are local in nature. Nonetheless, the large-scale
279 background circulation can still provide information in which areas strong winds are likely to
280 occur. Here, we apply the concept of atmospheric weather regimes (e.g. Hannachi et al., 2017)
281 to determine the characteristics of the large-scale circulation. Atmospheric weather regimes
282 are recurrent, dynamically relevant circulation patterns and allow the description of low-
283 frequency variability due to transitions between distinct regimes. Because of their preferred
284 occurrence locations, they potentially provide prediction and downscaling possibilities for
285 smaller scale weather events and extremes (Cassou et al. 2005).

286

287 To demonstrate the relation of specific regimes to preferred jet-stream patterns and storms,
288 we show weather regimes based on sea-level pressure fields from [the latest European Centre](#)
289 [for Medium-Range Weather Forecasts Reanalysis \(ERA5\)](#) (Hersbach et al. 2020) over the
290 North-Atlantic-Eurasian region (~~30~~ [30°N](#)-90°N, 90°W-90°E) for winters (December through
291 March, DJFM) from 1979-2020. The details of the applied regime analysis are described in
292 Crasemann et al. (2017). We identify 5 regime states (Fig. 1a): (1) Scandinavian/Ural blocking
293 (SCA-URAL BL), (2) the North Atlantic oscillation in the positive phase (NAO+), (3) blocking
294 over the North Atlantic (ATL BL), (4) North Atlantic trough (ATL TR) and (5) the NAO in its
295 negative phase (NAO-).



296



297

298 Figure 1: a) Weather regimes determined from ERA5 reanalysis data for December to March
 299 (DJFM). Shown are the regime patterns in terms of Sea-Level Pressure sea-level pressure

300 anomalies (shading), black contours indicate the climatology for DJFM, shown are isolines at
301 1000, 1005, 1010 and 1015 hPa. b) The ~~jet-stream~~ patterns ~~for the~~ associated ~~jet~~ with the
302 individual weather regimes, obtained by composites of zonal wind anomalies at ~~250hPa~~250
303 hPa (shading), black contours indicate the zonal wind climatology for DJFM, shown are
304 isolines at 20, 30, and 40 m/s. c) ~~Time-series-of-sequences-of-weather-regimes-(CL1-CL5)-The-date~~
305 ~~where-maximum-wind-speed-was-identified-over-land-for-storm-'Klaus'-is-marked-by-a-red-arrow.-d)~~
306 eXtreme WindStorms ~~storm~~XWS data base (Roberts et al. 2014) track for storm ~~'Klaus'~~Klaus
307 based on ERA5 data, identified with the method of Leckebusch et al. (2008). Storms are
308 defined by the exceedance of the local 98th percentile of near surface wind speed. Each dot
309 represents the position of the wind field center of storm "Klaus" for 6 ~~hr~~hour time steps from
310 22 January 2009, 06:00 to 26 January 2009, 00:00. The color of the dots shows the weather
311 regime of that date.

312
313 The characteristic patterns for the jet-stream associated with these weather regimes have
314 been obtained by compositing the zonal wind anomalies at 250 hPa over the days assigned
315 to each regime. The ~~5~~five ~~jet-regimes-stream~~ patterns (Fig. 1b) are very similar to those
316 obtained by previous studies (Dorrington and Strommen, 2020; Woollings et al., 2010; Franzke
317 et al., 2011).

318
319 The regime analysis assigns the atmospheric circulation of each day over the period 1979-
320 2020 to one specific cluster and enables a characterization of the large-scale background for
321 specific windstorm events. As one example, ~~Figs. 1b and Fig. 1c shows~~shows the ~~evolution-of-the~~
322 ~~prevailing-weather-regimes-over-the-winter-season-2008/2009.-From~~eastward movement of the
323 extreme storm Klaus from Jan 22 to Jan 26, 2009 ~~the-extreme-storm-'Klaus'-evolved.-'Klaus'~~
324 ~~moved-eastward~~ along an unusual southerly path. The storm ~~track and~~'Klaus' was characterized
325 by strong and record-breaking wind ~~speeds~~speed over northern Iberia and southern France
326 ~~(Liberato et al., 2011).~~ During the formation, intensification, and eastward movement of
327 ~~'Klaus'~~Klaus, the Atlantic trough weather regime associated with the central jet-stream
328 configuration prevails. ~~(Fig. 1c).~~ This central jet-stream pattern ~~sets~~sets the necessary large-
329 scale background flow for the development and movement of this extreme storm (Liberato et
330 al., 2011).

331
332 The concept of weather regimes enables the characterization of the large-scale atmospheric
333 circulation, in particular the jet-stream pattern, during extreme storm events. If changes in the
334 occurrence of these extremes can be related to an anomalous frequency of occurrence of a
335 specific weather regime, the use of these regime states offers ~~potentially~~potential predictability
336 of large- as well as small-scale wind impacts.

337 338 **2.5. ~~Storm's temporal~~ Temporal characteristics – ~~Seasonal Variability~~ of storms and seasonal** 339 **variability**

340 The occurrence of extreme wind ~~speeds~~speed and storms is subject to a strong seasonal
341 pattern in Europe. According to Young et al. (1999), windstorms occur 30% more frequently
342 in winter than in summer (see also Fig. S1). We compared the wind gusts from three reanalysis

343 products (ERA5 (Hersbach et al., 2020), COSMO-REA6 (Bollmeyer et al., 2015) and COSMO-
344 REA2 (Wahl et al., 2017)), to 145 German station observations (Kaspar et al., 2013). While a
345 direct comparison is difficult, qualitative statements on seasonality can be made with all data
346 sets. The number of occurrences of wind gusts is determined for certain wind speed intervals,
347 which are shown against the warning levels (WL) of the Deutscher Wetterdienst (DWD-
348 Reanalyses show an underestimation of the frequencies of the). The warning levels are defined by
349 6 different wind gusts compared speed thresholds: 14, 18, 25, 29, 33, and 39 m/s (Primo, 2016),
350 referring to the 4 WL (WL1- WL4), with WL2 and WL3 being divided into two intervals (DWD,
351 2021). Compared to observations, which become more extreme the wind gust frequencies are
352 underestimated in reanalyses. The higher the wind speed. Thereby gusts, the higher the
353 underestimation. Therefore, COSMO-REA2 shows a significantly better agreement with the
354 reference, especially for WL3 in summer and WL4 in winter. The benefit of the higher
355 resolution provided by regional reanalyses compared to their global counterparts is well
356 documented for near surface wind speeds speed (Niermann et al., 2019). Results shown in Fig.
357 S1 emphasize the importance of using high resolution models to represent extreme wind gusts
358 in reanalysis products.

359

360 Above 25 m/s there is a clear difference between summer and winter months, which becomes
361 stronger the higher wind speeds are speed is considered. In summer, wind speeds speed over
362 30 m/s de does not appear in the coarser reanalysis products ERA5 (~30km) and COSMO-
363 REA6 (~6km) at all and also for the high-resolution reanalysis COSMO-REA2 (2km) and the
364 point observations the occurrence of wind gusts of WL3 or WL4 in summer is smaller than in
365 winter by a factor of 10 to 100.

366

367 The intraseasonal intra-annual variability is not only visible in meteorological data but also in
368 loss data from insurance companies, (Hoeppel 2017, Franzke 2017), which shows the strong
369 impact of storms and especially winter storms on society and economic areas (Klawa and
370 Ulbrich, 2003). One The energy sector that is strongly affected by the occurrence of
371 windstorms, and especially their seasonal variability, is the energy sector. Due to the worldwide
372 effort to convert the energy system to renewable sources, the industry will have to deal more
373 with seasonal fluctuations in energy availability. The interest and the need for precise
374 knowledge of the wind conditions in various regions is therefore growing, as energy production
375 directly depends on it; for more details about wind-based energy production please see section
376 4.6.

377

378 **2.6. Convectively Winds induced winds by convective activity**

379 Most of the wind damage in temperate latitudes are is due to extra-tropical cyclones. However,
380 damage can also occur to structures, crops and forests from winds produced by convective
381 storms (e.g. Gatzen et al., 2020; Parodi et al., 2019); since our focus is more on extra-tropical
382 storms, we keep this part rather brief. The following conditions need to be met for convection
383 to occur (Wallace and Hobbs 2006): (1) The atmosphere needs to be conditionally unstable,
384 (2) there needs to be a reservoir of substantial moisture in the boundary layer, and (3) there

385 needs to be sufficient lifting due to low level convergence to cross the threshold to start the
386 instability.

387

388 Convective systems and storms can lead to severe wind ~~speeds~~ speed connected to tornadoes,
389 gust fronts and downbursts (Wallace and Hobbs 2006). Tornadoes are rapidly rotating air
390 systems which connect with the ground and can lead to devastatingly strong winds.
391 Downbursts are downward directed winds due to the negative buoyancy of the downdraft air.
392 Convective storms can also have gust fronts. The gust fronts form due to downdrafts in the
393 convective storm ~~by~~ forming a pool of cold, dense air which replaces the warmer, buoyant air
394 of the environment.

395

396 These downdrafts can lead to severe wind gust speeds at the surface (Bunkers and Hjelmfelt,
397 2021) with speeds of up to ~~150 km h⁻¹~~ 42 m/s. So far, relatively little attention has been paid to
398 wind damage to infrastructure, forests and agriculture from such events besides the studies
399 by ~~e.g.~~ Jim and Liu (1997) and Peterson (2000). Forest damage from thunderstorms ~~from~~ in
400 areas, which previously were rarely affected, such as eastern parts of Europe (~~e.g.~~ Nosnikau
401 et al., 2018; Sulik and Kejna, 2020) ~~are nowadays~~, but have experienced an increase in
402 convectively available potential energy and near surface moisture which can cause more
403 ~~commonly reported~~ thunderstorm activity (Taszarek et al. 2021). It is expected that
404 anthropogenic global warming will lead to an increase of ~~this type of~~ convective ~~activity~~ (storms
405 (Lepore et al. 2021; Taszarek et al. 2021; Diffenbaugh et al. 2013)).

406

407 Another type of convective storm is derechos, which are a clustering of downbursts, organized
408 by a line of thunderstorms (also called a squall line), that lead to widespread straight-line
409 winds, and can cause damaging winds. They occur frequently in the Great Plains area of the
410 USA (Ashley and Mote, 2005) but can occur around the world, including Central and Eastern
411 Europe (Gatzen et al., 2020). Some examples of the devastating impact of derechos on forests
412 are described in Goff et al. (2021), and Negrón-Juárez et al. (2010).

413

414 **3 Wind-surface interaction**

415 **3.1. The physics of fine scale interactions between surfaces and wind**

416 The characteristics of the wind ~~field and~~ speed and gustiness in a given environment are
417 dependent on surface characteristics, such as its roughness, all of which are highly influential
418 on the levels of damage caused ~~by wind~~. The momentum of the mean horizontal wind is
419 vertically transferred by turbulence, i.e. near the surface, large whirling air packages break up
420 ~~in~~ into smaller ones and their momentum dissipates into thermal energy or is absorbed by
421 roughness elements, such as trees and buildings. The strength of the wind is altered by
422 ~~orography~~ topography and the roughness of the surface (Stull, 2017; Kaimal and Finnigan,
423 1994); Finnigan et al. 2020). Thus, the damage level can vary dramatically at small scales
424 (Gromke and Ruck 2018; Forzieri et al. 2019).

425

426 Typically, the boundary layer above the ~~earth's~~Earth's surface is subdivided into three
427 sublayers: 1) a roughness sublayer that is characterized by the flow around obstacles and
428 varies locally and where mechanical turbulence dominates, 2) one or more inertial sublayers,
429 where the influence of the individual obstacles and surfaces is blended together and the
430 vertical energy fluxes are constant with height and 3) a mixing layer above, where the Coriolis
431 force gains influence and is often separated from the free atmosphere by a capping inversion
432 and an entrainment zone (Stull, 1988; Kaimal and Finnigan, 1994). The effect of buoyancy
433 and thermal stability is very important for the formation of strong winds, i.e. for cyclones and
434 thunderstorms. During storm events, high wind speed increases friction within the lower
435 boundary layer and also increases form drag by obstacles. The instability of the shear in the
436 flow created by the drag of the surface leads to turbulence, which affects the vertical exchange
437 of mass, momentum, and scalars. Thermal gradients near the surface are reduced or
438 disappear due to this mixing, which results in neutral stratification near the surface, i.e. thermal
439 stability need not be considered in the equations of the vertical wind profile (Stull, 1988).

440
441 As turbulent movements play a major role in the momentum transfer to the surface it is
442 important to regard shear forces and gustiness as the damaging characteristics of the wind
443 field (Gromke and Ruck 2018). For example, in forest ecosystems trees are blown down at a
444 mean wind ~~speeds~~ speed considerably lower than those estimated by pulling experiments
445 (Oliver and Mayhead, 1974; Milne, 1991). Boundary-layer eddies create a local increase in
446 wind speed and windshear close to the surface (Romanic und Hangan, 2020) and leading to
447 coherent eddies (Raupach et al., 1996). The loading due to these turbulent structures with
448 higher energy and momentum can be accounted for in a gust factor (Hale et al., 2015; Chen
449 et al., 2018; Holland et al., 2006; Usbeck et al., 2010). Since trees react to gusts like damped
450 harmonic oscillators (Mayer, 1987; Gardiner, 1992) there has been considerable debate about
451 whether the arrival frequencies of these coherent eddies could lead to resonant failure
452 (Gardiner, 1995; Peltola, 1996); however, this does not happen (Schindler and Mohr, 2019;
453 Schindler and Kolbe, 2020; Kamimura et al., 2022), probably due to the efficient damping of
454 trees (Spatz and Theckes, 2013). Besides the drag force of a plant (Rudnicki et al., 2004;
455 Queck et al., 2012; Vollsinger et al. 2005), the level of damage depends also on the acclimation
456 of plants to the wind (Telewski, 1995; Nicoll et al., 2019), which is a function of the maximum
457 wind speed (Bonnesoeur et al., 2016; Dêfossez et al., 2022). They are adapted to wind forces
458 and build stronger ~~root~~roots and wood structures depending on the main wind direction and
459 magnitude (Nicoll and Ray 1996; Tomczak et al. 2020).

460
461 Furthermore, the development of turbulence above and within the canopy is different between
462 naturally uneven aged woods and managed forests or plantations. Experiments showed that
463 the inflection of the wind profile (i.e. maximum gradient of wind speed) is weaker in
464 heterogeneous compared to homogeneous canopies, and that it occurs deeper within the
465 canopy, i.e. the displacement height is lower (Cionco, 1972; Belcher et al., 2012; Queck et al.
466 2016). Furthermore, homogeneous forests are more vulnerable than naturally uneven aged
467 woods (Everham and Brokaw, 1996; Mitchell 2013). Obviously, the adaptation to wind stress
468 is not restricted to single trees but extends to the structure of natural mixed woods too. The
469 characteristics of the tree (height, diameter, canopy size, wood properties), and the tree
470 resistance to uprooting and breakage are all affected by the level of wind exposure (Gardiner
471 et al. 2016). Recent experimental measurements of tree damage during a super typhoon

(Kamimura et al. (2022) has also shown that collisions between the crowns of individual trees and the crowns of their neighbours is extremely important in reducing tree movement during strong winds and contributing to their overall stability. These adaptations of plants to living in a windy environment must be considered when modelling the risk of wind damage to tree stands.

Large eddy simulations (LES) are used to better understand the complex current patterns and the acting wind forces near heterogeneous surfaces (Stoll et al. 2016, Takemi et al. 2020). These turbulence-resolving models include all the basic physical equations; however, they require considerable computer resources and are therefore unsuitable for operational use. Simplified mechanistic models (Holland et al. 2006, Gross et al. 2018, Duperat et al. 2021) parameterize the turbulence spectrum and operate on a larger spatial scale; thus, need less computational resources. Statistical approaches (Jung and Schindler 2015, Dupont 2016) focus on predicting critical thresholds at which wind damage occurs and are therefore efficient for operational damage prediction. The indices discussed in section 4 are based on empirical observations and have proven useful in a wide range of applications.

3.2. Mean wind and gust rates for different landscapes

The gustiness of the wind is critically important for assessing the likely impact of strong winds on forests, agriculture, and structures (e.g. Usbeck et al. 2010; Gardiner et al. 2016). The level of gustiness is known to be influenced by surface roughness (Table 1), the height above the ground, and wind speed (Ashcroft 1994; Verkaik 2000). Gust ratios are also affected by wind speed (see Born et al., 2012; their Fig. 2) and by the type of storm (Kramer and Marshall 1992; Harper et al. 2010).

Roughness Class	Aerodynamic roughness length (m)	Gust Ratio (3 s to 10 min)	Gust Ratio (3 s to 60 min)
1	0.003	1.36	1.44
2	0.01	1.42	1.49
3	0.03	1.48	1.56
4	0.1	1.58	1.66
5	0.3	1.74	1.85

Table 1. Wind rate (mean/gusts) for different landscapes. 3 s gust to 10 min and 60 min mean wind at 10 m height, by terrain category. From Ashcroft (1994). Roughness Classes: 1: off-sea wind onto flat coastal areas; 2: level grass plains, e.g. marsh; 3: standard category: fairly level terrain-mostly open fields with a few houses and buildings; 4: fairly level terrain with more hedges, trees and villages, farm buildings; 5: many trees and hedges, or fairly level wooded country or more open suburban areas.

4. Wind and storm related indices and critical thresholds

Wind and storm related indices and thresholds are a vital tool in assessing the likelihood and magnitude of damage. While there are many definitions for indices and thresholds, here we define an index as a number or a category, serving as an aggregated measure of a quality,

507 which can be reached by means of observation, arithmetic calculation, or different modelling
508 techniques. A threshold is defined here as a value taken or calculated from a numerical or a
509 categorical range, and when the threshold value is crossed, it indicates a significant increase
510 in the ~~probabilities~~probability for an event to take place or for a certain condition to be fulfilled.
511 Indices can be used to predict damage caused directly by wind or a storm, or when wind
512 modulates the damage caused by another process such as fire or drought. Since indices and
513 thresholds can be as effective as complex mechanistic models but more cost-effective, it is of
514 no surprise that there is a plethora of indices. There are general indices that are not bound to
515 a given ~~sector or~~ environment, but many of the indices and thresholds available require a
516 careful selection process according to the target ~~environment~~ecosystem. Below we provide an
517 extensive review of available indices, focusing on five ~~key~~ terrestrial ~~environments: forests,~~
518 ~~urban, transport, agriculture, and wind-based energy production-~~sectors.

519

520 **4.1. General storm indices—~~Storm:~~ scale and severity indices**

521 Classical wind scales are defined by phenomena caused by the interactions between wind
522 and the surface. A very prominent example is given by the Beaufort scale (Stull, 2017). It
523 classifies the effect of wind on wave generation, tree movement and the damage of buildings.
524 Similar scales exist for tornados, e.g., the Fujita scale and the Torro scale (Kirk, 2014), which
525 relates the tornado intensity to damage description. As short gusts and shear forces are very
526 important factors of storm risk, the Enhanced Fujita scale includes further information **on**
527 **derived maximal tangential 3s gust speeds (Fujita, 1981).** Recently an improved wind speed
528 scale and damage description has been suggested for Central Europe (Feuerstein et al.,
529 2011). Finally, The Saffir–Simpson hurricane wind scale (Ellis et al., 2020) is based on the
530 highest wind speed averaged over a one-minute interval 10 m above the surface. It can
531 provide some indication of the potential damage a hurricane will cause upon landfall.

532 Several storm severity indices have been developed to quantify the severity of a windstorm
533 regardless of the ~~environment type-~~land cover. These indices are used to identify severe winter-
534 storms and analyze their impacts and to investigate storm trends in past and future climate
535 conditions. They often include the cube of the wind speed, assuming a proportionality of the
536 dissipation rate of the wind kinetic energy to damage. A selection of these indices is presented
537 in Table S1.

538

539 From an historical context, one of the earliest storm severity indices was developed by Lamb
540 (1991) to grade and rank storms based on the greatest observed wind speed over land, the
541 area affected by damaging winds and the overall duration of occurrence of damaging winds.
542 Later, in a study by Klawa and Ulbrich (2003), the wind speed values were scaled with the
543 local 98th percentile. Based on this approach, Leckebusch et al. (2008) identified and tracked
544 windstorms in time and space and computed an event-based storm severity index that
545 quantifies the potential impact of a storm. This index considers the relation of the maximum
546 daily wind speed to a certain local percentile of ~~daily~~maximum ~~daily~~wind speed (e.g. the 95th
547 or 98th) as well as the affected area. For example, in their study they found a trend for an
548 increase in severity of storms during 1960–2000 and for 2070-2100 under anthropogenic
549 climate change conditions. Pinto et al. (2012) ~~extend~~extended this approach by taking into
550 account the exposure and including local population levels in a Loss Index, resulting in the

551 finding that the maximum storm losses for current climate conditions are likely to be exceeded
552 in the future. Additionally, Haylock (2011) used a storm severity index to identify the severest
553 storms for ~~72h~~72 hours storm footprints. This index considers the latitude and the excess of
554 the maximum wind speed over a ~~72h~~72 hour period taken from six-hourly values over a
555 threshold (e.g. the local 90th percentile of wind ~~speeds~~speed).

556 557 **4.2. Forests**

558 **4.2.1. Topographic indices**

559 Many topographic indices have been used for assessing the risk of wind damage to forests
560 (see Table S2). These indices can be based on elevation, slope characteristics such as
561 compass angle, aspect, and curvature, or are more complex such as TOPEX (Topographic
562 exposure; Quine and White, 1998) which was developed as part of a risk assessment method
563 (Windthrow Hazard Classification) to predict the height at which trees could be expected to be
564 first damaged (Miller, 1986). TOPEX is the sum of the angle to the horizon in the eight principal
565 points of the compass and can be calculated for different distances from the point of interest.
566 Furthermore, such indices can be used to create even more complex predictive systems. For
567 instance, when TOPEX is combined with elevation and aspect it produces a system called
568 DAMS (Detailed Aspect Method of Scoring; Quine and White, 1993) for predicting wind speed
569 variation in the landscape. This system is entirely based on topographic measures and
570 compares favorably with modelling systems based on solutions of the fluid equations (Suárez
571 et al., 1999).

572
573 The actual variation of wind speed with height above the ground is a function of the surface
574 roughness and the topography. Predicting variations of wind speed across flat surfaces is
575 relatively straight forward, especially for strong winds by using a measure of the aerodynamic
576 roughness of the surface and a logarithmic wind profile (Garratt, 1980; Stull, 1988). Even in
577 stable or unstable conditions the profile can be modified with the addition of the diabatic term
578 ψ_m (Kaimal and Finnigan, 1994; Panofsky and Dutton, 1984). Often the roughness of the
579 surface is simplified into different roughness classes (Troen and Petersen, 1989) to allow for
580 easier estimation of the surface roughness. However, when even-strong-winds flow over
581 topography the simple logarithmic profile breaks down and the shape of the wind profile
582 strongly varies between the upwind slope, the crest of the hill and the downwind slope, where
583 the flow may even separate (Belcher et al., 2012). Thus, one should not only calculate
584 topographic indices for the target locations but calculate also for the neighboring
585 environmentareas and assess the change in value between the target location and its
586 surroundings (Ruel et al., 1997; Schindler et al., 2012; Murshed and Reed, 2016).

587
588 We reviewed the literature, focusing on studies using topographic indices to assess and
589 predict damage caused by strong winds, as topographic indices are a common feature in
590 modelling wind damage in forests (Table S3). The most commonly used variables were (Fig.
591 2): elevation, slope, aspect and TOPEX. We assessed the usefulness of the four most
592 commonly used topographic indices in modelling forest damage according to their inclusion in
593 final models and according to the importance/influence metrics reported. We note that most


594 studies employed a multivariate modelling approach, thus, a certain variable may appear less
595 useful due to overlap in the variance explained with another variable, but not necessarily due
596 to the variable's lack of explanatory power (Scott and Mitchell, 2005). Furthermore, there are
597 other topographic indices that were not tested so far for their contribution in forest damage
598 prediction (e.g. see Florinsky, 2017).

599

600 Elevation was useful in about a third of the studies, and was particularly useful when the study
601 area was very large, encompassing an entire region, state or country (Díaz-Yáñez et al., 2019;
602 Kramer et al., 2001; Torun and Altunel, 2020; Mayer et al., 2005) or when there was a strong
603 gradient of elevation, preferably reaching above 900 m ~~as~~ above sea level (Krejci et al., 2018;
604 Pasztor et al., 2015; Torun and Altunel, 2020; Kramer et al., 2001; Mayer et al., 2005). ~~This~~
605 ~~relevance assessment for elevation is new to our best of knowledge.~~ The trend in the correlation
606 between elevation and forest damage was ~~both~~ found to be inconclusive, to be both positive
607 (Díaz-Yáñez et al., 2019; Krejci et al., 2018; Pasztor et al., 2015;) and negative (Mayer et al.,
608 2005; Albrecht et al., 2013), or only present for a certain range of elevation (Albrecht et al.,
609 2013; Torun and Altunel, 2020;). While there is an expectation for an increase in forest
610 damage with higher ~~altitudes~~ elevation due to an increase in wind speed (Machar et al., 2014),
611 diversity of trends can stem from the involvement of other topographic indices that may contain
612 ~~the~~ similar information (e.g. slope or TOPEX), and also due to varying levels of acclimation of
613 trees to the wind conditions present at different ~~altitudes~~ elevations (Gardiner 2021).


614

615 The slope was shown to be useful in about half of all articles, however it is difficult to observe
616 a clear relation to forest damage. In articles that identified a contribution of slope, the relation
617 of damage with slope was found to be either positive (Díaz-Yáñez et al., 2019) or negative
618 (Mayer et al., 2005; Morimoto et al., 2019; Schütz et al., 2006). But an important deciding
619 factor can be the aspect of the slope (useful in about 40% of all articles) as there is often an
620 interaction between the two (Suvanto et al., 2018, 2016; Díaz-Yáñez et al., 2019; Hanewinkel
621 et al., 2014). In this sense, the aspect likely indicates the forest's susceptibility to wind coming
622 from a certain direction, as in most cases of usefulness of aspect, the slope was also useful.
623 Finally, TOPEX was found by 77% of articles as useful, and when a trend was reported, all
624 studies reported higher ~~damages~~ damage or probabilities for forest damage being associated
625 with more exposed locations (Albrecht et al., 2013, 2012; Jung et al., 2016; Morimoto et al.,
626 2019; Mitchell et al., 2001; Taylor et al., 2019). One of the reasons for TOPEX's usefulness is
627 that it does not strongly overlap with the information contained in other wind-based variables
628 (Albrecht et al., 2019; Schindler et al., 2012). However, when TOPEX is calculated only for a
629 certain cardinal direction (e.g. west) it contains information that is very similar to aspect.



630

631



632

633 Figure 2. An assessment of the usefulness of the most commonly used topographic indices.

634

635 **4.2.2. Fine scale wind and surface interactions**

636 Interactions between the surface and the wind field are controlled by surface roughness,
 637 absolute wind velocity and atmospheric stability. The commonly used index related to
 638 surface interactions that is relevant for wind and storm damage is the critical wind speed
 639 (CWS). The CWS defines the threshold wind speed for overcoming the maximum resistance
 640 to stem breakage or uprooting of a tree (Gardiner et al., 2016; Peterson et al., 2019; Hale et
 641 al., 2015; Chen et al., 2018; Holland et al., 2006). CWS is a standard term in forest ecology.
 642 The typical averaging interval for CWS is a period of a few minutes, e.g. 3 minutes (Peltola
 643 and Kellomäki, 1993), 10 minutes (Dupont et al., 2015, Peltola et al. 1999) or 60 minutes
 644 (Hale et al., 2015). CWS is estimated either at a height of 10 m above the canopy or at the
 645 tree top at the stand edge.

646 One of the governing quantities to describe the interactions between wind forces and stem
 647 breakage or uprooting is the applied maximum bending moment (BMmax) (Quine et al.

2021), which is the sum of wind forces in the tree crown and the additional turning moment due to stem bending and deflection of the stem and crown of a tree (Peltola, 2006). ~~The amplitude of BM_{max} depends on wind speed and is a function of tree species, tree size, inter-tree spacing and tree location relative to any forest edges.~~ BM_{max} calculation refers typically to the mean bending moment (BM_{mean}) and a gust factor (see e.g., Gardiner et al. 1997). A tree uproots if its BM_{max} at the ground level exceeds the resistance of the root–soil plate, and a tree breaks if its BM_{max} at breast height (1.3 m) exceeds the critical value of the stem’s modulus of rupture (Peltola et al. 1999, Quine et al., 2020). The gust factor is parameterised by wind measurements (field or wind tunnel) and depends on the spacing/height ratio of tree stands and the location relative to the forest/stand edge (Gardiner et al. 1997; Quine et al., 2020). The wind measurements are taken from the top of the canopy, and the bending moment is typically determined from the level of zero-plane displacement (e.g., 0.8 of the tree height; Gardiner et al. (1997)). Nevertheless, measurements of the effects (Gardiner et al. 1997) as well as directly solved finite element models of the crown architecture (Ruy et al., 2022) have shown the influence of crown architecture on the maximum bending moment. Therefore, the gust factor used in the calculation of BM_{max} may need to be varied according to stand composition and tree type.

665

The probability of occurrence of CWS, as a measure of storm damage risk for specific forest stands, depends on the statistics of wind velocity, e.g., on hourly maximum synoptic winds (u_{max} : Usbeck et al., 2010, Chen et al., 2018) or maximum geostrophic wind speed (Blennow and Olofsson, 2008). CWS is used to parameterize impact models for the estimation of storm risk in forests such as ORCHIDEE-CAN (Chen et al., 2018), SWAN/ADCIRC (Akbar et al., 2017), GALES and HWIND (Peltola et al., 1999; Gardiner et al., 2000, 2008).

673

The key parameter in the calculation of CWS is the diameter at breast height (DBH_{1.3}), which is a standard parameter in forest inventories ~~and~~. DBH is commonly defined as the stem diameter ~~in~~ at 1.3 m above the ground (Peterson et al., 2019; Gardiner, 2021; Hale et al., 2015, Chen et al., 2018; Holland et al., 2006; Hanewinkel et al., 2014; Beck and Dotzek, 2010; Gardiner et al., 2008; Peltola, 2006). DBH is the most used structural parameter due to ~~the~~ its easy and practicable measurement and due to its widespread application in forest management (Liu et al., 2018). DBH is also used to derive other structural parameters like tree height and Leaf Area Index (LAI) which can also be derived from NDVI normalized difference vegetation index (NDVI) as a standard product of satellite remote sensing. These structural quantities are important both for statistical analysis and for the parameterization of storm risk models.

684

Other important parameters for calculating CWS are the mean drag coefficient (c_d) which is part of the equation of the drag force (Vogel, 1989; Akbar et al., 2017; Dupont et al., 2015), turbulence intensity, gust duration, ~~gust factor~~ (Hale et al., 2015; Chen et al., 2018), tree density (Peterson et al., 2019; Albrecht et al., 2015), tree height, crown projection area and crown volume (Peterson et al., 2019; Gardiner, 2010, 2021; Albrecht et al., 2015; Hale et al., 2015; Chen et al., 2018; Dupont et al. 2015; Peltola, 2006), and tree species (Hanewinkel et al., 2014). Additionally, the edge factor index describes the influence of a tree’s position relative to a forest edge, the shape of the forest edge and the width of any upwind gap (Chen et al., 2018; Gardiner et al., 2010; Peltola, 2006).

694

695 The severity of storm damage depends on the ability of a tree to resist the applied bending
696 moment from the wind and on the stability of the root soil complex (Nicoll et al., 2006;). If soil
697 water content is close to saturation the critical resistive moment of trees (BMcrit) can be
698 reduced significantly during storm events, which could become increasingly important with the
699 increasing frequency of heavy winter rain in temperate forests in the context of regional climate
700 change (Défossez et al., 2021).

701

702 The uncertainty of [critical wind speed CWS](#) results from the consecutive solving of analytic
703 equations including accumulated uncertainties of the different input quantities. Additional
704 uncertainties result from the differences in the models used. Sensitivity tests using GALE5
705 (Locatelli et al., 2017) and HWIND with a variation of the input parameters of +/-20% lead to
706 [a](#) more than 20% change in [critical wind speed CWS](#). For example, CWS is especially sensitive
707 to changes in [diameter of breast height DBH](#). The measurement uncertainty of the [diameter of](#)
708 [breast height DBH](#) ranges between 2 and 10% ~~dependent~~[depending](#) on the absolute diameter
709 (Qin et al., 2019). Applied in HWIND and GALE5 the variation of [diameter of breast height DBH](#)
710 of +/- 20% lead to changes of [Critical wind speed CWS](#) of +30% and -46% (Gardiner et al.,
711 2000). The most comprehensive analysis of wind risk model uncertainty was made by Locatelli
712 et al. (2017) who found that tree DBH, tree height and inter tree spacing were the most critical
713 factors.

714

~~715 Classical wind scales are defined by phenomena caused by the interactions between wind
716 and the surface. A very prominent example is given by the Beaufort scale (Stull, 2017). It
717 classifies the effect of wind on wave generation, tree movement and the damage of buildings.
718 Similar scales exist for tornadoes, e.g., the Fujita scale and the Torro scale (Kirk, 2014), which
719 relates the tornado intensity to damage description. As short gusts and shear forces are very
720 important factors of storm risk, the Enhanced Fujita scale includes further information to
721 [derived maximal tangential 3s gust speeds \(Fujita, 1981\)](#). Recently an improved wind speed scale
722 and damage description has been suggested for Central Europe (Feuerstein et al., 2011).
723 Finally, The Saffir Simpson hurricane wind scale (Ellis et al., 2020) is based on the highest
724 wind speed averaged over a one minute interval 10 m above the surface. It can provide some
725 indication of the potential damage a hurricane will cause upon landfall.~~

726

727 **4.3. Urban [areas](#)**

728 **[4.3.1](#) The urban boundary layer**

729 The small-scale interactions of the wind field with urban surfaces are significantly different
730 from natural surfaces due to high three-dimensional variability of impermeable artificial
731 obstacles (buildings). These differences lead to a higher mean surface roughness of the urban
732 surface (Grimmond and Oke, 1999; Oke et al., 2017) combined with a general attenuation of
733 the mean wind speed, [the wind speed averaged over some time period](#) (Chen et al., 2020),
734 as compared with more natural surfaces. The level of increase in roughness depends on the
735 morphology - density, size, and composition - of the obstacles along the flow direction. The
736 height of the roughness layer is 2-3 times the mean height of the buildings. Within this layer,

737 mechanical turbulence generation dominates, and average wind profiles can only be assumed
738 above the roughness layer, within the inertial sublayer. The averaged roughness of an urban
739 surface is described by roughness length z_0 within equations for vertical wind profiles. This
740 parameter serves as ~~an~~ a useful index for the prediction of turbulent impulse transfer and ~~also~~
741 for damage prediction, which are derived ~~based on the basis of~~ building height, areal fraction
742 and frontal area index (Grimmond and Oke, 1999). At finer scales, wind speed shows high
743 spatio-temporal variability. Thus, when using indices based on averaged wind speed, it is also
744 important to consider that due to the small-scale aerodynamic and thermal heterogeneities of
745 urban infrastructure (buildings and trees), the local magnitude of the wind speed is temporarily
746 larger than under rural conditions (Droste et al., 2018). The reasons for this anomaly are again
747 the inflexibility and impermeability of technical structures and buildings. These features cause
748 canalization of flows and stronger turbulence generation compared to natural surfaces. There
749 is also a diurnal-nocturnal distinction in the formation of local thermal wind systems, with street
750 canyon wind during the day and a nocturnal inflow to the urban heat island (Droste et al., 2018;
751 Lindén and Holmer, 2011). Thus, indices in urban areas should account for both spatial and
752 temporal heterogeneities.

753

754 **4.3.2 Indices for estimating damage to individual buildings**

755 Damage occurs either directly by wind pressure or indirectly by the impact at high speed of
756 objects and debris moved by the wind (Tamura, 2009). At the level of individual buildings, air
757 movement results in wind pressure on the building surface and an applied force ~~called the wind~~
758 ~~load.~~ ~~Damages.~~ Damage to buildings caused by extreme wind loads include resonance and
759 vibration induced damage, damage to roof tiles or sheet roofing, roof lift off and the collapse
760 of walls or entire houses.

761

762 The occurrence and type of ~~damages~~damage depend on the level of exposure as well as the
763 structural vulnerability of the individual buildings to ~~local~~-severe local winds. The European
764 wind loading code EN 1991-1-4 regulates how to adapt the structural design of buildings to
765 the local wind climate. The code defines basic wind velocities for different geographical wind
766 zones based on the 50-year return level of 10-~~minute~~ min wind ~~speeds~~speed at a 10 m height.
767 In Germany, for example, the basic wind velocities range from 22.5 m/s in wind zone 1 (inland
768 areas in southern Germany) up to 30 m/s in wind zone 4 (coastal areas). The basic wind
769 velocities are further adjusted based on the height above ground and the terrain roughness to
770 account for short term wind fluctuations. Terrain roughness is classified in five categories
771 ranging from coastal areas to cities with a high building density. Additionally, where
772 ~~orography~~topography (e.g. hills, cliffs etc.) increases wind velocities by more than 5% the effect
773 is taken into account using ~~an-orography~~ a topographic index, as the ratio of the mean wind
774 velocity at the height above the terrain to the mean wind velocity above flat terrain. Finally, the
775 wind ~~speeds-are~~speed is used to compute the local peak velocity pressure which is a
776 fundamental index for the determination of all wind loads for a specific building (Schmidt 2019).
777 Nonetheless, assigning critical wind speed thresholds to building damage is rather difficult
778 given the heterogeneity of buildings, topography and ~~environments~~land-cover.

779

780 **4.3.3 Storm loss models: estimating damage on a district level**

781 Often there is little to no information on the actual damage to individual buildings or small-
782 scale urban structures. Instead, storm loss models come into play, and they relate wind
783 ~~speeds~~speed to actual building damage data, usually by applying statistical modeling
784 techniques. In some cases, these models rely on the use of wind indices like the exceedance
785 of local wind ~~speeds~~speed over a critical threshold to calculate monetary loss. In other cases,
786 the model itself calculates a damage index. The purpose of storm loss models is, among other
787 things, to assess current risk to residential structures or to estimate expected losses in future
788 climate conditions. It is often assumed that the maximum daily ~~windgust~~ speed (24-hour
789 maximum) is the most influential factor compared to other wind parameters like daily mean
790 wind speed or wind direction and is commonly used in indices as well as in loss models (Donat
791 et al., 2011; Klawa & Ulbrich, 2003; Koks & Haer 2020; Leckebusch et al., 2008; Pardowitz et
792 al., 2016; Welker et al., 2021).

793

794 Building damage data on a district level is usually provided by insurance companies and is
795 analyzed in the form of the loss ratio, which is the amount of insured loss per day and district,
796 divided by the corresponding sum of insured value, or claim ratio, which is the number of
797 affected insurance contracts per day and district, divided by the corresponding total number
798 of insurance contracts (Prahl et al. 2015).

799

800 The functional relationships between wind and damage are usually referred to as damage
801 functions. As the relationship between damage and wind depends strongly on local
802 ~~condition~~conditions like building or city structure, there is no universal function or model and
803 instead a variety of different damage function formulations are in use. A detailed overview can
804 be found in Prahl et al. (2015). Power-law damage functions are common. Different exponents
805 for these functions can be found in the literature ranging from 2 to 12 (Münchener

806 Rückversicherungs-Gesellschaft. 1993; Heneka et al 2006; Prahl et al. 2012). Some damage
807 functions also assume an exponential form (Prahl et al., 2015).

808

809 Another type of model are probabilistic models which calculate the probability that a certain
810 loss threshold is exceeded (Pardowitz et al., 2016; Prahl et al., 2012). Some examples ~~for~~^{of}
811 existing models are shown in Table 2. Most models still need to be fitted to local conditions
812 and validated with existing damage data. Model selection depends on the available data.

813

$L(v_{max}) = 2.48 * 10^7 * exp(0.48v)$	Dorland et al. (1999)
$D(v) = \left(\frac{v_{max}}{v_{98}} - 1\right)^3$	Klawe & Ulbrich (2003)
$LR(v, f(v_{crit}), \Delta v) = \int_{-\infty}^v f(v_{crit})G(v)dv_{crit}$ $G(v) = \begin{cases} 0, & v < v_{crit} \\ D(v), & v_{crit} \leq v \leq v_{tot} \\ 1, & v_{tot} \leq v \end{cases}$ $D(v) = \left(\frac{v - v_{crit}}{v_{tot} - v_{crit}}\right)^2$	Heneka et al. (2006)
$P(LR > th) = \frac{exp(a + b * v)}{1 + exp(a + b * v)}$	Pardowitz et al. (2016)

814 Table 2: A selection of damage functions including exponential damage relationships
815 (Dorland et al. 1999), power law damage functions (Klawe & Ulbrich 2003, Heneka et al. 2006)
816 and probabilistic damage functions (Pardowitz et al. 2016). a , b denote coefficients, D a
817 damage index, $f(v_{crit})$ a normal distribution of the critical wind speed, G a damage ratio, L a
818 loss, LR a loss ratio, $P(LR > th)$ a probability that a certain loss threshold will be exceeded, th
819 a loss threshold, v a mean daily wind speed, v_{98} the 98th percentile of the local wind speed,
820 v_{crit} the critical wind speed at which buildings are assumed to suffer damage, (comparable to
821 the CWS used for trees), v_{max} the maximum daily gust speed, and v_{tot} the buildings total wind
822 speed at which maximum damage is reached.

823

824 4.4. Transport

825 Transport systems are the backbones of modern societies. Disruptions within the transport
826 systems can have serious cascading effects that can cause large costs. Weather in general,
827 and windstorms in particular, can affect all aspects and functions of transport systems
828 (Leviäkangas et al. 2011). However, relevant thresholds of wind speeds and their
829 impacts are different depending on the mode of transport. Vajda et al. (2014) identify three
830 wind gust thresholds of increasing magnitude, which they relate to general impacts and
831 consequences within different parts of the European transport system: (i) Wind gusts >17 m/s:
832 Adverse impacts on the transport system may start to occur, especially if the resilience of the
833 exposed part of the system is low, but disruptions are rather local. For example, some
834 windthrow of trees can occur along railways and roads, leading to local problems to road
835 and rail traffic. Furthermore, operation of smaller boats could be suspended due to reduced
836 maneuverability, (ii) Wind gusts > 25 m/s: Some adverse impacts can be expected, such as
837 windthrow and electricity cuts occurring on a larger scale. In addition, delays and cancellations
838 in air, rail, road traffic and disturbances of ferry traffic can be expected, and (iii) Wind gusts >
839 32 m/s adverse impacts are very likely to occur, windthrow of trees can be expected on a large
840 scale, leading to long lasting power failures and delays, and cancellation of rail and road traffic.
841 Furthermore, damages to traffic control devices and structures can occur, airports can
842 be closed, and ferries stay in harbour due to reduced visibility and high waves.

843

844 The effect of wind on road safety is not extensively explored in the literature (Theofilatos and
845 Yannis 2014). In general, the number of road vehicle crashes caused by strong wind is small
846 compared to the total number of crashes (Edwards, 1998). However, studies have identified
847 specific types of crashes which typically occur under strong wind conditions: overturning, side
848 slip and rotation crashes (Baker, 1986), with ~~high-sided lorries~~trucks, vans, or buses being
849 particularly affected ([Becker et al. 2022](#), Baker 1992). A critical rollover wind velocity of 20 m/s
850 was found for high-sided lorries in crosswind situations (~~Snaebjornsson~~[Snaebjornsson et al.](#)
851 2007). Particularly dangerous situations with strong crosswinds can occur on bridges (Wang
852 et al., 2014; Charuvisit et al., 2004). A vehicle overturning model is applied by the British
853 Meteorological Office (Hemingway et al. 2020). It estimates the risk of overturning based on
854 wind gust thresholds ranging from 23 to 45 m/s, depending on vehicle type, loading, driving
855 speed and wind direction. In addition to direct effects of high wind ~~speeds~~speed on road
856 vehicles, indirect effects like blocked roads due to falling trees or drifting snow can affect road
857 transport (Leviäkangas et al., 2011).

858

859 The most frequent impact of high wind speed on railway transport is the blockage of tracks
860 due to windthrow of trees or drifting snow, as well as loss of electricity due to damaged
861 overhead lines (Leviäkangas et al. ~~2011~~.2011), [an example of a compound event](#). Only in rare
862 cases, extreme gusts exceeding 40 m/s can blow trains off the track (Sprenger et al., 2017).
863 Mean winds above 17 m/s or wind gusts above 30 m/s have been identified as thresholds
864 relevant for wind induced ~~damages~~damage to railway transport (Thornes and Davis 2002).
865 Shaking of overhead cables can cause damage to masts and pantographs on trains.
866 Consequences of windthrow can be collisions of trains with fallen trees. Precursory measures
867 to prevent collisions are reduced traveling speeds or ~~canceling~~cancelling/limiting train services,
868 commonly leading to ~~wide-spread~~widespread delays.

869

870 The most common impacts on ports are delays due to the disruption of loading and unloading
871 procedures, as well as direct ~~damages~~damage to infrastructure. For example, maximum wind
872 ~~speeds~~speed recommended for crane operations are around 18 m/s, depending on the design
873 of the crane (TT Club et al. 2011). This can have effects on the overall efficiency of ports
874 (Garcia-Alonso et al., 2020). From 88 disruptive events affecting ports and their surrounding
875 seas in the UK between 1950 and 2014, 36% were attributed to wind storms and 12% to storm
876 surges, while the others were mainly related to human error and mechanical faults (Adam et
877 al. 2016).

878

879 In the case of inland waterway transport, there is generally no large impact of wind on vessels,
880 since they are sufficiently wide and stable (Leviäkangas et al. 2011). However, at specific
881 locations with high local wind ~~speeds~~speed due to topography or at locations which are difficult
882 to navigate, navigation of pushed convoys without bow thrusters may be suspended in case
883 of high wind ~~speeds~~speed. In addition to location-specific issues, the vulnerability of vessels to
884 strong wind is strongly dependent on the ~~vessels~~vessel's characteristics (Schweighofer, 2014).
885 For specific types of inland container vessel mean wind ~~speeds~~speed of 18 m/s can lead to
886 flooding of open cargo-holds due to heeling and rolling (Hofman and Bačkalov 2010) and
887 increase the risk of sliding of empty containers on the upper tiers.

888

889 In the case of deep-sea shipping, vessels like large container ships are rarely lost at sea.
890 However, high wind ~~speeds~~speed impose the danger of container losses (Allianz, 2019). The
891 global average annual loss of containers is estimated ~~to be~~ up to 10,000 per year (Frey and
892 DeVogelaere 2014). These numbers are low compared to a total number of more than 200
893 million containers transported per year, but each container lost at sea can lead to a significant
894 safety and environmental hazard. In contrast to container ships, losses of dry bulk carriers are
895 often related to heavy weather conditions (INTERCARGO, 2018). Forecasts of ocean surface
896 conditions are important for route planning to avoid areas affected by windstorms (Kite-Powell
897 2011).

898

899 Airplanes are affected by strong winds mainly during take-off and landing. Dangerous
900 situations related to wind are mainly caused by abrupt changes in wind speed due to wind
901 gusts, wind shear or microbursts (strong downward movements of air within and below
902 thunderstorms). In the USA, for example, 48% of weather-related aviation accidents are due
903 to adverse wind conditions, and of those wind-related accidents 34% are due to crosswinds
904 and 29% due to wind gusts (Jenama and Kumar, 2013). Therefore, for safety reasons,
905 separation distances between airplanes are increased under high-wind conditions.
906 Furthermore, depending on the wind direction, runways may need to be closed. At London
907 Heathrow, for example, tailwinds of more than ~~5 knots~~(2.6 m/s) and crosswinds above ~~25 knots~~
908 ~~(13 m/s)~~ are avoided by changing flight direction or runways (Pejovic et al. 2009). This can
909 lead to delays, diversions and cancellations of flights. At London Heathrow Airport, an
910 increase in wind speed of ~~1 knot~~(0.5 m/s) above the mean ~~will~~increases the probability of
911 delay by 8% (Pejovic et al. 2009).

912

913 **4.5. Agriculture**

914 ~~The agricultural sector is~~**4.5.1 Wind damage in the agriculture**

915 ~~Agricultural production levels are~~ crucial for the ~~European~~worldwide economy ~~and is a key~~
916 ~~employer in rural areas~~. Wind leads to substantial environmental, social, and economic losses
917 and has distinct impacts on agriculture: physical damage to crops and related infrastructure,
918 soil erosion including nutrient and soil carbon removal, dust storms, higher evapotranspiration
919 rates of plants, as well as negative impacts on flowering, pollinators and fruits (~~e.g.~~Torshizi et
920 al. 2020).

921

922 Wind can damage crops through various mechanisms. Most vegetables already react to low
923 wind ~~speeds~~speed of around 4 m/s with physiological adaptations that affect the quantity or
924 quality of the harvest (Rouse and Hodges, 2004). Most kinds of crops can also be directly
925 damaged by abrasion from windblown dust particles or rubbing leaves (Brandle et al., 2004).
926 In orchards, wind can cause a considerable loss by breaking branches or damaging the fruit
927 set (Gardiner et al., 2016). For cereals, lodging (i.e. flattening) is probably the most important
928 impact of wind (Berry et al. 2004). For instance, wheat yield is usually reduced about 25%
929 when fields are lodged (Baker et al., 2014), but the loss can reach up to 50-68% (Berry and
930 Spink, 2012) and ~~also~~the yield of other cereals can decrease by 35-50% under these
931 conditions (Rajkumara, 2008). In most cases lodging is caused by strong wind accompanied

932 by heavy rain, whereby the maximum wind speed is the critical parameter (Mohammadi et al.,
933 2020; Niu et al., 2016). The vulnerability of plants to lodging depends on many factors, for
934 example, excessive usage of nitrogen fertilizers increases lodging vulnerability of wheat (Berry
935 et al., 2019). It is therefore difficult to determine general threshold values for a critical wind
936 speedsspeed. However, typical lodging threshold wind speeds at 10 m above the ground for
937 maize, oilseed rape, oats and wheat can be assumed to be 11.5, 14.8, 15.1 and 16.5 m/s
938 respectively (Joseph et al., 2020; Baker et al., 2014).

939
940 In general, plants exposed to wind are shorter and have thicker leaves and mature plants are
941 less vulnerable to wind stress than younger plants (Brandle et al., 2004). Therefore, land users
942 must carefully balance between the investment in wind adaptation measures and yields
943 (Wiréhn et al. 2020). However, the careful selection of wind resistant varieties with short-
944 stems (Berry et al., 2014), climate resilient plants, or the use of cultivar mixtures can
945 significantly improve wind lodging stress resistance, as demonstrated in wheat (Kong et al.
946 2022). Field fruits react differently to wind exposure: vegetables in general have a very low
947 tolerance to wind stress, cucumber, pepper, and cabbage for example can be damaged by
948 even a low wind speedsspeed of around 5m5 m/s, corn and cotton are a bit more resistant than
949 most vegetables, but also susceptible to wind damage when wind speed exceeds 6m6 m/s
950 (Rouse & Hodges 2004). Overall, critical thresholds for damage linked to wind speeds
951 varyspeed varies substantially.

952
953 ~~Most of the studies that investigated climatic indices relevant for agriculture focused on indices~~
954 ~~related to temperature and precipitation (Kitsara et al., 2021; Sun et al., 2016; Tschurr et al., 2020),~~
955 ~~but a few took wind into consideration as well. Crespi et al. (2020) propose a set of 32 climate related~~
956 ~~hazard indicators for Europe. Only two of these indicators refer to wind: ‘mean wind speed’ and the~~
957 ~~‘extreme wind speed days’ index. Peña-Angulo et al. (2020) analyzed the trend of 125 climatic indices~~
958 ~~which are important for agriculture, five of which concern wind. For Europe, none of these five wind~~
959 ~~indicators showed a significant trend or influence. This may be because the reanalysis data used may~~
960 ~~not be ideal for such trend analyses. What adds to the uncertainty is that less well simulated~~
961 ~~phenomena such as convective storms could become more frequent and more severe, increasing the~~
962 ~~risk for damage from downbursts and straight-line winds.~~

963
964 Whether or not wind-related agricultural damage will increase under continued warming is
965 unclear. Peña-Angulo et al. (2020) found that none of the five metrics linked to wind speed
966 show a significant trend in either direction. However, the results are subject to considerable
967 uncertainty given that convective events, which are associated with downbursts and straight-
968 line winds, are poorly simulated in the current generation of global circulation models.

969 970 **4.5.2 Wind erosion, dust storms and agricultural drought**

971 In regions with open and sandy arable land, wind can cause wind erosion and dust storms.
972 Wind erosion refers to the loss of fertile topsoil, whereas dust storms are singular events where
973 strong winds displace huge amounts of soil in a short time. Dust storms are particularly

974 frequent in the so-called dust belt reaching from the north of Africa through the Middle east to
975 central Asia (Gholizadeh et al., 2021). However, soil loss due to wind erosion is also an
976 important issue in less erosion-prone areas such as Europe (Borrelli et al., 2017). While wind
977 is the main forcing factor, there are other climatic factors such as precipitation, soil moisture
978 and radiation which affects the soil surface and thus influence soil erosion (Barring et al. 2003).

979
980 The threshold values for the mean wind speed at which soil particles start to be dislodged vary
981 greatly depending on the type and condition of the soil (Shahabinejad et al., 2019). According
982 to Rouse and Hodges (2004) the minimum mean wind speed to create erosion is normally
983 about 5-6 m/s at 30 cm above the ground. Shahabinejad et al. (2019) found critical
984 thresholdCWS values of 5.7-8.9 m/s at 10 m height for soils in Iran. Plants can suffer from dust
985 storms due to loss of plant tissue through abrasion resulting in reduced photosynthesis and
986 burial of seedlings (Stefanski and Sivakumar, 2009). This can result in considerable economic
987 losses for farmers. For example, Gholizadeh et al. (2021) demonstrate that a dust storm
988 lasting one hour can reduce the annual income of farmers by up to 1.2%. Erosion reduces soil
989 fertility for long periods due to removal of soil containing essential nutrients. In many cases,
990 extreme drought conditions precede dust storms (Sivakumar, 2005; Sissakian et al., 2013),
991 as dry soil disaggregates faster and thus dislodgedislodges more easily enhancing erosion.
992 Wind erosion is thereby closely related to land use practices.

993
994 Physiological water stress can be enhanced by increased evapotranspiration, due to high wind
995 speedsspeed. The longer such wind conditions last, the more severe the risk as exemplified
996 forby a recent drought event in India (Masroor et al. (2020). Thus, wind can exacerbate drought
997 conditions and lead to crop failure. While wind speedsarespeed is not expected to increase as
998 a global average (McVicar et al., 2012), evapotranspiration likely will in many regions due to
999 the increased evaporationevaporative demand caused by higher air temperatures (Tomas-
1000 Burguera et al., 2020) and a reduced number of days with rainfall. The fact that some plants
1001 react to hot and windy weather conditions by closing their stomata, may balance some of the
1002 enhanced evapotranspiration deficit. However, this is at the expense of plant growth.

1003 1004 **4.5.3 Protection measures against wind**

1005 Because of the direct wind damagesdamage in agriculture, it is necessary or even
1006 indispensable to take countermeasures to minimize the risks. Such measures can be a better
1007 choice of location according to topographic features or using windbreaks. Windbreaks usually
1008 consist of natural barriers such as tree rows. The most important aspect of a windbreak is its
1009 height (Brandle et al. 2004). Indeed, windbreak effects on adjacent cropcrops result in a yield
1010 reduction due to water and light competition up to a distance of one to two windbreak heights,
1011 which is followed by a yield increase up to a distance 8–12 heights (Weninger et al. 2021). To
1012 moderate effects of wind flow around the windbreak, it should be at least ten times as broad
1013 as it is high (Brandle et al. 2004).

4.6. Wind-based energy production

Wind indices are of interest for estimating the wind potential and wind energy. Extreme wind events on different spatial and temporal scales, e. g. storms, gustiness or low-level jets, affect the energy production, the structural integrity and the stability/operational safety of wind turbines. ~~Even small-scale variations~~ Microscale variability in the wind field, ~~due to~~ occurs temporally (e.g., gustiness) and spatially (e.g., vertical wind shear). These variations of the wind field depend on the time of day, and thus on the stability of the atmospheric ~~stability, and stratification~~. There is also a dependence on the characteristics of the wind turbine site (land use, terrain, can affect the energy yield and the system safety). Microscale variations of the wind field influence both the wind potential and the operational reliability of a wind turbine.

Wind indices are typically defined as the ratio of the current values of a variable (~~wind speed or energy production-related~~) to the long-term mean. ~~In practice,~~ The variable is either related to the wind speed or to the wind ~~farm operators are interested in the current wind speeds, but also in an assessment of the annual wind energy yields in comparison to the long-term average energy production~~. Extreme wind events are directly related to wind ~~data~~speed-based indices. To identify the energy potential at a site, the Power Density Wind Index can be used (Katinas et al., 2018; Celik, 2003). It is based on parameters of wind speed frequency distribution. The Power Density Index results in significantly higher variations than the real energy production of the wind turbine at the location and should be applied carefully. In practice, both the current values of the wind speed are needed (control of the turbine) and the evaluation of the annual energy yield compared to the long-term average using wind indices (planning of turbines, financing)

When addressing wind climate at a location, including the occurrence of strong wind events, which includes both productive and destructive events, much attention was given to the connection between the wind climate and the wind energy potential (Carta and Mentado, 2007). In comparison to the wind speed-based indices, the production-based indices use the energy yield of turbines as input data. The Wind Energy Production Index can be based on a Wind Speed Index (Ritter et al., 2015) calculated from wind speed data by an additional application of a power curve (Hahn and Rohrig, 2003; Ding et al., 2005). Another possibility is the use of energy yield data of a wind turbine directly. The BDB index (BDB, 2021) describes the ratio of monthly reported energy yields from wind turbines in a region to the long-term mean yields of these wind turbines. High wind ~~speeds~~speed or wind shear due to storms or low-level jets need to be taken into account when calculating wind speed indices. However, the energy production-based indices, contain the effects of such events only when the wind turbine is working, i.e. until reaching the turbine cut-out wind speed. Due to their design, most systems switch off at a wind speeds~~speed~~ above 25 m/s (Christakos et al., 2016), but there are also slightly higher and lower shutdown wind ~~speeds~~speed values for different system types (Chauhan and Saini et al., 2014). An analysis showed that ~~high-impact~~ storms had a positive effect on the wind energy production for Southwestern Europe and the Iberian Peninsula (Gonçalves et al. 2020, 2021). As such, the highest values of wind energy production result for stormy weather conditions (Petrović and Bottasso, 2014). Climate change impacts on wind energy have been investigated for a few years (~~e.g.,~~ Pryor and Barthelmie, 2010; Moemken

1059 et al., 2018). The studies are mostly in agreement on a minimal effect of climate change on
1060 the wind energy production (~~e.g.,~~ Jung and Schindler, 2020).

1061
1062 Topographic effects are another example of small-scale effects on the wind field, leading to a
1063 local wind speed-up, separation, and reattachment. These processes can be studied by
1064 numerical models (Uchida and Ohya, 2003, 2008, 2011; Uchida and Li, 2018; Uchida, and
1065 Sugitani, 2020). Uchida and Kawashima (2019) defined two indices to evaluate the terrain-
1066 induced turbulence and the fatigue damage based on the measurement data and the design
1067 value. These studies indicated the need for further development of standards. A commonly
1068 used turbulence index is the effective turbulence for site-specific fatigue assessment of wind
1069 turbines (Slot et al., 2019). Additionally, the usage of the effective turbulence index significantly
1070 reduces the number of aero-elastic simulations needed for checking ~~if~~ the loads on major
1071 components of the wind turbine.

1072 1073 **4.7. Compound indices**

1074 ~~Wind~~ **Strong winds** often co-~~occurs~~**occur** with other phenomena and their co-occurrence
1075 affects the damage levels observed. This is an integral part of the compound event
1076 concept in which multiple phenomena or hazards form a complex causal chain of
1077 events that can lead to a more extreme impact than each phenomenon by itself
1078 (Zscheischler et al. 2018). A compound event is ~~characterised by~~**often associated with**
1079 **one driver (e.g. an ~~impact~~extreme cyclone)** which ~~is caused by a hazard~~**may cause multiple**
1080 **hazards (e.g. strong wind and heavy precipitation), but it can have more complex**
1081 **characteristics** (Zscheischler et al. 2020). ~~The hazard, in its turn, is caused by a driver. Finally,~~
1082 ~~a modulator influences the location, frequency and intensity of drivers and thereby hazards. Strong~~
1083 ~~wind can therefore either be a hazard itself or~~**For example, strong wind can also serve as** a
1084 modulator for hazards like drought and wildfire. **A full typology of compound events**
1085 **can be found in Zscheischler et al. (2020).**

1086 1087 **4.7.1 Precipitation.**

1088 Strong wind ~~speeds~~**speed** often co-~~occur~~**occurs** with heavy precipitation (Martius et al. 2016),
1089 causing multivariate compound events. Additionally, it is argued that wind and precipitation
1090 enhance the impact by extratropical cyclones, since cyclones with ~~a~~**extreme** precipitation
1091 **extreme** often have a longer lifetime than cyclones with only extreme wind speed (Messmer &
1092 Simmonds, 2021). Furthermore, the impact of such multivariate compound events is much
1093 higher than a hazard containing only wind or precipitation (Martius et al. 2016). In coastal
1094 ~~environments~~**areas**, even when wind is not considered as a hazard itself, wind together with
1095 heavy precipitation can cause storm surges and coastal flooding (~~e.g.~~Wahl et al. 2015;
1096 Couasnon et al., 2020). Furthermore, precipitation is important when saturating the soil prior
1097 to the occurrence of a windstorm. Soil water content is an index that governs the stability of
1098 the root sector of trees during storm events (Everham and Brokaw, 1996; Défossez et al.
1099 2021).

1100 1101 **4.7.2 Air Temperature.**

1102 Wind and low **air** temperatures are both drivers, causing wind chill as human health and
1103 agricultural hazards among other risks. Each driver, when acting by itself, would have caused

1104 less of an impact than the compound effect (Danielsson, 1996). Wind chill is a threat mainly
1105 in cold climates, where enhanced wind ~~speeds increase~~ ~~speed increases~~ the heat transfer from
1106 an object. Such heat loss can cause injuries and mortality both in animals and plants. Windchill
1107 can be calculated as the wind chill temperature, also called wind chill factor (Quayle and
1108 Steadman, 1998; Bluestein and Zecher, 1999), which is usually taken as the air temperature
1109 at which there would be an equivalent rate of heat loss. Also, low air temperatures can lead to
1110 the freezing of soil and ~~enhance~~ ~~enhances~~ the stability of trees against windthrow during
1111 windstorms (Pasztor et al., 2015). In contrast, trees in frozen soil are more likely to undergo
1112 stem breakage than uprooting (Everham and Brokaw, 1996; Peltola, 2006).

1113
1114 **4.7.3 Drought.** The impact of wind on drought is ~~relatively~~ ~~comparatively~~ small- compared with
1115 other drivers like temperature and (lack of) precipitation, but it has an effect in terms of the
1116 evapotranspiration. Wind is ~~only~~ ~~thus~~ included in some drought indices through
1117 evapotranspiration in the Penman or Penman-Monteith equation, such as in the Baumgartner
1118 index (Baumgartner et al., 1967). These indices are therefore short-term indices that operate
1119 on a scale of days and typically do not take into account the long-term impacts of drought on
1120 the risk of wind damage to forests. Drought can be considered as a pre-condition, that
1121 potentially amplifies the impact of winds. Csilléry et al. (2017) showed that long-term drought
1122 can increase the risk of wind damage on sites where drought can lead to a weakening of trees
1123 but can also decrease the risk of damage on normally extremely wet sites.

1124
1125 **4.7.4 Fire.** Indices used for assessing fire risk include often wind and topography to determine
1126 the rate of spread and damage caused by a wildfire. Wind and slope are viewed as the major
1127 factors influencing fire development (Byram 1959a, 1959b; Sharples 2008). The most used
1128 indices for fire risk are based on the Canadian Forest Fire Weather Index (FWI) system (Van
1129 Wagner, 1987) that uses information on fuel loading and meteorological conditions (rainfall,
1130 temperature, humidity, and wind speed) to predict the probability of a fire starting and then the
1131 probable spread of the fire. Humidity, wind speed and air temperature are used to calculate
1132 the day-to-day drying of the fuel load. The Initial Spread Index is then used to adjust the FWI
1133 as an exponential function of wind speed (doubles the FWI for every increase of wind speed
1134 by 19 km/h or 5.3 m/s). The spread of the fire will also be affected by the topography and, in
1135 particular, how the topography modifies ~~in~~ ~~the~~ wind speed and direction.


1136
1137 Wind can alter the angle of the fire toward unburnt fuel, extending the preheating range and
1138 increasing the rate of spread. Slope has a similar effect by affecting the distance between the
1139 flames and the fuel. Thus, typically the greatest rate of spread is found when an upslope is
1140 combined with upward winds and vice versa (Sharples 2008). Since topography influences
1141 wind traits, it can create a channeling effect enhancing fire intensity, but with the strength of
1142 the effect depending on the overlap between wind direction and landscape orientations
1143 (Barros 2012; Mansuy 2014). Kushal (1997) found in a review ~~of four articles~~ that a higher
1144 relative elevation, proximity to ridges and increased exposure to wind, all led to greater fire
1145 ~~damages~~ ~~damage~~ in forests. Additionally ~~in six articles~~, aspects that are associated with greater
1146 exposure to dry winds increased fire damage in forests, and damage was lower in aspects
1147 with cold and moist winds. ~~We are aware of~~ ~~There is~~ an index combining slope, aspect, and wind
1148 speed (wind-topo), but it had a rather low importance for the final model chosen for statistical
1149 interpretation (Masoudvaziri 2020).

4.8. Wind speed warning-levels used at national meteorological services and environmentsector-related critical thresholds

Advanced storm-warnings are crucial for the protection of property and lives. Meteorological services operate a structured warning system for windstorms and recommend appropriate protective measures and rules of conduct depending on the warning level: (e.g. Germany: (DWD, 2021), Ireland: (MetEireann, 2023) or Sweden: (SMHI, 2023)). The warnings will be published, when the event reaches a certain probability level to occur, can be well spatially located and especially when the warning criterion is met, such as wind speed or precipitation ~~exceed~~exceeding a certain threshold value. These threshold values are set individually by all meteorological services. ~~For example, the DWD uses 6 different gust wind speed thresholds: 14, 18, 25, 29, 33 and 39 m/s (Primo, 2016), referring to 4 warning levels (WL), WL2 and WL3 are divided into two intervals (DWD, 2021).~~ In some cases, the weather services already indicate possible consequences due to the wind ~~speeds~~speed, by warning of damage to infrastructure, forests, or energy systems at differing warning levels. There are even variants of weather forecasting systems that follow a more risk-based approach, i.e. the probabilities and consequences of extreme events are integrated into the forecasting system in order to achieve an improved warning management (Neal et al. 2014 or Kaltenberger et al. 2020). A Europe-wide overview of warnings and, in part, possible impacts is provided by Metealarm (www.meteoalarm.org), developed by EUMETNET (European Meteorological Network) provides relevant information on extreme weather events from 37 national meteorological services.


We collected many critical thresholds from the literature for the five sectors which are the focus of this manuscript. The vulnerability of each environmentsector to wind speed is illustrated in Table S1. Figure 3 provides a synthesis and comparison of thresholds from the five environmentsectors. The agriculture sector seems to be the most sensitive to wind, as negative effects are already noticeable at gustmean wind ~~speeds~~speed well below the first official warning level of the DWD (WL1 = 14 m/s); Rouse and Hodges, 2004.

At WL2 (18-29 m/s), initial restrictions must already be expected in all 5 sectors, but these are initially localized. In the forest, individual trees and areas may be affected; (Gardiner et al., 2010, 2013, 2016), buildings may show slight roof damage but no structural damage yet; (Feuerstein et al., 2011), in road traffic there may be some accidents and delays in train and air traffic. ~~For the~~ (Vajda et al., 2014). Concerning wind energy ~~sector~~, no damage is expected yet, but depending on the type of turbine, precautionary shutdowns of turbines may occur. For WL3 (29-39 m/s), the literature describes significant impacts in the forest, building, and transportation environmentsectors. Damage will be significant, and impacts are already affecting regional areas. Especially in the The influence of storms on transportation ~~sector, storms~~ can quickly impact society at regional to national levels. ~~Severe damage is described at the national level from WL4 onwards, including damage to wind turbines, massive building damage, or even the shutdown of entire transport sectors (air and railway). While forest, urban and transport are affected by wind speeds at the same order of magnitude (i.e. locally at WL2, regionally at WL3 and nationally at WL4), the wind energy sector shows strong impacts at higher wind speeds (when only damage is considered) and the agricultural sector at much lower wind speeds.~~



1194

1195 Severe damage is described at the national level from WL4 onwards, including damage to
 1196 wind turbines (Quaschnig et al., 2016), massive building damage (Feuerstein et al., 2011), or
 1197 even the shutdown of entire transport sectors (air and railway) (Vajda et al., 2014). While
 1198 forest, urban areas and transport are affected by wind speed at the same order of magnitude
 1199 (i.e. consequences for society are mostly locally at WL2, regionally at WL3 and nationally at
 1200 WL4), the energy production from wind is impacted at a much higher wind speed (when only
 1201 damage is considered) while agricultural productivity at much lower wind speed.



1202

1203 Figure 3: Critical ~~thresholds~~threshold ranges of wind speed (mean wind speed (averaging
 1204 interval 1 hour) and wind gusts) for five affected sectors. Warning levels (WL1-WL1-4 at speeds
 1205 of 14, 18, 29 and 39 m/s, respectively) of the DWD for wind gusts (Primo, 2016) are marked
 1206 on the right axis. For each sector different ranges of critical mean wind speed from the
 1207 literature are plotted, to show the mean wind speed (or gust), where impacts are expected.
 1208 Thresholds in the first three sectors (forest, urban areas and transport) refer to wind gusts at
 1209 10 m height, for agriculture we present the mean wind speed at 10 m height and the shown
 1210 thresholds for the energy sector refers to the wind speed measured at the height of the wind

1211 turbines. The upper bar (e.g. “National impact”) are left open as damage will occur also at higher wind
1212 speeds. mean wind speed measured at the height of the wind turbines (wind in hub height).
1213 The upper bar (e.g. “National impact”) is left open as damage will occur also at higher wind
1214 speed. The threshold ranges mean: (1) Forest local: Limited area of damage. Forest Regional:
1215 Damage level is meaningful for the affected forest and short-term forest planning and timber
1216 price. Forest National: Damage can occur across several countries. (2) Urban Marginal: Light
1217 objects, tiles can be lifted or come loose. Urban Noticeable: Heavier objects are lifted and first
1218 damage to individual building components are possible. Urban Major: Large vehicles overturn,
1219 roofs are severely damaged. (3) Transport Local disruption: Blocked roads through windthrow
1220 or sliding containers at ships. Transport Delays: Cancellation trough electricity cuts and
1221 increasing number of wind-related accidents. Transport Traffic stop: Damage of overhead
1222 cables and longer power failures as well as airport and harbor closures. (4) Agriculture erosion:
1223 Soil loss to wind erosion. Agriculture yield loss: Damage to leaves and yield loss due to lodged
1224 fields. (5) Energy cut-in speed: start of energy production. Energy Cut-out speed: Automatic
1225 shutdown of wind turbines.

1226

1227 5. Outlook & open research questions

1228 In this review we covered a wide range of topics dealing with wind damage to terrestrial
1229 ecosystems with an emphasis on studies dealing with ~~central~~Central Europe. To conclude, we
1230 address ~~trends~~issues of importance in the near future and topics that require further research.
1231 The most intriguing question in this field is how wind-related damage levels may change in
1232 future decades, given the strong dominance of decadal variability (e.g. Feser et al., 2015).
1233 Therefore, attention was given to identifying drivers of future changes in windstorms and
1234 cyclone characteristics which are particularly important for the predictability of present-day
1235 and long-term trends in socio-economic ~~damages~~damage (Koks and Haer 2020; Hoeppe
1236 2016; Franzke 2021). The key current drivers that contribute to future changes in storms are
1237 well known; many studies assume that the atmospheric moisture content will increase due to
1238 global warming (IPCC 2021). Idealized studies suggest that this increase in moisture will lead
1239 to a stronger circulation, more intense storms (including stronger winds and more rainfall) and,
1240 thus, to an expansion of the windstorm footprint (Catto et al. 2019). Additionally, studies show
1241 that the lower-tropospheric meridional air temperature gradient will decrease due to Arctic
1242 amplification, whereas the upper-tropospheric meridional air temperature gradient will
1243 increase due to the warming of the tropical upper troposphere and the cooling of the polar
1244 lower stratosphere (Lee et al. 2019). However, it is still uncertain how these contrasting forcing
1245 mechanisms will contribute to the future changes in storms quantitatively (Catto et al., 2019;
1246 their Fig. 2). The recently extended ERA5 reanalysis product could enable further studies to
1247 deal with wind-related ~~damages~~damage in the past, present and future, reducing uncertainties.
1248 Indeed, increasing the resolution of climate models may improve their capacity to quantify
1249 statistical storm properties. CMIP6 models already indicate a general improvement in future
1250 storm tracking (Priestley et al. 2020b; Harvey et al. 2020). As a result, more accurate
1251 projections of wind and storm damage based on future emission scenarios and climate change
1252 may be attainable in the future. According to a recent study, winter-storm-related wind gusts
1253 could increase towards the 2nd half of the 21st century in Germany (Jung und Schindler 2021).
1254 This demonstrates the need for more studies in damage analysis.

1255

Methodologically, the usage of the indices described here in damage analysis has many advantages, but their creation can be time consuming, and their usage may lead to statistical pitfalls. ~~We emphasize the importance of choosing~~ For example, it is important to choose indices while trying to avoid an overlap in variability explained by different topographic indices (e.g. Mitchel 2001). In this sense, we are lacking a clear methodology that can ~~address~~ select the most ~~efficient manner to select suitable~~ indices in advance. ~~Such a methodology needs to be developed,~~ especially when many indices are easily available. There are three main approaches: 1) a hypothesis-based approach, where typically only few variables are used in the analyses because these variables can be well explained and justified due to past research, incorporation of expert knowledge in the development of indices using co-design and familiarity with the study site (Gebhardt et al. 2019, Merz et al. 2020), 2) a computational approach, where a feature-selection algorithm (e.g. genetic algorithm) is first used to trim down the number of independent variables before performing an analysis, and 3) an exploratory approach with little limitation on the number of independent variables used, where one can examine, for example, if a certain group of indices is more useful than another (e.g. gust-related indices vs. topographic indices) in achieving accurate models according to a given evaluation metric (e.g. coefficient of determination or area under the curve). The choice of method is dependent on the specific research goals, but also on the skillset and computational resources available, for instance, an exploratory analysis including many variables on a large area may demand access to high-performance computing. Furthermore, when modelling on a large spatial scale to evaluate in which, it is important to choose analysis tools that test and quantify the homogeneity of the relation between indices and damage variables across the different sub-regions ~~certain groups of indices are useful. A possible solution can be to first assess the relative contribution of an index or a group of indices according to their relative contribution to a given metric (e.g. R^2 , AUC) when running models with all possible combinations of explanatory variables in the study site.~~ Thus, ~~enabling us to assess their contribution~~ taking into account that key parameters may change within the study area, such as a standalone ~~the topography or the vegetation structure, altering the relations between an independent variable (more useful for simple logistic regression) or their relative contribution when using many explanatory variables at the same time (e.g. and storm damage. Finally, when using machine learning algorithms)~~ analyzing socio-economic impacts, the availability of data is often a limiting factor, and these limitations shape the selection and analysis approach.

We identify that the area most in need of new indices for wind-related damage analysis are compound events. Damage from extreme climatic events most commonly occurs through interactions between different hazards (Zscheischler et al., 2020). The main challenge is to handle the different time scales of each factor, for example, a storm may last from hours to days, but drought can last years. Therefore, we require indices that incorporate a multitude of factors that are very site specific, as both the topography and the land cover can strongly modify these interactions. Another important challenge is the inclusion of non-climate drivers related to exposure and vulnerability in the compound indices. Concerning the five sectors dealt with here, we present sector-specific outlooks:

1299 ~~We lack information in several key areas.~~**5.1 Forest.** In a forest setting, there are very few
1300 measurements of tree damage due to storms (Kamimura et al., 2022) and very few studies of
1301 the dynamic nature of damage at the time scale of a storm. Such studies are required to
1302 understand damage initiation and propagation during storms (Dupont et al., 2015). In addition,
1303 predicting airflow over complex terrain is still difficult when there are steep slopes and multiple
1304 changes in vegetation height (Finnigan et al., 2020). Similarly, there is a need for improvement
1305 of land surface information, and in particular, the acquirement of highly resolved 3D
1306 distributions of vegetation elements at the landscape scale to enable the creation of fine scale
1307 maps for risk assessment. ~~Recent~~To this end, it is often difficult to assess damage or risk in
1308 the most relevant spatial scale. The recent developments in remote sensing techniques
1309 (terrestrial and airborne laser scanning) promise effective assessments of surfaces structures
1310 (Favorskaya and Jain, 2017).~~However~~, may prove useful for many of these issues. However,
1311 it may take much time to achieve a sufficient level of data collection, for example, terrestrial
1312 laser scanning is accurate but confined to small areas, and an effective assessment for larger
1313 areas using airborne laser scanning and satellite data are not at a sufficient resolution and
1314 need further development. Another consequence is that we still lack in monitoring and
1315 modelling the small-scale variability in the interactions of the wind field with the surface. The
1316 main research questions for the future are: How does the structure of a forest canopy influence
1317 the turbulence within and above the canopy? And, as they grow, stems, roots and canopies
1318 acclimate to the wind forces, so, what is the optimal cultivation and canopy structure to reduce
1319 damage (Dèfossez et al., 2022)?~~These questions and points made above are important in forests~~
1320 ~~but also in agricultural setting even though the surface is typically smoother than in forests.~~

1321

1322 **5.2 Urban.** In urban settings, storm, and loss indices as well as damage functions do not
1323 usually consider differences in the exposure and vulnerability of different building types or
1324 types of urban areas. To further assess wind damage risks on a smaller spatial scale
1325 investigations of individual building damage or damage to specific types of neighborhoods are
1326 needed, together with modelling of urban ~~environments~~areas. However, damage data at a fine
1327 spatial scale is difficult to obtain, and it is a priority to improve the documentation of urban
1328 damage to support the development of new indices. The availability of data, such as the spatial
1329 extent of wind damage to individual buildings or green spaces, is key in developing mitigation
1330 strategies. For example, wind channeling as a function of wind speed and direction needs to
1331 be reliably simulated during the development phase of new building projects.

1332

1333 ~~Furthermore, it will be~~**5.3 Transport.** Studies addressing wind effects on transport usually focus
1334 on direct effects in a particular part of the transport system. Results from such studies can
1335 strongly depend on the region, data and methodologies used for the study. Studies with a
1336 more unified approach addressing wind effects on transport on a broader scale could lead to
1337 more comparable results. Furthermore, little research is available that takes into account
1338 cascading effects that propagate through different parts of the transport system. In general, it
1339 remains unclear how climate change and resulting changes in the wind extremes will affect
1340 the transport system. Studies addressing this question should not only consider future
1341 changes in wind extremes, but also potential changes of the transport system as part of
1342 climate change mitigation measures. Such measures could make the transport system more
1343 vulnerable to extreme winds. For example, a shift from road to rail transport to reduce CO₂

emissions could lead to a higher vulnerability to wind-related tree fall, because single storm events can lead to a collapse of rail transport over whole countries for periods of several days.

5.4 Agriculture. The future of the agriculture sector is closely linked to the global challenge of feeding a still growing population, which is expected to reach 9.7 billion people by 2050 (UN World Population Prospects 2022). In response to this challenge, the awareness for sustainable and efficient agricultural practices has been gradually increasing. Wind damage in agriculture landscapes is thereby a growing concern due to the potential change in frequency and intensity of wind events as the climate continues to warm (Seneviratne et al. 2021). In order to optimize crop yields and reduce waste, the relationship between wind damage and crop yields needs to be investigated in more detail to quantify this impact. For example, understanding and better short-term prediction of wind events are key to improved crop management. Furthermore, there is a lack of simple indices incorporating soil properties and their tendency to lead to soil erosion and nutrient loss, or wind erosion, as such events can be a major challenge for farmers. There is much space to develop new practices to mitigate wind damage in agriculture by using vegetation and cover crops reduce wind damage. Since the positioning of vegetation (e.g. trees as windbreaks) alter the small-scale interactions of the wind field with the soil and crops, a more accurate positioning of vegetation would be supported by the creation or adaptation of existing compound indices or modeling platforms. Such manipulations of the surface cover can be highly flexible in the spatial scale of the wind-field modification, thus providing a good counter measure to different types of vulnerability in agricultural sector.

5.5 Renewable Energy. It is important to follow the influence of climate change projections on wind energy production. With the increase in the reliance on renewable energy, it will be important to reduce uncertainties in wind potential and the risk for technical and safety issues in the operation of the wind turbines. Furthermore, while we know where turbines are located and their characteristics, it would be important that the data on the turbines wind field and the energy generated were accessible for scientific projects and to the private sector. Currently much of the data is not made publicly available. For instance, we especially lack wind data at hub height for the evaluation of numerical models. Other key challenges, that are similar to other sectors, are the acquirement of high spatial resolution measurement, and past and future modeling of the wind field over heterogeneous surfaces and complex terrain.

In conclusion, predicting and assessing the damage caused by wind and storms is a complex matter but there are effective and simple methodologies to support assessment and decision making. In the light of the future uncertainties, it is vital to continue developing tools to prepare for the next calamities that are bound to occur.

6. Acknowledgements:

This publication is the outcome of a working group of the project ClimXtreme, funded by the German Bundesministerium fuer Bildung und Forschung (AK, DG, DH, DN, HL, JG). AZ was funded by Project QuWind100 (German Federal Ministry for Economic Affairs and Energy, BMWi, based on a decision by the German Bundestag, funding code 0325940A). CF was supported by the Institute for Basic Science (IBS), Republic of Korea, under IBS-R028-D1 and by Pusan National University research grant 2021-Research Fund of Korea (NRF-2022M3K3A1097082). JGP thanks the AXA Research Fund for their

1388 support. [NBNB's](#) work was carried out within the framework of the Hans-Ertel-Centre for
1389 Weather Research. This research network of universities, research institutes and the
1390 Deutscher Wetterdienst is funded by the Bundesministerium für Verkehr und Digitale
1391 Infrastruktur (grant no. 4818DWDP3A). JGP is a member of the editorial board of the journal
1392 Natural Hazards and Earth System Sciences.

1393

1394 7. References

- 1395 ○ Adam, E. F., Brown, S., Nicholls, R. J., and Tsimplis, M.: A systematic assessment of
1396 maritime disruptions affecting UK ports, coastal areas and surrounding seas from 1950 to
1397 2014. *Nat. Hazard.*, 83(1), 691-713, <https://doi.org/10.1007/s11069-016-2347-4>, 2016.
- 1398 ○ Akbar, M. K., Kanjanda, S., and Musinguzi, A.: Effect of Bottom Friction, Wind Drag
1399 Coefficient, and Meteorological Forcing in Hindcast of Hurricane Rita Storm Surge Using
1400 SWAN + ADCIRC Model, *J. Marine Sci. Eng.*, 5, 38, <https://doi.org/10.3390/jmse5030038>,
1401 2017.
- 1402 ○ [Albrecht, A. T., Fortin, M., Kohnle, U., and Ningre, F.: Coupling a tree growth model with](#)
1403 [storm damage modeling - conceptual approach and results of scenario simulations. *Environ.*](#)
1404 [Mod. Software. 69, 63–76, 2015.](#)
- 1405 ○ [Albrecht, A.](#), Hanewinkel, M., Bauhus, J., and Kohnle, U.: How does silviculture affect storm
1406 damage in forests of south-western Germany? Results from empirical modeling based on
1407 long-term observations, *Eur. J. Forest Res.*, 131, 229–247, [https://doi.org/10.1007/s10342-](https://doi.org/10.1007/s10342-010-0432-x)
1408 [010-0432-x](#), 2012.
- 1409 ~~○ Albrecht, A., Kohnle, U., Hanewinkel, M., and Bauhus, J.: Storm damage of Douglas fir~~
1410 ~~unexpectedly high compared to Norway spruce, *Ann. Forest Sci.*, 70, 195–207,~~
1411 ~~<https://doi.org/10.1007/s13595-012-0244-x>, 2013.~~
- 1412 ~~○ Albrecht, A. T., Fortin, M., Kohnle, U., and Ningre, F.: Coupling a tree growth model with~~
1413 ~~storm damage modeling - conceptual approach and results of scenario simulations. *Environ.*~~
1414 ~~Mod. Software, 69, 63–76, 2015.~~
- 1415 ○ ~~Albrecht, A. T.~~, Jung, C., and Schindler, D.: Improving empirical storm damage models by
1416 coupling with high-resolution gust speed data, *Agricul. Forest Meteor.*, 268, 23–31,
1417 <https://doi.org/10.1016/j.agrformet.2018.12.017>, 2019.
- 1418 ○ [Albrecht, A., Kohnle, U., Hanewinkel, M., and Bauhus, J.: Storm damage of Douglas-fir](#)
1419 [unexpectedly high compared to Norway spruce, *Ann. Forest Sci.*, 70, 195–207,](#)
1420 [https://doi.org/10.1007/s13595-012-0244-x, 2013.](#)
- 1421 ○ Allianz: Safety and shipping review 2019 - An annual review of trends and developments in
1422 shipping losses and safety, Allianz Global Corporate & Specialty, Munich, Germany,
1423 [https://www.agcs.allianz.com/content/dam/onemarketing/agcs/agcs/reports/AGCS-Safety-](https://www.agcs.allianz.com/content/dam/onemarketing/agcs/agcs/reports/AGCS-Safety-Shipping-Review-2019.pdf)
1424 [Shipping-Review-2019.pdf](#), last access: 14 December 2021, 2019.
- 1425 ○ Ashcroft, J.: The relationship between the gust ratio, terrain roughness, gust duration and
1426 the hourly mean wind speed, *J. Wind Eng. Ind. Aerodyn.*, 53, 331–355, doi:10.1016/0167-
1427 [6105\(94\)90090-6](#), 1994.
- 1428 ○ Ashley, W. S. and Mote, T. L.: Derecho hazards in the United States, *Bull. Am. Meteorol.*
1429 *Soc.*, 86, 1577–1592, doi:10.1175/BAMS-86-11-1577, 2005.
- 1430 ○ Baker, C. J.: A simplified analysis of various types of wind-induced road vehicle accidents,
1431 *J. Wind Eng. Ind. Aerodyn.*, 22, 69-85, [https://doi.org/10.1016/0167-6105\(86\)90012-7](https://doi.org/10.1016/0167-6105(86)90012-7),
1432 1986.

- 1433 ○ Baker, C. J., and Reynolds, S.: Wind-induced accidents of road vehicles, *Accid. Anal. Prev.*,
1434 24, 559-575, [https://doi.org/10.1016/0001-4575\(92\)90009-8](https://doi.org/10.1016/0001-4575(92)90009-8), 1992.
- 1435 ○ Baker, C. J.; Sterling, M.; Berry, P.: A generalised model of crop lodging. *J. Theor. Biology*,
1436 363, 1–12. DOI: 10.1016/j.jtbi.2014.07.032, 2014.
- 1437 ○ Barring, L., Jönsson, P., Mattsson, J.O. and Åhman, R., 2003: Wind erosion on arable land
1438 in Scania, Sweden and the relation to the wind climate—a review. *Catena*, 52, 173-190.
- 1439 ○ Barros, A. M. G., Pereira, J. M. C. and Lund, U. J.: Identifying geographical patterns of
1440 wildfire orientation: A watershed-based analysis, *For. Ecol. Manage.*, 264, 98–107,
1441 doi:10.1016/j.foreco.2011.09.027, 2012.
- 1442 ○ [Bartkowski, B., Schepanski, K., Bredenbeck, S., & Müller, B. \(2022\). Wind erosion in](#)
1443 [European agricultural landscapes: More than physics. *People and Nature*, doi:](#)
1444 [10.1002/pan3.10418](#)
- 1445 ○ Batke, S. P., Jocque, M., and Kelly, D. L.: Modelling hurricane exposure and wind speed on
1446 a mesoclimate scale: A case study from Cusuco NP, Honduras, *PloS One*, 9, e91306,
1447 <https://doi.org/10.1371/journal.pone.0091306>, 2014.
- 1448 ○ Baumgartner, A., Klemmer, L., Raschke, E. and Waldmann., G.: Waldbrände in Bayern
1449 1950 bis 1959., *Mitteilungen aus der Staatsforstverwaltung Bayerns*, 36, 57-79, 1967.
- 1450 ○ BDB (2021): WEA-Standortdaten, Betreiber-Datenbasis (BDB),
1451 <http://www.btrdb.de/sto.html>, last access: 21 December 2021.
- 1452 ○ Beck, V. and Dotzek, N.: Reconstruction of Near-Surface Tornado Wind Fields from Forest
1453 Damage. *J. Appl. Meteor. Clim.*, 49, 1517–1537, <https://doi.org/10.1175/2010JAMC2254.1>,
1454 2010.
- 1455 ○ [Becker, N., Rust, H.W., and Ulbrich, U.: Weather impacts on various types of road crashes:](#)
1456 [a quantitative analysis using generalized additive models. *Eur. Transp. Res. Rev.* 14, 37.,](#)
1457 [2022.](#)
- 1458 ○ Belcher, S. E., Harman, I. N., and Finnigan, J. J.: The Wind in the Willows: Flows in Forest
1459 Canopies in Complex Terrain, *Annu. Rev. Fluid. Mech.*, 44, 479–504,
1460 <https://doi.org/10.1146/annurev-fluid-120710-101036>, 2012.
- 1461 ○ [Berry, P. M.; Baker, C. J.; Hatley, D.; Dong, R.; Wang, X.; Blackburn, G. A. et al.:](#)
1462 [Development and application of a model for calculating the risk of stem and root lodging in](#)
1463 [maize. *Field Crops Res.*, 262, 108037. DOI: 10.1016/j.fcr.2020.108037, 2021.](#)
- 1464 ○ ~~Berry, P. Spink, J.: Predicting yield losses caused by lodging in wheat. In *Field Crops Res.*,~~
1465 ~~137, 19–26. DOI: 10.1016/j.fcr.2012.07.019, 2012~~
- 1466 ○ ~~Berry, P., S. Kendall, Z. Rutterford, S. Orford, S. Griffiths: Historical analysis of the effects~~
1467 ~~of breeding on the height of winter wheat (*Triticum aestivum*) and consequences for lodging,~~
1468 ~~*Euphytica*, 203: 375–383. DOI: 10.1007/s10681-014-1286-y, 2014.~~
- 1469 ○ ~~Berry, P.; Blackburn, A.; Sterling, M.; Miao, Y.; Hatley, D.; Gullick, D. et al.: A multi-~~
1470 ~~disciplinary approach for the precision management of lodging risk. In John V. Stafford (Ed.):~~
1471 ~~Precision agriculture '19. Papers presented at the 12th European Conference on Precision~~
1472 ~~Agriculture, Montpellier, France, 8-11 July 2019. [12th European Conference on Precision](#)~~
1473 ~~[Agriculture. Montpellier, France, 08-11-07, 2019.](#) Wageningen: Wageningen Academic~~
1474 ~~Publishers, 969–975, 2019.~~
- 1475 ○ [Berry, P. M.; Berry, P., S. Kendall, Z. Rutterford, S. Orford, S. Griffiths: Historical analysis of](#)
1476 [the effects of breeding on the height of winter wheat \(*Triticum aestivum*\) and consequences](#)
1477 [for lodging. *Euphytica*, 203: 375-383. DOI: 10.1007/s10681-014-1286-y, 2014.](#)
- 1478 ○ [Berry, P. M.; Spink, J.: Predicting yield losses caused by lodging in wheat. In *Field Crops*](#)
1479 [Res., 137, 19–26. DOI: 10.1016/j.fcr.2012.07.019, 2012](#)

- 1480 ~~○ Baker, C. J.; Hatley, D.; Dong, R.; Wang, X.; Blackburn, G. A. et al.: Development and~~
 1481 ~~application of a model for calculating the risk of stem and root lodging in maize. *Field Crops*~~
 1482 ~~*Res.*, 262, 108037. DOI: 10.1016/j.fcr.2020.108037, 2021.~~
- 1483 ○ Bittelli, M., Ventura, F., Campbell, G., Snyder, R., Gallegati, F., and Pisa, P.: Coupling of
 1484 heat, water vapor, and liquid water fluxes to compute evaporation in bare soils, *J. Hydrol.*,
 1485 362, 191–205, <https://doi.org/10.1016/j.jhydrol.2008.08.014>, 2008
 - 1486 ○ Bjerknes, J.: Life cycle of cyclones and the polar front theory of atmospheric circulation.
 1487 *Geophys. Publik.*, 3, 1-18, 1922.
 - 1488 ○ Blender, R., Fraedrich, K. and Lunkeit, F.: Identification of cyclone-track regimes in the North
 1489 Atlantic. *Quart. J. Roy. Meteor. Soc.*, 123, 727-741, 1997.
 - 1490 ○ Blender, R., Raible, C. and Lunkeit, F.: Non-exponential return time distributions for vorticity
 1491 extremes explained by fractional Poisson processes. *Q. J. Roy. Meteorol. Soc.*, 141, 249-
 1492 257, 2015.
 - 1493 ○ Blennow, K. and Olofsson, E.: The probability of wind damage in forestry under a changed
 1494 wind climate, *Clim. Change*, 87, 347–360, <https://doi.org/10.1007/s10584-007-9290-z>,
 1495 2008.
 - 1496 ○ Bluestein, M., and Zecher, J.: A New Approach to an Accurate Wind Chill Factor, *Bull. Amer.*
 1497 *Meteor. Soc.*, 80, 9, 1893-1900, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0477(1999)080<1893:ANATAA>2.0.CO;2)
 1498 [0477\(1999\)080<1893:ANATAA>2.0.CO;2](https://doi.org/10.1175/1520-0477(1999)080<1893:ANATAA>2.0.CO;2), 1999.
 - 1499 ○ Bollmeyer, C., Keller, J. D., Ohlwein, C., Wahl, S., Crewell, S., Friederichs, P., Hense, A.,
 1500 Keune, J., Kneifel, S., Pscheidt, I., Redl, S., and Steinke, S.: Towards a high-resolution
 1501 regional reanalysis for the European CORDEX domain, *Q. J. Roy. Meteor. Soc.*, 141, 1-15,
 1502 <https://doi.org/10.1002/qj.2486><https://doi.org/10.1002/qj.2486>, 2015.
 - 1503 ○ Bonnesoeur, V., Constant, T., Moulia, B. and Fournier, M.: Forest trees filter chronic wind-
 1504 signals to acclimate to high winds. *New Phytologist*, 210, 850–860,
 1505 <https://doi.org/10.1111/nph.13836>, 2016.
 - 1506 ○ Born K., Ludwig P., Pinto J. G.: Wind gust estimation for Mid-European winter storms:
 1507 Towards a probabilistic view. *Tellus A* 64:17471. doi:10.3402/tellusa.v64i0.17471, 2012.
 - 1508 ○ Borrelli, P.; Lugato, E.; Montanarella, L.; Panagos, P.: A New Assessment of Soil Loss Due
 1509 to Wind Erosion in European Agricultural Soils Using a Quantitative Spatially Distributed
 1510 Modelling Approach. *Land Degrad. Develop.*, 28, 335–344. DOI: 10.1002/ldr.2588, 2017.
 - 1511 ○ Brandle, J., Hodges, L. and Zhou, X.: Windbreaks in North American Agricultural Systems
 1512 (*Agronomy and Horticulture - Faculty Publications*, 389), last accessed on 11/3/2021, 2004.
 - 1513 ○ Brasseur O.: Development and application of a physical approach to estimating wind gusts.
 1514 *Mon. Wea. Rev.*, 129: 5–25, 2001
 - 1515 ○ Bunkers, M. and Hjelmfelt, M.: Chapter Three - Thunderstorm downbursts: Windstorms and
 1516 blowdowns, In *Plant Disturbance Ecology (Second Edition)*, edited by E. Johnson and K.
 1517 Miyanishi, pp. 65–115, Academic Press, San Diego., [https://doi.org/10.1016/B978-0-12-](https://doi.org/10.1016/B978-0-12-818813-2.00003-4)
 1518 [818813-2.00003-4](https://doi.org/10.1016/B978-0-12-818813-2.00003-4)<https://doi.org/10.1016/B978-0-12-818813-2.00003-4>, 2021.
 - 1519 ○ Byram, G.: Combustion of forest fuels, in *Forest fire: control and use*, edited by K. P. Davis,
 1520 pp. 61–89, McGraw-Hill, New York., 1959a.
 - 1521 ○ Byram, G.: Forest fire behaviour, in ‘*Forest Fire: Control and Use*’, Ed. Davis, K., McGraw-
 1522 Hill Book Comp. Inc., New York, 584 pp., 1959b.
 - 1523 ○ Carta, J. and Mentado, D.: A continuous bivariate model for wind power density and wind
 1524 turbine energy output estimations. *Energy Conversion Manag.*, 48, 420-432,
 1525 <https://doi.org/10.1016/J.ENCONMAN.2006.06.019>, 2007.

- 1526 ○ Cassou, C., Terray, L., and Phillips, A.: Tropical Atlantic influence on European heat waves,
1527 J. Clim., 18, 2805-2811, doi:10.1175/JCLI3506.1, 2005.
- 1528 ○ Catto, J., Ackerley, D., Booth, J. et al. The Future of Midlatitude Cyclones. Curr Clim Change
1529 Rep 5, 407–420. <https://doi.org/10.1007/s40641-019-00149-4>, 2019.
- 1530 ○ CEN. (2010). EN 1991-1-4 (2005): Eurocode 1: Actions on structures—Part 1-4: General
1531 actions—Wind actions.
- 1532 ○ Christakos, K., Cheliotis, I., Varlas, G., and Steeneveld, G.-J.: Offshore Wind Energy
1533 Analysis of Cyclone Xaver over North Europe, Energy Procedia, 94, 37–44,
1534 <https://doi.org/10.1016/j.egypro.2016.09.187>, 2016.
- 1535 ○ Celik, A.: Assessing the suitability of wind speed probability distribution functions based on
1536 wind power density, Renewable Energy, 28, 1563–1574, [https://doi.org/10.1016/S0960-
1537 1481\(03\)00018-1](https://doi.org/10.1016/S0960-1481(03)00018-1), 2003.
- 1538 ○ Charuvisit, S., Kimura, K., and Fujino, Y.: Effects of wind barrier on a vehicle passing in
1539 the wake of a bridge tower in cross wind and its response, J. Wind Eng. Ind. Aerodyn., 92,
1540 609-639, <https://doi.org/10.1016/j.jweia.2004.03.006>, 2004.
- 1541 ○ Chauhan, A. and Saini, R.: Statistical analysis of wind speed data using Weibull distribution
1542 parameters, in: 2014 1st International Conference on Non-Conventional Energy (ICONCE
1543 2014), 160–163, <https://doi.org/10.1109/ICONCE.2014.6808712>, 2014.
- 1544 ○ Chang, E., Lee, S. and Swanson, K.: Storm track dynamics. J. Climate, 15, 2163-2183,
1545 2002.
- 1546 ○ Chen, Y.-Y., Gardiner, B., Pasztor, F., Blennow, K., Ryder, J., Valade, A., Naudts, K., Otto,
1547 J., McGrath, M. J., Planque, C., and Luysaert, S.: Simulating damage for windstorms in the
1548 land surface model ORCHIDEE-CAN (revision 4262), Geosci. Model Develop., 11, 771–
1549 791, <https://doi.org/10.5194/gmd-11-771-2018>, 2018.
- 1550 ○ Chen, X., Jeong, S., Park, H., Kim, J., and Park, C.-R.: Urbanization has stronger impacts
1551 than regional climate change on wind stilling: a lesson from South Korea, Environ. Res.
1552 Lett., 15, 054016, <https://doi.org/10.1088/1748-9326/ab7e51>, 2020.
- 1553 ○ Chmielewski, T., Szer, J. and Bobra, P.: Derecho windstorm in Poland on 11–12 August
1554 2017: results of the post-disaster investigation, Environ. Hazards, 1–21,
1555 doi:10.1080/17477891.2020.1730154, 2020.
- 1556 ○ Cionco, R.: A wind-profile index for canopy flow. Bound.-Layer Meteor. 3, 255–263,
1557 <https://doi.org/10.1007/BF02033923>, 1972.
- 1558 ○ Clark, P. and Gray, S.: Sting jets in extratropical cyclones: A review. Q. J. Roy. Meteorol.
1559 Soc., 144, 943-969, 2018.
- 1560 ○ Crasemann, B., Handorf, D., Jaiser, R., Dethloff, K., Nakamura, T., Ukita, J., and Yamazaki,
1561 K.: Can preferred atmospheric circulation patterns over the North-Atlantic-Eurasian region
1562 be associated with Arctic sea ice loss? Polar Sci., 14, 9-20, doi:10.1016/j.polar.2017.09.002,
1563 2017.
- 1564 ~~○ Crespi, A., Terzi, S., Cocuccioni, S., Zebisch, M., Berckmans, J. and Füssel, H.M.: 2020. Climate-~~
1565 ~~related hazard indices for Europe. ETC-CCA Technical Paper, 1, European Topic Centre on Climate~~
1566 ~~Change—Impacts, Vulnerability and Adaptation (ETC/CCA).~~
1567 ~~https://doi.org/10.25424/cmcc/climate_related_hazard_indices_europe_2020~~
- 1568 ○ Csilléry, K., Kunstler, G., Courbaud, B., Allard, D., Lassegues, P., Haslinger, K., Gardiner,
1569 B., Lassègues, P., Haslinger, K. and Gardiner, B.: Coupled effects of wind-storms and
1570 drought on tree mortality across 115 forest stands from the Western Alps and the Jura
1571 mountains, Glob. Chang. Biol., 23, 5092–5107, doi:10.1111/gcb.13773, 2017.
- 1572 ○ Couasnon, A., Eilander, D., Muis, S., Veldkamp, T., Haigh, I., Wahl, T., Winsemius, H., and
1573 Ward, P.: Measuring compound flood potential from river discharge and storm surge

- 1574 extremes at the global scale and its implications for flood hazard, *Nat. Hazards Earth Syst.*
 1575 *Sci.* 20, 489–504, <https://doi.org/10.5194/nhess-20-489-2020>, 2020.
- 1576 ○ Cullen, S.: Trees and wind: a practical consideration of the drag equation velocity exponent
 1577 for urban tree risk management, *Arboriculture Urban Forestry*, 31, 101, 2005.
- 1578 ○ Cvitan, L.: Determining wind gusts using mean hourly wind speed. *Geofizika*, 20, 63-74,
 1579 2003.
- 1580 ○ Dacre, H.: [A review of extratropical cyclones: observations and conceptual models over the](#)
 1581 [past 100 years. *Weather*, 75, 4–7, 2020](#)
- 1582 ○ Dacre, H. and Gray, S.: The spatial distribution and evolution characteristics of North
 1583 Atlantic cyclones. *Mon. Wea. Rev.*, 137, 99–115, 2009
- 1584 ~~○ Dacre, H.: [A review of extratropical cyclones: observations and conceptual models over the](#)~~
 1585 ~~[past 100 years. *Weather*, 75, 4–7, 2020](#)~~
- 1586 ○ ~~Dacre, H.~~, and Pinto, J. G.: Serial clustering of extratropical cyclones: a review of where,
 1587 when and why it occurs. *npj Clim. Atmos. Sci.* 3, 48, [https://doi.org/10.1038/s41612-020-](https://doi.org/10.1038/s41612-020-00152-9)
 1588 00152-9, 2020
- 1589 ○ Dalezios, N. R., Gobin, A., Tarquis Alfonso, A. M. and Eslamian, S.: Agricultural Drought
 1590 Indices: Combining Crop, Climate, and Soil Factors, in *Handbook of Drought and Water*
 1591 *Scarcity: Principles of Drought and Water Scarcity*, edited by S. Eslamian and F. A.
 1592 Eslamian, pp. 73–89, CRC Press., 2017.
- 1593 ○ Danielsson, U.: Windchill and the risk of tissue freezing, *J. Appl. Physiol.*, 81, 6, 2666-2673,
 1594 <https://doi.org/10.1152/jappl.1996.81.6.2666>, 1996.
- 1595 ○ Decker, L.: Windthrow risk assessment of Douglas-fir stands in Denmark - A comparison
 1596 study of Douglas-fir (*Pseudotsuga menziesii*) and Norway spruce (*Picea abies*), PhD.
 1597 Thesis, [https://naturstyrelsen.dk/media/249959/windthrow-risk-assessment-of-douglas-fir-](https://naturstyrelsen.dk/media/249959/windthrow-risk-assessment-of-douglas-fir-stands-in-denmark.pdf)
 1598 [stands-in-denmark.pdf](https://naturstyrelsen.dk/media/249959/windthrow-risk-assessment-of-douglas-fir-stands-in-denmark.pdf), 2018.
- 1599 ○ Défosse, P., Rajaonalison, F. and Bosc, A.: [How wind acclimation impacts *Pinus pinaster*](#)
 1600 [growth in comparison to resource availability. *Forestry*, 95, 118–129,](#)
 1601 <https://doi.org/10.1093/forestry/cpab028>, 2022.
- 1602 ○ Défosse, P., Veylon, G., Yang, M., Bonnefond, J., Garrigou, D., Trichet, P. and Danjon, F.:
 1603 Impact of soil water content on the overturning resistance of young *Pinus Pinaster* in sandy
 1604 soil, *For. Ecol. Manage.*, 480, 118614, doi:10.1016/j.foreco.2020.118614, 2021.
- 1605 ~~○ Défosse, P., Rajaonalison, F. and Bosc, A.: [How wind acclimation impacts *Pinus pinaster*](#)~~
 1606 ~~[growth in comparison to resource availability. *Forestry*, 95, 118–129,](#)~~
 1607 ~~<https://doi.org/10.1093/forestry/cpab028>, 2022.~~
- 1608 ○ Díaz-Yáñez, O., Mola-Yudego, B. and González-Olabarria, J.: Modelling damage
 1609 occurrence by snow and wind in forest ecosystems. *Ecological Modelling*, 408, 108741,
 1610 2019.
- 1611 ○ Diffenbaugh, N., Scherer, M. and Trapp, R.: Robust increases in severe thunderstorm
 1612 environments in response to greenhouse forcing. *Proc. Nat. Acad. Sci.*, 110, 16361-16366,
 1613 2013.
- 1614 ○ Ding, M., Wu, Y.-C., and Zhang, L.-J.: Study on the algorithm to the probabilistic distribution
 1615 parameters of wind speed in wind farms. *Proc. Chinese Soc. Elec. Eng.*, 25, 107-110,
 1616 http://en.cnki.com.cn/Article_en/CJFDTOTAL-ZGDC200510019.htm, last access: 17
 1617 December 2021, 2005.
- 1618 ○ Donat, M., Pardowitz, T., Leckebusch, G., Ulbrich, U., and Burghoff, O.: High-resolution
 1619 refinement of a storm loss model and estimation of return periods of loss-intensive storms

- 1620 over Germany. *Nat. Hazards Earth Syst. Sci.*, 11, 2821–2833. [https://doi.org/10.5194/nhess-](https://doi.org/10.5194/nhess-11-2821-2011)
1621 [11-2821-2011](https://doi.org/10.5194/nhess-11-2821-2011)<https://doi.org/10.5194/nhess-11-2821-2011>, 2011.
- 1622 ○ Dorland, C., Tol, R., and Palutikof, J.: Vulnerability of the Netherlands and Northwest Europe
1623 to storm damage under climate change. *Clim. Change*, 43, 513–535.
1624 <https://doi.org/10.1023/A:1005492126814><https://doi.org/10.1023/A:1005492126814>, 1999.
- 1625 ○ Dorrington, J., and Strommen, K.: Jet speed variability obscures Euro-Atlantic regime
1626 structure, *Geophys. Res. Lett.*, 47, e2020GL087907, doi:10.1029/2020GL087907, 2020.
- 1627 ○ Droste, A., Steeneveld, G., and Holtslag, A.: Introducing the urban wind island effect,
1628 *Environ. Res. Lett.*, 13, 094007, <https://doi.org/10.1088/1748-9326/aad8ef>, 2018.
- 1629 ○ Duperat, M., Gardiner, B., Ruel, J.-C.: Testing an individual tree wind damage risk model in
1630 a naturally regenerated balsam fir stand: potential impact of thinning on the level of risk.
1631 *Forestry: An International Journal of Forest Research* 94:141–150,
1632 <https://doi.org/10.1093/forestry/cpaa023>, 2021
- 1633 ○ Dupont, S.: A simple wind–tree interaction model predicting the probability of wind
1634 damage at stand level. *Agricultural and Forest Meteorology* 224:49–63,
1635 <https://doi.org/10.1016/j.agrformet.2016.04.014>, 2016.
- 1636 ○ Dupont, S., Ikonen, V.-P., Väisänen, H., and Peltola, H.: Predicting tree damage in
1637 fragmented landscapes using a wind risk model coupled with an airflow model, *Can. J. For.*
1638 *Res.*, 45, 1065–1076, <https://doi.org/10.1139/cjfr-2015-0066>, 2015.
- 1639 ○ DWD: Warnkriterien, Deutscher Wetterdienst (DWD):
1640 https://www.dwd.de/DE/wetter/warnungen_aktuell/kriterien/warnkriterien.html, last access:
1641 15 December 2021.
- 1642 ○ Economou, T., Stephenson, D., Pinto, J. G., Shaffrey, L. and Zappa, G.: Serial clustering of
1643 extratropical cyclones in a multi-model ensemble of historical and future simulations. *Quart.*
1644 *J. Roy. Meteor. Soc.*, 141, 3076–3087, 2015
- 1645 ○ Edwards, J.: The relationship between road accident severity and recorded weather, *J.*
1646 *Saf. Res.*, 29, 249–262, [https://doi.org/10.1016/S0022-4375\(98\)00051-6](https://doi.org/10.1016/S0022-4375(98)00051-6), 1998.
- 1647 ○ Ellis, J., Harris, M., Román-Rivera, M., Ferguson, J., Tereszkievicz, P., and McGill, S.:
1648 Application of the Saffir-Simpson Hurricane Wind Scale to Assess Sand Dune Response to
1649 Tropical Storms, *J. Marine Sci. Eng.*, 8, 670, <https://doi.org/10.3390/jmse8090670>, 2020.
- 1650 ○ Evans, J., Dasey, T., Rhoda, D., Cole, R., Wilson, F., and Williams, E.: Weather sensing
1651 and data fusion to improve safety and reduce delays at major west coast airports,
1652 Massachusetts Institute of Technology, Lincoln Laboratory, Project Report ATC-290, 30,
1653 [https://www.ll.mit.edu/sites/default/files/publication/doc/2018-12/Evans_1999_ATC-](https://www.ll.mit.edu/sites/default/files/publication/doc/2018-12/Evans_1999_ATC-290_WW-10288.pdf)
1654 [290_WW-10288.pdf](https://www.ll.mit.edu/sites/default/files/publication/doc/2018-12/Evans_1999_ATC-290_WW-10288.pdf), last access: 14 December 2021, 1999.
- 1655 ○ Everham, E. and Brokaw, N.: Forest damage and recovery from catastrophic wind. *Bot.*
1656 *Rev.*, 62, 113–185, <https://doi.org/10.1007/BF02857920>, 1996.
- 1657 ○ Favorskaya MN, Jain LC. 2017. Innovations in Remote Sensing of Forests. In: Handbook
1658 on Advances in Remote Sensing and Geographic Information Systems. Intelligent Systems
1659 Reference Library, vol 122. Springer, Cham. [https://doi.org/10.1007/978-](https://doi.org/10.1007/978-3-319-52308-8_1)
1660 [3https://doi.org/10.1007/978-3-](https://doi.org/10.1007/978-3-319-52308-8_1) 319-52308-8_1
- 1661 ○ Feldstein, S. and Franzke, C.: Atmospheric teleconnection patterns. In *Nonlinear and*
1662 *stochastic climate dynamics*, pp. 54–104. Cambridge University Press, 2017.
- 1663 ○ Feser, F., Barcikowska, M., Krueger, O., Schenk, F., Weisse, R., and Xia, L: Storminess
1664 over the North Atlantic and Northwestern Europe - a review. *Q. J. R. Meteorol. Soc.* 141,
1665 350–382, 2015.
- 1666 ○ Feuerstein, B., Groenemeijer, P., Dirksen, E., Hubrig, M., Holzer, A. and Dotzek, N.:
1667 Towards an improved wind speed scale and damage description adapted for Central

- 1668 Europe. Atmos. Res., 100, 547–564.
 1669 <https://doi.org/10.1016/j.atmosres.2010.12.026><https://doi.org/10.1016/j.atmosres.2010.12.02>
 1670 [6](https://doi.org/10.1016/j.atmosres.2010.12.026), 2011.
- 1671 ○ Fink, A., Brücher, T., Ermert, V., Krüger, A., and Pinto, J. G.: The European storm Kyrill in
 1672 January 2007: synoptic evolution, meteorological impacts and some considerations with
 1673 respect to climate change, *Nat. Hazards Earth Syst. Sci.*, 9, 405–423,
 1674 <https://doi.org/10.5194/nhess-9-405-2009>, 2009.
 - 1675 ○ Finnigan, J., Ayotte, K., Harman, I., Katul, G., Oldroyd, H., Patton, E., Poggi, D., Ross, A.
 1676 and Taylor, P.: Boundary-Layer Flow Over Complex Topography, *Boundary-Layer*
 1677 *Meteorol.*, 177(2–3), 247–313, doi:10.1007/s10546-020-00564-3, 2020.
 - 1678 ○ Flather, R.: Storm surges, in: *Encyclopedia of Ocean Sciences*, edited by: Steele, J.,
 1679 Thorpe, S., and Turekian, K., Academic, San Diego, California, 2882–2892, 2001
 - 1680 ○ Florinsky, I.: An illustrated introduction to general geomorphometry. *Prog. Phys. Geography*,
 1681 41, 723-752, 2017.
 - 1682 ○ Forzieri, G., Pecchi, M., Girardello, M., Mauri, A., Klaus, M., Nikolov, C., Rüetschi, M.,
 1683 Gardiner, B., Tomaščík, J., Small, D., Nistor, C., Jonikavicius, D., Spinoni, J., Feyen, L.,
 1684 Giannetti, F., Comino, R., Wolynski, A., Pirotti, F., Maistrelli, F., Ionut, S., Lucas-Stephanie,
 1685 W., Stefan, K., Zieba-Kulawik, K., Strejczek-Jazwinska, P., Mokroš, M., Stefan, F., Krejci,
 1686 L., Haidu, I., Nilsson, M., Wezyk, P., Chirici, G., Cescatti, A., and Beck, P.: A spatially-explicit
 1687 database of wind disturbances in European forests over the period 2000-2018, *Earth Sys.*
 1688 *Sci. Data*, 1–23, <https://doi.org/10.5194/essd-2019-141>, 2019.
 - 1689 ○ Franzke, C.: Persistent regimes and extreme events of the North Atlantic atmospheric
 1690 circulation. *Phil. Trans. Roy. Soc.*, 371, 20110471, 2013.
 - 1691 ○ Franzke, C.: [Impacts of a Changing Climate on Economic Damages and Insurance.](#)
 1692 [Economics of Disasters and Climate Change, 1, 95-110. doi:10.1007/s41885-017-0004-3,](#)
 1693 [2017.](#)
 - 1694 ○ [Franzke, C.:](#) Towards the development of economic damage functions for weather and
 1695 climate extremes, *Ecol. Econ.*, 189, 107172, 2021.
 - 1696 ○ Franzke, C., Woollings, T. and Martius, O.: Persistent circulation regimes and preferred
 1697 regime transitions in the North Atlantic. *J. Atmos. Sci.*, 68, 2809-2825, 2011.
 - 1698 ○ Frey, O. and DeVogelaere, A.: The Containerized Shipping Industry and the Phenomenon
 1699 of Containers Lost at Sea. *Marine Sanctuaries Conservation Series*, ONMS-14. 1-51,
 1700 <https://sanctuaries.noaa.gov/science/conservation/lostcontainers.html>, last access: 14
 1701 December 2021, 2014.
 - 1702 ○ Fujita, T.: Tornadoes and Downbursts in the Context of Generalized Planetary Scales, *J.*
 1703 *Atmos. Sci.*, 38, 1511–1534, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0469(1981)038<1511:TADITC>2.0.CO;2)
 1704 [0469\(1981\)038<1511:TADITC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1981)038<1511:TADITC>2.0.CO;2), 1981.
 - 1705 ○ Ganbat, G., Seo, J., Han, J. et al.: A theoretical study of the interactions of urban breeze
 1706 circulation with mountain slope winds. *Theor. Appl. Clim.* 121, 545–555,
 1707 <https://doi.org/10.1007/s00704-014-1252-6>, 2015.
 - 1708 ○ Garcia-Alonso, L., Moura, T. G. Z., and Roibas, D.: The effect of weather conditions on port
 1709 technical efficiency, *Mar. Policy*, 113, 103816,
 1710 <https://doi.org/10.1016/j.marpol.2020.103816>, 2020.
 - 1711 ○ Gardiner, B.: Mathematical modelling of the static and dynamic characteristics of plantation
 1712 trees. In Franke, J. and Roeder, A.: *Mathematical Modelling of Forest Ecosystems*,
 1713 Sauerländers Verlag, 40–61, 1992.
 - 1714 ○ Gardiner, B.: The interactions of wind and tree movement in forest canopies, in: *Wind and*
 1715 *Trees*, edited by: Coutts, M. and Grace, J., Cambridge University Press, 1995.

- 1716 ○ Gardiner, B.: Wind damage to forests and trees: a review with an emphasis on planted and
1717 managed forests, 26, 248–266, <https://doi.org/10.1080/13416979.2021.1940665>, 2021.
- 1718 ○ Gardiner, B., Berry, P. and Moulia, B.: Review: Wind impacts on plant growth, mechanics
1719 and damage. Plant Science, 245, 94-118,
1720 <https://doi.org/doi:10.1016/j.plantsci.2016.01.006>, 2016.
- 1721 ~~○ Gardiner, B., Peltola, H., and Kellomäki, S.: Comparison of two models for predicting the~~
1722 ~~critical wind speeds required to damage coniferous trees, *Ecol. Model.*, 129, 1–23,~~
1723 ~~[https://doi.org/10.1016/S0304-3800\(00\)00220-9](https://doi.org/10.1016/S0304-3800(00)00220-9), 2000.~~
- 1724 ~~○ Gardiner, B., Byrne, K., Hale, S., Kamimura, K., Mitchell, S., Peltola, H., and Ruel, J. C.: A~~
1725 ~~review of mechanistic modelling of wind damage risk to forests, 81, 447–463,~~
1726 ~~<https://doi.org/10.1093/forestry/cpn022>, 2008.~~
- 1727 ○ Gardiner, B., Blennow, K., Carnus, J. M., Fleischer, P., Ingemarson, F., Landmann, G.,
1728 Lindner, M., Marzano, M., Nicoll, B., Orazio, C., Peyron, J., Reviron, M., Schelhaas, M.,
1729 Schuck, A., Spielmann, M., and Usbeck, T.: Destructive storms in European forests: past
1730 and forthcoming impacts., Final report to European Commission-DG Environment.
1731 European Forest Institute, <https://library.wur.nl/WebQuery/wurpubs/fulltext/162053> 2010.
- 1732 ○ Gardiner, B., Byrne, K., Hale, S., Kamimura, K., Mitchell, S., Peltola, H., and Ruel, J.-C.: A
1733 review of mechanistic modelling of wind damage risk to forests, 81, 447–463,
1734 <https://doi.org/10.1093/forestry/cpn022>, 2008.
- 1735 ○ Gardiner, B., Peltola, H., and Kellomäki, S.: Comparison of two models for predicting the
1736 critical wind speed required to damage coniferous trees, *Ecol. Model.*, 129, 1–23,
1737 [https://doi.org/10.1016/S0304-3800\(00\)00220-9](https://doi.org/10.1016/S0304-3800(00)00220-9), 2000.
- 1738 ○ Gardiner, B., Schuck, A., Schelhaas, M.-J., Orazio, C., Blennow, K. and Nicoll, B.: Living
1739 with Storm Damage to Forests: What Science Can Tell Us, edited by B. Gardiner, A. Schuck,
1740 M.-J. Schelhaas, C. Orazio, K. Blennow, and B. Nicoll, European Forest Institute, Joensuu.,
1741 2013.
- 1742 ~~○ Gardiner, B., Berry, P. and Moulia, B.: Review: Wind impacts on plant growth, mechanics and~~
1743 ~~damage. *Plant Science*, 245, 94–118, <https://doi.org/doi:10.1016/j.plantsci.2016.01.006>,~~
1744 ~~2016.~~
- 1745 ○ Garratt, J.: Surface influence upon vertical profiles in the atmospheric near-surface layer.
1746 *Quart. J. Roy. Meteorol. Soc.*, 106, 803–819, doi:10.1256/smsqj.45010, 1980.
- 1747 ○ Gatzen, C., Fink, A., Schultz, D., and Pinto, J. G.: An 18-year climatology of derechos in
1748 Germany, *Nat. Hazards Earth Syst. Sci.*, 20, 1335–1351, [https://doi.org/10.5194/nhess-20-](https://doi.org/10.5194/nhess-20-1335-2020)
1749 [1335-2020](https://doi.org/10.5194/nhess-20-1335-2020), 2020.
- 1750 ○ Gholizadeh, H.; Zoghipour, M., Torshizi, M.; Nazari, M., and Moradkhani, N.: Gone with the
1751 wind: Impact of soil-dust storms on farm income. *Ecol. Econ.*, 188, 107133. DOI:
1752 10.1016/j.ecolecon.2021.107133, 2021.
- 1753 ○ Goff, T., Nelson, M., Liknes, G., Feeley, T., Pugh, S. and Morin, R.: Rapid assessment of
1754 tree damage resulting from a 2020 windstorm in Iowa, USA, *Forests*, 12, 1–13,
1755 doi:10.3390/f12050555, 2021.
- 1756 ○ Gonçalves, A., Liberato, M., and Nieto, R.: Wind Energy Assessment during High-Impact
1757 Winter Storms in the Iberian Peninsula, *Environ. Sci. Proc.*, 4, 28,
1758 <https://doi.org/10.3390/ecas2020-08132>, 2020.
- 1759 ○ Gonçalves, A., Liberato, M., and Nieto, R.: Wind Energy Assessment during High-Impact
1760 Winter Storms in Southwestern Europe, *Atmosphere*, 12, 509,
1761 <https://doi.org/10.3390/atmos12040509>, 2021.

- 1762 ○ Grams, C., Binder, H., Pfahl, S., Piaget, N. and Wernli, H.: Atmospheric processes triggering
 1763 the Central European floods in June 2013, *Nat. Hazards Earth Syst. Sci.*, 14, 1691–1702,
 1764 doi:10.5194/nhess-14-1691-2014, 2014.
- 1765 ○ Grimmond, C. and Oke, T.: Aerodynamic Properties of Urban Areas Derived from Analysis
 1766 of Surface Form, *J. App. Meteor. Clim.*, 38, 1262–1292, [https://doi.org/10.1175/1520-0450\(1999\)038<1262:APOUAD>2.0.CO;2](https://doi.org/10.1175/1520-0450(1999)038<1262:APOUAD>2.0.CO;2), 1999.
- 1767 ○ Gromke, C. and Ruck, B.: On Wind Forces in the Forest-Edge Region During Extreme-Gust
 1768 Passages and Their Implications for Damage Patterns, *Bound.-Layer Meteor.*, 168, 269–
 1769 288, <https://doi.org/10.1007/s10546-018-0348-4>, 2018.
- 1771 ○ Gross, G.: A windthrow model for urban trees with application to storm “Xavier.”
 1772 *Meteorologische Zeitschrift* 27 (2018), Nr. 4 27:299–308, <https://doi.org/10.15488/4901>,
 1773 2018.
- 1774 ○ Gyakum, J. and Danielson, R.: Analysis of meteorological precursors to ordinary and
 1775 explosive cyclogenesis in the western North Pacific. *Mon. Wea. Rev.*, 128, 851-863, 2000.
- 1776 ○ Hahn, B. and Rohrig, K.: ISET-Wind-Index Assessment of the Annual Available Wind
 1777 Energy, *Proc. Europ. Wind Energy Conf. Exh. EWEC2003*, Madrid, Spain, 16-19 Jun 2003,
 1778 <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.196.1675&rep=rep1&type=pdf>,
 1779 last access: 21 December 2021, 2003
- 1780 ○ Hale, S., Gardiner, B., Peace, A., Nicoll, B., Taylor, P., and Pizzirani, S.: Comparison and
 1781 validation of three versions of a forest wind risk model, *Environ. Model. Software*, 68, 27–
 1782 41, <https://doi.org/10.1016/j.envsoft.2015.01.016>, 2015.
- 1783 ~~○ Hanewinkel, M., Zhou, W. and Schill, C.: A neural network approach to identify forest stands
 1784 susceptible to wind damage. *Forest Ecol. Manag.*, 196, 227–243, 2004.~~
- 1785 ~~○ Hanewinkel, M., Kuhn, T., Bugmann, H., Lanz, A., and Brang, P.: Vulnerability of uneven-
 1786 aged forests to storm damage, *Forestry*, 87, 525–534,
 1787 <https://doi.org/10.1093/forestry/cpu008>, 2014.~~
- 1788 ○ ~~Hanewinkel, M., Cullmann, D. a., Schelhaas, M.-J., Nabuurs, G.-J., Zimmermann, N:~~
 1789 ~~Climate change may cause severe loss in the economic value of European forest land. *Nat.*
 1790 ~~*Clim. Chang.* 3, 203–207. <https://doi.org/10.1038/nclimate1687>, 2013.~~~~
- 1791 ○ Hanewinkel, M., Kuhn, T., Bugmann, H., Lanz, A., and Brang, P.: Vulnerability of uneven-
 1792 aged forests to storm damage, *Forestry*, 87, 525–534,
 1793 <https://doi.org/10.1093/forestry/cpu008>, 2014.
- 1794 ○ Hanewinkel, M., Zhou, W. and Schill, C.: A neural network approach to identify forest stands
 1795 susceptible to wind damage. *Forest Ecol. Manag.*, 196, 227-243, 2004.
- 1796 ○ Hannachi, A., Straus, D., Franzke, C., Corti, S., and Woollings, T.: Low-frequency
 1797 nonlinearity and regime behavior in the Northern Hemisphere extratropical atmosphere,
 1798 *Rev. Geophys.*, 55, 199-234, doi:10.1175/2009JCLI3087.1, 2017.
- 1799 ○ Harper, B., Kepert, J. and Ginger, J.: Wind speed time averaging conversions for tropical
 1800 cyclone conditions, in *AMS 28th Conference on Hurricanes and Tropical Meteorology*, pp.
 1801 1–2, Orlando, FL., 2008.
- 1802 ○ Harvey, B., Cook, P., Shaffrey, L. and Schiemann, R.: The response of the northern
 1803 hemisphere storm tracks and jet streams to climate change in the CMIP3, CMIP5, and
 1804 CMIP6 climate models. *J. Geophys. Res.*, 125, e2020JD032701, 2020.
- 1805 ○ Haylock, M.: European extra-tropical storm damage risk from a multi-model ensemble of
 1806 dynamically-downscaled global climate models. *Nat. Haz. Earth Sys. Sci.*, 11, 2847–2857.
 1807 <https://doi.org/10.5194/nhess-11-2847-2011><https://doi.org/10.5194/nhess-11-2847-2011>,
 1808 2011.

- 1809 ○ Held, I.M.: Momentum transport by quasi-geostrophic eddies. *J. Atmos. Sci.*, 32, 1494-1497,
1810 1975.
- 1811 ○ Held, I.: 100 years of progress in understanding the general circulation of the atmosphere.
1812 *Meteorological Monographs*, 59, 6-1, 2019.
- 1813 ○ Held, I.M. and Hou, A.Y.: Nonlinear axially symmetric circulations in a nearly inviscid
1814 atmosphere. *J. Atmos. Sci.*, 37, 515-533, 1980.
- 1815 ○ Hemingway, R., and Robbins, J.: Developing a hazard-impact model to support impact-
1816 based forecasts and warnings: The Vehicle OverTurning (VOT) Model, *Meteorol. Appl.*,
1817 27, <https://doi.org/10.1002/met.1819>, 2020.
- 1818 ○ Heneka P, Hofherr T, Ruck B, Kottmeier C: Winter storm risk of residential structures—model
1819 development and application to the German state of Baden-Württemberg. *Nat. Haz. Earth*
1820 *Sys.*, 6, 721–733, 2006.
- 1821 ○ Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas,
1822 J., Peubey, C., Radu, R., Schepers, D., Simmons, A., et al.: The ERA5 global reanalysis,
1823 *Q. J. Roy. Meteor. Soc.*, 146, 1999-2049,
1824 <https://doi.org/10.1002/qj.3803><https://doi.org/10.1002/qj.3803>, 2020.
- 1825 ○ Hewson, T. and Neu, U.: Cyclones, windstorms and the IMILAST project. *Tellus A*, 67,
1826 27128, 2015.
- 1827 ○ Hoeppe, P.: Trends in weather related disasters—Consequences for insurers and society.
1828 *Weather and climate extremes*, 11, 70-79, 2016.
- 1829 ○ Hofman, M., and Bačkalov, I.: Risk-based analysis of inland vessel stability, In Proceedings
1830 of the 11th International Ship Stability Workshop, Wageningen, 67-72,
1831 [http://www.shipstab.org/files/Proceedings/ISSW/ISSW_2010_Wageningen_The_Netherla](http://www.shipstab.org/files/Proceedings/ISSW/ISSW_2010_Wageningen_The_Netherlands/ISSW_2010_s03-p2.pdf)
1832 [nds/ISSW_2010_s03-p2.pdf](http://www.shipstab.org/files/Proceedings/ISSW/ISSW_2010_Wageningen_The_Netherlands/ISSW_2010_s03-p2.pdf), last access: 14 December 2021, 2010.
- 1833 ○ Holland, A., Riordan, A., and Franklin, E.: A Simple Model for Simulating Tornado Damage
1834 in Forests, *J. App. Meteor. Clim.* 45, 1597–1611, <https://doi.org/10.1175/JAM2413.1>, 2006.
- 1835 ○ Holton, J. and Hakim, G.: *An Introduction to Dynamic Meteorology*, Vol. 9780123848, Wiley,
1836 2012.
- 1837 ○ Hoskins B. and Valdes P.: On the existence of storm tracks. *J. Atmos. Sci.*, 47, 1854–1864,
1838 1990.
- 1839 ○ IPCC: Summary for Policymakers. In: *Climate Change 2021: The Physical Science Basis.*
1840 *Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental*
1841 *Panel on Climate Change [Masson-Delmotte, V., et al. (eds.)].*
1842 <https://www.ipcc.ch/report/ar6/wg1/#SPM>, 2021.
- 1843 ○ INTERCARGO: Bulk Carrier Casualty Report - Years 2008 to 2017 and the trends,
1844 International Organization of Dry Cargo Shipowners, London, United Kingdom,
1845 [https://www.intercargoo.org/wp-content/uploads/2018/05/bulk-carrier-casualty-report-](https://www.intercargoo.org/wp-content/uploads/2018/05/bulk-carrier-casualty-report-2017.pdf)
1846 [2017.pdf](https://www.intercargoo.org/wp-content/uploads/2018/05/bulk-carrier-casualty-report-2017.pdf), last access: 14 December 2021, 2018.
- 1847 ○ Jenamani, R., and Kumar, A.: Bad weather and aircraft accidents—global vis-à-vis Indian
1848 scenario. *Curr. Sci.*, 104, 316-325, <https://www.jstor.org/stable/24089632>, last access: 14
1849 December 2021, 2013.
- 1850 ○ Jim, C. and Liu, H.: Storm damage on urban trees in Guangzhou, China, *Landsc. Urban*
1851 *Plan.*, 38, 45–59, doi:10.1016/S0169-2046(97)00018-2, 1997.
- 1852 ○ Joseph, G., Mohammadi, M., Sterling, M., Baker, C., Gillmeier, S., Soper, D. et al.:
1853 Determination of crop dynamic and aerodynamic parameters for lodging prediction. *J. Wind*
1854 *Eng. Industrial Aerodyn.*, 202, 104169. DOI: 10.1016/j.jweia.2020.104169, 2020.

- 1855 ~~○ Jung, C., Schindler, D., Albrecht, A. and Buchholz, A.: The role of highly resolved gust~~
 1856 ~~speed in simulations of storm damage in forests at the landscape scale: A case study from~~
 1857 ~~southwest Germany. Atmosphere, 7, 7, 2016.~~
- 1858 ○ Jung, C. and Schindler, D.: Introducing a new approach for wind energy potential
 1859 assessment under climate change at the wind turbine scale. Energy Conversion Manag.,
 1860 225, 113425, <https://doi.org/10.1016/j.enconman.2020.113425>, 2020.
- 1861 ○ Jung, C. and Schindler, D.: Does the winter storm-related wind gust intensity in Germany
 1862 increase under warming climate? – A high-resolution assessment. In Wea. Clim. Extremes
 1863 33, 100360. DOI: 10.1016/j.wace.2021.100360, 2021.
- 1864 ○ Jung, C., Schindler, D., Albrecht, A. and Buchholz, A.: The role of highly-resolved gust
 1865 speed in simulations of storm damage in forests at the landscape scale: A case study from
 1866 southwest Germany. Atmosphere, 7, 7, 2016.
- 1867 ○ Jung, C., Schindler, D.: Statistical Modeling of Near-surface Wind Speed: A Case Study
 1868 from Baden-Wuerttemberg (Southwest Germany). Austin Journal of Earth Science Austin J
 1869 Earth Science:1006, 2015.
- 1870 ○ Kaimal, J. and Finnigan, J.: Atmospheric boundary layer flows: Their structure and
 1871 measurement, Oxford University Press, 1994.
- 1872 ○ Finnigan, J., Ayotte, K., Harman, I., Katul, G., Oldroyd, H., Patton, E., Poggi, D., Ross, A.
 1873 and Taylor, P.: Boundary-Layer Flow Over Complex Topography, Boundary-Layer
 1874 Meteorol., 177(2–3), 247–313, doi:10.1007/s10546-020-00564-3, 2020.
- 1875 ○ Kaltenberger, R., Schaffhauser, A. & Staudinger, M. (2020) “What the weather will do” –
 1876 results of a survey on impact-oriented and impact-based warnings in European NMHSs.
 1877 Advances in Science and Research, 17, 29–38. <https://doi.org/10.5194/asr-17-29-2020>
- 1878 ○ Kamimura, K., Nanko, K., Matsumoto, A., Ueno, S. and Gardiner, B.: Tree dynamic
 1879 response and survival in a category-5 tropical cyclone: The case of super typhoon Trami.
 1880 Sci. Adv., 8, 1–11, 2022.
- 1881 ○ Kaspar, F., Müller-Westermeier, G., Penda, E., Mächel, H., Zimmermann, K., Kaiser-Weiss,
 1882 A., and Deutschländer, T.: Monitoring of climate change in Germany – data, products and
 1883 services of Germany’s National Climate Data Centre, Adv. Sci. Res., 10, 99–106,
 1884 <https://doi.org/10.5194/asr-10-99-2013>, 2013.
- 1885 ○ Katinas, V., Gecevicius, G., and Marciukaitis, M.: An investigation of wind power density
 1886 distribution at location with low and high wind ~~speeds~~speed using statistical model, Applied
 1887 Energy, 218, 442–451, <https://doi.org/10.1016/j.apenergy.2018.02.163>, 2018.
- 1888 ○ Kelman, I.: Defining Risk, FloodRiskNet Newsletter, Issue 2,
 1889 <http://www.ilankelman.org/abstracts/kelman2003frn.pdf>, 2003.
- 1890 ○ Kirk, P.: An updated tornado climatology for the UK: 1981–2010, Weather, 69, 171–175,
 1891 <https://doi.org/10.1002/wea.2247>, 2014.
- 1892 ○ Kite-Powell, H.: The value of ocean surface wind information for maritime commerce, Mar.
 1893 Technol. Soc. J., 45(1), 75-84, <https://doi.org/10.4031/MTSJ.45.1.13>, 2011.
- 1894 ○ ~~Kitsara, G., van der Schriek, T., Varotsos, K., and Giannakopoulos, C.: Future changes in climate~~
 1895 ~~indices relevant to agriculture in the Aegean islands (Greece). Euro-Mediterr. J. Environ. Integr., 6,~~
 1896 ~~1. DOI: 10.1007/s41207-020-00233-4, 2021.~~
- 1897 ○ Klawa, M., and Ulbrich, U.: A model for the estimation of storm losses and the identification
 1898 of severe winter storms in Germany, Nat. Hazard Earth Sys, 3, 725-732,
 1899 <https://doi.org/10.5194/nhess-3-725-2003><https://doi.org/10.5194/nhess-3-725-2003>, 2003.
- 1900 ○ Koks, E. and Haer, T.: A high-resolution wind damage model for Europe, Sci. Rep., 10,
 1901 6866, <https://doi.org/10.1038/s41598-020-63580-w>, 2020

- 1902 ○ Kong, X., Peng, P., Li, L., Zhang, K., Hu, Z., Wang, X. and Zhao, G., 2022. Wind lodging-
 1903 associated yield loss is reduced by wheat genetic diversity. *European Journal of Agronomy*,
 1904 133, p.126441.
- 1905 ○ König, K. and Höfle, B.: Full-Waveform Airborne Laser Scanning in Vegetation Studies—A
 1906 Review of Point Cloud and Waveform Features for Tree Species Classification. *Forests*, 7,
 1907 198, <https://doi.org/10.3390/f7090198>, 2016.
- 1908 ○ Kramer, M., Hansen, A., Taper, M. and Kissinger, E.: Abiotic controls on long-term
 1909 windthrow disturbance and temperate rain forest dynamics in southeast Alaska. *Ecology*,
 1910 82, 2749-2768, 2001.
- 1911 ○ Krayner, W. and Marshall, R.: Gust factors applied to hurricane winds, *Bull. Am. Meteorol.*
 1912 *Soc.*, 73, 613–617, doi:10.1175/1520-0477(1992)073<0613:gfathw>2.0.co;2, 1992.
- 1913 ○ Krejci, L., Kolejka, J., Vozenilek, V. and Machar, I.: Application of GIS to empirical windthrow
 1914 risk model in mountain forested landscapes. *Forests*, 9, 96, 2018.
- 1915 ○ Kushla, J. and Ripple, W.: The role of terrain in a fire mosaic of a temperate coniferous
 1916 forest, *For. Ecol. Manage.*, 95, 97–107, doi:10.1016/S0378-1127(97)82929-5, 1997.
- 1917 ○ Lamb, H.: *Historic storms of the North Sea, British Isles, and Northwest Europe*. Cambridge
 1918 University Press, 1991.
- 1919 ○ Leckebusch, G., Renggli, D. and Ulbrich, U.: Development and application of an objective
 1920 storm severity measure for the Northeast Atlantic region. *Met. Zeitschrift*, 17, 575–587.
 1921 <https://doi.org/10.1127/0941-2948/2008/0323> <https://doi.org/10.1127/0941-2948/2008/0323>,
 1922 2008.
- 1923 ○ Lee, S., Furtado, J. and Charlton-Perez, A.: Wintertime North American weather regimes
 1924 and the Arctic stratospheric polar vortex. *Geophys. Res. Lett.*, 46, 14892– 14900.
 1925 <https://doi.org/10.1029/2019GL085592>, 2019.
- 1926 ○ [Lepore, C., Abernathey, R., Henderson, N., Allen, J. T., & Tippett, M. K.: Future global
 1927 convective environments in CMIP6 models. *Earth's Future*, 9, e2021EF002277.
 1928 <https://doi.org/10.1029/2021EF002277>, 2021.](https://doi.org/10.1029/2021EF002277)
- 1929 ○ Leviäkangas, P., Tuominen, A., Molarius, R., and Kojo, H. et al.: Extreme weather impacts
 1930 on transport systems, VTT Working Papers 168, VTT Technical Research Centre of Finland,
 1931 <https://www.vttresearch.com/sites/default/files/pdf/workingpapers/2011/W168.pdf>, last
 1932 access: 14 December 2021, 2011.
- 1933 ○ Li, C. and Wettstein, J.: Thermally driven and eddy-driven jet variability in reanalysis. *J.*
 1934 *Climate*, 25, 1587-1596, 2012.
- 1935 ○ Liberato M., Pinto J. G., Trigo I., and Trigo R.: Klaus - an exceptional winter storm over
 1936 Northern Iberia and Southern France. *Weather* 66:330–334. doi:10.1002/wea.755, 2011
- 1937 ○ Lindén, J. and Holmer, B.: Thermally induced wind patterns in the Sahelian city of
 1938 Ouagadougou, Burkina Faso, *Theor. Appl. Clim.*, 105, 229–241,
 1939 <https://doi.org/10.1007/s00704-010-0383-7>, 2011.
- 1940 ○ Liu, R., Liu, S., Yang, X., Lu, H., Pan, X., Xu, Z., Ma, Y., and Xu, T.: Wind Dynamics Over
 1941 a Highly Heterogeneous Oasis Area: An Experimental and Numerical Study, *J. Geophys.*
 1942 *Res.* 123, 8418-8440 <https://doi.org/10.1029/2018JD028397>, 2018.
- 1943 ○ Locatelli, T., Tarantola, S., Gardiner, B., Patenaude, G.: Variance-based sensitivity analysis
 1944 of a wind risk model - Model behaviour and lessons for forest modelling, *Environ. Model.*
 1945 *Softw.* 87, <https://doi.org/10.1016/j.envsoft.2016.10.010>, 84–109, 2017.
- 1946 ○ Ma, Q., Lembo, V. and Franzke, C.: The Lorenz energy cycle: trends and the impact of
 1947 modes of climate variability. *Tellus A*, 73,1-15, 2021.
- 1948 ○ Machar, I., Pechanec, V., Brus, J., Kilianova, H., Kirchner, K.: Forest management at the
 1949 upper treeline in Jeseníky Mountains (Czech Republic). In *International Multidisciplinary*

- 1950 Scientific GeoConference-SGEM; SGEM, Ed.; STEF92 Technology Ltd.: Albena, Bulgaria,
1951 2014; pp. 361–36
- 1952 ○ Maeden, G.: Tornadoes in Britain: Their intensities and distribution in space and time. *J.*
1953 *Meteor.*, 1, 242–251, 1976.
- 1954 ○ Mailier, P., Stephenson, D., Ferro, C. and Hodges, K.: Serial clustering of extratropical
1955 cyclones. *Mon. Wea. Rev.*, 134, 2224–2240, 2006.
- 1956 ○ Martius, O., Pfahl, S., and Chevalier, C.: A global quantification of compound precipitation
1957 and wind extremes, *Geophys. Res. Lett.*, 43, 7709–7717,
1958 <https://doi.org/10.1002/2016GL070017>, 2016.
- 1959 ○ Masoudvaziri, N., Ganguly, P., Mukherjee, S. and Sun, K.: Impact of geophysical and
1960 anthropogenic factors on wildfire size: a spatiotemporal data-driven risk assessment
1961 approach using statistical learning, *Stoch. Environ. Res. Risk Assess.*, doi:10.1007/s00477-
1962 021-02087-w, 2021.
- 1963 ○ Masroor, M., Rehman, S., Avtar, R., Sahana, M., Ahmed, R. and Sajjad, H.: Exploring
1964 climate variability and its impact on drought occurrence: evidence from Godavari Middle
1965 sub-basin, India. *Wea. Clim. Extremes*, 30, 100277, 2020.
- 1966 ○ Mayer, H. Wind-induced tree sways. *Trees*, 1, 195–206,
1967 <https://doi.org/10.1007/BF01816816>, 1987.
- 1968 ○ Mayer, P., Brang, P., Dobbertin, M., Hallenbarter, D., Renaud, J., Walthert, L. and
1969 Zimmermann, S.: Forest storm damage is more frequent on acidic soils. *Ann. Forest Sci.*,
1970 62, 303-311, 2005.
- 1971 ○ McGinley, J.: A Diagnosis of Alpine Lee Cyclogenesis. *Mon. Wea. Rev.*, 110, 1271-1287,
1972 1982.
- 1973 ○ McVicar, T., Roderick, M., Donohue, R., Li, L., van Niel, T., Thomas, A. et al.: Global review
1974 and synthesis of trends in observed terrestrial near-surface wind ~~speeds~~speed: Implications
1975 for evaporation. *J. Hydrology*, 416-417, 182–205. DOI: 10.1016/j.jhydrol.2011.10.024, 2012.
- 1976 ○ [Merz, B., Kuhlicke, C., Kunz, M., Pittore, M., Babeyko, A., Bresch, D.N., Domeisen, D.I.,
1977 Feser, F., Koszalka, I., Kreibich, H. and Pantillon, F.: Impact forecasting to support
1978 emergency management of natural hazards. *Reviews of Geophysics*, 58\(4\),
1979 \[p.e2020RG000704\]\(#\), 2020.](#)
- 1980 ○ Messmer, M. and Simmonds, I.: Global analysis of cyclone-induced compound precipitation
1981 and wind extreme events, *Wea. Clim. Extremes*, 32, 100324,
1982 <https://doi.org/10.1016/j.wace.2021.100324>, 2021.
- 1983 ○ [MetEireann warning levels: <https://www.met.ie/met-eireann-warning-system-explained>, last
1984 access: 10.01.2023](#)
- 1985 ○ Miller, K. F.: Windthrow hazard in conifer plantations, *Irish For.*, 43, 66–78, 1986.
- 1986 ○ Milne, R.: Dynamics of swaying of *Picea sitchensis*, *Tree Physiology*, 9, 383–399,
1987 <https://doi.org/10.1093/treephys/9.3.383>, 1991.
- 1988 ○ Mitchell, S.: Wind as a natural disturbance agent in forests: a synthesis. *Forestry*. 86, 147–
1989 157, <https://doi.org/10.1093/forestry/cps058>, 2013.
- 1990 ○ Mitchell, S., Hailemariam, T. and Kulis, Y.: Empirical modeling of cutblock edge windthrow
1991 risk on Vancouver Island, Canada, using stand level information. *Forest Ecology Manag.*,
1992 154, 117-130, 2001.
- 1993 ○ Mitchell-Wallace, K., Jones, M., Hillier, J. and Foote, M.: Natural catastrophe risk
1994 management and modelling: A practitioner's guide. John Wiley & Sons, 2017.
- 1995 ○ Moemken, J., Reyers, M., Feldmann, H., and Pinto, J. G.: Future Changes of Wind Speed
1996 and Wind Energy Potentials in EURO-CORDEX Ensemble Simulations, *J. Geophys. Res.*
1997 123, 6373–6389, <https://doi.org/10.1029/2018JD028473>, 2018.

- 1998 ○ Mohammadi, M., Finnan, J., Sterling, M. and Baker, C.: A calibrated oat lodging model
1999 compared with agronomic measurements, *F. Crop. Res.*, 255(June), 107784,
2000 doi:10.1016/j.fcr.2020.107784, 2020.
- 2001 ○ Morimoto, J., Nakagawa, K., Takano, K., Aiba, M., Oguro, M., Furukawa, Y., Mishima, Y.,
2002 Ogawa, K., Ito, R., Takemi, T., Nakamura, F., and Peterson, C.: Comparison of vulnerability
2003 to catastrophic wind between Abies plantation forests and natural mixed forests in northern
2004 Japan. *Forestry*. 92, 436-443, <https://doi.org/10.1093/forestry/cpy045>, 2019.
- 2005 ○ Münchener Rückversicherungs-Gesellschaft: Munich Re: Winterstürme in Europa (II) –
2006 Schadenanalyse 1999 – Schadenspotentiale, 1999.
- 2007 ○ Murshed, S. and Reed, E.: Mapping of the vulnerability of forest resources due to extreme
2008 winter storms in the state of Baden-Württemberg in Germany. *AutoCarto*, 128-140, 2016.
- 2009 ○ Negrón-Juárez, R., Chambers, J., Guimaraes, G., Zeng, H., Raupp, C., Marra, D. M.,
2010 Ribeiro, G., Saatchi, S., Nelson, B. and Higuchi, N.: Widespread Amazon forest tree
2011 mortality from a single cross-basin squall line event, *Geophys. Res. Lett.*, 37, 1–5,
2012 doi:10.1029/2010GL043733, 2010.
- 2013 ○ [Neal, R.A., Boyle, P., Grahame, N., Mylne, K. & Sharpe, M. \(2014\) Ensemble based first
2014 guess support towards a risk-based severe weather warning service: ensemble severe
2015 weather forecasts. *Meteorological Applications*, 21, 563–577.
2016 <https://doi.org/10.1002/met.1377>](https://doi.org/10.1002/met.1377)
- 2017 ○ Neu, U., Akperov, M., Bellenbaum, N., Benestad, R., Blender, R., Caballero, R., Coccozza,
2018 A., Dacre, H., Feng, Y., Fraedrich, K., Grieger, J., Gulev, S., Hanley, J., Hewson, T., Inatsu,
2019 M., Keay, K., Kew, S., Kindem, I., Leckebusch, G., Liberato, M., Lionello, P., Mokhov, I.,
2020 Pinto, J. G., Raible, C., Reale, M., Rudeva, I., Schuster, M., Simmonds, I., Sinclair, M.,
2021 Sprenger, M., Tilinina, N., Trigo, I., Ulbrich, S., Ulbrich, U., Wang, X. and Wernli, H.:
2022 IMILAST: a community effort to intercompare extratropical cyclone detection and tracking
2023 algorithms. *Bull. Amer. Meteor. Soc.*, 94, 529–547, 2013.
- 2024 ○ Nicoll, B. and Ray, D.: Adaptive growth of tree root systems in response to wind action and
2025 site conditions. *Tree Physiology* 16, 891–898, [https://doi.org/10.1093/treephys/16.11-
2026 12.891](https://doi.org/10.1093/treephys/16.11-12.891), 1996.
- 2027 ~~○ Nicoll, B., Gardiner, B., Rayner, B. and Peace, A.: Anchorage of coniferous trees in relation
2028 to species, soil type, and rooting depth, *Can. J. For. Res.*, 36, 1871–1883, doi:10.1139/x06-
2029 072, 2006.~~
- 2030 ○ ~~Nicoll, B.~~ Connolly, T. and Gardiner, B. : Changes in Spruce Growth and Biomass Allocation
2031 Following Thinning and Guying Treatments. *Forests*, 10, 253, [https://www.mdpi.com/1999-
2032 4907/10/3/253](https://www.mdpi.com/1999-4907/10/3/253), 2019.
- 2033 ○ [Nicoll, B., Gardiner, B., Rayner, B. and Peace, A.: Anchorage of coniferous trees in relation
2034 to species, soil type, and rooting depth, *Can. J. For. Res.*, 36, 1871–1883, doi:10.1139/x06-
2035 072, 2006.](https://doi.org/10.1139/x06-072)
- 2036 ○ Niermann, D., Borsche, M., Kaiser-Weiss, A. and Kaspar, F.: Evaluating renewable-energy-
2037 relevant parameters of COSMO-REA6 by comparison with satellite data, station
2038 observations and other reanalyses. *Meteorol. Z.*, 28, 347-360,
2039 <https://doi.org/10.1127/metz/2019/0945><https://doi.org/10.1127/metz/2019/0945>, 2019.
- 2040 ○ Nissen K., Leckebusch G., Pinto J. G., Renggli D., Ulbrich S., and Ulbrich U.: Cyclones
2041 causing windstorms in the Mediterranean: characteristics, trends and links to large-scale
2042 patterns. *Nat. Haz. Earth Sys. Sci.*, 10, 1379–1391. doi:10.5194/nhess-10-1379-2010,
2043 2010.

- 2044 ○ Niu, L., Feng, S., Ding, W. and Li, G.: Influence of Speed and Rainfall on Large-Scale Wheat
 2045 Lodging from 2007 to 2014 in China. *PloS one*, 11, e0157677. DOI:
 2046 10.1371/journal.pone.0157677, 2016.
- 2047 ○ Nosnikau, V., Gardiner, B., Donis, J. and Fedarovich, L.: The Overview of Existing
 2048 Silvicultural Approaches Under the Influence of Adverse Weather Phenomena and
 2049 Recommendations for Increasing the Forest Resilience to High Wind Events in Belarus,
 2050 World Bank: GFDRR, Minsk., 2018.
- 2051 ○ Oke, D., Mills, D., Christen, D., and Voogt, J.: *Urban Climates*, Cambridge University Press,
 2052 Cambridge, 582 pp., 2017.
- 2053 ○ Oliver, H. and Mayhead, G.: Wind Measurements in a Pine Forest During a Destructive
 2054 Gale. *Forestry*. 47, 185–194, <https://doi.org/10.1093/forestry/47.2.185>, 1974.
- 2055 ○ Panofsky, H. and Dutton, J.: *Atmospheric Turbulence*, Wiley, New York., 1984.
- 2056 ○ Pardowitz, T., Osinski, R., Kruschke, T., and Ulbrich, U.: An analysis of uncertainties and
 2057 skill in forecasts of winter storm losses. *Nat. Hazards Earth Syst. Sci.*, 16, 2391–2402.
 2058 ~~<https://doi.org/10.5194/nhess-16-2391-2016>~~<https://doi.org/10.5194/nhess-16-2391-2016>,
 2059 2016.
- 2060 ○ Parker, D.: Secondary frontal waves in the North Atlantic region: a dynamical perspective
 2061 of current ideas. *Quart. J. Roy. Meteor. Soc.*, 124, 829–856, 1998.
- 2062 ○ Parodi, A., Lagasio, M., Maugeri, M., Turato, B. and Gallus, W.: Observational and
 2063 modelling study of a major downburst event in Liguria: The 14 October 2016 case,
 2064 *Atmosphere*, 10, 1–17, doi:10.3390/ATMOS10120788, 2019.
- 2065 ○ Pasztor, F., Matulla, C., Zuvela-Aloise, M., Rammer, W., and Lexer, M.: Developing
 2066 predictive models of wind damage in Austrian forests. *Ann. Forest Sci.*, 72, 289–301,
 2067 <https://doi.org/10.1007/s13595-014-0386-0>, 2015.
- 2068 ○ Patton, E. G. and Finnigan, J. J.: Canopy Turbulence, *Handb. Environ. Fluid Dyn. Vol. One*,
 2069 311–328, doi:10.1201/b14241-28, 2012.
- 2070 ○ Pejovic, T., Williams, V. A., Noland, R. B., and Toumi, R.: Factors Affecting the Frequency
 2071 and Severity of Airport Weather Delays and the Implications of Climate Change for Future
 2072 Delays, *Transp. Res. Rec.*, 2139(1), 97–106, <https://doi.org/10.3141/2139-12>, 2009.
- 2073 ○ Peltola, H.: Swaying of trees in response to wind and thinning in a stand of Scots pine,
 2074 *Bound.-Layer Meteorol*, 77, 285–304, <https://doi.org/10.1007/BF00123529>, 1996.
- 2075 ○ ~~Peltola, H.: [Mechanical stability of trees under static loads, *Am. J. Bot.*, 93, 1501–1511,](https://doi.org/10.3732/ajb.93.10.1501)~~
 2076 ~~<https://doi.org/10.3732/ajb.93.10.1501>, 2006.~~
- 2077 ○ [Peltola, H.](https://doi.org/10.1139/x99-029), Kellomäki, S., Väisänen, H., and Ikonen, V.-P.: A mechanistic model for
 2078 assessing the risk of wind and snow damage to single trees and stands of Scots pine,
 2079 Norway spruce, and birch, *Can. J. For. Res.* 29: 647–661, <https://doi.org/10.1139/x99-029>,
 2080 1999.
- 2081 ○ ~~[Peltola, H.: Mechanical stability of trees under static loads, *Am. J. Bot.*, 93, 1501–1511,](https://doi.org/10.3732/ajb.93.10.1501)~~
 2082 ~~<https://doi.org/10.3732/ajb.93.10.1501>, 2006.~~
- 2083 ○ Peña-Angulo, D.; Reig-Gracia, F.; Domínguez-Castro, F.; Revuelto, J.; Aguilar, E.; Schrier,
 2084 G.; Vicente-Serrano, S. M.: ECTACI: European Climatology and Trend Atlas of Climate
 2085 Indices (1979–2017). *J. Geophys. Res. Atmos.*, 125, 9. DOI: 10.1029/2020JD032798, 2020.
- 2086 ○ Peterson, C.: Catastrophic wind damage to North American forests and the potential impact
 2087 of climate change, *Sci. Total Environ.*, 262, 287–311, doi:10.1016/S0048-9697(00)00529-
 2088 5, 2000.

- 2089 ○ Peterson, C., Ribeiro, G., Negrón-Juárez, R., Marra, D., Chambers, J., Higuchi, N., Lima,
2090 A. and Cannon, J.: Critical wind ~~speeds~~speed suggest wind could be an important
2091 disturbance agent in Amazonian forests. *Forestry*. 92, 444-459, 2019.
- 2092 ○ Petrović, V. and Bottasso, C.: Wind turbine optimal control during storms, *J. Phys.: Conf.*
2093 *Ser.*, 524, 012052, <https://doi.org/10.1088/1742-6596/524/1/012052>, 2014.
- 2094 ~~○ Pinto, J. G., Spanghel T., Ulbrich U., and Speth P.: Sensitivities of a cyclone detection
2095 and tracking algorithm: individual tracks and climatology. *Meteorol. Zeitschrift*, 14, 823–838.
2096 [doi:10.1127/0941-2948/2005/0068](https://doi.org/10.1127/0941-2948/2005/0068), 2005~~
- 2097 ○ Pinto, J., Fröhlich, E., Leckebusch, G. and Ulbrich, U.: Changing European storm loss
2098 potentials under modified climate conditions according to ensemble simulations of the
2099 ECHAM5/MPI-OM1 GCM. *Nat. Haz. Earth Sys. Sci.*, 7, 165–175.
2100 <https://doi.org/10.5194/nhess-7-165-2007>, 2007.
- 2101 ○ Pinto, J. G., Gómara, I., Masato, G., Dacre, H., Woollings, T. and Caballero, R.: Large-scale
2102 dynamics associated with clustering of extratropical cyclones affecting Western Europe. *J.*
2103 *Geophys. Res.*, 119, 13704–13719, <https://doi.org/10.1002/2014JD022305>, 2014
- 2104 ○ Pinto, J. G., Karremann, M., Born, K., Della-Marta, P., and Klawa, M.: Loss potentials
2105 associated with European windstorms under future climate conditions. *Climate Res.*, 54, 1–
2106 20. <https://doi.org/10.3354/cr01111>, 2012
- 2107 ○ Pinto J. G., Spanghel T., Ulbrich U., and Speth P.: Sensitivities of a cyclone detection and
2108 tracking algorithm: individual tracks and climatology. *Meteorol. Zeitschrift*, 14, 823–838.
2109 [doi:10.1127/0941-2948/2005/0068](https://doi.org/10.1127/0941-2948/2005/0068), 2005
- 2110 ~~○, Karremann, M., Born, K., Della Marta, P., and Klawa, M.: Loss potentials associated with
2111 European windstorms under future climate conditions. *Climate Res.*, 54, 1–20.
2112 <https://doi.org/10.3354/cr01111>, 2012~~
- 2113 ~~○ Pinto, J., Gómara, I., Masato, G., Dacre, H., Woollings, T. and Caballero, R.: Large scale
2114 dynamics associated with clustering of extratropical cyclones affecting Western Europe. *J.*
2115 *Geophys. Res.*, 119, 13704–13719, <https://doi.org/10.1002/2014JD022305>, 2014~~
- 2116 ○ ~~Prahl, B., Rybski, D., Kropp, J., Burghoff, O. and Held, H.: Applying stochastic small scale
2117 damage functions to German winter storms: Damage functions for German storm loss.
2118 *Geophys. Res. Lett.*, 39, 6. <https://doi.org/10.1029/2012GL050961>, 2012.~~
- 2119 ○ Prahl, B., Rybski, D., Burghoff, O. and Kropp, J.: Comparison of storm damage functions
2120 and their performance. *Nat. Haz. Earth Sys. Sci.*, 15, 769–788.
2121 <https://doi.org/10.5194/nhess-15-769-2015>, 2015.
- 2122 ○ Prahl, B., Rybski, D., Kropp, J., Burghoff, O. and Held, H.: Applying stochastic small-scale
2123 damage functions to German winter storms: Damage functions for German storm loss.
2124 *Geophys. Res. Lett.*, 39, 6. <https://doi.org/10.1029/2012GL050961>, 2012.
- 2125 ○ Priestley, M., Ackerley, D., Catto, J., Hodges, K., McDonald, R., and Lee, R.: An Overview
2126 of the Extratropical Storm Tracks in CMIP6 Historical Simulations, *J. Climate*, 33, 6315–
2127 6343, 2020b.
- 2128 ○ Priestley, M., Dacre, H., Shaffrey, L., Schemm, S, and Pinto, J. G.: The role of secondary
2129 cyclones and cyclone families for the North Atlantic storm track and clustering over western
2130 Europe. *Q J R Meteorol Soc.*, 146, 1184–1205. <https://doi.org/10.1002/qj.3733>, 2020a.
- 2131 ~~○ Priestley, M., Ackerley, D., Catto, J., Hodges, K., McDonald, R., and Lee, R.: An Overview
2132 of the Extratropical Storm Tracks in CMIP6 Historical Simulations, *J. Climate*, 33, 6315–
2133 6343, 2020b.~~
- 2134 ○ Primo, C.: Wind gust warning verification, *Adv. Sci. Res.*, 13, 113-120,
2135 <https://doi.org/10.5194/asr-13-113-2016>, 2016.

- 2136 ○ Pryor, S. and Barthelmie, R. : Climate change impacts on wind energy: A review, *Renewable*
2137 *Sus. Energy Rev.*, 14, 430–437, <https://doi.org/10.1016/j.rser.2009.07.028>, 2010.
- 2138 ○ Quaschnig, V.: *Understanding renewable energy systems.* Routledge,
2139 <https://doi.org/10.4324/9781315769431>, 2016.
- 2140 ○ Quayle, R., and Steadman, R.: The Steadman Wind Chill: An Improvement over Present
2141 Scales, *Wea. Forecasting*, 13, 4, 1187-1193, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0434(1998)013<1187:TSWCAI>2.0.CO;2)
2142 [0434\(1998\)013<1187:TSWCAI>2.0.CO;2](https://doi.org/10.1175/1520-0434(1998)013<1187:TSWCAI>2.0.CO;2), 1998.
- 2143 ○ Queck, R., Bernhofer, C., Bienert, A., and Schlegel, F.: *The TurbEFA Field Experiment—*
2144 *Measuring the Influence of a Forest Clearing on the Turbulent Wind Field, Bound.-Layer*
2145 *Meteor.*, 160, 397–423, <https://doi.org/10.1007/s10546-016-0151-z>, 2016.
- 2146 ○ Queck, R., Bienert, A., Maas, H.-G., Harmansa, S., Goldberg, V., and Bernhofer, C.: Wind
2147 fields in heterogeneous conifer canopies: parameterisation of momentum absorption using
2148 high-resolution 3D vegetation scans, *Eur. J. Forest Res.*, 131, 165–176,
2149 <https://doi.org/10.1007/s10342-011-0550-0>, 2012.
- 2150 ~~○ Queck, R., Bernhofer, C., Bienert, A., and Schlegel, F.: *The TurbEFA Field Experiment—*~~
2151 ~~*Measuring the Influence of a Forest Clearing on the Turbulent Wind Field, Bound. Layer*~~
2152 ~~*Meteor.*, 160, 397–423, <https://doi.org/10.1007/s10546-016-0151-z>, 2016.~~
- 2153 ○ Qin, L., Liu, Q., Zhang, M., and Saeed, S.: Effect of measurement errors on the estimation
2154 of tree biomass, *Can. J. For. Res.*, 49, 1371–1378, <https://doi.org/10.1139/cjfr-2019-0034>,
2155 2019.
- 2156 ○ Quine, C., Gardiner, B., and Moore, J., 2020: Wind disturbance in forests: The process of
2157 wind created gaps, tree overturning, and stem breakage. *Plant Disturbance Ecology*.
2158 Elsevier, New York, pp. 117–184. <https://doi.org/10.1016/b978-0-12-818813-2.00004-6>
- 2159 ○ Quine, C. and White, I.: Revised windiness scores for the windthrow hazard classification:
2160 the revised scoring method, Research Division of the Forestry Authority., 1993.
- 2161 ○ Quine, C. and White, I.: The potential of distance-limited topex in the prediction of site
2162 windiness, *Forestry*, 71, 325–332, 1998.
- 2163 ○ Radinovic, D.: Mediterranean cyclones and their influence on the weather and climate.
2164 PSMP Rep Series 24, WMO, Geneva, Switzerland, 131 pp, 1987
- 2165 ○ Raupach, M., Finnigan, J., Brunet, Y., 1996. Coherent eddies and turbulence in vegetation
2166 canopies: the mixing-layer analogy. *Bound.-Layer Meteorol.* 78, 351–382.
- 2167 ○ Rajkumara, S.: Lodging in cereals—a review. *Agricultural Rev.*, 29, pp.55-60, 2008.
- 2168 ○ Ritter, M., Shen, Z., López Cabrera, B., Odening, M., and Deckert, L.: Designing an index
2169 for assessing wind energy potential, *Renewable Energy*, 83, 416–424,
2170 <https://doi.org/10.1016/j.renene.2015.04.038>, 2015.
- 2171 ○ Roberts, J. F., Champion, A. J., Dawkins, L. C., Hodges, K. I., Shaffrey, L. C., Stephenson,
2172 *D. B., Stringer, M. A., Thornton, H. E., and Youngman, B. D.: The XWS open access*
2173 *catalogue of extreme European windstorms from 1979 to 2012, Nat. Hazards Earth Syst.*
2174 *Sci.*, 14, 2487–2501, <https://doi.org/10.5194/nhess-14-2487-2014>, 2014.
- 2175 ○ Romanic, D. and Hangan, H.: Experimental investigation of the interaction between near-
2176 surface atmospheric boundary layer winds and downburst outflows, *J. Wind Eng. Ind.*
2177 *Aerodynamics*, 205, 104323, <https://doi.org/10.1016/j.jweia.2020.104323>, 2020.
- 2178 ○ Rouse, R. and Hodges, L.: Windbreaks. *Agronomy and Horticulture Department (Agronomy*
2179 *and Horticulture - Faculty Publications*, 427), 2004.
- 2180 ○ Rudnicki, M., Mitchell, S., and Novak, M.: Wind tunnel measurements of crown streamlining
2181 and drag relationships for three conifer species, *Can. J. For. Res.*, 34, 666–676,
2182 <https://doi.org/10.1139/x03-233>, 2004.

- 2183 ○ Ruel, J., Pin, D., Spacek, L., Cooper, K., and Benoit, R.: The estimation of wind exposure
2184 for windthrow hazard rating: comparison between Strongblow, MC2, Topex and a wind
2185 tunnel study. *Forestry*. 70, 253–266, <https://doi.org/10.1093/forestry/70.3.253>, 1997.
- 2186 ○ Sanders F, and Gyakum J.: Synoptic-dynamic climatology of the ``bomb. *Mon Wea Rev.*
2187 108:1589–606. 1980.
- 2188 ○ Schindler, D., Grebhan, K., Albrecht, A., Schönborn, J., and Kohnle, U.: GIS-based
2189 estimation of the winter storm damage probability in forests: A case study from Baden-
2190 Wuerttemberg (Southwest Germany). *Int. J. Biometeor.*, 56, 57–69,
2191 <https://doi.org/10.1007/s00484-010-0397-y>, 2012.
- 2192 ○ Schindler, D. and Kolbe, S.: Assessment of the Response of a Scots Pine Tree to Effective
2193 Wind Loading. *Forests*. 11, 145. <https://doi.org/10.3390/f11020145>, 2020.
- 2194 ○ Schindler, D. and Mohr, M.: No resonant response of Scots pine trees to wind excitation.
2195 *Agricultural Forest Meteorol.*, 265, 227–244, <https://doi.org/10.1016/j.agrformet.2018.11.021>,
2196 2019.
- 2197 ○ ~~Schindler, D. and Kolbe, S.: Assessment of the Response of a Scots Pine Tree to Effective~~
2198 ~~Wind Loading. *Forests*. 11, 145. <https://doi.org/10.3390/f11020145>, 2020.~~
- 2199 ○ Schmidt, P.: *Lastannahmen - Einwirkungen auf Tragwerke: Grundlagen und Anwendung*
2200 nach EC 1. Springer Vieweg, 2019.
- 2201 ○ Schultz, D., Bosart, L., Colle, B., Davies, H., Dearden, C., Keyser, D., Martius, O., Roebber,
2202 P., Steenburgh, W., Volkert, H. and Winters, A.: Extratropical cyclones: a century of
2203 research on meteorology’s centerpiece. *Meteor. Monographs*, 59, 16-1, 2019.
- 2204 ○ Schütz, J., Götz, M., Schmid, W., and Mandallaz, D.: Vulnerability of spruce (*Picea abies*)
2205 and beech (*Fagus sylvatica*) forest stands to storms and consequences for silviculture. *Eur.*
2206 *J. Forest Res.*, 125, 291-302, <https://doi.org/10.1007/s10342-006-0111-0>, 2006.
- 2207 ○ Schweighofer, J.: The impact of extreme weather and climate change on inland waterway
2208 transport, *Nat. Hazard.*, 72, 23-40, <https://doi.org/10.1007/s11069-012-0541-6>, 2014.
- 2209 ○ Schwierz, C., Köllner-Heck, P., Zenklusen Mutter, E., Bresch, D., Vidale, P.-L., Wild, M.,
2210 and Schär, C.: Modelling European winter windstorm losses in current and future climate.
2211 *Clim. Change*, 101, 485–514. <https://doi.org/10.1007/s10584-009-9712-1>
2212 <https://doi.org/10.1007/s10584-009-9712-1>, 2010.
- 2213 ○ Scott, R. and Mitchell, S.: Empirical modelling of windthrow risk in partially harvested stands
2214 using tree, neighbourhood, and stand attributes. *Forest Ecology Manag.*, 218, 193-209,
2215 <https://doi.org/10.1016/j.foreco.2005.07.012>, 2005.
- 2216 ○ Seneviratne, S.I., X. Zhang, M. Adnan, W. Badi, C. Dereczynski, A. Di Luca, S. Ghosh, I.
2217 Iskandar, J. Kossin, S. Lewis, F. Otto, I. Pinto, M. Satoh, S.M. Vicente-Serrano, M. Wehner,
2218 and B. Zhou, 2021: Weather and Climate Extreme Events in a Changing Climate. In *Climate*
2219 *Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth*
2220 *Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte,*
2221 *V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I.*
2222 *Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O.*
2223 *Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United*
2224 *Kingdom and New York, NY, USA, pp. 1513–1766, doi:10.1017/9781009157896.013.*
- 2225 ○ Shahabinejad, N., Mahmoodabadi, M., Jalalian, A., and Chavoshi, E.: In situ field
2226 measurement of wind erosion and threshold velocity in relation to soil properties in arid and
2227 semiarid environments. *Environ. Earth Sci.*, 78, 710. DOI: 10.1007/s12665-019-8508-5,
2228 2019.

- 2229 ○ Shapiro, M. and Keyser, D.: Fronts, jet-streams and the tropopause. In Extratropical
 2230 cyclones (pp. 167-191). American Meteorological Society, Boston, MA, 1990.
- 2231 ○ Sharples, J.: Review of formal methodologies for wind-slope correction of wildfire rate of
 2232 spread, *Int. J. Wildl. Fire*, 17, 179–193, 2008.
- 2233 ○ Sissakian, V., Al-Ansari, N. and Knutsson, S.: Sand and dust storm events in Iraq. *J. Nat.*
 2234 *Sci.*, 5, 1084-1094, 2013.
- 2235 ○ Sivakumar, M.: Impacts of sand storms/dust storms on agriculture. In *Natural Disasters and*
 2236 *Extreme Events in Agriculture*. Berlin, Heidelberg: Springer, pp. 159–177, 2005.
- 2237 ○ [SMHI warning levels: https://www.smhi.se/en/weather/sweden-weather/warnings](https://www.smhi.se/en/weather/sweden-weather/warnings), [last](https://doi.org/10.01.2023)
 2238 [access: 10.01.2023](https://doi.org/10.01.2023)
- 2239 ○ Snæbjörnsson, J., Baker, C., and Sigbjörnsson, R.: Probabilistic assessment of road vehicle
 2240 safety in windy environments, *J. Wind Eng. Ind. Aerodyn.*, 95, 1445-1462.,
 2241 <http://doi.org/10.1016/j.jweia.2007.02.020>, 2007.
- 2242 ○ Spatz, H.-C. and Theckes, B.: Oscillation damping in trees. *Plant Science*, 207, 66–71,
 2243 <https://doi.org/10.1016/j.plantsci.2013.02.015>, 2013.
- 2244 ○ Sprenger, M., Schmidli, J., and Egloff, L.: The Laseyer wind storm-case studies and a
 2245 climatology, *Meteorol. Z.*, 27(1), 15-32, <https://doi.org/10.3929/ethz-b-000239023>, 2018.
- 2246 ○ Stefanski, R.; Sivakumar, M.: Impacts of sand and dust storms on agriculture and potential
 2247 agricultural applications of a SDSWS. In *IOP Conf. Ser.: Earth Environ. Sci.* 7, p. 12016.
 2248 DOI: 10.1088/1755-1307/7/1/012016, 2009.
- 2249 ○ de Souza, C., Dias-Júnior, C., Tóta, J. and de Abreu Sá, L.: An empirical-analytical model
 2250 of the vertical wind speed profile above and within an Amazon forest site. *Met. Apps*,
 2251 <https://doi.org/10.1002/met.1543>, 2015.
- 2252 ○ [Stoll, R., Gibbs, J.A., Salesky, S.T., Anderson, W., Calaf, M.: Large-Eddy Simulation of the](https://doi.org/10.1007/s10546-020-00556-3)
 2253 [Atmospheric Boundary Layer. *Boundary-Layer Meteorol* 177:541–581,](https://doi.org/10.1007/s10546-020-00556-3)
 2254 [https://doi.org/10.1007/s10546-020-00556-3, 2020.](https://doi.org/10.1007/s10546-020-00556-3)
- 2255 ○ Stull, R.: An introduction to boundary layer meteorology. Springer,
 2256 <https://link.springer.com/book/10.1007/978-94-009-3027-8>, 1988.
- 2257 ○ Stull, R.: *Practical Meteorology: An Algebra-based Survey of Atmospheric Science*, Black
 2258 and White ed. edition., Sundog Publishing, LLC, Vancouver, 942 pp., 2017.
- 2259 ○ Suárez, J., Gardiner, B. and Quine, C.: A comparison of three methods for predicting wind
 2260 [speeds](#) [speed](#) in complex forested terrain, *Meteorol. Appl.*, 6, 3290342, 1999.
- 2261 ○ Sulik, S. and Kejna, M.: The origin and course of severe thunderstorm outbreaks in Poland
 2262 on 10 and 11 August, 2017, *Bull. Geogr. Phys. Geogr. Ser.*, 18, 25–39, doi:10.2478/bgeo-
 2263 2020-0003, 2020.
- 2264 ○ ~~Sun, Q., Miao, C., and Duan, Q.: Extreme climate events and agricultural climate indices in China:~~
 2265 ~~CMIP5 model evaluation and projections. *Int. J. Climatol.*, 36, 43–61. DOI: 10.1002/joc.4328, 2016.~~
- 2266 ○ Suvanto, S., Henttonen, H., Nöjd, P., and Mäkinen, H.: Forest susceptibility to storm
 2267 damage is affected by similar factors regardless of storm type: Comparison of thunder
 2268 storms and autumn extra-tropical cyclones in Finland. *Forest Ecology Manag.*, 381, 17–28,
 2269 <https://doi.org/10.1016/j.foreco.2016.09.005>, 2016.
- 2270 ○ Suvanto, S., Henttonen, H., Nöjd, P., and Mäkinen, H.: High-resolution topographical
 2271 information improves tree-level storm damage models. *Can. J. Forest Res.*, 48, 721–
 2272 728z, <https://doi.org/10.1139/cjfr-2017-0315>, 2018.
- 2273 ○ Takano, K., Nakagawa, K., Aiba, M., Oguro, M., Morimoto, J., Furukawa, Y., Mishima, Y.,
 2274 Ogawa, K., Ito, R., and Takemi, T.: Projection of impacts of climate change on windthrows
 2275 and evaluation of potential adaptation measures in forest management: A case study from

- 2276 empirical modelling of windthrows in Hokkaido, Japan, by Typhoon Songda (2004). *Hydrol.*
 2277 *Res. Lett.*, 10, 132–138, <https://doi.org/10.3178/hrl.10.132>, 2016.
- 2278 ○ [Takemi, T., Yoshida, T., Horiguchi, M., Vanderbauwhede, W.: Large-Eddy-simulation](#)
 2279 [analysis of airflows and strong wind hazards in urban areas. *Urban Climate* 32:100625,](#)
 2280 <https://doi.org/10.1016/j.uclim.2020.100625>, 2020.
- 2281 ○ Tamura, Y.: Wind-induced damage to buildings and disaster risk reduction. 23, 2009.
- 2282 ○ Taszarek, M., Pilguy, N., Orlikowski, J., Surowiecki, A., Walczakiewicz, S., Pilorz, W.,
 2283 Piasecki, K., Pajurek, L. and Pórolniczak, M.: Derecho evolving from a Mesocyclone-A
 2284 Study of 11 August 2017 severe weather outbreak in Poland: Event analysis and high-
 2285 resolution simulation, *Mon. Weather Rev.*, 147, 2283–2306, doi:10.1175/MWR-D-18-
 2286 0330.1, 2019.
- 2287 ○ [Taszarek, M., Allen, J.T., Brooks, H.E., Pilguy, N. and Czernecki, B.: Differing trends in](#)
 2288 [United States and European severe thunderstorm environments in a warming climate.](#)
 2289 [*Bulletin of the American Meteorological society*, 102, E296-E322, 2021.](#)
- 2290 ○ Taylor, A., Dracup, E., MacLean, D., Boulanger, Y., and Endicott, S.: Forest structure more
 2291 important than topography in determining windthrow during Hurricane Juan in Canada's
 2292 Acadian Forest. *Forest Ecology Manag.*, 434, 255–263,
 2293 <https://doi.org/10.1016/j.foreco.2018.12.026>, 2019.
- 2294 ○ Telewski, F.: Wind-induced physiological and developmental responses in trees. In Grace,
 2295 J. and Coutts, M.: *Wind and Trees*, Cambridge University Press, 237–263,
 2296 <https://doi.org/10.1017/CBO9780511600425.015>, 1995.
- 2297 ○ Theofilatos, A., and Yannis, G.: A review of the effect of traffic and weather characteristics
 2298 on road safety, *Accid. Anal. Prev.*, 72, 244-256, <https://doi.org/10.1016/j.aap.2014.06.017>,
 2299 2014.
- 2300 ○ Thornes, J., and Davis, B.: Mitigating the impact of weather and climate on railway
 2301 operations in the UK, In *ASME/IEEE Joint Railroad Conference* (pp. 29-38). IEEE,
 2302 <https://doi.org/10.1109/RRCON.2002.1000089>, 2002.
- 2303 ○ Tomas-Burguera, M., Vicente-Serrano, S., Peña-Angulo, D., Domínguez-Castro, F.,
 2304 Noguera, I., and El Kenawy, A.: Global Characterization of the Varying Responses of the
 2305 Standardized Precipitation Evapotranspiration Index to Atmospheric Evaporative Demand.
 2306 *J. Geophys. Res.*, 125, 17. DOI: 10.1029/2020JD033017, 2020.
- 2307 ○ Tomczak, A., Jelonek, T., Grzywiński, W., Mania, P., Tomczak, K. and Pazdrowski, W.: The
 2308 Effects of Wind Exposure on Scots Pine Trees: Within-Stem Variability of Wood Density and
 2309 Mechanical Properties. *Forests* 11, 1095, <https://doi.org/10.3390/f11101095>, 2020.
- 2310 ○ Torshizi, M., Miri, A., Shahriari, A., Dong, Z., and Davidson-Arnott, R.: The effectiveness of
 2311 a multi-row Tamarix windbreak in reducing aeolian erosion and sediment flux, Niatak area,
 2312 Iran. *J. Environ. Manag.*, 265, 110486. DOI: 10.1016/j.jenvman.2020.110486, 2020.
- 2313 ○ Torun, P. and Altunel, A.: Effects of environmental factors and forest management on
 2314 landscape-scale forest storm damage in Turkey. *Ann. Forest Sci.*, 77, 1-13,
 2315 <https://doi.org/10.1007/s13595-020-00945-1>, 2020.
- 2316 ○ Trigo I., Davies T., and Bigg G.: Objective climatology of cyclones in the Mediterranean
 2317 region. *J. Clim.*, 12, 1685–1696, 1999.
- 2318 ○ Troen, I. and Petersen, E.: *European Wind Atlas*, 1st ed., Risø National Laboratory,
 2319 Roskilde., 1989.
- 2320 ○ [Tschurr, F., Feigenwinter, I., Fischer, A., and Kotlarski, S.: Climate Scenarios and Agricultural Indices:](#)
 2321 [A Case Study for Switzerland. *Atmosphere*, 11, 535. DOI: 10.3390/atmos11050535, 2020.](#)

- 2322 ○ TT Club, ICHCA International, Port Equipment Manufacturers Association: Recommended
 2323 minimum safety specifications for quay container cranes, TT Club (in collaboration with
 2324 ICHCA International and PEMA), [https://www.ttclub.com/-/media/files/tt-club/port-](https://www.ttclub.com/-/media/files/tt-club/port-equipment-information-papers/recommended-minimum-safety-features-for-quay-container-cranes.pdf)
 2325 [equipment-information-papers/recommended-minimum-safety-features-for-quay-](https://www.ttclub.com/-/media/files/tt-club/port-equipment-information-papers/recommended-minimum-safety-features-for-quay-container-cranes.pdf)
 2326 [container-cranes.pdf](https://www.ttclub.com/-/media/files/tt-club/port-equipment-information-papers/recommended-minimum-safety-features-for-quay-container-cranes.pdf), last access: 14 December 2021, 2011.
- 2327 ○ ~~Uchida, T. and Ohya, Y.: and Kawashima, Y.: New Assessment Scales for Evaluating the~~
 2328 ~~Degree of Risk of Wind Turbine Blade Damage Caused by Terrain-Induced Turbulence,~~
 2329 ~~Energies, 12, 2624, <https://doi.org/10.3390/en12132624>, 2019.~~
- 2330 ○ ~~Uchida, T. Large eddy simulation of turbulent airflow over complex terrain, J. Wind Eng.~~
 2331 ~~Indus. Aerodynamics, 91, 219–229, [https://doi.org/10.1016/S0167-6105\(02\)00347-1](https://doi.org/10.1016/S0167-6105(02)00347-1), 2003.~~
- 2332 ○ Uchida, T. and Li, G.: Comparison of RANS and LES in the Prediction of Airflow Field over
 2333 Steep Complex Terrain, Open J. Fluid Dyn., 08, 286,
 2334 <https://doi.org/10.4236/ojfd.2018.83018>, 2018.
- 2335 ○ Uchida, T. and Ohya, Y.: [Large-eddy simulation of turbulent airflow over complex terrain, J.](https://doi.org/10.1016/S0167-6105(02)00347-1)
 2336 [Wind Eng. Indus. Aerodynamics, 91, 219–229, \[https://doi.org/10.1016/S0167-\]\(https://doi.org/10.1016/S0167-6105\(02\)00347-1\)](https://doi.org/10.1016/S0167-6105(02)00347-1)
 2337 [6105\(02\)00347-1, 2003.](https://doi.org/10.1016/S0167-6105(02)00347-1)
- 2338 ○ ~~Uchida, T. and Ohya, Y.: Micro-siting technique for wind turbine generators by using large-~~
 2339 ~~eddy simulation, J. Wind Eng. Indus. Aerodyn., 96, 2121–2138,~~
 2340 ~~<https://doi.org/10.1016/j.jweia.2008.02.047>, 2008.~~
- 2341 ○ Uchida, T. and Ohya, Y.: Latest Developments in Numerical Wind Synopsis Prediction
 2342 Using the RIAM-COMPACT® CFD Model—Design Wind Speed Evaluation and Wind Risk
 2343 (Terrain-Induced Turbulence) Diagnostics in Japan, Energies, 4, 458–474,
 2344 <https://doi.org/10.3390/en4030458>, 2011.
- 2345 ○ ~~Uchida, T. and Kawashima, Y.: New Assessment Scales for Evaluating the Degree of Risk~~
 2346 ~~of Wind Turbine Blade Damage Caused by Terrain-Induced Turbulence, Energies, 12,~~
 2347 ~~2624, <https://doi.org/10.3390/en12132624>, 2019.~~
- 2348 ○ ~~Uchida, T.~~ and Sugitani, K.: Numerical and Experimental Study of Topographic Speed-Up
 2349 Effects in Complex Terrain, Energies, 13, 3896, <https://doi.org/10.3390/en13153896>, 2020.
- 2350 ○ Ulbrich U., Leckebusch G., and Pinto J. G.: Extra-tropical cyclones in the present and future
 2351 climate: a review. Theor. Appl. Climatol., 96: 117–131 doi:10.1007/s00704-008-0083-8,
 2352 2009
- 2353 ○ [United Nations Department of Economic and Social Affairs, Population Division. World](https://doi.org/10.1007/s00704-008-0083-8)
 2354 [Population Prospects 2022: Summary of Results. UN DESA/POP/2022/TR/NO. 3. \(2022\)](https://doi.org/10.1007/s00704-008-0083-8)
- 2355 ○ Usbeck, T., Wohlgemuth, T., Pfister, C., Volz, R., Beniston, M., and Dobbertin, M.: Wind
 2356 speed measurements and forest damage in Canton Zurich (Central Europe) from 1891 to
 2357 winter 2007: Wind Speed Measurements and Forest Damage in Canton Zurich, Int. J.
 2358 Climatol., 30, 347–358, <https://doi.org/10.1002/joc.1895>, 2010.
- 2359 ○ Vajda, A., Tuomenvirta, H., Juga, I., Nurmi, P., Jokinen, P., and Rauhala, J.: Severe weather
 2360 affecting European transport systems: the identification, classification and frequencies of
 2361 events, Nat. Hazard., 72, 169-188, <https://doi.org/10.1007/s11069-013-0895-4>, 2014.
- 2362 ○ Van Der Schrier, G., Tank, A., Van Den Besselaar, E., and Swarinoto, Y.: Observed Trends
 2363 and Variability in Climate Indices Relevant for Crop Yields in Southeast Asia. J. Climate 29,
 2364 2651–2669. DOI: 10.1175/JCLI-D-14-00574.1, 2016.
- 2365 ○ van Oldenborgh, G., Krikken, F., Lewis, S., Leach, N., Lehner, F., Saunders, K., van Weele,
 2366 M., Haustein, K., Li, S., Wallom, D., Sparrow, S., Arrighi, J., Singh, R. K., van Aalst, M.,
 2367 Philip, S., Vautard, R., and Otto, F.: Attribution of the Australian bushfire risk to
 2368 anthropogenic climate change, Nat. Hazards Earth Syst. Sci., 21, 941–960,
 2369 <https://doi.org/10.5194/nhess-21-941-2021>, 2021.

- 2370 ○ Van Wagner, C. and Forest, P.: Development and structure of the Canadian forest fire
2371 weather index system. Can. For. Serv., Forestry Tech. Rep., 1987.
- 2372 ~~○ Vitolo, R., Stephenson, D., Cook, I. and Mitchell-Wallace, K.: Serial clustering of intense
2373 European storms. Meteorol. Z., 18, 411-424, 2009.~~
- 2374 ○ Verkaik, J. W.: Evaluation of Two Gustiness Models for Exposure Correction Calculations,
2375 J. Appl. Meteorol., 39, 1613–1626, doi:10.1175/1520-
2376 0450(2000)039<1613:EOTGMF>2.0.CO;2, 2000.
- 2377 ○ Vitolo, R., Stephenson, D., Cook, I. and Mitchell-Wallace, K.: Serial clustering of intense
2378 European storms. Meteorol. Z. 18, 411-424, 2009.
- 2379 ○ Vogel, S.: Drag and Reconfiguration of Broad Leaves in High Winds. J. Exper. Botany, 40,
2380 941–948, 1989.
- 2381 ○ Vollsinger, S., Mitchell, S. J., Byrne, K. E., Novak, M. D. and Rudnicki, M.: Wind tunnel
2382 measurements of crown streamlining and drag relationships for several hardwood species,
2383 Can. J. For. Res., 35(5), 1238–1249, doi:10.1139/x05-051, 2005.
- 2384 ~~○ Wahl, T., Jain, S., Bender, J., Meyers, S., and Luther, M.: Increasing risk of compound
2385 flooding from storm surge and rainfall for major US cities, Nature Clim. Change, 5, 1093–
2386 1097, <https://doi.org/10.1038/nclimate2736>, 2015.~~
- 2387 ○ Wahl, S., Bollmeyer, C., Crewell, S., Figura, C., Friederichs, P., Hense, A., Keller, J. D., and
2388 Ohlwein, C.: A novel convective-scale regional reanalysis COSMO-REA2: Improving the
2389 representation of precipitation. Meteorol. Z., 26, 345-361, [https://doi.org/
2390 10.1127/metz/2017/0824](https://doi.org/10.1127/metz/2017/0824), 2017.
- 2391 ○ Wahl, T., Jain, S., Bender, J., Meyers, S., and Luther, M.: Increasing risk of compound
2392 flooding from storm surge and rainfall for major US cities, Nature Clim. Change, 5, 1093–
2393 1097, <https://doi.org/10.1038/nclimate2736>, 2015.
- 2394 ○ Wang, B., Xu, Y., Zhu, L., and Li, Y.: Crosswind effect studies on road vehicle passing by
2395 bridge tower using computational fluid dynamics, Eng. Appl. Comput. Fluid Mech., 8, 330-
2396 344, <https://doi.org/10.1080/19942060.2014.11015519>, 2014.
- 2397 ○ Wallace, J. and Hobbs, P.: Atmospheric science: an introductory survey (Vol. 92). Elsevier,
2398 2006.
- 2399 ○ Welker, C., Rössli, T., and Bresch, D.: Comparing an insurer’s perspective on building
2400 damages with modelled damages. Nat. Haz. Earth Sys. Sci., 21, 279–299.
2401 <https://doi.org/10.5194/nhess-21-279-2021><https://doi.org/10.5194/nhess-21-279-2021>, 2021.
- 2402 ○ Weninger, T., Scheper, S., Lackóová, L., Kitzler, B., Gartner, K., King, Cornelis, W., Strauss,
2403 P., and Michel, K.: Ecosystem services of tree windbreaks in rural landscapes—a systematic
2404 review. Environ. Res. Lett., <https://doi.org/10.1088/1748-9326/ac1d0d>., 2021.
- 2405 ○ Wiréhn, L., Käyhkö, J., Neset, T.S. and Juhola, S.: 2020. Analysing trade-offs in adaptation
2406 decision-making—agricultural management under climate change in Finland and Sweden.
2407 Reg. Environ. Change, 20, 1-14.
- 2408 ○ Woollings, T., Hannachi, A., and Hoskins, B.: Variability of the North Atlantic eddy-driven
2409 jet-stream, Q. J. R. Meteorol. Soc., 136, 856-868, doi:10.1002/qj.625, 2010.
- 2410 ○ World Shipping Council: Containers Lost At Sea – 2020 Update, Washington, D.C., United
2411 States, <https://www.bifa.org/news/articles/2020/jul/containers-lost-at-sea-2020-update>, last
2412 access: 14 December 2021, 2020.
- 2413 ○ Wurman, J., Kosiba, K., White, T., and Robinson, P.: Supercell tornadoes are much stronger
2414 and wider than damage-based ratings indicate, Proc. Nat. Acad. Sci., 118,
2415 <https://doi.org/10.1073/pnas.2021535118>, 2021.
- 2416 ○ Wyngaard, J.: Turbulence in the Atmosphere, 1st ed., Cambridge University Press, 2010.

- 2417 ○ Young, I.: Seasonal variability of the global ocean wind and wave climate, *Int. J. Climatol*,
 2418 19, 931-950, [https://doi.org/10.1002/\(SICI\)1097-0088\(199907\)19:9<931::AID-JOC412>3.0.CO;2-](https://doi.org/10.1002/(SICI)1097-0088(199907)19:9<931::AID-JOC412>3.0.CO;2-0)
 2419 [https://doi.org/10.1002/\(SICI\)1097-0088\(199907\)19:9<931::AID-JOC412>3.0.CO;2-0](https://doi.org/10.1002/(SICI)1097-0088(199907)19:9<931::AID-JOC412>3.0.CO;2-0),
 2420 1999.
- 2421 ○ Zappa G., Shaffrey L., Hodges K., Sansom P., and Stephenson D.: A multimodel
 2422 assessment of future projections of North Atlantic and European extratropical cyclones in
 2423 the CMIP5 climate models. *J. Clim.*, 26, 5846–5862, 2013.
- 2424 ~~○ Zscheischler, J., Westra, S., van den Hurk, B., Seneviratne, S., Ward, P., Pitman, A.,
 2425 AghaKouchak, A., Bresch, D., Leonard, M., Wahl, T., and Zhang, X.: Future climate risk
 2426 from compound events, *Nature Clim. Change*, 8, 469–477, [https://doi.org/10.1038/s41558-](https://doi.org/10.1038/s41558-018-0156-3)
 2427 [018-0156-3](https://doi.org/10.1038/s41558-018-0156-3), 2018.~~
- 2428 ○ Zscheischler, J., Martius, O., Westra, S., Bevacqua, E., Raymond, C., Horton, R. M., van den
 2429 Hurk, B., AghaKouchak, A., Jézéquel, A., Mahecha, M. D., Maraun, D., Ramos, A., Ridder,
 2430 N., Thiery, W., and Vignotto, E.: A typology of compound weather and climate events, *Nat.*
 2431 *Rev. Earth Environ.*, 1, 333–347, <https://doi.org/10.1038/s43017-020-0060-z>, 2020.
- 2432 ○ Zscheischler, J., Westra, S., van den Hurk, B., Seneviratne, S., Ward, P., Pitman, A.,
 2433 AghaKouchak, A., Bresch, D., Leonard, M., Wahl, T., and Zhang, X.: Future climate risk
 2434 from compound events, *Nature Clim. Change*, 8, 469–477, [https://doi.org/10.1038/s41558-](https://doi.org/10.1038/s41558-018-0156-3)
 2435 [018-0156-3](https://doi.org/10.1038/s41558-018-0156-3), 2018.
- 2436 ○