

设置了格式: 字体: (默认) + 西文标题 (Times New Roman)

# Quantifying unequal urban resilience to rainfalls across China from location-aware big data

Jiale Qian<sup>1,2</sup>, Yunyan Du<sup>1,2,\*</sup>, Jiawei Yi<sup>1,2</sup>, Fuyuan Liang<sup>1</sup>, Nan Wang<sup>1,2</sup>, Ting Ma<sup>1,2</sup>, Tao Pei<sup>1,2</sup>

<sup>1</sup>State Key Laboratory of Resources and Environmental Information System, Institute of Geographical Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China.

<sup>2</sup>University of Chinese Academy of Sciences, Beijing 100049, China.

Correspondence to: Yunyan Du (duyy@lreis.ac.cn)

**Abstract.** The disaster-relevant authorities could make an uninformed decision due to the lack of a clear picture of the urban resilience to adverse natural events. Previous studies seldom examine the near-real-time human dynamics, which are critical

10 to disaster emergency response and mitigation, in response to the development and evolution of mild and frequent rainfall events a natural hazard. In this study, we used the aggregated Tencent location request data (TLR) to examine the variations in collective human activities in response to rainfalls in 346 cities in China. Then two resilience metrics, rainfall threshold and response sensitivity, were introduced to report a comprehensive study of the urban resilience to rainfalls across the mainland China. Our results show that, on average, a 1-mm increase in rainfall intensity is associated with a 0.49% increase of the human 15 activity anomalies. In the cities of northwestern and southeastern China, human activity anomalies are affected more by rainfall intensity and rainfall duration, respectively. Our results highlight the unequal urban resilience to rainfalls across China, showing current heavy rain warning standards underestimate the impacts of heavy rains on the residents in the northwest arid region and the central underdeveloped areas, and overestimate the impacts on the residents in the southeastern coastal area. An overhaul of current heavy rain alert standards therefore is needed to better serve the residents in our study area.

## 20 1 Introduction

Heavy rains with intense precipitation have become more frequent in the context of global climate changes (Myhre et al., 2019; Ogie et al., 2018), and pose significant threats to urban residents, mainly due to the uncoordinated watershed management and undersized infrastructures (Chan et al., 2018; Dewan, 2015; Nahiduzzaman et al., 2015; Song et al., 2019). China is frequently perplexed by urban flooding, particularly in summer when the Asian monsoon brings heavy rains to inland China. It is

25 estimated that 55.15 million people are affected by floods in China in 2017 alone and the direct economic loss is approximately 214 billion Chinese Yuan, which significantly exceeds that of the 2017 typhoon disasters (5.879 million, 34.62 billion). In addition to threatening human daily activities and cities' normal operation (Aerts et al., 2014; Grinberger and Felsenstein, 2016; Kasmalkar et al., 2020; Owrangi et al., 2014), the ever-increasing rainstorms endlessly challenge cities' flood resistance capacity and relevant authorities' real-time decisions in response to such adverse events. Urban decision makers have learned

设置了格式: 字体: (默认) + 西文标题 (Times New Roman), 非突出显示

设置了格式: 字体: (默认) + 西文标题 (Times New Roman)

30 that city management and planning would significantly benefit from a better understanding of urban resilience(Bertilsson et al., 2019; de Bruijn, 2004; O'Sullivan et al., 2012).

Urban resilience refers to the ability of an urban system to prepare for, respond to, and recover from adverse events(Ambelu et al., 2017; Hong et al., 2021a; Liao, 2012; Meerow et al., 2016). Biologists, psychologists, engineers, and geographers have all made their own contributions to the urban resilience studies(Adger et al., 2005; Brusberg and Shively, 2015; Olsson et al.,

35 2015; Ouyang et al., 2012; Poulin and Kane, 2021; Shiferaw et al., 2014). Over the past 20 years, geographers have heavily relied satellite imagery to assess disaster-related resilience as satellites have been providing ever-increasing information about the Earth at a relatively low cost(Mpandeli et al., 2019; Stefan et al., 2016; Tellman et al., 2021). For example, satellite-based emergency mapping systems have been developed to monitor the inundation and recovery processes of the 2005 Switzerland flood(Buehler et al., 2006); assess the damage, restoration, and reconstruction induced by the 2010 Haiti earthquake(Honey et 40 al., 2010); and evaluate the changes of power supply before and after Hurricane Maria in 2017(Román et al., 2019). Emergency rescuers can use high-resolution images to closely monitor ongoing natural disasters and coordinate disaster relief. However, it is almost impossible to extract near real-time human dynamics over the evolution of a disaster from satellite images and such information is very important in disaster mitigation and reduction(Ghaffarian et al., 2018; Liu et al., 2015).

45 Location-aware big data such as the smartphone call records, signaling data, and social media posts, have been widely used to infer real-time human activities(Wang et al., 2020, 2019; Yi et al., 2019; Yue et al., 2017), estimate disaster-induced losses(Kryvasheyev et al., 2016; Liu et al., 2019b), monitor resettlement and restoration(Martín et al., 2020a, 2020b; Wang and Taylor, 2018; Yabe et al., 2020), and study disaster-related resilience(Hong et al., 2021a; Huang and Ling, 2018; Kasmalkar et al., 2020; Zou et al., 2018). Urban residents would adjust their activities when their living environments are 50 socially and physically impacted by an adverse event and such adjustment could be inferred from location-aware big data. In other words, the changes of human social activities extracted from location-aware big data (hereafter referred to as social activities unless specified otherwise) could be used to study the resilience capacity of an urban system in response to an adverse event. For example, Hong et al.,(2021b)(Hong, Bonezak et al. 2021) quantified changes of mobility behaviour before, during, and after the Hurricane Harvey using smartphone geolocation data, and analysed the spatial variable of community resilience capacity which was defined as the function of the magnitude of impact and time-to-recovery.

55 Human social activities may change in response to a disaster and quite a few studies have been conducted to evaluate the urban resilience to destructive disasters(Qiang and Xu, 2020; Takagi et al., 2021; Walch, 2018). For example, significant social and geographic differences were found in the use of social medias across the 126 counties impacted by Hurricane Sandy(Zou et al., 2018) and the 76 counties in Texas and Louisiana affected by Hurricane Harvey(Zou et al., 2019), illustrating how residents prepare for, respond to, and recover from such catastrophic events.(2019)

60 Human social activities may also change in response to mild yet frequent adverse natural events, such as urban floodsrainstorms. Unlike Hurricanes, dwellers are usually not mobilized by relevant authorities to prepare for and resettle after such events. Instead, nearly 90% of flood-related tweets in a city are released during heavy rains(Wang et al., 2020). Consequently, social mediashuman activities mainly show how an urban system respond to but not prepare for and recover from such adverse natural

设置了格式: 字体: (默认) + 西文标题 (Times New Roman)

events (Qian et al., 2022; Zhang et al., 2022). Furthermore, the mild adverse natural events usually affect a large geographic scale and for a much longer time (Qian et al., 2022; Zhang et al., 2022). As a result, urban resilience to mild and frequent adverse events refers to the ability of an urban system to respond to adverse events. Furthermore, urban resilience urban resilience to mild and frequent adverse events could significantly differ from that of destructive disasters and may show significant spatiotemporal variations due to the areal difference in local natural settings, socioeconomic status, and infrastructure completeness (Adger et al., 2005; Guan and Chen, 2014; Östh et al., 2015; Zou et al., 2019, 2018). Study of such regional inequality is therefore of great value to disaster relief and mitigation.

However, previous resilience metric, which mainly focuses on unique disaster event, was not suitable for the assessing in large scale. Two resilience metrics were introduced into this study from other fields. The sensitivity is a widely used tool for understanding resilience in different regions in many other weather events, such as heat wave and air pollution (Hong et al., 2021b; Wang et al., 2021). For example, Zheng et al. (2019) defined the links between the city-level happiness index calculated from the social media data and the daily local air quality metric as the perception sensitivity and explored its spatial variation. However, response sensitivity is not yet to be studied for rainstorm events through analysing the relation between the city-scale human activity response metric and rainstorm event index. Another index, rainfall threshold, is commonly used to study rainfall events that have resulted in landslides (Marra et al., 2016; Naidu et al., 2018). In this study, rainfall threshold, which is defined as the minimum rainfall index that corresponds to significant urban human activity response anomaly, is introduced to the study of the urban resilience. Two metrics can effectively depict the urban resilience in different focuses.

In this study, we proposed a method for measuring and evaluating how urban systems respond to heavy rains as reflected in location-aware big data. We extracted human social-activities from the Tencent location request (TLR) data in 346 cities across mainland China from May to August 2017 and used two indicators - rainfall threshold and response sensitivity – to quantify the urban resilience across our study area. We found significant regional inequality of urban resilience in mainland China and the inequality could be explained by the variations in the regional natural, socioeconomic, and infrastructure variables. The findings from this study provide a new perspective and method to quantify urban resilience to the frequent yet no so destructive adverse events across a large geographic scale. Practically, our findings suggest an urgent need to revise current unified rainstorm warning standards to better serve the residents.

## 2 Data and methods

### 90 2.1 Data

We collected the Tencent location requested (TLR) data from May 1<sup>st</sup> to August 31<sup>st</sup>, 2017 from Tencent's big data portal. Tencent, with over 700 million users, is the largest social media platform in China. A Tencent user may check in the platform for a variety of purposes such as location-based searching, navigation, location sharing and so on. The dataset we downloaded has an hourly temporal resolution and a 1 km x 1 km spatial resolution. The data has been proven as a reliable proxy for

设置了格式: 字体: (默认) + 西文标题 (Times New Roman)  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman)  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman), (中文 Times New Roman) 非突出显示  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman)  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman), 非突出显示  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman)  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman)  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman), 非突出显示  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman)  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman)  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman), 非突出显示  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman)  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman)

95 collective human activities in many studies from multiple dimensions of time and space(Liu et al., 2019b; Ma, 2018; Qian et al., 2021b).

We collected the Version06 Global Precipitation Measurement (GPM) Integrated Multi-satellite Retrievals (IMERG) 30-min precipitation dataset(Levizzani et al., 2020, p. 1). This dataset has a spatial resolution of  $0.1^\circ \times 0.1^\circ$  and has been evaluated and widely used in related studies(Liu et al., 2019a; Yi et al., 2019). We used this dataset to extract the characteristics of rainfall


100 events of interest.

We collected six natural, socioeconomic, or infrastructure indicators to help explain the variations in urban resilience, including the annual precipitation in China since 1980calculated by aggregating GPM data in 2017, population density, gross domestic product, green coverage rate, drainage network density, and per capita area of paved roads. The precipitation data was obtained from <https://www.resde.en/data.aspx?DATAID=229> and the other indicators from the 2017 China City Statistical Yearbook.

## 105 2.2 Methods

Fig. 1 shows the data process and analysis flow chart of this study. We first proposed a Multilevel Human Activity Anomaly Detection (MHAAD) methodological framework to detect and characterize the TLR anomalies in response to rainfall events. The framework has two major parts. In the first part, we identified the grids with a stable TLR number and then the anomalies from the time series TLR of each grid. We then used the two-sided Welch's t-test and probability density function (PDF) 110 method to detect whether human activity anomalies are triggered by a rainfall event or not. Rainfall indices were extracted for the grids with a stable TLR number and selected by their importance as shown in the random forest model. We then explored the multi-level relationship between rainfall characteristics and human activity anomalies. We proposed two indicators -rainfall threshold and response sensitivity- to describe urban resilience. Lastly, we assessed the association between the urban resilience with the natural, socio-economic, and infrastructure explanatory variables.

设置了格式: 字体: (默认) +西文标题 (Times New Roman)






Fig. 1 A flow chart showing the data analysis process in this study.

### 2.2.1 TLR anomaly detection

We first identified all grids with a stable TLR number (hereafter referred to as stable grids unless stated otherwise) using the method Qian et al. (2021a) proposed. The stable grids are the regions with stable human activity and rhythm in urban area.

In total, 832,630 out of 9,600,903 grids across our study area are identified as stable grids. The number of smartphone users and TLR vary significantly by grid. We therefore normalized the TLR using the median interquartile normalization method (Geller et al., 2003) to make the TLR in the stable grids comparable. We then employed the Seasonal Hybrid Extreme Studentized Deviate (S-H-ESD) method (Vallis et al., 2014) to detect anomalies from the gridded TLR time series. The S-H-ESD method could be denoted by the following additive model:

$$T_s = T + S + R$$

(1)

设置了格式: 字体: (默认) + 西文标题 (Times New Roman)  
 设置了格式: 字体: (默认) + 西文标题 (Times New Roman)

设置了格式: 字体: (默认) + 西文标题 (Times New Roman), 非突出显示  
 设置了格式: 字体: (默认) + 西文标题 (Times New Roman), 非突出显示  
 设置了格式: 字体: (默认) + 西文标题 (Times New Roman)  
 设置了格式: 字体: (默认) + 西文标题 (Times New Roman)

where  $T$ ,  $S$ , and  $R$  represent the trend, seasonality, and residual components, respectively.

The S-H-ESD method has two major steps. First, the piecewise median method is used to fit and remove the long-term trend.

Then the Seasonal-Trend decomposition using Loess (STL) method is employed to remove seasonality (Cleveland et al., 1990).

130 We then used the Generalized Extreme Studentized Deviate (G-ESD) statistic (Rosner, 1975) to identify the significant anomalies in the residuals. In this study, we used a piecewise combination of the biweekly medians to model the underlying trend, which shows little changes in the TLR time series. The significance level  $\alpha$  is set to 0.05 and the number of anomalies is set to no more than 25% of the total observations.

We first then extracted the total numbers of the grids with positive (PTLR) and negative anomalies (NTLR) by city, respectively

135 and then examined the variations in the PTLR/NTLR time series over the periods with rains and without rains to identify whether a rainfall event triggers collective human activity anomalies. The two-sided Welch's t-test was used for the significance test. Human activity anomalies usually happen shortly before or after the peak rainfall intensity and last as a spell instead of the entire rainfall event. We therefore employed a moving time window method to find the period with the largest accumulative rainfall and used the period to detect the statistical significance of the change in related to the PTLR/NTLR time 140 series in a raining and non-raining period. In this study, we used a 6-hour moving window, which is half of the average duration of all rainfall events of interest in this study (see Supplementary Fig. 3).

## 2.2.2 Feature selection of rainfall indices

The GPM data were first resampled to the same spatiotemporal resolutions as that of the TLR using the nearest-neighbor 145 interpolation method. We then extracted the hourly rainfall intensity for each city/hour, i.e., the average hourly GPM precipitation within the stable grids of the city.

In this study, a rainfall event is defined as a precipitation process that lasts for at least 3 hours and with no rains preceding for at least one hour. The number of rainfall events in each city is normally distributed. We selected the 346 cities with at least 40 (the top 5% quantile) rainfall events for this study (Supplementary Fig. 1 and 2).

150 Every rainfall event is described with three rainfall indices (the 1-hour peak intensity, 6-hour peak intensity, and cumulative rainfall) and two temporal indices (the duration and peak hour) (Supplementary Table 1). From these indices, we used the random forest model (RF) to calculate the importance score (mean decrease accuracy) for each indicator. The importance score shows the global importance over all the out-of-bag cross validated predictions. The random forest model is robust and less susceptible to multicollinearity as it averages all predictions for a given feature variable and is more efficient in terms of feature 155 selection than the multi-linear regression (Pal, 2005; Strobl et al., 2007). We then identified the most important indicator that triggers human activity anomalies (Liaw and Wiener, 2001).

设置了格式: 字体: (默认) +西文标题 (Times New Roman)

### 2.2.3 Anomaly detection at event scale

We first extracted the total numbers of the grids with positive (PTLR) and negative anomalies (NTLR) by city, respectively and then examined the variations in the PTLR/NTLR time series over the periods with rains and without rains to identify 160 whether a rainfall event triggers collective human activity anomalies. The two-sided Welch's t test was used for the significance test. Human activity anomalies usually happen shortly before or after the peak rainfall intensity and last as a spell instead of the entire rainfall event. We therefore employed a moving time window method to find the period with the largest accumulative rainfall and used the period to detect the statistical significance of the change in related to the PTLR/NTLR time series in a raining and non-raining period. In this study, we used a 6-hour moving window, which is half of the average duration 165 of all rainfall events of interest in this study (see Supplementary Fig. 3).

### 2.2.4.3 Quantifying the rainfall threshold and response sensitivity

In this study, we used two indices, rainfall threshold and response sensitivity, to quantitatively characterize urban resilience. The rainfall threshold is the peak intensity of the rainfall event which triggers collective human activity anomalies. We first 170 used a linear binary classifier to examine the paired values of the peak intensity and duration to determine whether a rainfall event brings more or less rain than the threshold to trigger collective human activity anomalies (Fig. 2a). The Fisher Discriminant Analysis method was then applied to identify the discriminant function to minimize the classification errors (Mika et al., 1999). The rainfall thresholds associated with different rainfall durations are directly extracted from Fig. 2a.




Fig. 2 The schematic diagram (a) and contingency table (b) of the binary classifier for Beijing.

175

Three quantitative indices, the probability of detection (*POD*), false alarm rate (*FAR*), and critical success index (*CSI*) are used to evaluate the performance of the threshold identification method based on the contingency table (Fig. 2b):

$$POD = \frac{H}{H + M}$$

(2)

$$FAR = \frac{FA}{H + FA}$$

$$CSI = \frac{H}{H + M + FA} = \frac{1}{POD^{-1} + (1 - FAR)^{-1} - 1}$$

(3)  
(4)

设置了格式: 字体: (默认) + 西文标题 (Times New Roman)  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman)

180

where H, FA, M represent the percent of hits, false alarms, and misses, respectively. All three indices range from 0 to 1. A value of 1 for the POD and FAR indicates a perfect hit and a 100% false positive rate. A higher CSI is associated with a higher POD and a lower FAR value.

185 The second index, the response sensitivity, is defined as the rate of abnormal change in collective human activities triggered by a rainfall event. We first selected the rainfall events with precipitation above the threshold and that trigger human activity anomalies, i.e., those represented with red crosses and are above the decision boundary in Fig. 2a. Then we defined two indicators cPTLR/cNTLR to describe the rate of abnormal change in collective human activities (Supplementary Fig. 4). The cPTLR/cNTLR were calculated as the difference between the mean of the two time series, i.e., the PTLR/NTLR in the 6-hour raining time window and non-raining time window. A multiple linear regression model was then constructed between the 190 cPTLR and the rainfall intensity/duration for each city. In the end, we calculated the marginal city-specific partial derivatives of cPTLR with respect to the peak intensity and the duration, respectively. The response sensitivity index is calculated as the average of the regression coefficients of the peak intensity and the duration. The adjusted  $R^2$  was also calculated to assess the model accuracy.

195 Finally, we separately classified the rainfall threshold and response sensitivity indices of the 346 cities into three classes using the Jenks natural breaks classification method, which clusters data into different groups by seeking minimum variance within a class and maximum variance between classes (McDougall and Temple-Watts, 2012). In this study, we used the 6-hour rainfall threshold index in line with the time window in Supplementary Fig. 4. In total, there are nine different combinations between the threshold and response sensitivity indices.

## 2.2.5.4 Quantifying the relationships between resilience and urban characteristics

200 We then examined the relationships between the two urban resilience indices and the annual rainfall, population density, gross domestic product, green coverage rate, drainage network density, and per capita area of paved roads. The Kendall, Pearson, and Spearman correlation coefficients and multi-linear regression were used to measure the correlation between rainfall threshold, response sensitivity, and the city characteristics at the city level, respectively.

## 3. Results


### 205 3.1 Collective human activities in response to rainfall events

The gridded TLR could increase (positive anomaly) or decrease (negative anomaly) in response to rains (Supplementary Fig.

5). Counting the overall TLR changes by city would not show how rains impact collective human activities (Yi et al., 2019).

设置了格式: 字体: (默认) + 西文标题 (Times New Roman)  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman)

In this study, instead, we calculated the changes of the total numbers of the grids showing positive (cPTLR) and negative anomalies (cNTLR) by city during a raining period in related to those over the non-raining period, respectively (Fig. 3a) to 210 illustrate how collective human activities change in response to rains.



设置了格式: 字体: (默认) + 西文标题 (Times New Roman)  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman)

Fig. 3 (a) The schematic definitions of the human activity anomaly associated with a specific rainfall event used in this study. (b) The relative proportions and numbers of the rainfall events with various cPTLR and cNTLR values. (c) The probability density distribution curves of the events with significantly increased cPTLR/cNTLR. (d) Box plots by peak rainfall intensity groups per 0.5mm. Trend lines are shown for the OLS regression (black dotted), 0.25 (purple), median (red), 0.75 quantile (blue) range. The numbers in Fig. 3(d) show the numbers of rainfall events with specific peak intensity as shown along the x axis.

The city-level collective human activities jump to an excited state (Fig. 3b) with a significantly increased number of the grids 220 exhibiting positive anomalies, in response to 55.11% of the daytime rains (Supplementary Fig. 64) whereas nighttime rains show no significant impact on the collective human activities. About 93.2% (i.e., 10,710) of the rainfall events in this study are associated with a greater change of the number of the grids with positive anomalies than that of the grids with negative anomalies (i.e., |cPTLR| > |cNTLR|). Around 59.7% of the 10,710 (i.e., 6394) rains show an increased number of the grids with

positive anomaly by city. Furthermore, 35.3% of the 6,394 (i.e., 2257) rainfall events associated with excited-state human activities show a significant increased number of the grids with positive anomalies, which we believed could be attributed to heavy rains. We noticed that a small number (103, 13.19%  $|cPTLR| \leq |cNTLR|$ ) of heavy rains brought by typhoons trigger the city-level collective human activities to a dispirited state, with a significantly increased number of grids exhibiting NTLR (Fig. 3c) as compared to the non-typhoon rains. Accordingly, we excluded all typhoon-related rainfall events from this study. The higher rainfall intensity could trigger more excited-state collective human activities (Fig. 3d). The 1-hour peak intensity values of the rainfall events associated with excited-state human activities are positively correlated with the corresponding cPTLR values (fitting slope = 0.49%, p value <0.001). However, the cPTLR slope against rainfall is affected by the divergence of the peak intensity anomaly. Quantile regression results show that the cPTLR slope coefficient estimates gradually increase from 0.17% for the lower 25% quantile to 0.84% for the higher 75% quantile (p value <0.01). In other words, the cPTLR growth rate generally increases from the less-anomaly to the sensitive-anomaly rainfall events with respect to the increasing magnitude of the rainfall peak intensity.




Fig. 4. (a) The importance of the five rainfall indicators obtained by the random forest model. The peak hour is the most important covariate that triggers human activity anomalies and has the highest decrease accuracy value of 114.5. (b) The peak hour thresholds identified from the differences of the peak hour PDF values between the rainfall events with and without significant collective human activity anomalies. When the peak hours are between 8:59 am and 6:50 pm, the PDF values of the events with human activity anomalies are higher than those without. In other words, daytime rains are more likely to trigger human activity anomalies. There are 11,491 daytime rains (i.e., from 8:59 am to 6:50 pm) out of the total 20,860 rainfall events (ratio: 55.11%).

设置了格式: 字体: (默认) + 西文标题 (Times New Roman)  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman)

设置了格式: 字体: (默认) + 西文标题 (Times New Roman), 小五  
设置了格式: 字体: (默认) + 西文标题 (Times New Roman), 小五, 加粗

设置了格式: 字体: (默认) + 西文标题 (Times New Roman)




Fig. 3 (a) The schematic definitions of the human activity anomaly associated with a specific rainfall event used in this study. (b) The relative proportions and numbers of the rainfall events with various cPTLR and cNTLR values. (c) The probability density distribution curves of the events with significantly increased cPTLR/cNTLR. (d) Box plots by peak rainfall intensity groups per 0.5mm. Trend lines are shown for the OLS regression (black dotted), 0.25 (purple), median (red), 0.75 quantile (blue) range. The numbers in Fig. 3(d) show the numbers of rainfall events with specific peak intensity as shown along the x axis.

### 3.2 Regional inequality of urban resilience

We derived two indicators - rainfall threshold and response sensitivity - from the perspective of public's social response to rains to evaluate the urban resilience. The threshold is defined as the intensity of an event that triggers a city into an undesirable state (Liao, 2012). In this study, we defined the threshold of a rainfall event with a specific duration as the minimum rainfall intensity that corresponds to significant urban TLR anomalies. As shown in Supplementary Fig. 6a4a, the peak rain intensity and rain duration are the second and third important characteristics that trigger human activity anomalies, respectively. We extracted the rainfall thresholds for each city using binary classification models (supplementary Fig. 76) and the results show that the rainfall thresholds drop with increased rainfall duration across all 346 cities in China (Fig. 45 a-c). The 3-, 6-, and 12-hour rainfall thresholds all show significant spatial autocorrelation and a pattern of gradual decrease from the southeast coast


设置了格式: 字体: (默认) +西文标题 (Times New Roman)  
设置了格式: 字体: (默认) +西文标题 (Times New Roman)

设置了格式: 字体: (默认) +西文标题 (Times New Roman)  
设置了格式: 字体: (默认) +西文标题 (Times New Roman)

to the northwest inland. However, with increased rainfall durations, the average rainfall thresholds of all 346 cities decrease from 4.24 to 2.75, whereas their standard deviations decrease from 3.55 to 2.45. Such results indicate the public's response to the short rainfall events varies greatly in different cities and tends to be more consistent with increased rainfall durations.

The impacts of the peak rain intensity and duration on human activity vary across our study area as shown by the slopes of the decision boundary of different cities (Fig. 4d5d). In the arid and semi-arid northwestern China, the slope is close to zero, showing the public response is mainly affected by peak intensity. Residents in the northwestern China may adjust their activities in response to the rain peak intensity as rains in this area seldom last long. By contrast, the slope is high for the southeastern region, indicating the public's response is more affected by rainfall duration. The wet southeastern China usually receives frequent and heavy rains and residents already have adopted to it. Consequently, residents in this area may change their activities in response to rainfall duration more than the peak intensity.


Results of the binary classification are solid as shown by the anomaly detection POD, FAR, and CSI values for different rainfall durations based on rainfall thresholds (Supplementary Fig. 87). More specifically, the POD values for different rainfall durations in the 346 cities range from 0.71 to 1.00, the FAR values from 0 to 0.46, and the CSI values from 0.48 to 1.



设置了格式: 字体: (默认) +西文标题 (Times New Roman)  
设置了格式: 字体: (默认) +西文标题 (Times New Roman)

**Fig. 4-5** Spatial distribution of the rainfall threshold that triggers human activity anomalies for rainfall events lasting three (a), six (b), and 12 (c) hours, respectively and slope (d) of the decision boundary. The isohyets show the 2017 annual rainfall.

The other urban resilience indicator, the response sensitivity, is defined as the rate of the collective human activity anomalies triggered by a rainfall event and was extracted from the multiple regression analysis. The response sensitivity is low in the southeast coast and high in the northwest inland, showing an opposite trend as that of the rainfall threshold (Fig. 5a6a). The higher response sensitivity in the northwest suggests that the residents in this area tend to change their activities more significantly in response to rains. By contrast, activities of the residents in the southeast are not significantly impacted by rains. Such findings are consistent with those derived from the rain threshold indicator. The accuracy of the regression model (the adjusted  $R^2$ ) also shows a similar trend as that of the response sensitivity (Fig. 5b6b), which indicate that the response mode of collective human activities to rainfall in the southeast coastal area is more complex.



**Fig. 5-6** Spatial distribution of the response sensitivity (a) and adjusted  $R^2$  (b) in multiple linear regression. The isohyets show the 2017 annual rainfall.

The associations between the two urban resilience indices also exhibit a significant pattern across our study area (Fig. 67). Cities located in the area with over 1600mm annual precipitation are mainly categorized into type HL (threshold  $> 10.29\text{mm}$  and response sensitivity  $< 0.003$ ), and surrounded by the types ML and HM cities. The cities located in the areas with less than 400mm annual precipitation are mainly classified into LH, MH, and LL types, indicating the annual precipitation has a significant impact on the human activities in different cities.

The LH type cities have a low rainfall threshold ( $< 3.25\text{mm}$ ) and high response sensitivity ( $> 0.025$ ). These cities are mainly found in the northwest fragile region (mainly including the Yili prefecture in Xinjiang province) and the central underdeveloped China (including Shaanxi, Shanxi, and Hebei provinces). Such cities may have underdeveloped infrastructure and weaker rains could trigger human activity anomalies.

设置了格式: 字体: (默认) +西文标题 (Times New Roman)  
设置了格式: 字体: (默认) +西文标题 (Times New Roman)




Fig. 6-7 Spatial distribution of urban resilience. The basemap colors indicate the 2017 annual rainfall.

### 3.3 Associations between the urban resilience with the urban indicators

Results of multiple regression analysis show that variations in urban resilience by city could be explained by the variations in a variety of natural, socioeconomic, and urban infrastructure indicators. Fig. 7a-8a shows the relationship between rainfall thresholds and the explanatory indicators. The 3-hour, 6-hour, and 12-hour rainfall thresholds all show similar correlations with the indicators (Supplementary Table 2, 3, 4), and we only show the correlations between the 6-hour rainfall and the indicators in this section.

All six explanatory variables are significantly correlated with the rainfall threshold ( $p < 0.05$ ). About 42% of the variations in the rainfall threshold could be explained by the variations in the explanatory variables as shown by the  $R^2$  (Supplementary Table 3). Among all explanatory variables, the annual rainfall has the highest coefficient value of 0.43, indicating the variations in the threshold are most affected by the annual rainfall. In other words, residents living in regions with different annual precipitation amount are more likely to accordingly adjust their daily activities once the rainfall is over a specific threshold. Other explanatory variables are also positively correlated with the threshold variable as shown by the positive [correlation regression](#) coefficients ranging from 0.21 to 0.10, except the per capita area of paved road, which is negatively correlated with the threshold. In fact, the per capita area of paved roads is the only indicator showing a negative correlation. Increased per

设置了格式: 字体: (默认) +西文标题 (Times New Roman)

设置了格式: 字体: (默认) +西文标题 (Times New Roman)

capita area of paved road weakens rainwater infiltration capacity and increases surface runoffs, which is more likely to cause traffic congestion and trigger human activity anomalies even the rainfall amount is below a lower threshold.




Fig. 7-8 Regression coefficients between the six explanatory variables and the 6-hour rainfall threshold (a), response sensitivity (b).

320 The horizontal lines mean 95% confidence interval.

Multi-regression analysis shows that the response sensitivity is negatively correlated with all explanatory variables, except for the per capita area of paved roads, which show a positive [regression correlation coefficient](#) (Fig. 7b8b, [Supplementary Table 5](#)). All correlations are statistically significant ( $p < 0.05$ ) ([Supplementary Table 5](#)). About 31% of the variations in the response 325 sensitivity could be explained by the variations of the explanatory variables as shown by an  $R^2$  value of 0.31.

#### 4. Conclusions and discussions

Residents in different cities may adjust and change their activities in different ways in response to [rainfalls](#). Such activity changes and adjustment, also known as urban resilience, could be characterized and studied from location-aware big data. In this study, we used the Tencent aggregated location request data to examine the changes of collective human activities in major 330 cities in China in response to [rainfalls](#) over 2017 summer. Our results show that rainfall time, peak intensity, and duration are the three most important indicators that determine whether a rain would trigger human activity anomalies or not. We also proposed two indices, the rainfall threshold, and response sensitivity, to describe the urban resilience, which show significant spatial variations across our study area. Furthermore, the unequal urban resilience could be explained by a series of explanatory variables.

设置了格式: 字体: (默认) +西文标题 (Times New Roman)

设置了格式: 字体: (默认) +西文标题 (Times New Roman)

335 We believe this paper provides a new perspective to studying urban resilience and that the results bridge the knowledge gap between heavy rains, collective human activities, and urban resilience. Such knowledge is of great significance to urban planning, traffic management, and emergency response.

We also believed this study has three other contributions regarding urban resilience research. First, this study expanded the research framework of urban resilience in response to high-frequency yet mild adverse events and such a framework could be 340 used to study the variations in urban resilience across a large area. Previous studies mainly focused on urban resilience to a specific adverse event such as a typhoon or a hurricane. For example, Zou et al. (2018) used a normalized ratio index to assess the regional variations in urban resilience to Hurricane Sandy. In this study, we examined urban resilience to rains over a relatively long period and across a large area. Such a study would better show how residents in different regions change their activities in different ways in response to rains with different duration, peak intensity, and accumulative rainfall.

345 Secondly, our research analyzed the impacts of different features of the rainfall events on human activities. Previous studies often simply characterized a disaster using its threat levels. For example, Zou et al. (2018) used the average hurricane track kernel density and its wind speed to define the threat levels. In this study, instead, we extracted five major elements of a rainfall event and employed the random forest model to study the impacts of different elements on collective human activities.

Thirdly, we used the rainfall threshold to quantify urban resilience and the thresholds is valuable for the authorities to revise 350 heavy rain alerts. Conventionally, heavy rain alerts are usually based on rainfall intensity and precipitation only and seldom consider the areal difference of infrastructures. According to current Chinese Standard, Chinese authorities would issue a blue, yellow, and orange alert when precipitation is or will be over 50 mm in 12, 6, and 3 hours and if the rain might not stop (Mendiondo, 2005). A red alert would be issued when precipitation is or will be over 100 mm in 3 hours and if the rain might not stop. Results from this study show that it is not appropriate to apply such a unified alert standard to different groups

355 of cities across China. In the resilience fragile areas (Fig. 67), for instance, a rainfall event with 3.25mm precipitation per hour (i.e., 19.5mm in six hours) already triggers significant human activity anomalies. As a result, the national heavy rain alert standard significantly underestimates the impacts of rainfall on the residents in the northwestern and central China.

Tab. 1 shows the precipitation thresholds of different city groups that trigger human activity anomalies and the 6-hour accumulative precipitation based on which a heavy rain warning should be issued. For example, for the LH cities, a rainfall 360 intensity of 1.8 mm and 10.8mm 6-hour accumulative precipitation should trigger a yellow heavy rain warning. Such a value of the 6-hour accumulative precipitation is much lower than current Chinese heavy rain yellow warning standard (50 mm in 6 hour). By contrast, for the HH cities, the rainfall threshold and 6-hour accumulative precipitation that trigger human activity anomalies is 22.34 mm and 134.05 mm, respectively. The accumulative precipitation is much higher than the heavy rain yellow warning standard. In other words, it is amateurish to issue a heavy rain warning when the 6-hour accumulative precipitation is 365 50 mm for the HH cities. Results from this study therefore are of great value for the authorities who revise heavy rain alerts across China to help local residents be better prepared for such adverse events.

The study could by further studied. Rather than all the residents of a city, the Tencent location request dataset is generated by over one billion monthly active users. The Tencent dataset's aggregate geotagged human activities may underestimate the

设置了格式: 字体: (默认) +西文标题 (Times New Roman)

370 effects of rainstorms on infrequent users, particularly the elderly and children. To address this limitation and further investigate human responses to various weather events, our future studies would aim to integrate multisource geospatial datasets. Furthermore, identifying disaster types such as rainstorm, waterlogging, and flood from social media data and then analysing regional response variation of large-scale human activity in different disasters can improve deep understanding of urban resilience.

375 **Tab. 1 Urban resilience indecies from this study for different city groups and the difference from current Chinese yellow heavy rain alter standard.**

Type	Threshold(mm)	Response sensitivity	Alarm standard(mm)	Yellow warning(mm)	Difference from yellow warning
LH	1.8	0.71	10.8	50	-39.2
LM	2.08	0.51	12.46	50	-37.54
LL	2.23	0.39	13.35	50	-36.65
MH	3.61	0.63	21.68	50	-28.32
ML	4.99	0.41	29.94	50	-20.06
MM	5.03	0.49	30.19	50	-19.81
HL	17.5	0.42	104.97	50	54.97
HM	18.94	0.48	113.64	50	63.64
HH	22.34	0.64	134.05	50	84.05

#### Data availability

380 The information can be made available upon request to the corresponding author.

设置了格式: 字体: (默认) +西文标题 (Times New Roman)  
设置了格式: 字体: (默认) +西文标题 (Times New Roman)

#### Author contributions

J.Q., Y.D., J.Y., and F.L. conceived and designed the study and methods; J.Q., J.Y., Y.D., N.W., T.M., and T.P. analyzed the data; J.Q., F.L., Y.D. and J.Y. wrote the paper, and all coauthors contributed to the interpretation of the results and to the text. All authors read the manuscript and approved the submission.

设置了格式: 字体: (默认) +西文标题 (Times New Roman), 非加粗, 字体颜色: 自动设置  
设置了格式: 字体: (默认) +西文标题 (Times New Roman)  
设置了格式: 字体: (默认) +西文标题 (Times New Roman), 非加粗

#### Competing interests

385 The contact author has declared that neither they nor their co-author has any competing interests.

设置了格式: 字体: (默认) +西文标题 (Times New Roman)  
设置了格式: 字体: (默认) +西文标题 (Times New Roman), 非加粗, 字体颜色: 自动设置

## Financial support

设置了格式: 字体: (默认) +西文标题 (Times New Roman)

This research was jointly supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (Grant No. XDA19040501), the National Key Research and Development Program of China (Grant No. 2017YFC1503003), the 390 National Science Foundation of China (42176205 and 41901395).

设置了格式: 字体: (默认) +西文标题 (Times New Roman), 非加粗, 字体颜色: 自动设置

设置了格式: 字体: (默认) +西文标题 (Times New Roman)

## References

设置了格式: 字体: (默认) +西文标题 (Times New Roman)

- Adger, W.N., Hughes, T.P., Folke, C., Carpenter, S.R., Rockström, J., 2005. Social-Ecological Resilience to Coastal Disasters. *Science* 309, 1036–1039. <https://doi.org/10.1126/science.1112122>
- Aerts, J.C.J.H., Botzen, W.J.W., Emanuel, K., Lin, N., Moel, H. de, Michel-Kerjan, E.O., 2014. Evaluating Flood Resilience Strategies for Coastal Megacities. *Science* 344, 473–475. <https://doi.org/10.1126/science.1248222>
- Ambelu, A., Birhanu, Z., Tesfaye, A., Berhanu, N., Muhumuza, C., Kassahun, W., Daba, T., Woldemichael, K., 2017. Intervention pathways towards improving the resilience of pastoralists: A study from Borana communities, southern Ethiopia. *Weather and Climate Extremes* 17, 7–16. <https://doi.org/10.1016/j.wace.2017.06.001>
- Bertilsson, L., Wiklund, K., de Moura Tebaldi, I., Rezende, O.M., Verôl, A.P., Miguez, M.G., 2019. Urban flood resilience – A multi-criteria index to integrate flood resilience into urban planning. *Journal of Hydrology* 573, 970–982. <https://doi.org/10.1016/j.jhydrol.2018.06.052>
- Brusberg, M.D., Shively, R., 2015. Building drought resilience in agriculture: Partnerships and public outreach. *Weather and Climate Extremes, USDA Research and Programs on Extreme Events* 10, 40–49. <https://doi.org/10.1016/j.wace.2015.10.003>
- Buehler, Y.A., Kellenberger, T.W., Small, D., Itten, K.I., 2006. Rapid mapping with remote sensing data during flooding 2005 in Switzerland by object-based methods: a case study, in: *Geo-Environment and Landscape Evolution II: Monitoring, Simulation, Management and Remediation*. Presented at the GEO-ENVIRONMENT 2006, WIT Press, Rhodes, Greece, pp. 391–400. <https://doi.org/10.2495/GEO060391>
- Chan, F.K.S., Griffiths, J.A., Higgitt, D., Xu, S., Zhu, F., Tang, Y.-T., Xu, Y., Thorne, C.R., 2018. “Sponge City” in China—A breakthrough of planning and flood risk management in the urban context. *Land Use Policy* 76, 772–778. <https://doi.org/10.1016/j.landusepol.2018.03.005>
- Cleveland, R., Cleveland, W., McRae, J.E., Terpenning, I.J., 1990. STL: A seasonal-trend decomposition procedure based on loess (with discussion). *undefined*.
- de Bruijn, K.M., 2004. Resilience and flood risk management. *Water Policy* 6, 53–66. <https://doi.org/10.2166/wp.2004.0004>
- Dewan, T.H., 2015. Societal impacts and vulnerability to floods in Bangladesh and Nepal. *Weather and Climate Extremes, SI: IGBP APN* 7, 36–42. <https://doi.org/10.1016/j.wace.2014.11.001>
- Geller, S.C., Gregg, J.P., Hagerman, P., Rocke, D.M., 2003. Transformation and normalization of oligonucleotide microarray data. *Bioinformatics* 19, 1817–1823. <https://doi.org/10.1093/bioinformatics/btg245>
- Ghaffarian, S., Kerle, N., Filatova, T., 2018. Remote Sensing-Based Proxies for Urban Disaster Risk Management and Resilience: A Review. *Remote Sensing* 10, 1760. <https://doi.org/10.3390/rs10111760>
- Grinberger, A.Y., Felsenstein, D., 2016. Dynamic agent based simulation of welfare effects of urban disasters. *Computers, Environment and Urban Systems* 59, 129–141. <https://doi.org/10.1016/j.compenvurbsys.2016.06.005>
- Guan, X., Chen, C., 2014. Using social media data to understand and assess disasters. *Nat Hazards* 74, 837–850. <https://doi.org/10.1007/s11069-014-1217-1>
- Honey, M., Brink, S., Chang, S., Davidson, R., Amyx, P., Pyatt, S., Mills, R., Eguchi, R., Bevington, J., Panjwani, D., Hill, A., Adams, B., 2010. Uncovering Community Disruption Using Remote Sensing: An Assessment of Early Recovery in Post-earthquake Haiti (Technical Report). Disaster Research Center.
- Hong, B., Bonczak, B.J., Gupta, A., Kontokosta, C.E., 2021a. Measuring inequality in community resilience to natural disasters using large-scale mobility data. *Nat Commun* 12, 1870. <https://doi.org/10.1038/s41467-021-22160-w>

- Hong, B., Bonczak, B.J., Gupta, A., Kontokosta, C.E., 2021b. Measuring inequality in community resilience to natural disasters using large-scale mobility data. *Nat Commun* 12, 1870. <https://doi.org/10.1038/s41467-021-22160-w>
- Huang, W., Ling, M., 2018. System resilience assessment method of urban lifeline system for GIS. *Computers, Environment and Urban Systems* 71, 67–80. <https://doi.org/10.1016/j.compenvurbysys.2018.04.003>
- 435 Kasmalkar, I.G., Serafin, K.A., Miao, Y., Bick, I.A., Ortolano, L., Ouyang, D., Suckale, J., 2020. When floods hit the road: Resilience to flood-related traffic disruption in the San Francisco Bay Area and beyond. *Science Advances* 6, eaba2423. <https://doi.org/10.1126/sciadv.eaba2423>
- Kryvasheyeu, Y., Chen, H., Obradovich, N., Moro, E., Hentenryck, P.V., 2016. Rapid assessment of disaster damage using social media activity. *Science Advances* 2, e1500779–e1500779. <https://doi.org/10.1126/sciadv.1500779>
- 440 Levizzani, V., Kidd, C., Kirschbaum, D.B., Kummerow, C.D., Nakamura, K., Turk, F.J., 2020. Satellite Precipitation Measurement. *Springer Nature*.
- Liao, K.-H., 2012. A Theory on Urban Resilience to Floods--A Basis for Alternative Planning Practices. *E&S* 17, art48. <https://doi.org/10.5751/ES-05231-170448>
- Liaw, A., Wiener, M., 2001. Classification and Regression by RandomForest. *Forest* 23.
- 445 Liu, Y., Liu, X., Gao, S., Gong, L., Kang, C., Zhi, Y., Chi, G., Shi, L., 2015. Social Sensing: A New Approach to Understanding Our Socioeconomic Environments. *Annals of the Association of American Geographers* 105, 512–530.
- Liu, Z., Du, Y., Yi, J., Liang, F., Ma, T., Pei, T., 2019a. Quantitative Association between Nighttime Lights and Geo-Tagged Human Activity Dynamics during Typhoon Mangkhut. *Remote Sensing* 11, 2091. <https://doi.org/10.3390/rs11182091>
- 450 Liu, Z., Du, Y., Yi, J., Liang, F., Pei, T., 2019b. Quantitative estimates of collective geo-tagged human activities in response to typhoon Hato using location-aware big data. *International Journal of Digital Earth* 1–21. <https://doi.org/10.1080/17538947.2019.1645894>
- Ma, T., 2018. Multi-Level Relationships between Satellite-Derived Nighttime Lighting Signals and Social Media-Derived Human Population Dynamics. *Remote Sensing* 10, 1128. <https://doi.org/10.3390/rs10071128>
- 455 Marra, F., Nikolopoulos, E.I., Creutin, J.D., Borga, M., 2016. Space-time organization of debris flows-triggering rainfall and its effect on the identification of the rainfall threshold relationship. *Journal of Hydrology, Flash floods, hydro-geomorphic response and risk management* 541, 246–255. <https://doi.org/10.1016/j.jhydrol.2015.10.010>
- Martín, Y., Cutter, S.L., Li, Z., 2020a. Bridging Twitter and Survey Data for Evacuation Assessment of Hurricane Matthew and Hurricane Irma. *Nat. Hazards Rev.* 21, 04020003. [https://doi.org/10.1061/\(ASCE\)NH.1527-6996.0000354](https://doi.org/10.1061/(ASCE)NH.1527-6996.0000354)
- 460 Martín, Y., Cutter, S.L., Li, Z., Emrich, C.T., Mitchell, J.T., 2020b. Using geotagged tweets to track population movements to and from Puerto Rico after Hurricane Maria. *Popul Environ* 42, 4–27. <https://doi.org/10.1007/s11111-020-00338-6>
- McDougall, K., Temple-Watts, P., 2012. THE USE OF LIDAR AND VOLUNTEERED GEOGRAPHIC INFORMATION TO MAP FLOOD EXTENTS AND INUNDATION. *ISPRS Ann. Photogramm. Remote Sens. Spatial Inf. Sci.* 1–4, 251–256. <https://doi.org/10.5194/isprsaannals-1-4-251-2012>
- 465 Meerow, S., Newell, J.P., Stults, M., 2016. Defining urban resilience: A review. *Landscape and Urban Planning* 147, 38–49. <https://doi.org/10.1016/j.landurbplan.2015.11.011>
- Mendiondo, E.M., 2005. FLOOD RISK MANAGEMENT OF URBAN WATERS IN HUMID TROPICS: EARLY WARNING, PROTECTION AND REHABILITATION 14.
- 470 Mika, S., Ratsch, G., Weston, J., Scholkopf, B., Mullers, K.R., 1999. Fisher discriminant analysis with kernels, in: *Neural Networks for Signal Processing IX: Proceedings of the 1999 IEEE Signal Processing Society Workshop (Cat. No.98TH8468)*. Presented at the *Neural Networks for Signal Processing IX: Proceedings of the 1999 IEEE Signal Processing Society Workshop (Cat. No.98TH8468)*, pp. 41–48. <https://doi.org/10.1109/NNSP.1999.788121>
- Mpandeli, S., Nhamo, L., Moeletsi, M., Masupha, T., Magidi, J., Tshikolomo, K., Liphadzi, S., Naidoo, D., Mabhaudhi, T., 2019. Assessing climate change and adaptive capacity at local scale using observed and remotely sensed data. *Weather and Climate Extremes* 26, 100240. <https://doi.org/10.1016/j.wace.2019.100240>
- 475 Myhre, G., Alterskær, K., Stjern, C.W., Hodnebrog, Ø., Marelle, L., Samset, B.H., Sillmann, J., Schaller, N., Fischer, E., Schulz, M., Stohl, A., 2019. Frequency of extreme precipitation increases extensively with event rareness under global warming. *Scientific Reports* 9, 16063. <https://doi.org/10.1038/s41598-019-52277-4>
- Nahiduzzaman, K.M., Aldosary, A.S., Rahman, M.T., 2015. Flood induced vulnerability in strategic plan making process of 480 Riyadh city. *Habitat International* 49, 375–385. <https://doi.org/10.1016/j.habitatint.2015.05.034>

- Naidu, S., Sajinkumar, K.S., Oommen, T., Anuja, V.J., Samuel, R.A., Muraleedharan, C., 2018. Early warning system for shallow landslides using rainfall threshold and slope stability analysis. *Geoscience Frontiers, Reliability Analysis of Geotechnical Infrastructures* 9, 1871–1882. <https://doi.org/10.1016/j.gsf.2017.10.008>
- Ogie, R.I., Holderness, T., Dunn, S., Turpin, E., 2018. Assessing the vulnerability of hydrological infrastructure to flood damage in coastal cities of developing nations. *Computers, Environment and Urban Systems* 68, 97–109. <https://doi.org/10.1016/j.compenvurbsys.2017.11.004>
- 485 Olsson, L., Jerneck, A., Thoren, H., Persson, J., O'Byrne, D., 2015. Why resilience is unappealing to social science: Theoretical and empirical investigations of the scientific use of resilience. *Science Advances* 1, e1400217. <https://doi.org/10.1126/sciadv.1400217>
- 490 Östh, J., Reggiani, A., Galiazzo, G., 2015. Spatial economic resilience and accessibility: A joint perspective. *Computers, Environment and Urban Systems* 49, 148–159. <https://doi.org/10.1016/j.compenvurbsys.2014.07.007>
- O'Sullivan, J.J., Bradford, R.A., Bonaiuto, M., De Dominicis, S., Rotko, P., Aaltonen, J., Waylen, K., Langan, S.J., 2012. Enhancing flood resilience through improved risk communications. *Nat. Hazards Earth Syst. Sci.* 12, 2271–2282. <https://doi.org/10.5194/nhess-12-2271-2012>
- 495 Ouyang, M., Dueñas-Osorio, L., Min, X., 2012. A three-stage resilience analysis framework for urban infrastructure systems. *Structural Safety* 36–37, 23–31. <https://doi.org/10.1016/j.strusafe.2011.12.004>
- Wrangi, A.M., Lannigan, R., Simonovic, S.P., 2014. Interaction between land-use change, flooding and human health in Metro Vancouver, Canada. *Nat Hazards* 72, 1219–1230. <https://doi.org/10.1007/s11069-014-1064-0>
- 500 Pal, M., 2005. Random forest classifier for remote sensing classification. *International Journal of Remote Sensing* 26, 217–222. <https://doi.org/10.1080/01431160412331269698>
- Poulin, C., Kane, M.B., 2021. Infrastructure resilience curves: Performance measures and summary metrics. *Reliability Engineering & System Safety* 216, 107926. <https://doi.org/10.1016/j.ress.2021.107926>
- 505 Qian, J., Du, Y., Yi, J., Liang, F., Huang, S., Wang, X., Wang, N., Tu, W., Pei, T., Ma, T., 2022. Regional geographical and climatic environments affect urban rainstorm perception sensitivity across China. *Sustainable Cities and Society* 87, 104213. <https://doi.org/10.1016/j.scs.2022.104213>
- 510 Qian, J., Liu, Z., Du, Y., Liang, F., Yi, J., Ma, T., Pei, T., 2021a. Quantify city-level dynamic functions across China using social media and POIs data. *Computers Environment and Urban Systems* 85. <https://doi.org/10.1016/j.compenvurbsys.2020.101552>
- Qian, J., Liu, Z., Du, Y., Wang, N., Yi, J., Sun, Y., Ma, T., Pei, T., Zhou, C., 2021b. Multi-level Inter-regional Migrant Population Estimation Using Multi-source Spatiotemporal Big Data: A Case Study of Migrants in Hubei Province During the Outbreak of COVID-19 in Wuhan, in: Shaw, S.-L., Sui, D. (Eds.), *Mapping COVID-19 in Space and Time: Understanding the Spatial and Temporal Dynamics of a Global Pandemic, Human Dynamics in Smart Cities*. Springer International Publishing, Cham, pp. 169–188. [https://doi.org/10.1007/978-3-030-72808-3\\_9](https://doi.org/10.1007/978-3-030-72808-3_9)
- 515 Román, M., Stokes, E., Shrestha, R., Wang, Z., Schultz, L., Carlo, E., Sun, Q., Bell, J., Molthan, A., Kalb, V., ji, C., Seto, K., Mcclain, S., Enenkel, M., 2019. Satellite-based assessment of electricity restoration efforts in Puerto Rico after Hurricane Maria. *PLOS ONE* 14, e0218883. <https://doi.org/10.1371/journal.pone.0218883>
- Rosner, B., 1975. On the Detection of Many Outliers. *null* 17, 221–227. <https://doi.org/10.1080/00401706.1975.10489305>
- 520 Shiferaw, B., Tesfaye, K., Kassie, M., Abate, T., Prasanna, B.M., Menkir, A., 2014. Managing vulnerability to drought and enhancing livelihood resilience in sub-Saharan Africa: Technological, institutional and policy options. *Weather and Climate Extremes, High Level Meeting on National Drought Policy* 3, 67–79. <https://doi.org/10.1016/j.wace.2014.04.004>
- Song, J., Chang, Z., Li, W., Feng, Z., Wu, J., Cao, Q., Liu, J., 2019. Resilience-vulnerability balance to urban flooding: A case study in a densely populated coastal city in China. *Cities* 95, 102381. <https://doi.org/10.1016/j.cities.2019.06.012>
- 525 Stefan, V., Fabio, G.-T., Josh, L., Jan, K., Brenda, J., 2016. Global Trends in Satellite-based Emergency Mapping. *Science* 353, 247–252.
- Strobl, C., Boulesteix, A.-L., Zeileis, A., Hothorn, T., 2007. Bias in random forest variable importance measures: Illustrations, sources and a solution. *BMC Bioinformatics* 8, 25. <https://doi.org/10.1186/1471-2105-8-25>
- Tellman, B., Sullivan, J.A., Kuhn, C., Kettner, A.J., Doyle, C.S., Brakenridge, G.R., Erickson, T.A., Slayback, D.A., 2021. Satellite imaging reveals increased proportion of population exposed to floods. *Nature* 596, 80–86. <https://doi.org/10.1038/s41586-021-03695-w>

- Vallis, O., Hochenbaum, J., Kejariwal, A., 2014. A Novel Technique for Long-Term Anomaly Detection in the Cloud. Presented at the 6th {USENIX} Workshop on Hot Topics in Cloud Computing (HotCloud 14).
- Wang, B., Loo, B.P.Y., Zhen, F., Xi, G., 2020. Urban resilience from the lens of social media data: Responses to urban flooding in Nanjing, China. *Cities* 106, 102884. <https://doi.org/10.1016/j.cities.2020.102884>
- 535 Wang, J., Meng, B., Pei, T., Du, Y., Zhang, J., Chen, S., Tian, B., Zhi, G., 2021. Mapping the exposure and sensitivity to heat wave events in China's megacities. *Science of The Total Environment* 755, 142734. <https://doi.org/10.1016/j.scitotenv.2020.142734>
- Wang, N., Du, Y., Liang, F., Yi, J., Wang, H., 2019. Spatiotemporal Changes of Urban Rainstorm-Related Micro-Blogging Activities in Response to Rainstorms: A Case Study in Beijing, China. *Applied Sciences* 9, 4629. <https://doi.org/10.3390/app9214629>
- 540 Wang, Y., Taylor, J.E., 2018. Coupling sentiment and human mobility in natural disasters: a Twitter-based study of the 2014 South Napa Earthquake. *Nat Hazards* 92, 907–925. <https://doi.org/10.1007/s11069-018-3231-1>
- Yabe, T., Tsubouchi, K., Fujiwara, N., Sekimoto, Y., Ukkusuri, S.V., 2020. Understanding post-disaster population recovery patterns. *Journal of The Royal Society Interface* 17, 20190532. <https://doi.org/10.1098/rsif.2019.0532>
- 545 Yi, J., Du, Y., Liang, F., Pei, T., Zhou, C., 2019. Anomalies of dwellers' collective geotagged behaviors in response to rainstorms: a case study of eight cities in China using smartphone location data. *Natural Hazards and Earth System Sciences* 19, 2169–2182.
- Yue, Y., Zhuang, Y., Yeh, A.G.O., Xie, J.-Y., Ma, C.-L., Li, Q.-Q., 2017. Measurements of POI-based mixed use and their relationships with neighbourhood vibrancy. *International Journal of Geographical Information Science* 31, 658–675. <https://doi.org/10.1080/13658816.2016.1220561>
- 550 Zhang, R., Chen, Y., Zhang, X., Ma, Q., Ren, L., 2022. Mapping homogeneous regions for flash floods using machine learning: A case study in Jiangxi province, China. *International Journal of Applied Earth Observation and Geoinformation* 108, 102717. <https://doi.org/10.1016/j.jag.2022.102717>
- Zheng, S., Wang, J., Sun, C., Zhang, X., Kahn, M.E., 2019. Air pollution lowers Chinese urbanites' expressed happiness on social media. *Nature Human Behaviour* 3, 237–243. <https://doi.org/10.1038/s41562-018-0521-2>
- 555 Zou, L., Lam, N., Cai, H., Qiang, Y., 2018. Mining Twitter Data for Improved Understanding of Disaster Resilience. *Annals of the Association of American Geographers* 108, 1422–1441. <https://doi.org/10.1080/24694452.2017.1421897>
- Zou, L., Lam, N.S.N., Shams, S., Cai, H., Meyer, M.A., Yang, S., Lee, K., Park, S.-J., Reams, M.A., 2019. Social and geographical disparities in Twitter use during Hurricane Harvey. *International Journal of Digital Earth* 12, 1300–1318. <https://doi.org/10.1080/17538947.2018.1545878>
- 560

Hong, B., B. J. Bonczak, A. Gupta and C. E. Kontokosta (2021). "Measuring inequality in community resilience to natural disasters using large-scale mobility data." *Nature communications* 12(1): 1–9.

设置了格式: 字体: (默认) +西文标题 (Times New Roman), 德语 (德国), 不检查拼写或语法

带格式的: 正文, 定义网格后不调整右缩进, 不对齐到网格

设置了格式: 字体: (默认) +西文标题 (Times New Roman)