

---

1                    Characteristics of cloud-to-ground lightning (CG) and  
2                    differences between +CG and -CG stroke in China regarding  
3                    China National Lightning Detection Network

4     **Ruijiao Jiang<sup>a</sup>, Guoping Zhang<sup>a,\*</sup>, Shudong Wang<sup>a</sup>, Bing Xue<sup>a</sup>, Zhengshuai Xie<sup>b</sup>,**  
5     **Tingzhao Yu<sup>a</sup>, Kuoyin Wang<sup>a</sup>, Jin Ding<sup>a</sup>, Xiaoxiang Zhu<sup>a</sup>**

6     <sup>a</sup> Public Meteorological Service Center, China Meteorological Administration, Beijing 100081, China

7     <sup>b</sup> Weather Modification Centre, China Meteorological Administration, Beijing 100081, China

8     \*Corresponding author. E-mail address: zhanggp@cma.gov.cn

9     **Abstract**

10            The lightning location system consisting of multiple ground-based stations is an  
11            effective means of lightning observation. The dataset from CNLDN (China National  
12            Lightning Detection Network) in 2016-2022 is employed to analyze the temporal and  
13            spatial lightning distributions and the differences between +CG (positive cloud-to-  
14            ground lightning) and -CG (negative cloud-to-ground lightning) stroke in China. On  
15            the **annual** scale, lightning activity is most prevalent during the summer months (June,  
16            July, and August), accounting for 72.6 % of the year. Spring sees more lightning than  
17            autumn, and winter has only a small amount in southeastern coastal areas. During the  
18            day, the frequency of lightning peaks at 15:00-17:00 CST (China Standard Time) and  
19            is lowest at 8:00-10:00 CST. For the period with high CG stroke frequency (summer of  
20            a year or afternoon of a day), the proportion of +CG stroke and the discharge peak  
21            current is relatively small. Winter in a year and morning or midnight in a day  
22            corresponds to a greater +CG stroke proportion and discharge current. Spatially, low  
23            latitude, undulating terrain, seaside, and humid surface are favorable factors for  
24            lightning occurrence. Thus, the southeast coastland has the largest lightning stroke  
25            density, while the northwest deserts and basins and the western and northern Tibetan  
26            Plateau, with altitudes over 6000 meters, have almost no lightning. The proportion of  
27            +CG stroke and the peak current are low in the southern region with high density but  
28            diverse in other regions. The Tibetan Plateau leads to the diversity of lightning activity  
29            in China and lays the foundation for studying the impact of surface elevation on  
30            lightning. Results indicate that the +CG stroke proportion on the eastern and southern  
31            Tibetan Plateau is up to 15 %, larger than the plain regions. The **peak current** of -CG  
32            stroke is positively correlated with altitude, but +CG stroke shows a negative  
33            correlation, resulting in a large difference in **peak current** between +CG and -CG on the  
34            plain and approach on the plateau.

---

35 **Keywords:** China, CNLDN, Lightning characteristics, +CG stroke, Peak current

36 1. Introduction

37 Most lightning is generated mainly through mesoscale or small scale severe  
38 convective weather, with occasionally occurring in tropical cyclones, volcanic  
39 eruptions, and dust storms (Rakov and Uman, 2003). Lightning, a violent long-distance  
40 transient discharge phenomenon, could cause severe disasters such as human and  
41 animal casualties, forest fires, and electronic and communication equipment  
42 interruptions. Lightning is also associated with extreme weather events such as heavy  
43 rainfall, hail, and strong winds. These events can cause damage to infrastructure, crops  
44 and property, and pose a threat to public safety. Therefore, the timely and accurate  
45 monitoring of lightning serves as an effective approach for the development of lightning  
46 science and scientifically mitigating the hazards of lightning strikes.

47 Lightning discharge emits electromagnetic spectrums with a broad range,  
48 providing an essential avenue for lightning detection. The very low frequency / low  
49 frequency (VLF/LF, 20-300 kHz) band radiation is mainly produced by the cloud-to-  
50 ground lightning (CG) return strokes, intracloud (IC) K-changes, and other discharge  
51 processes with a large spatial scale (Preston and Tolver, 1989; Schulz et al., 2016;  
52 Cummins et al., 1998). VLF/LF electromagnetic waves can propagate along the ground  
53 surface or be reflected between the surface and ionosphere propagation, with the  
54 superiority of long propagation distance (hundreds to thousands of kilometers) and slow  
55 attenuation. This frequency range thus is suitable for large-scale lightning detection and  
56 is currently the most commonly used target detection band for ground-based lightning  
57 location systems (Wang et al., 2020).

58 Representative examples of modern lightning location systems working in  
59 VLF/LF band are mainly the U.S. National Lightning Detection Network (NLDN)  
60 (Rudlosky and Fuelberg, 2010), Los Alamos Sferic Array (LASA) (Shao et al., 2006),  
61 European Cooperation for Lightning Detection (EUCLID) (Schulz et al., 2016),  
62 Vaisala's Global Lightning Dataset (GLD360) (Said et al., 2013), World Wide  
63 Lightning Location Network (WWLLN) (Rudlosky and Shea, 2013), etc. The three  
64 nationwide detection networks in China are China National Lightning Detection  
65 Network (CNLDN), operated by the Meteorological Observation Centre of China  
66 Meteorological Administration (CMA), the Lightning Location System (LLS) of the  
67 State Grid Corporation of China, and the Three-Dimensional Lightning Location  
68 System (3D-LLS) deployed by the Institute of Electrical Engineering of Chinese  
69 Academy of Sciences (CAS). There are also small-scale and refined detection systems  
70 in local China areas, such as the Beijing Lightning Network (BLNET) established by  
71 the Institute of Atmospheric Physics of CAS, the Guangdong-Hongkong-Macao  
72 Lightning Location System (GHMLLS) deployed by the meteorological departments  
73 of Guangdong Province, Hongkong, and Macao.

74 China spans a wide range of latitudes from north to south, and significant terrain

---

75 changes from east to west. The western and northern parts of the Tibetan Plateau have  
76 large uninhabited areas with altitudes above 6000 m. The above factors pose challenges  
77 for the installation of lightning detectors and the improvement of the accuracy of  
78 locating algorithms. Currently, most of the analyses of lightning characteristics in China  
79 are based on lightning imagers on satellites and the World-Wide Lightning Location  
80 Network (WWLLN) (Ma et al., 2005; You et al., 2019; Ma et al., 2021). However, the  
81 Optical Transient Detector (OTD) on Microlab-1 and Lightning Imaging Sensor (LIS)  
82 on TRMM (Tropical Rainfall Measuring Mission) satellite discontinued updates, and  
83 the Chinese Lightning Mapping Imager (LMI) on FY-4A is not oriented to the China  
84 area all year round. Meanwhile, the detection efficiency of the satellite sensor and  
85 WWLLN is relatively low and not valuable for analyzing the difference between  
86 positive cloud-to-ground lightning (+CG) and negative cloud-to-ground lightning (-  
87 CG). CNLDN is nowadays the most widely used system by the meteorological  
88 departments in China and has accumulated observational data for many years. Currently,  
89 there has been no study evaluating the detection performance of CNLDN on a national  
90 scale. Based on some localized assessments, the detection efficiency for lightning flash  
91 and stroke in Beijing was reported to be 49.4 % and 36.5 % (Srivastava et al., 2017). In  
92 Jiangsu, the detection efficiency for lightning flash was documented as 61.4 % (Min-  
93 Xue et al., 2015). In our previous study, we calculated the lightning density in the  
94 vicinity of the Canton Tower in Guangdong using optical observation by the Tall-Object  
95 Lightning Observatory in Guangzhou (TOLOG), which is currently regarded as a  
96 nearly no-miss observation (Wu et al., 2019; Jiang, 2021). The result was that the CG  
97 flash density was  $20 \text{ km}^{-2}\cdot\text{yr}^{-1}$  within a 3 km radius of the Canton Tower. The detection  
98 efficiency of lightning flash for CNLDN was correspondingly only 33.7 % in the same  
99 region. Despite the relatively lower detection efficiency of CNLDN compared to other  
100 advanced international systems, it still holds significant importance in analyzing the  
101 lightning characteristics in nationwide China.


102 This study makes use of CNLDN data from 2016-2022 to analyze the lightning  
103 climate over China by dividing the continental region into four blocks. We also focus  
104 on comparing the differences between +CG and -CG stroke regarding temporal and  
105 spatial distribution. Furthermore, China's extensive geographical expanse, spanning a  
106 wide range of latitudes and longitudes, coupled with its intricate topography, provides  
107 a unique opportunity for investigating the correlation between lightning occurrences  
108 and geographic factors.

## 109 2. Data source

110 CNLDN was initially developed in 2007 by the National Space Science Center  
111 (NSSC) of CAS and is now operated by the Meteorological Observation Centre of  
112 CMA. It carries out the monitoring and early warning of strong convective weather in  
113 thunderstorms. The CNLDN system comprised 435 sub-stations (as of 2020), each  
114 equipped with a lightning detector, and a central data processing station located at the

115 National Meteorological Information Center. The baseline distance typically ranges  
116 from tens to hundreds of kilometers. Although with some blind areas in the desert  
117 regions of Xinjiang and the uninhabited high-altitude areas of Tibet, CNLND can  
118 generally achieve nationwide lightning detection. The distribution of sub-stations is  
119 illustrated in Fig. 1.

120 The network uses a time-of-arrival (TOA) technique, with a GPS timing error of  
121 fewer than 20 ns (in clear sky conditions), to detect VLF/LF pulses of CG return strokes.  
122 A lightning flash may consist of several CG strokes, and the stroke locations within  
123 each flash always do not overlap, causing striking disasters in different areas. In this  
124 study, we analyze lightning stroke characteristics in inland China using the CNLND  
125 dataset from 2016-2022, downloaded from the CMA big data cloud platform. For each  
126 stroke, we can obtain information on the time of occurrence, latitude, longitude, peak  
127 current value, and the number of reporting sensors.



128  
129 **Fig. 1. CNLND sites distribution and altitude distribution map of China with labeled names of each**  
130 **province and municipality. The alphabetical labels (a, b, c, d, etc.) in the figure indicate the positions of the**  
131 **mentioned locations referred to in the text.**

### 132 3. CG characteristics in China

133 China's climate features are greatly influenced by its wide latitudinal span,  
134 significant terrain disparity, complex topography, and ocean currents (Ren et al., 2012).

135 Lightning is a fundamental meteorological element, and its long-term accumulation  
 136 characteristics are closely linked to China's climate. The spatial distribution of lightning  
 137 in China is determined by a combination of factors, including atmospheric circulation,  
 138 topography, distance from the sea, latitude, etc.

### 139 3.1 CG stroke distribution in China

140 Previous studies have often divided China into four major geographical regions,  
 141 each with relatively uniform climatic characteristics. These regions are Southern China  
 142 (Region-I), Northern China (Region-II), Northwestern China (Region-III), and the  
 143 Qinghai-Tibet region of China (Region-IV), as illustrated in Fig. 2. The Qinling  
 144 Mountains (a) - Huaihe River (b) line, which roughly coincides with the 0 °C isotherm  
 145 and 800 mm annual precipitation line in January, serves as the boundary between  
 146 Region-I and Region-II. The Daxing'an Mountains (c) - Yinshan Mountains (d) - Helan  
 147 Mountains (e), which divide the monsoon and non-monsoon regions and the 400 mm  
 148 annual precipitation line, serve as the boundaries between Region-II and Region-III.  
 149 The boundary between Region-IV and Regions I-II-III is approximately the line  
 150 between China's first and second steps in terrain.


151 **Tab. 1 Statistics on the annual average numbers of return strokes, the stroke densities, and the peak current**  
 152 **values of the four regions**

	Region-I	Region-II	Region-III	Region-IV
-CG stroke number ( $10^3 \cdot \text{yr}^{-1}$ )	3653	499	127	129
+CG stroke number ( $10^3 \cdot \text{yr}^{-1}$ )	388	151	37	18
stroke number ( $10^3 \cdot \text{yr}^{-1}$ )	4041	650	166	147
Mean stroke density ( $\text{km}^{-2} \cdot \text{yr}^{-1}$ )	1.68	0.34	0.05	0.07
Mean peak current of -CG (-kA)	-41	-39	-37	-45
Mean peak current of +CG (kA)	47	54	65	66

153 The statistical results of the annual average number of -CG return stroke, +CG  
 154 return stroke, all types of stroke, as well as the mean stroke density of four regions are  
 155 shown in Tab. 1. The data indicates a sequential decrease in lightning activity across  
 156 the four regions. Fig. 2 displays the distribution of annual average CG stroke density  
 157 from 2016-2022. When calculating lightning density, it is necessary to determine the  
 158 size of grid cells. According to the research of Diendorfer (2008), lightning is a highly  
 159 stochastic phenomenon, and if we require an uncertainty of less than  $\pm 20\%$  per grid  
 160 cell, there should be more than 80 events in it. Therefore, we establish the grid size as  
 161  $0.25^\circ \times 0.25^\circ$ , ensuring that the results are within the confidence interval for almost all  
 162 regions, except part of areas in Xinjiang and Tibet. Lightning primarily occurs in  
 163 convective precipitation and, to a lesser extent, in stratus cloud precipitation. Generally,  
 164 the spatial distribution of lightning stroke in Fig. 2 is consistent with the distribution of  
 165 annual average precipitation in China, as illustrated in Fig. 3 in Jin et al. (2021).

166 Region-I has the highest concentration of CG stroke, with an average density of  
 167  $1.68 \text{ km}^{-2} \cdot \text{yr}^{-1}$ . The leap line of lightning density corresponds well with the 0 °C

168 isotherms in January, the 800 mm annual equivalent precipitation line, and the eastern  
 169 dividing line of the first and second terrain steps. The climate in Region-I is mainly  
 170 influenced by the tropical/subtropical monsoon. The southeast monsoon from the  
 171 Pacific Ocean and the southwest monsoon from the Indian Ocean make the summer hot  
 172 and humid and prone to thunderstorms. In particular, the monsoon influence is more  
 173 pronounced in coastal areas with abundant water vapor and thermal conditions. In the  
 174 mountainous regions of Hainan, Guangdong, Fujian, and Zhejiang, where the rolling  
 175 topography lifts the warm and humid air masses, thunderstorm activity is most frequent,  
 176 resulting in high lightning density. Although the Sichuan Basin and Yunnan are far from  
 177 the coastline, they are located at the eastern and southern windward slopes of the  
 178 Tibetan Plateau, which benefits the generation of thunderstorm activities due to the  
 179 topographic uplift. The distribution in Region-I is similar with the observation results  
 180 by Xia et al. (2015).



181  
 182 **Fig. 2. 2016-2022 annual average CG stroke density distribution in China. Region-I: Southern China;**  
 183 **Region-II: Northern China; Region-III: Northwestern China; Region-IV: Qinghai-Tibet region of China.**  
 184 **The grid size is  $0.25^\circ \times 0.25^\circ$ .**

185 Region-II has a temperate monsoon climate, with summer influenced by the  
 186 southeast monsoon carrying temperate marine air mass or **degenerative** tropical marine  
 187 air mass, making summer warm and rainy. Most areas have CG stroke density between  
 188  $0.1-1 \text{ km}^{-2}\cdot\text{yr}^{-1}$ , slightly lower than Region-I. The lightning activity in Region-II is also

---

189 greater in the seaside area than inland areas. Shanxi is located in a mountainous region,  
190 and the undulating terrain makes it a high incidence area for thunderstorm activity.  
191 Region-II has the most extensive plain, Northeastern China Plain (f), surrounded by the  
192 Daxing'an Mountains (c) - Xiaoxing'an Mountains (g) - Changbai Mountains (h). The  
193 landform is conducive to the southeast monsoon reaching the inland areas of Region-II  
194 and forming summer thunderstorms. Jilin is only a dozen kilometers from the Sea of  
195 Japan, facilitating the entry of Japanese warm air currents. Therefore, despite its high  
196 latitude, thunderstorm activity is relatively intense in Region-II.

197 Region-III, which includes Xinjiang, northern Gansu, and most land of Inner  
198 Mongolia, has a temperate continental climate. There are significant differences in  
199 lightning distribution characteristics within this region. The southern and central parts  
200 of Region-III consist primarily of vast deserts and gobies. The Tibetan Plateau blocks  
201 the humid South Asian monsoon, and its arid surface cannot produce abundant water  
202 vapor, resulting in few thunderstorms. However, the Tianshan Mountains (i), Kunlun  
203 Mountains (j), and Altay Mountains (k), located in the hinterland of the Eurasian  
204 continent, are provided with water vapor for thunderstorm generation through the  
205 westerly circulation that transports evaporated water vapor from the Atlantic Ocean and  
206 the Eurasian continent. As a result, the northern mountainous areas occupy almost all  
207 the lightning activity in Xinjiang. The southeastern monsoon flowing through Region-  
208 II can also bring some thunderstorm processes to the eastern and central mountainous  
209 regions in Inner Mongolia during summer.

210 Region-IV's primary landmass is the Tibetan Plateau, which includes Tibet,  
211 Qinghai, southern Xinjiang, and western Sichuan. It has a highland mountain climate,  
212 and the overall geomorphic distribution trend increases from east to west (Ma et al.,  
213 2021). The uninhabited areas above 4500 meters in elevation in the west and north of  
214 Region-IV are icy all year round, covered by snow and glaciers. The Qaidam Basin (l)  
215 in Qinghai is a closed, huge, interrupted basin, where dry sinking airflow from the  
216 northern edge of the plateau in summer leads to water shortage. Consequently, there are  
217 few thunderstorm activities in these areas, and the distribution of sub-stations is sparse,  
218 making them the regions with the lowest lightning density detected in China, with CG  
219 stroke density less than  $10^{-3} \text{ km}^{-2} \cdot \text{yr}^{-1}$ . In contrast, the southern Himalayas Mountains  
220 (m), near the Yarlung Tsangpo River Grand Canyon (n), have a relatively low altitude,  
221 opening a "gap" for the influx of abundant water vapor from the Bay of Bengal.  
222 However, this narrow plain area, located in Mêdog County (o), has very high  
223 precipitation but low lightning density, which can also be concluded from the  
224 observations of TRMM. The remaining moisture continues northward across this plain,  
225 causing most thunderstorms between the east-west Himalayas Mountains and Tanggula  
226 Mountains (p). The thunderstorms on the east side of the plateau are mainly influenced  
227 by the low vortex and the shear line, which is usually stable at around  $32.5^{\circ}\text{N}$  (You et  
228 al., 2019; Qie et al., 2003). The high lightning density area is precisely located on the  
229 south side of the shear line.

---

## 230 3.2 Differences between +CG and -CG stroke

231 Based on the different polarities of neutralized charges in thunderclouds, CG can  
232 be classified into two types: +CG and -CG. Generally, +CG has a lower occurrence  
233 probability, accounting for only about 10 % of all CG, but it is characterized by a larger  
234 charge transfer, which results in a more significant hazard (Preston and Tolver, 1989;  
235 Carey and Buffalo, 2007). Studies have suggested that thunderstorms dominated by  
236 +CG are more likely to result in tornadoes and hail, particularly if the dominant phase  
237 lasts for tens of minutes. This may be related to changes in the charge distribution  
238 structure within thunderstorm clouds during extreme weather events (Williams, 1985).  
239 Previous research has been conducted on the comparative analysis of +CG and -CG in  
240 specific regions (Nag et al., 2014; Rakov and Uman, 2003). Based on these findings,  
241 this study aims to investigate further the spatial and temporal variability of +CG and -  
242 CG stroke in China, taking into account the complex climatic and geographical factors  
243 that influence lightning activity.


### 244 3.2.1 Comparison of temporal distribution of +CG and -CG stroke

245 The geographical and climatic features differ considerably across the four regions.  
246 Therefore, we analyze the temporal distribution of lightning stroke separately for each  
247 region.

248 Fig. 3 illustrates the monthly average CG stroke frequency distribution in China  
249 over a seven-year period from 2016 to 2022. The stroke frequency varies significantly  
250 across the four regions, with Region-I having the highest frequency, followed by  
251 Region-II and Region-III, and Region-IV having the lowest. But the stroke frequency  
252 shows similar fluctuations throughout the year among the regions, with August having  
253 the highest frequency and December having the lowest. Lightning activity is also scarce  
254 in November, January, and February, with a sudden surge in March and a gradual  
255 increase in the following months. Based on the seasonal classification, lightning  
256 activity is most active in summer (June, July, and August), accounting for 72.6 % of the  
257 year. In other seasons, lightning is more frequent in spring (18.4 %) than in autumn  
258 (8.6 %), but much less frequent than in summer. This is mainly because the summer  
259 monsoon affecting China starts to form during April and May, while the cold and dry  
260 winter monsoon starts to build up and push southward from September, making  
261 thunderstorm activity in spring and autumn mainly concentrated in southern areas,  
262 particularly coastal areas. In winter, most regions in China are controlled primarily by  
263 cold high pressure, resulting in very little lightning, with only a small amount occurring  
264 in southeastern coastal areas, accounting for just 0.4 % of the year. Overall, lightning  
265 stroke distribution follows a seasonal trend that advances from south to north and then  
266 retreats southward, which is consistent with the trend of the summer monsoon. The  
267 above proportions of four seasons closely resemble the statistical results conducted in  
268 the US by Holle and Cummins (2010).




269 Furthermore, the proportion of +CG stroke in different months is calculated and  
 270 represented by the gray line in Fig. 3. To ensure the reliability of the analysis, months  
 271 with less than 50 +CG strokes are excluded to avoid the impact of outliers. Results  
 272 indicate an evident inverse relationship between the proportion of +CG stroke and the  
 273 frequency of stroke. Notably, the months with the highest incidence of CG stroke,  
 274 namely July and August, exhibit the lowest proportion of +CG stroke across the four  
 275 regions. During this period, Region-I and Region-IV, located at lower latitudes, exhibit  
 276 a proportion of +CG stroke of approximately 10 %, while Region-II and Region-III  
 277 display proportions between 10 % and 30 %. Winter in four regions have rare  
 278 thunderstorms but demonstrates the highest proportion of +CG stroke, even reaching  
 279 over 50 % in December in Region-I, Region-II and Region-IV. In other months, some  
 280 irregular fluctuations are observed, Region-I and Region-II show significantly high  
 281 proportions of +CG stroke in February and April-May, respectively.



282  
 283 **Fig. 3. Annual scale variation of the frequency distribution of +CG and -CG stroke. The gray line**  
 284 **represents the proportion of +CG stroke.**

285 Tab. 1 also displays the mean peak current values of two types of CG stroke in the  
 286 four regions. The results reveal that the +CG stroke current in Region-III and Region-  
 287 IV exhibits a notably higher magnitude compared to other two regions, and the disparity  
 288 between +CG and -CG is more pronounced. While the average peak current values of  
 289 both types are found to be in close proximity in Region-I. The value fluctuation in  
 290 different months is illustrated in Fig. 4. To avoid outliers, no box is drawn when the

291 stroke count is less than 50. Overall, the distribution range is wider in winter than in  
 292 other seasons, and Region-II has a wider current distribution interval than other regions.  
 293 The average peak current value of each month is indicated by a white cross, and the  
 294 variation trend is shown by a red line. The results indicate that +CG strokes generally  
 295 have a higher peak current than -CG strokes. The peak current and the proportion of  
 296 +CG stroke exhibit similar trends, with a higher proportion and stronger peak current  
 297 in winter and a lower proportion and weaker peak current in summer. **In Region-I and**  
 298 **Region-II, the seasonal fluctuations of +CG peak current are more pronounced than -**  
 299 **CG peak current, with the peak current of +CG stroke falling even below -CG stroke**  
 300 **in summer and early autumn in Region-I.** The trends of +CG and -CG stroke peak current  
 301 are more consistent in Region-III and Region-IV.




302  
 303 **Fig. 4. Annual scale variation of the peak current distribution of +CG and -CG stroke. The red line**  
 304 **represents the average peak current value of each month.**

305 Fig. 5 illustrates the hour-by-hour frequency and proportion variation of +CG  
 306 stroke throughout the day. The frequency of both +CG and -CG strokes show noticeable  
 307 and consistent fluctuations. The active period for lightning activity is concentrated in  
 308 the late afternoon to wee hours, which coincides with the maximum accumulation of  
 309 radiative heating and vapor conducive to the development of convection, particularly  
 310 during summer thunderstorms in China. **Stroke frequency peaks at 15:00 to 16:00 CST**  
 311 **(China Standard Time, UTC+8 h) in Region-I and Region-II in the east of China and 1-**  
 312 **2 hours later in Region-III and Region-IV in the west of China. After nightfall, lightning**

313 activity gradually weakens due to the decline in unstable energy, dropping to a trough  
 314 at 8:00-10:00 CST the following day. The monthly and hourly variations in lightning  
 315 frequency are similar to the observations in Europe by EUCLID (Poelman et al., 2015)  
 316 and in US by NLDN (Koehler, 2020; Holle and Cummins, 2010).


317 The proportion of +CG stroke is inversely correlated with the total number of CG  
 318 stroke in a day in Region-I, Region-II, and Region-IV, as shown in Fig. 5. The maxima  
 319 of the +CG stroke proportion coincides with the lowest frequency at 8:00-10:00 CST  
 320 in all four regions, but the minima appear 2-3 hours earlier than the frequency peak.  
 321 Region-I and Region-IV at low latitudes have maximum proportions in the morning,  
 322 while Region-II and Region-III at high latitudes have maximum proportions in the  
 323 evening. **Additionally, the proportion of +CG stroke is lower in Region-I and Region-**  
 324 **IV, with values of less than 20 %, than in Region-II and Region-III, with values of over**  
 325 **20 %.** These findings demonstrate a close relationship between thunderstorm  
 326 characteristics and geographical features such as latitude, topography, and sea distance.



327  
 328 **Fig. 5. Daily scale variation of the frequency distribution of +CG and -CG stroke. The gray line represents**  
 329 **the proportion of +CG stroke. The time zone is CST (UTC+8 h).**

330 The hourly distribution of the peak current value and its average is illustrated in  
 331 Fig. 6. Region-II and Region-III, situated at higher latitudes, exhibit a broader range of  
 332 peak current distribution. Despite this, their variation remains relatively stable, with a  
 333 slightly larger current for +CG stroke in Region-III than Region-II. In Region-1, there  
 334 is a significant decrease in current during the midday and afternoon periods, with a

335 more pronounced change observed in +CG strokes than in -CG strokes. This  
 336 phenomenon leads to a situation where the absolute peak current of the two types of  
 337 stroke can even be reversed. Conversely, Region-IV, characterized by the intricate  
 338 terrain resulting from the Tibetan Plateau, showcases complex current variations.  
 339 Specifically, there is an increase in peak current for +CG strokes and a decrease for -  
 340 CG strokes since 8:00 CST in this region. Consequently, a substantial disparity in peak  
 341 current between the two types emerges around the daytime hours.




342  
 343 **Fig. 6. Daily scale variation of the peak current distribution of +CG and -CG stroke. The red line represents**  
 344 **the average peak current value of each hour. The time zone is CST (UTC+8 h).**

### 345 3.2.2 Comparison of spatial distribution of +CG and -CG stroke


346 The geographical features of China contribute to the complex distribution of +CG  
 347 and -CG strokes across various regions, which is reflected in the variability of the ratio  
 348 of +CG and -CG strokes. Fig. 7 illustrates the spatial distribution of the proportion of  
 349 +CG stroke, with gray areas indicating grids with less than 50 CG strokes accumulated  
 350 over a seven-year period. These grids are mainly located in the central, western, and  
 351 northern parts of Tibet and the western and southern parts of Xinjiang. Region-I, which  
 352 has the highest density of CG stroke, has a low proportion of +CG stroke, at less than  
 353 10 %. Conversely, the other three regions have a higher proportion of +CG stroke,  
 354 particularly the North China Plain and adjacent Inner Mongolia, as well as some parts  
 355 of Xinjiang, where the +CG proportion can reach 30-40 %. The proportion of +CG

356 stroke in Shanxi and Shaanxi, both located in Region-II, is lower than in other regions  
 357 of the same area. Overall, regions with lower CG stroke density tend to have a higher  
 358 proportion of +CG stroke, and high latitudes correspond to a higher +CG stroke  
 359 proportion.




360  
 361 **Fig. 7. Distribution of the proportion of +CG stroke in China. The gray grids have a CG stroke number of**  
 362 **less than 50 in seven years and thus are not calculated. The grid size is  $0.25^\circ \times 0.25^\circ$ .**

363 Based on Fig. 8, it can be inferred that the spatial distribution of the peak current  
 364 values for both +CG and -CG stroke is generally similar, with lower peak current values  
 365 observed in southeast China where lightning activity is more frequent, and higher peak  
 366 current values found in other inland areas. Notably, the peak current in Mêdog County  
 367 (o) is especially high, with most areas with an average of over 100 kA. Besides,  
 368 southern Gansu, southern Shaanxi, and the intersection of Guizhou, Hunan, and  
 369 Guangxi are also recognized as high-value regions, where the proportion of +CG stroke  
 370 is also relatively high. Therefore, it can be concluded that a high proportion of +CG  
 371 stroke typically corresponds to larger peak current values in terms of temporal and  
 372 spatial scales.



373



374

375  
376  
377


**Fig. 8. Distribution of the average peak current values of (a) -CG stroke and (b) +CG stroke in China. The gray grids have a -CG stroke or +CG stroke number of less than 50 in seven years and thus are not calculated. The grid size is  $0.25^\circ \times 0.25^\circ$ .**

378  
379  
380  
381  
382  
383  
384  
385

The proportion of +CG stroke in different altitude layers is calculated, as shown by the gray line in Fig. 9. Below 4500 meters altitude, the proportion increases with altitude, ranging from 7 % to over 15 %. A sub-peak is observed at 1500 meters, which is caused by the high proportion region of +CG strokes in Xinjiang and Inner Mongolia. However, above 4500 meters altitude, which mainly comprises the uninhabited areas of the western and northern Tibetan Plateau, the proportion of +CG stroke decreases rapidly. It is worth noting that only 91 CG strokes were observed above 6000 meters of altitude during the seven-year period and are not included in the statistics.

386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397

The box plot in Fig. 9 shows the peak current distribution of +CG and -CG strokes at different altitudes. The distribution of +CG peak current narrows with increasing altitude. Interestingly, the average peak current of -CG strokes shows a slight positive correlation with altitude, whereas +CG strokes exhibit a negative correlation with altitude. The opposite trend of the two types of lightning strokes leads to a large difference in their peak current at low altitudes and coincidence at high altitudes. The Tibetan Plateau is primarily responsible for the intricate lightning activity versus altitude over China. As the "third pole" of the Earth, the charge structure of thunderstorm clouds on the Tibetan Plateau always has some special characteristics due to the high-altitude ground surface (Li et al., 2013; Qie et al., 2005). Furthermore, its influence on the uplift and obstruction of water vapor can also affect the climatic characteristics of other regions.



398  
399

**Fig. 9. The peak current distribution of +CG and -CG stroke and the proportion of +CG versus altitude**

400

#### 4. Conclusion

401

China is primarily located in temperate and subtropical zones, with climate subject

---

402 to a variety of factors, including cold and warm monsoons, the interplay of land and  
403 sea, and varied topography. As a result, there are frequent convective weathers and a  
404 high prevalence of lightning activities. This paper utilizes the dataset from a ground-  
405 based lightning location system, CNLDN, which serves as the most extensively  
406 deployed national lightning detection system in China, to analyze the CG characteristics  
407 in China over the past seven years. The spatial and temporal distribution of +CG and -  
408 CG strokes exhibit regular patterns in terms of their frequency, proportion, and peak  
409 current.

410 The results indicate that there is more CG in southern regions than in northern  
411 regions, more in mountainous areas than in plains at the same latitude, more in humid  
412 areas than in arid areas, and more in coastal areas than in inland areas within the same  
413 climate zone. The southeast coastland of China has the highest CG stroke density, while  
414 the northwest deserts and basins as well as the east and north Tibetan Plateau have the  
415 lowest density. The monsoon system plays a critical role in lightning activities in  
416 southern and Northern China, while the Tibetan Plateau contributes to the variability of  
417 lightning activities in Northwestern China and the Qinghai-Tibet region. Overall, the  
418 distribution of lightning activity across China is consistent with the precipitation  
419 distribution observed at a climatic scale.

420 In general, +CG strokes have a lower occurrence rate than -CG strokes, but they  
421 carry higher currents and are more destructive. The spatial and temporal distribution of  
422 +CG and -CG stroke also varies significantly due to different storm structures. In terms  
423 of **annual** scale, the lightning activity follows a seasonal pattern, with the highest  
424 frequency occurring during summer (72.6 %), followed by spring (18.4 %) and autumn  
425 (8.6 %), and the least frequent in winter (0.4 %). In spring, autumn, and winter,  
426 lightning is mainly concentrated in the southeastern coastal areas. The percentage of  
427 +CG stroke is always inversely correlated with lightning frequency. High stroke  
428 frequency in summer generally corresponds to a low proportion of +CG stroke, while  
429 low frequency in winter corresponds to a high proportion of +CG stroke. **The proportion**  
430 **of +CG stroke in December in most regions reaches up to 50 %**. The average peak  
431 current of return stroke is strongly correlated with the proportion of +CG stroke and  
432 also follows a seasonal pattern of being high in winter and low in summer. The seasonal  
433 fluctuations of +CG stroke are stronger than -CG stroke in Southern China and Northern  
434 China, where the average intensity of +CG stroke in summer is even below -CG stroke.  
435 On the hourly scale, lightning is active in the late afternoon, with a peak between **15:00-**  
436 **17:00 CST**, and drops to a trough the following day between 8:00-10:00 CST. The  
437 proportion of +CG stroke throughout the day follows an inverse trend with the  
438 frequency of stroke in most regions, but the minimum proportion occurs 2-3 hours  
439 earlier than the maximum frequency. The highest proportion of +CG stroke at low  
440 latitudes always occurs in the morning, while at high latitudes, it tends to occur at  
441 midnight. The changes in peak current during the day at high latitudes are not  
442 significant. In Southern China, the peak current of +CG and -CG stroke drops  
443 significantly at noon and afternoon, with +CG dropping even lower than -CG.



---

444 The distribution of the +CG stroke proportion exhibits significant spatial  
445 variability. In Southern China, where the density of CG is the highest, the +CG stroke  
446 proportion is the lowest, at less than 10 %. In contrast, the high-latitude regions such as  
447 the North China Plain, Inner Mongolia, and northern and central Xinjiang have a much  
448 higher proportion of 30-40 %. The proportion of +CG stroke below 4500 meters is  
449 positively correlated with altitude and drops sharply after exceeding 4500 meters in the  
450 western and northern regions of the Tibetan Plateau. The spatial distribution of peak  
451 current values of +CG and -CG stroke is consistent, and a higher proportion of +CG  
452 stroke is generally associated with greater peak current for both types. As latitude  
453 increases, the current distribution widens. The peak current of +CG shows a slight  
454 decrease with increasing altitude, while the peak current of -CG increases with altitude.  
455 Consequently, there is a significant difference in discharge intensities between the two  
456 types at low altitudes, but they tend to be similar at higher altitudes.

457 The lightning location system sites cannot be evenly distributed due to geographic  
458 factors, thus bringing about errors in lightning distribution analysis. The observation  
459 from the Lightning Mapping Imager (LMI) on the FY-4A satellite will be used to correct  
460 the distribution deviations by ground-based data in our following research. Given the  
461 vast size of China, a simple division into four regions may be too crude to study the  
462 influence of geographic and climatic factors on CG characteristics in depth. Therefore,  
463 a more detailed division will be necessary for future studies.

## 464 Acknowledgments:

465 This study is supported by the Key Technologies Research and Development  
466 Program of China (2020YFB1600103). We appreciate the Meteorological Observation  
467 Centre of CMA and the Institute of Electrical Engineering of CAS for their data support.  
468 We also thank the reviewers and editors for their valuable suggestions for this study.

## 469 Reference

470 Carey, L. D. and Buffalo, K. M.: Environmental control of cloud-to-ground lightning polarity in  
471 severe storms, *Monthly Weather Review*, 135, 1327---1353, 2007.

472 Cummins, K. L., Murphy, M. J., Bardo, E. A., Hiscox, W. L., Pyle, R. B., and Pifer, A. E.: A  
473 combined TOA/MDF technology upgrade of the US National Lightning Detection Network, *Journal of*  
474 *Geophysical Research: Atmospheres*, 103, 9035-9044, 1998.

475 Diendorfer, G.: Some comments on the achievable accuracy of local ground flash density values,  
476 2008.

477 Holle, R. L. and Cummins, K. L.: Monthly distributions of US NLDN cloud-to-ground lightning,  
478 3rd International Lightning Meteorology Conference, 21-22,

479 Jiang, R.: Observation and Simulation of CG Lightning Activity Characteristics in the Regions with  
480 Tall Structures, University of Chinese Academy of Sciences, 2021.

481 Jin, H., Chen, X., Wu, P., Song, C., and Xia, W.: Evaluation of spatial-temporal distribution of

---

482 precipitation in mainland China by statistic and clustering methods, *Atmospheric Research*, 262, 105772,  
483 2021.

484 Koehler, T. L.: Cloud-to-ground lightning flash density and thunderstorm day distributions over the  
485 contiguous United States derived from NLDN measurements: 1993–2018, *Monthly Weather Review*,  
486 148, 313-332, 2020.

487 Li, Y., Zhang, G., Wen, J., Wang, D., Wang, Y., Zhang, T., Fan, X., and Wu, B.: Electrical structure  
488 of a Qinghai–Tibet Plateau thunderstorm based on three-dimensional lightning mapping, *Atmospheric*  
489 *Research*, 134, 137-149, 2013.

490 Ma, M., Tao, S., Zhu, B., and Lü, W.: Climatological distribution of lightning density observed by  
491 satellites in China and its circumjacent regions, *Science in China Series D: Earth Sciences*, 48, 219-229,  
492 2005.

493 Ma, R., Zheng, D., Zhang, Y., Yao, W., Zhang, W., and Cuomu, D.: Spatiotemporal Lightning  
494 Activity Detected by WWLLN over the Tibetan Plateau and Its Comparison with LIS Lightning, *Journal*  
495 *of Atmospheric and Oceanic Technology*, 38, 511-523, 2021.

496 Min-Xue, F., Yu, Z., Min, Y. U., and Yan-Hua, Z.: Evaluation of Jiangsu ADTD Lightning Location  
497 System Data in 2013, *Science Technology and Engineering*, 2015.

498 Nag, A., Rakov, V. A., and Cummins, K. L.: Positive Lightning Peak Currents Reported by the U.S.  
499 National Lightning Detection Network, *IEEE Transactions on Electromagnetic Compatibility*, 56, 404-  
500 412, 2014.

501 Poelman, D., Schulz, W., Diendorfer, G., and Bernardi, M.: The European lightning location system  
502 EUCLID-Part 2: Observations, *Natural Hazards & Earth System Sciences Discussions*, 3, 2015.

503 Preston and Tolver, S.: The lightning discharge, *Philosophical Magazine Series 1*, 31, 443-445, 1989.

504 Qie, X., Toumi, R., and Zhou, Y.: Lightning activity on the central Tibetan Plateau and its response  
505 to convective available potential energy, *Chinese Science Bulletin*, 48, 296-299, 2003.

506 Qie, X., Zhang, T., Chen, C., Zhang, G., Zhang, T., and Wei, W.: The lower positive charge center  
507 and its effect on lightning discharges on the Tibetan Plateau, *Geophysical research letters*, 32, 2005.

508 Rakov, V. A. and Uman, M. A.: *Lightning: Physics and Effects*, Lightning: Physics and Effects 2003.

509 Ren, G., Ding, Y., Zhao, Z., Zheng, J., Wu, T., Tang, G., and Xu, Y.: Recent progress in studies of  
510 climate change in China, *Advances in Atmospheric Sciences*, 29, 958-977, 2012.

511 Rudlosky, S. D. and Fuelberg, H. E.: Pre-and postupgrade distributions of NLDN reported cloud-  
512 to-ground lightning characteristics in the contiguous United States, *Monthly Weather Review*, 138, 3623-  
513 3633, 2010.

514 Rudlosky, S. D. and Shea, D. T.: Evaluating WWLLN performance relative to TRMM/LIS,  
515 *Geophysical Research Letters*, 40, 2344-2348, 2013.

516 Said, R., Cohen, M., and Inan, U.: Highly intense lightning over the oceans: Estimated peak currents  
517 from global GLD360 observations, *Journal of Geophysical Research: Atmospheres*, 118, 6905-6915,  
518 2013.

519 Schulz, W., Diendorfer, G., Pedeboy, S., and Poelman, D. R.: The European lightning location  
520 system EUCLID–Part 1: Performance analysis and validation, *Natural Hazards and Earth System*  
521 *Sciences*, 16, 595-605, 2016.

522 Shao, X.-M., Stanley, M., Regan, A., Harlin, J., Pongratz, M., and Stock, M.: Total lightning  
523 observations with the new and improved Los Alamos Sferic Array (LASA), *Journal of Atmospheric and*

---

524 Oceanic Technology, 23, 1273-1288, 2006.

525 Srivastava, A., Tian, Y., Qie, X., Wang, D., Sun, Z., Yuan, S., Wang, Y., Chen, Z., Xu, W., and Zhang,  
526 H.: Performance assessment of Beijing Lightning Network (BLNET) and comparison with other  
527 lightning location networks across Beijing, Atmospheric Research, 197, 76-83, 2017.

528 Wang, J., Huang, Q., Ma, Q., Chang, S., He, J., Wang, H., Zhou, X., Xiao, F., and Gao, C.:  
529 Classification of VLF/LF Lightning Signals Using Sensors and Deep Learning Methods, Sensors (Basel,  
530 Switzerland), 20, 2020.

531 Williams, E. R.: Large - scale charge separation in thunderclouds, Journal of Geophysical Research:  
532 Atmospheres, 90, 6013-6025, 1985.

533 Wu, S., Weitao, L., Qi, Q., Wu, B., Chen, L., Su, Z., Jiang, R., and Zhang, C.: Characteristics of  
534 Downward Cloud-to-ground Lightning Flashes Around Canton Tower Based on Optical Observations,  
535 Journal of Applied Meteorological Science, 2019.

536 Xia, R., Zhang, D.-L., and Wang, B.: A 6-yr cloud-to-ground lightning climatology and its  
537 relationship to rainfall over central and eastern China, Journal of Applied Meteorology and Climatology,  
538 54, 2443-2460, 2015.

539 You, J., Zheng, D., Zhang, Y., Yao, W., and Meng, Q.: Duration, spatial size and radiance of lightning  
540 flashes over the Asia-Pacific region based on TRMM/LIS observations, Atmospheric Research, 223, 98-  
541 113, 2019.

542