



1 **Brief communication on the NW Himalayan towns; slipping towards potential disaster**

2 Yaspal Sundriyal¹, Vipin Kumar², Neha Chauhan¹, Sameeksha Kaushik¹, Rahul Ranjan³, Mohit
3 K. Punia⁴

4 ¹Department of Geology, HNB Garhwal University, Srinagar, India

5 ²Department of Geology, Doon University, Dehradun, India

6 ³Oslo Metropolitan University, Oslo, Norway

7 ⁴National Geotechnical Facility, Dehradun, India

8 *Correspondence: v.chauhan777@gmail.com

9 **ABSTRACT**


10 The NW Himalaya has been one of most affected terrains of Himalaya subjected to frequent
11 disastrous landslides owing to active tectonics and multiple precipitation sources. This article
12 aims at two towns (Joshimath and Bhatwari) of the Uttarakhand in the NW Himalaya (India),
13 which have been witnessing subsidence for decades. In the last 1-2 weeks, Joshimath has
14 witnessed widespread cracks in more than 500 houses that has created the social unrest. The
15 hillslopes accommodating both the towns comprise highly jointed gneisses with schistose
16 interlayers rockmass, subsidence in road, broken retaining wall, holes, displacing boulders, and
17 cracks in the houses. Recently, such slope instability phenomena have increased that is leading to
18 social movements in the region seeking government action for possible evacuation and
19 rehabilitation. Present study has involved continuum modeling-based slope stability simulation
20 to determine the response of these hillslopes under various loading conditions; gravity, rainfall,
21 building load, domestic discharge, and seismic load. Results revealed that the displacement in
22 these hillslopes might reach up to 20-25 m that will further aggravate the situation. Occurrence
23 of frequent extreme rainfalls in these towns and three major earthquakes i.e., 1 Sep. 1803
24 (M_w 7.8), 20 Oct. 1991 (M_w 6.8), and 29 Mar. 1999 (M_w 6.6) having hypocentral distance less
25 than 30 km make such study more viable for decision making.

26 **Keywords:** NW Himalaya; Landslides; Human Population; Slope Stability

27 The NW Himalaya has been one of most affected terrains of Himalaya subjected to frequent
28 disastrous landslides (Martha et al., 2015; Gupta et al. 2017; Kumar et al. 2021b). Continent-
29 collisional orogeny caused active tectonics and multiple precipitation sources i.e., Indian
30 Summer Monsoon (ISM) and Western Disturbance caused extreme rainfall are main causes of



1 such landslides (Petley, 2010; Dimri et al. 2015). The NW Himalaya has accommodated ~51 %
2 of all the landslides in India during the years 1800-2011 (Parkash 2011). This article aims at two
3 towns (Joshimath and Bhatwari) of the Uttarakhand in the NW Himalaya, which have been
4 witnessing subsidence for decades. Notably, Uttarakhand has been subjected to four known
5 major flood events in last 5 decades, which are as follows; 24-26th August 1894, 20th July 1970,
6 16-17th June 2013, 7th Feb. 2021 (Ziegler et al. 2014; Sundriyal et al. 2015). Apart from floods,
7 three major earthquakes have also caused widespread damage in the region that are as follows; 1
8 Sep. 1803 (M_w 7.8), 20 Oct. 1991 (M_w 6.8), and 29 Mar. 1999 (M_w 6.6) (Bilham, 2019).
9 Joshimath town (Fig. 1) had a human population of 16709 as per last census (2011) of Govt. of
10 India and might have reached at least 18630 as per 11.5 % population growth
11 (<https://main.mohfw.gov.in/>, data retrieved on 10/08/2022).





1 **Fig. 1.** Field observation of Joshimath town. (a) Location of Joshimath town with inset highlighting the
2 position in the NW Himalaya. CS 1 to CS 4 mark the position of 2D slope sections used for the slope
3 stability evaluation; (b) Rockmass exposure; (c) Natural drainage (Nala) through hillslope. Position of
4 soil sampling site (S-1); (d) Cracks and displacement of road; (e) Bulging and failure of retaining wall;
5 (f-g) Displacing hillslope material and hence pavements; (h-i) Cracks in houses. Picture (f-i) courtesy:
6 Mr. Atul Sati.

7 This town has witnessed widespread cracks in more than 500 houses, even in temples in the last
8 1-2 weeks that is creating the social unrest. Various media reports involving interviews of
9 experts aim at the potential role of infrastructural projects in promoting instability in this region
10 ([https://timesofindia.indiatimes.com/city/dehradun/uttarakhand-joshimath-land-subsidence-](https://timesofindia.indiatimes.com/city/dehradun/uttarakhand-joshimath-land-subsidence-cracks-sinking-news-live-updates-8-january-2023/liveblog/96843072.cms)
11 [cracks-sinking-news-live-updates-8-january-2023/liveblog/96843072.cms](https://timesofindia.indiatimes.com/city/dehradun/uttarakhand-joshimath-land-subsidence-cracks-sinking-news-live-updates-8-january-2023/liveblog/96843072.cms)). The town is situated
12 along a narrow gorge at the confluence of two major rivers; Dhauliganga and Alaknanda. The
13 gorge owes its existence to exhumation along Main Central Thrust (MCT) fault passing through
14 the southern side of town (Supp. Fig. 1). This orographic barrier results in ~3672 mm/month
15 rainfall in the town (Supp. Fig. 2). In a recent study by Agarwal et al. (2022), the town belongs to
16 a hotspot, which is experiencing changing climate and rapid anthropogenic influence. Field
17 observation of the town revealed that hillslope of town comprises highly jointed gneisses with
18 schistose interlayers rockmass, subsidence in road, broken retaining wall, holes, displacing
19 boulders, and cracks in the houses (Fig. 1). Loose hillslope material comprising sandy-gravel soil
20 (Fig. 1c), exposed in the eastern part of town points towards the possibility of paleo-landslide on
21 which most of the town is located. The last big earthquake (29 Mar. 1999: M_w 6.6) occurred
22 within 26 km hypocentral distance from this town.

23 Another town that is included in this article is Bhatwari (Fig. 2). As per census 2011, it had a
24 population of 1268 that is 1.025 % lower than that of year 1991. Though population migration
25 from hilly terrain to plain area is becoming common phenomena in the NW Himalaya owing to
26 socio-economic priorities, decreased human population of this town might be due to such slope
27 instability problems also. Similar to the Joshimath town, Bhatwari is also situated along a narrow
28 gorge made by Bhagirathi River valley. The hillslope of the town comprises gneissic rockmass
29 and the MCT-I (locally termed as Vaikrita Thrust: VT) and MCT-II (locally termed as Munsiyari
30 Thrust: MT) faults pass through the northern and southern side, respectively of this region.
31 (Supp. Fig. 1). Two tributaries merging into Bhagirathi River bound the hillslope from northern
32 and southern side. Bhatwari town is also situated in the orographic barrier setting and hence



1 receives ~3732 mm/month rainfall (Supp. Fig. 2). Field observation revealed exposed scarps in
2 the upper part of town, cracks in the road, and tilting houses (Fig. 2b-h). Big (4-5 ft.) boulders
3 and active slope failure in the loose material at the slope toe indicate the possibility of paleo-
4 landslide on which Bhatwari town is located (Fig. 2i-j). Yadav et al. (2020) have also noted the
5 material displacement in the range of 12-22 mm/yr. during the years 2006-2016 at this location.
6 The last big earthquakes (1 Sep. 1803: M_w 7.8, 20 Oct. 1991: M_w 6.8) occurred within 22 km
7 hypocentral distance from this town.



8 **Fig. 2.** Field observation of Bhatwari town. (a-b) Location of Bhatwari town with inset highlighting the
9 position in the NW Himalaya. CS 1 to CS 4 mark the position of 2D slope sections used for the slope
10 stability evaluation; (c-d) Exposed scarp. S-1 refers to soil sampling site; (e) Cracks and displacement of
11 road; (f-g) Tilting houses/shops; (h) Active slope failure at slope toe; (i) Big Gneissic boulder implying
12 paleo-landslide site.

13 These two towns were selected because both of these are subjected to similar litho-tectonic
14 conditions and precipitation pattern (owing to orographic barrier setting). Since both of these
15 locations comprise surficial cracks, displacing boulders, exposed scarp, possibility of the
16 existence of paleo-landslide cannot be ignored. Though a detailed geophysical survey is being
17 planned for future prospects to understand the subsurface regime at these locations, present study
18 has involved continuum (Finite Element Method) based slope stability simulation.




1 This simulation was performed to determine the response of hillslopes that accommodate these
2 towns, under various loading conditions; gravity, rainfall, building load, domestic discharge, and
3 seismic load. Rainfall infiltration (RF) is based on extreme rainfall of 122 mm/day (18th Oct.
4 2021) and 124 mm/day (26th July 2010) at the Joshimath and Bhatwari, respectively (Supp. Fig.
5 2. Data source: GPM IMERG Final Precipitation (Huffman et al., 2020), spatial resolution: ~ 1
6 km). Domestic discharge (DD) infiltration and Building load (B) are based on the Indian
7 Standard (IS) code: 2470 (Part 1)-1985 and Indian Standard (IS) code: 875 (Part 2)-1987,
8 respectively. Domestic discharge refers to liquid waste of a house, whereas building/house load
9 refers to uniformly distributed load of houses. Notably, domestic discharge and building load are
10 consistent in nature, whereas the rainfall infiltration is relatively random. Seismic loads are based
11 on 20 Oct. 1991 (Mw 6.8) Uttarkashi earthquake and 29 Mar. 1999 (Mw 6.6) Chamoli
12 earthquake acceleration history (Supp. Fig. 3. Data source: Dept of Earthquake Eng., Indian Inst.
13 of Technology, Roorkee, India through COSMOS web portal, <http://www.cosmos-eq.org/>,
14 retrieved on 03 Aug. 2022) used for Bhatwari and Joshimath, respectively. This simulation
15 method has been used for various hillslopes of the NW Himalaya owing to flexibility in slope
16 geometry solution and acceptable approximation of displacement output (Kumar et al. 2021b).
17 Details of slope configuration, material models, and input parameters have been summarized in
18 Supplementary Annexure I. For both the towns, four 2D slope sections (CS 1 to CS 4) were
19 taken to minimize the uncertainty caused by subjectivity associated with single slope section
20 (Fig. 1, 2). A configuration model is shown in Supp. Fig. 4. Though 3D slope stability could
21 have been more informative, subsurface regime is yet to be explored in future prospects.

22 Stability evaluation results for both the locations are discussed using the total (horizontal &
23 vertical component) displacement in the top 5-10 m material layer along the slope surface.
24 Displacement profiles for Joshimath revealed that slope may yield 1-4 m displacement under
25 gravity conditions that might increase up to 6 m under combined effect of rainfall (RF)
26 infiltration, building/house load (B), and domestic discharge (DD) (CS 1 JM to CS 4 JM in Fig.
27 3). The CS 1 and CS 3 sections of Joshimath town yield relatively higher displacement under
28 these loads, particularly near slope toe. Under seismic load conditions, displacement increase
29 rapidly in the range of 8-22 m, particularly at the knickpoints of slope surface and near the slope
30 toe. Relatively higher seismic displacement at the slope toe is attributed to stress accumulation at
31 slope toe and narrow geomorphic setting resulting in constructive seismic interference, as also



1 noted in other case studies involving earthquake induced landslides Meunier et al. (2008).
2 Notably, slope crest region comprises debris layer that must have affected topographic
3 amplification and hence relatively lower displacement, as also noted in recent study in the SE
4 Carpathians (Kumar et al. 2021a).

5 **Fig. 3.** Results of the
6 Slope stability
7 evaluation. CS 1 JM to
8 CS 4 JM refer to
9 Joshimath Slope
10 sections, whereas CS 1
11 BT to CS 4 BT refer to
12 slope sections of
13 Bhatwari town hillslope.
14 Dn, RF, B, DD, and St
15 refer to Dynamic,
16 Rainfall infiltration,
17 Building load, Domestic
18 discharge, and Static
19 load, respectively



20
21
22
23
24
25
26
27



1 Similar to Joshimath town hillslope, displacement profiles for the Bhatwari town hillslope show
2 dominance of higher displacement (up to ~25 m) at the slope toe under seismic load conditions
3 (Fig. 3). Notably, under combined effect of rainfall, domestic discharge, and building load, CS 1
4 and CS 2 sections of Bhatwari town slope comprises pockets of higher displacement (5-20 m),
5 particularly near slope toe. This could be attributed to thicker debris deposit and cluster of human
6 settlement at the location of these slope sections (CS 1 and CS 2).

7 Field observations of surface cracks, tilting houses, displacing hillslope material and predictive
8 findings of more displacement in case of local (domestic discharge, building load) and external
9 loads (rainfall, earthquake) emphasize the critical situation of such NW Himalayan towns.
10 Notably, recent social movements in Joshimath town seeking government action for possible
11 evacuation, rehabilitation, and suspension of infrastructural projects (assumed as a major reason
12 of hillslope instability) (<https://timesofindia.indiatimes.com/city/dehradun/joshimath-sinking-thousands-of-fearful-residents-hit-road-in-protest/articleshow/96488198.cms>) make such study
13 more viable for decision making.
14

15 As far as the cause of slope instability in Joshimath is concerned, a committee report (No.
16 142/28-5/44/76 dated 8th April 1976) submitted jointly by 17 experts headed by Late M. Mishra
17 considered the paleo-landslide material in this region as a major cause of instability. Google
18 Earth imagery of June, 2014 (Supp. Fig. 5) also indicates growing slope toe failure, possibly
19 leading to retrogression. However, contribution of anthropogenic impacts involving population
20 increase, continuing rapid construction to cater high tourist influx (Agarwal et al. 2022) and
21 infrastructural projects can't be denied. Unlike Joshimath, Bhatwari town is witnessing
22 population decrease and lack of tourist influx. Therefore, natural conditions i.e., loose material of
23 paleo-landslide coupled with excessive rainfall dominate as the causative factors.

24 Notably, Joshimath and Bhatwari towns occupy a cornerstone to the religious and political
25 landscape of the NW Himalaya. With the emergent threats of cracks, this paper suggests a need
26 for a robust study to develop an integrated approach involving detailed scientific study and
27 people's grievances against/about the development projects. Further, there is an urgent need to
28 develop an integrative framework that combines the social composition of the region, various
29 forms of resilience/adaptation practices and robust scientific inquiries. Often it is assumed that



1 scientific factors that influence landscape are well understood and explained to the public. This
2 invariably miscommunicates the potential threat causing irreversible socio-economic loss.

3 **Code availability.** Slope stability analysis simulations are performed using the RS2 (v.11.012)
4 software (Rocscience Inc.). Figures are prepared using CorelDRAW 2021 (commercial software)
5 and JASP v.0.15 (open source).

6 **Data availability.** Rainfall datasets are openly accessible at
7 <https://giovanni.gsfc.nasa.gov/giovanni/> (NASA, 2022). The earthquake acceleration history is
8 available at <http://www.cosmos-eq.org/>.

9 **Author contribution:** YP, VK conceived the idea in view of increasing social problems at these
10 sites. VK, NC, and SK performed the field work and collected soil samples. MKP performed the
11 laboratory analysis. VK, NC, MP and SK interpreted the laboratory data and performed the
12 numerical simulation. RR performed a social anthropological survey. All authors contributed to
13 the writing of the final draft.

14 **Competing interests:** The authors declare that they have no conflict of interest.

15 **Financial support:** Authors are thankful for the financial support by the Department of Science
16 and Technology, Government of India, New Delhi [Ref. No. DST/CCP/MRDP/187/2019(G)].

17 ACKNOWLEDGEMENT

18 Authors are thankful to Mr. Atul Sati (Social activist) and Prof. S.P. Sati for the constructive
19 discussion and Misra Committee (1976) report, respectively.

20 REFERENCES

- 21 Agarwal, S., Kumar, V., Kumar, S., Sundriyal, Y., Bagri, D. S., Chauhan, N., ... & Rana, N. (2022).
22 Identifying potential hotspots of land use/land cover change in the last 3 decades, Uttarakhand, NW
23 Himalaya. doi.org/10.31223/X5VK9F.
- 24 Bilham, R. (2019). Himalayan earthquakes: a review of historical seismicity and early 21st century slip
25 potential. Geological Society, London, Special Publications, 483(1), 423-482.
- 26 Dimri, A. P., Niyogi, D., Barros, A. P., Ridley, J., Mohanty, U. C., Yasunari, T., & Sikka, D. R. (2015).
27 Western disturbances: a review. Reviews of Geophysics, 53(2), 225-246.
- 28 Gupta, V., Jamir, I., Kumar, V., & Devi, M. (2017). Geomechanical characterisation of slopes for
29 assessing rockfall hazards in the Upper Yamuna Valley, Northwest Higher Himalaya, India.
30 Himalayan Geology, 38(2), 156-170.



- 1 Huffman, G. J., Bolvin, D. T., Braithwaite, D., Hsu, K. L., Joyce, R. J., Kidd, C., ... & Xie, P. (2020).
- 2 Integrated multi-satellite retrievals for the global precipitation measurement (GPM) mission
- 3 (IMERG). In *Satellite precipitation measurement* (pp. 343-353). Springer, Cham.
- 4 Kumar, V., Cauchie, L., Mreyen, A. S., Micu, M., & Havenith, H. B. (2021a). Evaluating landslide
- 5 response in a seismic and rainfall regime: a case study from the SE Carpathians, Romania. *Natural*
- 6 *Hazards and Earth System Sciences*, 21(12), 3767-3788.
- 7 Kumar, V., Jamir, I., Gupta, V., & Bhasin, R. K. (2021b). Inferring potential landslide damming using
- 8 slope stability, geomorphic constraints, and run-out analysis: a case study from the NW Himalaya.
- 9 *Earth Surface Dynamics*, 9(2), 351-377.
- 10 Martha, T. R., Roy, P., Govindharaj, K. B., Kumar, K. V., Diwakar, P. G., & Dadhwal, V. K. (2015).
- 11 Landslides triggered by the June 2013 extreme rainfall event in parts of Uttarakhand state, India.
- 12 *Landslides*, 12(1), 135-146.
- 13 Meunier, P., Hovius, N., & Haines, J. A. (2008). Topographic site effects and the location of earthquake
- 14 induced landslides. *Earth and Planetary Science Letters*, 275(3-4), 221-232.
- 15 Parkash, S. (2011). Historical records of socio-economically significant landslides in India, *J. S. Asia*
- 16 *Disast. Stud.*, 4, 177-204.
- 17 Petley, D. N. (2010). On the impact of climate change and population growth on the occurrence of fatal
- 18 landslides in South, East and SE Asia. *Quarterly Journal of Engineering Geology and*
- 19 *Hydrogeology*, 43(4), 487-496.
- 20 Sundriyal, Y. P., Shukla, A. D., Rana, N., Jayangondaperumal, R., Srivastava, P., Chamyal, L. S., ... &
- 21 Juyal, N. (2015). Terrain response to the extreme rainfall event of June 2013: Evidence from the
- 22 Alaknanda and Mandakini River Valleys, Garhwal Himalaya, India. *Episodes Journal of*
- 23 *International Geoscience*, 38(3), 179-188.
- 24 Yadav, R. K., Gahalaut, V. K., Gautam, P. K., Jayangondaperumal, R., Sreejith, K. M., Singh, I., ... &
- 25 Sati, S. P. (2020). Geodetic Monitoring of Landslide Movement at two sites in the Garhwal
- 26 Himalaya. *Himalayan Geology*, 41(1), 21-30.
- 27 Ziegler, A. D., Wasson, R. J., Bhardwaj, A., Sundriyal, Y. P., Sati, S. P., Juyal, N., ... & Saklani, U.
- 28 (2014). Pilgrims, progress, and the political economy of disaster preparedness—the example of the
- 29 2013 Uttarakhand flood and Kedarnath disaster. *Hydrological Processes*, 28(24), 5985-5990.