# <u>Invited perspectives:</u> <u>Current challenges to face knowns and</u> unknowns in natural hazard risk modelling: an insurer perspective<u>An</u> insurer's perspective on the knowns and unknowns to face in natural hazard risk modelling

Madeleine-Sophie Déroche<sup>1</sup>
 <sup>1</sup>AXA Group Risk Management, Paris, 75008, France
 *Correspondence to*: Madeleine-Sophie Déroche (madeleinesophie.deroche@axa.com)

Abstract. All a long the Property and Casualty (P&C) insurance value chain, a wide vast range of data feed an equally wide vast range of models to estimate the losses for the varying probabilities and magnitudes of all the underwritten event risks, be

- 10 they natural hazards, financial, cyber. In effect, beyond reflecting and supporting the current understanding and knowledge of risks, these data and models reflect and support the assessment of situations that have yet to be experienced. These data and models in effect reflect and support the current understanding and knowledge of risks as well as the assessment of situations that have yet to be experienced. The (re)insurance market's current body of knowledge on natural hazards loss modelling results from over 30 years' research involving private companies like (re)insurers, brokers, and modelling firms and academia
- 15 [OR] academics/academic researchers in various disciplines (atmospheric sciences, geosciences, civil engineering studies, and data sciences, to name <u>but</u> a few <u>disciplines</u>) involving academic researchers and private companies like (re)insures, brokers and modelling firms. This paper highlights the need to conduct an in-depth review of the existing loss modelling framework created in the early 1990s to capture the increasing complexity of the three risk drivers – exposure, hazard, and vulnerability – as well as their interconnections.

## 20 1 Introduction

25

The mission of Property and Casualty (P&C) insurers is to effectively protect clients' property and activities while ensuring the solvency of their company. Even though insurers develop an ever-increasing number of products to respond to specific client needs, P&C insurance products fall underare composed of two business segments: individual insurance or the retail business for home or car owners and corporate insurance or <u>c</u>Commercial business for corporate clients. Insurance protection goes beyond risk transfer (i.e., the payment of a premium against future claims); it also encompasses prevention actions such

as reinforcing customers' risk a wareness and proposing a dapted protective solutions.

For example, in <u>c</u>-ommercial business, technical risk experts perform on-site visits to evaluate the state of buildings and identify potential vulnerabilities to natural hazards. The objective is to assess how natural hazards could generate damage

either to the buildings themselves (e.g. storage warehouses, data centers, shopping centers) or to their contents (e.g. machinery,

- 30 production chains, stock), and if such damage could cause business interruption (e.g. employees / clients being unable to access the building for N days resulting in a loss of turnover or profits). Prevention measures like elevating goods or machinery in the event of flooding are then suggested or imposed post-assessment to reduce the risk and adjust the premium. In the retail business with its mass of clients, protection actions have to be taken globally instead of individually. For
- 35 insurance premium rates were lowered for buildings made of brick to encourage brick constructions instead of wood thus reducing the fire risk in London.

example, after the Great Fire of London in 1666 that destroyed most of the city's buildings, made of wood at that time,

To achieve their mission of protection, it is essential for insurers to identify and quantify the risks associated with the underwritten policies. All along the P&C insurance value chain, a vast range of data feed an equally vast range of models to estimate the losses for the varying probabilities and magnitudes of all the underwritten risks

- 40 , be they natural hazards, financial, or cyber. These models serve to support decision-making from the actual underwriting and , be they natural hazards, financial, or cyber. These models serve to support decision-making from the actual underwriting and pricing of an individual or corporate's policy to the setting and optimization of the reinsurance programs at the insurance company level. The regulatory environment also plays a significant role in validating the models (re)insurers use to assess risks. Regulations require (re)insurers to assess the extreme losses of all their risks to determine their
- 45 minimum level of economic capital to ensure the (re)insurance companies' solvency in the event of intensely severe years. The European Solvency 2 regulation is a case in point. (Re)insurers with Europe-based headquarters are required to annually project their losses for a 200-year return period shock along with the associated risk management actions such as the purchasing of reinsurance covers. This estimated amount of loss determines the level of capital (re)insurers have to bear in their owned funds to resist such a shock if it were to occur in the following year. The models used to assess this loss require
- 50 approval and any change is monitored by regulatory authorities.

Sound and adaptive risk assessment and management are built over time through a continuous reassessment of insurers' understanding of: the "known knowns", what we know we know; the "known unknowns", what we know we do not know; and the "unknown unknowns", what we don't even know we do not know (Girard, 2009). This reassessment process induces a knowledge cycle: data supports the current understanding and knowledge of a risk, that is, what we know we know. On the

55 basis\_of this understanding, models are <u>also</u> built to support assessing situations that have yet to be experienced such as extreme events, that is, what we know we don't know. <u>However, the occurrence of unknown unknowns triggered</u> by natural and organizational issues that insurers either ignore or have yet to understand, points to a dire need to upgrade data collection, modelling methods and tools to perpetually enhance the view of risk and further insights for the decision-making process.

Prior to focusi	Prior to focusing on the scientific and technical advances made to keep refining what we know about the risk drivers, exposure.										
hazard, and vi	ulnerability	, and how to i	ncreas	se insurers	' prepared	iness fo	or the unkno	wn, it is impo	rtant to	recall	how the
reassessment	process	engendered	the	natural	hazard	loss	modeling	framework	from	the	outset.
		-					-				

65

# 2 Natural hazard modelling: a brief overview

# 2.1 The co-influence of (re)insurance market and natural hazard modelling

The actual assessing of natural event related costs has greatly evolved over the past 30 years. At first, so-called catastrophe models focused on the modelling of extreme losses to assess therisk of a portfolio (i.e. large ensembles of insured buildings).
Before the 1990s, catastrophe modelling consisted in extrapolating the loss experience to estimate extreme losses. This loss experience was usually limited and recorded as a total amount of loss per event and per insurer or per event and for the whole (re)insurance market. The data were thus too coarse to capture the three risk drivers' individual impact on the losses are generated by a storm surge versus wind in the case of a tropical cyclone), the vulnerability (e.g., how effective the flood)

- 75 defenses, building codes are). As a result, while the data and the resulting modelling failed to take into account individual effects when assessing extreme losses, it did reflect the state of what was known by insurers and public authorities at that time. Hurricane Andrew in 1992 and its unexpected impact was a game changer for modelling natural hazard-generated losses (Grossi et al. 2005; Mitchell-Wallace, 2017). According to McChristian (2012), before Hurricane Andrew, the loss assessment for an event of that strength was \$4 to \$5bn. This is 3 times lower than Hurricane Andrew's actual
- 80 loss at \$15 bn. Insurers underestimated their exposure as well as their exposure's vulnerability to such an event. McChristian (2012) also indicates that though recent loss history was adjusted to reflect current macro-economic trends, it has failed to capture the increasing population over coastal areas. In the aftermath of Hurricane Andrew, a collective realization grew for the need to both separately characterize the three drivers of the risk exposure, hazard and vulnerability and model their interconnections. Catastrophe modelling therefore evolved from a statistical extrapolation to a framework
- 85 divided into 4 components as shown in Figure 1; one component by risk driver (exposure, hazard and vulnerability), and one component that contains the insurance policies' financial conditions and its modelling. The occurrence of natural disasters, in particular those with a strong impact for the (re)insurance market, continues to feed research insofar that the research is in turn integrated into the hazard and vulnerability components of the loss modelling framework every 2 to 5 years. This is how, the successive 1999 occurrence of the two extreme European winter windstoms
- 90 Lothar and Martin triggered the introduction of the serial clustering effect in modelling the frequency of European winter windstorms (Mitchell-Wallace, 2017). The serial clustering effect refers to the higher probability that two extreme winter windstorms occur in a short period of time, under particular atmospheric conditions (Vitolo et al., 2009; Pinto et al., 2013; Priestley et al., 2017). Prior to these windstorms, the assumption used to calculate the occurrence probability of

a mis en forme : Non Surlignage

Code de champ modifié

3

1	European winter windstorms followed the Poisson distribution and thus failed to allow for the modelling of successive events.			
95	As shown by, (Priestley et al., 2018), the clustering effect has a significant impact on the estimation of yearly aggregated losses			
	and therefore on the sizing of reinsurance covers.			
	Within the reinsurancemarket, the use of catastrophe models - developed internally or licensed through third-party vendors -			
	has grown in the aftermath of Hurricane Andrew. For insurers to cede their risk they must provide their exposure information			
	to reinsurers so they can conduct a loss assessment prior to estimating the reinsurance premium corresponding to the accepted			
100	risk. Today, catastrophe models continue to be used primarily to set reinsurance programs.			
	The implementation of regulation has prompted insurers to use catastrophe models, mainly licensed by third-party vendors, as			
	tools to assess the risk, define the risk appetite, and set the solvency capital requirement. For example, the Solvency 2 regulation			
	implemented in Europe in 2016 requires (re)insurers with Europe-based headquarters to annually assess their loss			
	for a 200-year return period shock. (Re)Insurers conduct this assessment for all the risks they are			
105	exposed to. They then aggregate these estimated losses to determine the total potential loss and the			
	economic capital they have to bear in their owned funds. To achieve this assessment, (re)insurers have two options:			
	either to use the so-called Standard Formula, calibrated on market exposure, or to develop an internal view of their risk that			
	requires regulator approval. Most (re)insurers choosing to develop their own view of natural hazard risk use one or several			
	models licensed to third-party vendors; others develop their own suite of models. When (re)insurers opt for using third-party			
110	models, model evaluation becomes a necessary activity for assessing the model's strengths and limitations leads to gaining in			
	understanding and in taking ownership of the model. Model adjustments may be defined and applied to the models' loss			
	estimation to address identified limitations (e.g. a non-modeled peril). (Re)insurance companies a lso invest in the development			
	of in-house models either on scopes where no third-party vendors model is a vailable or to gain in flexibility and transparency.			
	In the past few years, both the scope and use of catastrophe models have evolved. Indeed, to estimate			
115	the insurance premium of an average risk, insurers are now as interested in capturing small frequent events as large rare ones			
	. As more and more data have been included in the			
	calibration of the catastrophe models, they do capture much more than just extreme loss events. In the hazard module, the full			
	spectrum of events (i.e. moderate/intense; frequent/rareevents) is considered. In the vulnerability module, vulnerability curves			
	cover the entire range of hazard intensity. As for the modelling scope, catastrophe models even exist for man-			
120	madeperils such as cyber and terrorism. To reflect these evolutions, we will use the term 'natural hazard models/modelling'			
	as it allows for greater precision on the model's targeted scope and reaffirms the use of these models for other purposes than			
	the analysis of extreme events.			
1	2.2 Natural hazard modelling framework			

# The loss modelling framework is composed of 4 components, namely exposure, hazard, vulnerability and financial components 125 (Figure 1). The description below provides a brief introduction. Greater details on the different components can be found in (Mitchell-Wallace, 2017).

Code de champ modifié

a mis en forme : Anglais (États-Unis)

The exposure component contains the insurance portfolio's information: the buildings' location, and their key physical properties (e.g. structure, occupancy, year of construction...). The hazard component contains a synthetic catalogue of several tens of thousands of events that represent the range of possible and plausible events for a given natural hazard (e.g., Asia

130 typhoon, US ground shaking, Europe severe convective storms, ranging from small frequent events to extreme rare events. Each event is characterized by a footprint (i.e., the maximum intensity over event duration) and an annual occurrence probability. The vulnerability component is composed of vulnerability curves that translate the hazard's intensity into a building damage ratio. Ideally, there is one vulnerability curve for every combination of a building's physical properties. Finally, the financial component contains the insurance contract's financial data: the sum insured corresponding to the

135 coverage (building, content, business interruption), the deductibles and limits, as well as the coinsurance programs or reinsurance treaties, if any.

For every event of the hazard component and for every building in the insurance portfolio, the 3-step loss modelling process consists in:

Step 1. Intersecting the building's location with the event footprint to obtain the location's hazard intensity value.

Step 2. Taking into account the hazard intensity value and the building's physical properties and using the vulnerability curve reflecting the building's characteristics to derive the corresponding damage ratio.

Step 3. Applying the damage ratio to the insured value of the building, as specified in the financial module, to provide a loss amount prior applying the financial conditions to the loss amount to get the ultimate loss borne by the insurance company.

145

140

The primary outputs of natural hazard models are exceedance probability distributions representing the probability to exceed a certain amount of loss. Two distributions are commonly used: the one for the annual maximum loss - Occurrence Exceedance Probability (OEP)-, and the one for the annual aggregated loss - Aggregate Exceedance Probability (AEP). The Annual Average Loss (AAL) is also frequently used for budget planning for instance. Analyses of building losses are

150 aggregated at granularities going from the building level to the portfolio level to characterize the probability to exceed an amount of loss.\_This granularity is set in function of an analysis' objective, i.e. policy underwriting or portfolio management.

The loss modelling process is supported by a platform that contains (i) the data of each component stored in a specific format (e.g. csv or netcdf file, digital precision...) and (ii) the code functions that process data and estimate the losses. Until the early

- 155 2010s, the loss modelling process could only be performed on proprietary platforms. Launched in 2010, the OASIS initiative's ambition was to provide an open-source loss modelling platform to further transparency and to expand the use of natural hazard modelling beyond the (re)insurance market. From a business perspective, integrating such a process in daily operational activities requires the run time to take no more than a few hours. As an example of volumes at stake, assuming we have a catalogue of around 30,000 events and a portfolio
- of 5 million buildings, there will be 150 billion computations to run and as much data to be stored temporarily 160

А	a mis en forme : Non Surlignage
1	a mis en forme : Non Surlignage
1	a mis en forme : Non Surlignage
A	a mis en forme : Non Surlignage
-	a mis en forme : Non Surlignage
	a mis en forme : Non Surlignage
{	a mis en forme : Non Surlignage
$\supset$	a mis en forme : Non Surlignage
$\langle \rangle$	a mis en forme : Non Surlignage
Y	a mis en forme : Non Surlignage

in a constrained IT environment with limited storage space and a memory limit\_To keep to the expected run time and given the volumes considered, the loss modelling platform is to be rationalized and optimized, even if it results in a drop of formatting flexibility and data precision within the 4-components.

There is therefore a gap between the quality and the sophistication of the modelling produced by research and the derivative

- 165 data compiled in the loss modelling framework's 4-components. For example, the severity of natural events is captured in the hazard component through the use of hazard footprints defined as the maximum hazard value (e.g., windspeed, flood depth, peak ground acceleration) at each grid cell of the considered area over the duration of the event. The information relative to the event's duration and to the hazard value's evolution over time however are lost, even though both of these parameters affect the damage assessment of a building.
- 170 The 4-component loss modelling framework makes it easier to identify the areas where, component by component, a moreindepth investigation is needed to refine data collection and modelling. The next section focuses on three of the loss modelling framework's components where (i) a thorough and systematic data collection needs to be put in place, and (ii) the loss modelling framework requires investment to upgrade it and tailor it to respond to insurers' business needs.

175 3 Current challenges in modelling natural hazards

The (re)insurance market's current body of knowledge on natural hazards loss modelling results from over 30 years' research involving private companies like (re)insurers, brokers, and modelling firms and a cademic researchers in a tmospheric sciences, engineering studies, and data sciences, geosciences, civil to name but a few disciplines (Ward et al., 2020). The learning curve has been steep, closely linked to the increase of computer power (e.g., enabling the 180 (Ward et al., 2020). The learning curve has been steep, closely linked to the increase of computer power (e.g., enabling the al., 2020). The learning curve has been steep, closely (Ward et linked to the increase of computer power (e.g., enabling the development and implementation of millions of possible climatic or seismic scenarios) and the collection of increasingly granular observational data (e.g. hazard, claims, geocoded exposure).

185 A significant gap exists today between what research produces in terms of data and models and what insurers ultimately have access to through the natural hazard loss modelling framework. This reflects the gap between the loss modelling framework's initial goal\_providing aggregated loss estimates for extreme events-, and insurers current needs that is accurately assessing the impact of natural hazards on their business, by conducting sensitivity tests on modelling parameters to evaluate their impact on (re)insurance premiums for instance.

190 **3.1** 

Exposure

component

a mis en forme : Non Surlignage

a mis en forme : Non Surlignage

a mis en forme : Anglais (États-Unis)

a mis en forme : Non Surlignage

6

	Infough an increasing use of natural nazards models, insurers have realized boundata quality and its completeness reduces the
	uncertainty in the modelling. Over the past 5 to 10 years, insurers have significantly improved the collection process of
195	information characterizing their exposure, namely the coordinates of the location of the buildings as well as the buildings'
	physical properties. As mentioned previously, exposure data in the loss modelling process is used (i) to estimate the hazard's
	severity at the location of the building and (ii) to select the suitable damage curve. The more precise the exposure data, the
	more accurate the loss evaluation will be. However, as some elements are particularly difficult to get at the time of underwriting
	individual insurance, the systematic extraction and completion of the data remains a challenge and any missing information
200	needs to be completed once the policy is underwritten either in the exposure database or at a later stage in the modelling.
	The increasing volume and precision of geographical information captured by satellites allowed for the development of
	performant geocoding tools supporting the completion of the
	exposure database.

With an address, it is possible to get the geolocation, the structure of the building, number of floors <u>and</u> even the 205 roof type, all critical drivers of damage\_ for different perils (Ehrlich and Tenerelli, 2013; Castagno and Atkins, 2018; Kang et al., 2018; Schorlemmer et al., 2020). This progress in characterizing buildings' properties along with geolocations, was a major advancement enabling insurers to visualize and analyze their accumulation to natural hazard risk.

# <u>To</u>

210 fill the gap when critical information is missing in the exposure database, assumptions are made by using either other data sources to complete the exposure database (e.g. exposure disaggregation to fill in buildings' geolocation) or generic vulnerability curves defined as the weighted average of specific vulnerability curves in the loss modelling process . Any omission on the properties of a building's construction induces an uncertainty on that given building(s)'s exposure that . Any omission on the properties of a building's construction induces an uncertainty on that given building(s)'s exposure that 215 Any omission on the properties of а building's construction induces an uncertainty on that given building(s)'s exposure that can be quantified through sensitivity tests that assess varying combinations of building's construction properties. The impact of inferring а geolocation might however be greater, depending on the peril in question, as for flood and severe convective storms. Testing the impact on losses of a disaggregation scheme requires running the model using several versions of disaggregated portfolios, 220 which is inconceivable today notably because of run time constraints. The disaggregation technique could also provide a solution to modelling the impact of natural hazard on movable exposure. Today, motor and marine exposures are modelled like buildings'. The geolocation used is the car owner's address or the vessel's home port as specified in the policy contract. Disaggregating the motor or marine exposure multiple times would give different vehicle locations and hence capture a range of potential losses

А	a mis en forme : Non Surlignage
1	a mis en forme : Non Surlignage
	a mis en forme : Non Surlignage
· · · · ·	a mis en forme : Non Surlignage

#### 225 3.2 Hazard component

An ever-growing a mount of data on the hazard component has been made accessible, refined, and maintained. A multitude of An ever-growing a mount of data on the hazard component has been made accessible, refined, and maintained. A multitude of An ever-growing a mount of data on the hazard component has been made accessible, refined, and maintained. A multitude of An ever-growing a mount of data on the hazard component has been made accessible, refined, and maintained. A multitude of An ever-growing amount of data on the hazard component has been made accessible, refined, and maintained, and an ever-growing amount of data on the hazard component has been made accessible, refined, and maintained. A multitude of types of data, from observations to model simulations or a mixture of both, substantially support the development of hazard catalogues and their validation. Hazard modelling sets out to characterize, via a hazard

- events catalogue, the full spectrum of severity and frequency of hazards on a specific geographical area. A review of hazard modelling approaches by peril can be found in (Ward et al., 2020). Beyond the perpetual enhancement necessary to complete and refine the view of the risk and to adapt to an ever-evolving environment, uncertainties persist in being only partially
- 235 quantified due to (i) IT constraints and (ii) the information loss perpetuated by simplifying assumptions to derive data compiled in the loss modelling framework. Resolving these two sources of uncertainties would enable insurers to heighten their understanding of the risk and make sounder business decisions.

Uncertainties in the hazard component come from the input data and the modelling parameters used to generate the stochastic event catalogue. For example, (Kaczmarska et al., 2018) quantifies how in changing flooding parameters the loss estimates are

- 240 impacted (Winter et al., 2018) go a step further notably in identifying and quantifying uncertainties present in the production of the hazard events catalogue. Such an analysis first requires running the production of the hazard catalogue several times to test different sets of parameters and secondly running the loss simulation engine multiple times. Including the quantification of uncertainties is costly both in terms of computer power and runtime but should be systematized as a modelling best practice. As mentioned in Section 2.2, the information relative to the event's duration and to the hazard value's evolution
- 245 however are lost when generating the events' footprint, i.e. the maximum value of hazard intensity over the duration of the event. In (re)insurance policies an event's duration is a metric used, within the hours clause, to specify that the (re)insurer will cover all the financial losses accumulated in a defined number of hours, varying depending on the peril. If financial losses are still recorded surpassing the defined number of hours, it will be counted as a second and separate event and activate a double reimbursement from the reinsurer. According to how the reinsurance program is defined, the insurer may have to pay additional
- 250 fees to get a cover for the second event. Analyzing the impact of the hours clause on the ultimate loss would therefore be beneficial for the (re)insurance market. The loss modelling framework must evolve to allow for more flexibility and more completeness.

# 3.3 Vulnerability component

When a natural event occurs, damage results from the rupture of one or several of the building's components, the level of the rupture depending on the hazard's severity and the components' vulnerability. In the aftermath of the event, reconstruction costs are assessed based on the current material prices and labor costs. However, in post-disaster situations, reconstruction

C	Code de champ modifié
a	mis en forme : Non Surlignage
a	mis en forme : Non Surlignage
_	
. <b>C</b>	Code de champ modifié
- (C	Code de champ modifié

a mis en forme : Non Surlignage

a mis en forme : Non Surlignage

	costs may be significantly higher due to a post-event demand surge and inflation. This effect is called post-loss amplification	
	(PLA) and is modelled using a sigmoid function whose calibration remains difficult as (i) it has been observed subsequent to	
	very extreme events and (ii) reconstruction costs or claims available in the historical record includes the PLA effect. As the	
260	PLA may have a substantial impact on the ultimate amount of loss paid by insurers, further research is needed to analyze and	
	model this effect.	
	Systematic data collection of damage information and its associated hazard magnitude is therefore vital to	
	characterize the impact of natural hazards on buildings and to improve the calibrations <u>not only</u> of the buildings'	
	destruction rate but also of the reconstruction costs in the vulnerability component. New technologies such as drones and	
265	satellites provide alternative ways to access impacted areas to collect detailed and granular measurements within a few hours	
	or days of an event's occurrence (Chesnel et al., 2007; Kakooei and Baleghi, 2017). While there has been a substantial increase	
	in the availability of observational data over the past two decades (Yu et al., 2018), further investments should be made to	
	systematically collect: (i) the event's level of <u>hazard</u> severity at <u>the building's</u> location	
	(i.e. values of the relevant hazards' variables leading to the building's damage), (ii) the building's level of damage	
270	and the prevention measures if any (concurrently recording all relevant information on the building itself) and (iii) the level	
	$of\ associated\ repair\ costs\ (including\ information\ on\ loss\ a\ djustments\ a\ nd\ e\ conomic\ metrics\ such\ a\ s\ post\ event\ in\ flation). This$	
	data collection effort should be a joint public and private sector undertaking to build up a core common knowledge.	
	A point of attention is the need for data collectors to coordinate and use the common definitions of the damages' scale to avoid	
	A point of attention is the need for data collectors to coordinate and use the common definitions of the damages' scale to avoid	
275	A point of attention is the need for data collectors to coordinate and use the common definitions of the damages' scale to avoid	
	A point of attention is the need for data collectors to coordinate and use the common definitions of the damages' scale to avoid	
	A point of attention is	
	the need for data collectors to coordinate and use the common definitions of the damages' scale to avoid duplicating and	
	overlapping datasets that are incomparable. Research initiatives dedicated to gathering various data sources already exist at	
280	country  level.  One  such  example  is the  HOWAS  database  for  flood  damage  in  Germany  (Kreibich  et  al., 2017; Kellermann  et  al.,	
	$al., 2020). Could this type of work be extended to the whole of Europe or even more globally? \\ \underline{The PERILS initiative is worth}$	
	mentioning as it is an example of the (re)insurance market's claims data collection initiative. When an event's loss estimation	
	exceeds a defined threshold, the PERILS organization collects claims from the (re)insurers taking part in the consortium.	
	While this data is aggregated at CRESTA level, the initial estimates of the loss ratios are fundamental	
285	to establish market's loss benchmarks and derive market vulnerability curves for instance.	
	While the challenges set out in this section indicate how to improve what we know we don't know, they also highlight the	
	potential limitations of the current loss modelling framework and its simulation platform.	
	The shortcomings of the current loss modelling framework herein described point to the need for an in-depth review of the	
	The shortcomings of the current	

 $290 \quad loss \, modelling \, framework \, herein \, described \, point \, to \, the \, need \, for \, an \, in-depth \, review \, of \, the \, framework \, to \, improve \, and \, increase$ 

a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage

insurers' understanding of natural hazard risk particularly in an ever more connected environment that is described in the next section. From an insurer's perspective, in a context of growing focus on natural hazard impacts, data collection, modelling flexibility, and transparency have become core strategic elements to enhance and gain confidence in its assessment of natural hazard risk. To achieve modelling flexibility and transparency will require in-depth changes to absorb the high amount of data and to incorporate uncertainty quantifications. Data collection, especially for damages and claims, is a societal matter

4 <u>Future challenges and further</u> needs

300 Since the building of the loss modelling framework in the 1990s, clients have become more interconnected (Gereffi et al. 2001), and the correlations between natural hazards and regions have also become better understood and quantified (Steptoe et al., 2018; Zscheischler et al., 2020; Tilloy et al., 2020). This section explores 3 elements that would advance natural risk assessment and would support insurers' in their ambition to more accurately project and plan out their business activities related to natural hazards.

305 <u>4.1 Introducing a 5<sup>th</sup> component to quantify the uncertainty</u>

and should be tackled collectively.

As stated in previous sections, the assessment of uncertainty all along the modelling chain constitutes the loss modelling framework's notable shortcoming and the one that requires further investigation. To a certain extent, uncertainty is inherent to modelling and is partly captured in the loss modelling framework today through (i) the primary uncertainty, that is the assumptions and the simulation of the hazard catalogue, and

310 (ii) the secondary uncertainty, that is the damage and loss assessment. Including the quantification of uncertainties in the loss modelling framework is costly both in terms of computer power and runtime. However, given the rapid evolution of IT, computer power and run time should not be an issue for long and the question will then be how to implement a comprehensive uncertainty quantification scheme. While (Beven et al., 2018) suggest a framework to deal with epistemic uncertainty in natural hazard modelling.recent work like (Noacco et al., 2019; KC et al.,

- 315 2020) has been carried out to address quantifying uncertainty with appropriate methods and tools. How about introducing a specific "uncertainty component" that would deal with the multiple datasets from the different components and propagate the quantification all along the loss modelling process? Along with their systematic quantification, we are convinced that uncertainties' management and communication around it
- will evolve and that insurers will take ownership of this management and make it a tool to enhance the modelling (Thompson 320 and Warmink, 2016; Doyle et al., 2019).

a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
a mis en forme : Non Surlignage
Code de champ modifié
Code de champ modifié
a mis en forme : Non Surlignage
a mis en forme : Exposant

Code de champ modifié

_	a mis en forme : Non Surlignage
{	a mis en forme : Non Surlignage
$\geq$	a mis en forme : Non Surlignage
Ň	a mis en forme : Non Surlignage
Ň	Code de champ modifié
N	a mis en forme : Non Surlignage

# 4.2 Supply chain modelling

	With globalization, clients around the world have become increasingly interconnected and dependent on each other within so-	
	called Global Value Chains (Gereffi et al., 2001; Baldwin and Lopez-Gonzalez, 2015; Phillips, 2018). This dependency	 Code de champ modifié
	became apparent with the 2011 floods in Thailand when Thailand's brutal interruption of microprocessor production led to a	
325	$\underline{halt\ in\ global\ production, a\ global\ shortage\ of\ microprocessors, and\ consequently, a\ loss\ in\ benefits\ for\ companies\ p\ roducing}$	
	chips, hard disc drives and other electronic devices. (Chopra and Sodhi, 2014; Haraguchi and Lall, 2015).	 Code de champ modifié
	From an insurer's perspective, suppliers' defaulting in their deliveries due to the occurrence of a natural hazard is not insurable,	
	as it is not quantifiable with the current modelling that fails to capture this connection between suppliers and their client	
	producers. Supply chain data has improved. (Tiwari et al., 2018; Beorchia and Crook, 2020) and needs to be analyzed further	 Code de champ modifié
330	and incorporated into natural hazard loss modelling. This could provide a source of opportunities for insurers to deliver new	
	services to customers while continuing to contribute to advancing research in visualizing and measuring the levels of	
	complexity (volume, direction and intensity of interconnections).	
	The interconnections between hazards or between clients have yet to be captured even in the latest loss modelling framework.	
	It remains siloed by hazard and region and pmits supply chain information. Failing to integrate these interactions	 a mis en forme : Non Surlignage
335	may result in instilling a bias in our understanding of the underlying risk. A deeper review of the loss modelling framework is	
	to be conducted to reflect on this new and complex reality.	
	4.3 Forward-looking scenario: modelling the future of natural hazard risks	
	Natural hazard models have been primarily developed to overcome the limited historical loss record and to assess extreme	
	losses driven by exposure, hazard and vulnerability in the present. They are now envisaged as tools to assess	
340	the future of natural hazard climate risks.	
540	To perform this analysis, insurers need to not only project the plausible future scenarios of hazard events (information provided	
	by climate model simulations) but also to project the evolution of exposure and vulnerability. In this context, the most pressing	
	questions global insurers need to respond to are: how to gather future projections of population growth or decline and/or wealth	
	worldwide? How will building codes evolve?	
345	(Cremen et al., 2022) perform a thorough review of the available literature and provide initial answers to these questions. Such	 Code de champ modifié
	a review is particularly enlightening to enhance the simple initial assumptions that were made, especially for exposure growth	
	and vulnerability. Furthermore, as vulnerability, is a crucial element in adapting to climate change impacts, further	
	investigations on the implementation of prevention measures and the quantification of the resulting risk reduction are needed.	
	Finally, while this forward-looking analysis is necessary, its outcomes should be taken with great caution. As (Fiedler et al.,	 Code de champ modifié
350	2021) highlight, uncertainty around future exposure, hazard or vulnerability projections,	
	, is significant and compounds the uncertainty already present in the	 a mis en forme : Non Surlignage

#### loss modelling framework

#### **5** Conclusion

To date, models have evolved through the incorporation of new information, without ever undergoing an in-depth transformation. Modifications have stemmed from the observation of the growing number of interconnections – and mutual impacts – at multiple levels: between insured customers and their suppliers and interactions and cross impacts between the disasters-causing natural phenomena. Though this make-do approach has served, it no longer suffices. In today's world where complex intrinsic interconnections exist between natural hazards, exposure, and vulnerability, models fail to reflect this reality. They are in want of an in-depth transformation. Only then will they convey and advance the new

- 360 level of understanding insurers need to cultivate and enable the design and testing of new products and protection mechanisms. <u>As said in (Baum, 2015), "threats are rarely completely unknown or unquantifiable". Sometimes what we do not know is already present in the data or the model but it has yet to be understood or analyzed.</u> We propose reflecting on how to bring together a transdisciplinary research team composed not only of IT, data science, and the geosciences, but also civil engineers, urban planning sciences, and socio-economic sciences to investigate the opportunities to build global loss models for natural
- 365 hazards that would deal with the complexity of the interactions of both natural elements and the customer ecosystem. This would enable insurers to better anticipate the needs of their customers while being better equipped to cope not only with uncertainty but also the unknown.

# References

Baldwin, R. and Lopez-Gonzalez, J.: Supply-chain Trade: A Portrait of Global Patterns and Several Testable Hypotheses, World Econ, 38, 1682–1721, https://doi.org/10.1111/twec.12189, 2015.

Baum, S. D.: Risk and resilience for unknown, unquantifiable, systemic, and unlikely/catastrophic threats, Environ Syst Decis, 35, 229–236, https://doi.org/10.1007/s10669-015-9551-8, 2015.

 Beorchia, A. and Crook, T. R.: Bloomberg Supply Chain Analysis: A Data Source for Investigating the Nature, Size, and Structure of Interorganizational Relationships, in: Research Methodology in Strategy and Management, edited by: Crook, T.
 R., Lê, J., and Smith, A. D., Emerald Publishing Limited, 73–100, https://doi.org/10.1108/S1479-838720200000012017, 2020.

Beven, K. J., Aspinall, W. P., Bates, P. D., Borgomeo, E., Goda, K., Hall, J. W., Page, T., Phillips, J. C., Simpson, M., Smith, P. J., Wagener, T., and Watson, M.: Epistemic uncertainties and natural hazard risk assessment – Part 2: What should constitute good practice?, Nat. Hazards Earth Syst. Sci., 18, 2769–2783, https://doi.org/10.5194/nhess-18-2769-2018, 2018.

Castagno, J. and Atkins, E.: Roof Shape Classification from LiDAR and Satellite Image Data Fusion Using Supervised Learning, Sensors, 18, 3960, https://doi.org/10.3390/s18113960, 2018. Chesnel, A.-L., Binet, R., and Wald, L.: Object oriented assessment of damage due to natural disaster using very high resolution images, in: 2007 IEEE International Geoscience and Remote Sensing Symposium, Barcelona, Spain, 3736–3739, https://doi.org/10.1109/IGARSS.2007.4423655, 2007.

Chopra, S. and Sodhi, M. S.: Reducing the Risk of Supply Chain Disruptions, SUPPLY CHAIN MANAGEMENT, 9, 2014.

385 Cremen, G., Galasso, C., and McCloskey, J.: Modelling and quantifying tomorrow's risks from natural hazards, Science of The Total Environment, 817, 152552, https://doi.org/10.1016/j.scitotenv.2021.152552, 2022.

Doyle, E. E. H., Johnston, D. M., Smith, R., and Paton, D.: Communicating model uncertainty for natural hazards: A qualitative systematic thematic review, International Journal of Disaster Risk Reduction, 33, 449–476, https://doi.org/10.1016/j.ijdrr.2018.10.023, 2019.

390 Ehrlich, D. and Tenerelli, P.: Optical satellite imagery for quantifying spatio-temporal dimension of physical exposure in disaster risk assessments, Nat Hazards, 68, 1271–1289, https://doi.org/10.1007/s11069-012-0372-5, 2013.

Fiedler, T., Pitman, A. J., Mackenzie, K., Wood, N., Jakob, C., and Perkins-Kirkpatrick, S. E.: Business risk and the emergence of climate analytics, Nat. Clim. Chang., 11, 87–94, https://doi.org/10.1038/s41558-020-00984-6, 2021.

Gereffi, G., Humphrey, J., Kaplinsky, R., and Sturgeon\*, T.J.: Introduction: Globalisation, Value Chains and Development, IDS Bulletin, 32, 1–8, https://doi.org/10.1111/j.1759-5436.2001.mp32003001.x, 2001.

Girard, J.: A Leader's Guide to Knowledge Management: Drawing on the Past to Enhance Future Performance, 1st ed, Business Expert Press, https://doi.org/10.4128/9781606490198, 2009.

Grossi, P., Kunreuther, H., and Windeler, D.: An Introduction to Catastrophe Models and Insurance, in: Catastrophe Modeling: A New Approach to Managing Risk, vol. 25, edited by: Grossi, P. and Kunreuther, H., Kluwer Academic Publishers, Boston, 23–42. https://doi.org/10.1007/0-387-23129-3 2. 2005.

400

Haraguchi, M. and Lall, U.: Flood risks and impacts: A case study of Thailand's floods in 2011 and research questions for supply chain decision making, International Journal of Disaster Risk Reduction, 14, 256–272, https://doi.org/10.1016/j.ijdrr.2014.09.005, 2015.

Kaczmarska, J., Jewson, S., and Bellone, E.: Quantifying the sources of simulation uncertainty in natural catastrophe models,
 Stoch Environ Res Risk Assess, 32, 591–605, https://doi.org/10.1007/s00477-017-1393-0, 2018.

Kakooei, M. and Baleghi, Y.: Fusion of satellite, aircraft, and UAV data for automatic disaster damage assessment, International Journal of Remote Sensing, 38, 2511–2534, https://doi.org/10.1080/01431161.2017.1294780, 2017.

Kang, J., Körner, M., Wang, Y., Taubenböck, H., and Zhu, X. X.: Building instance classification using street view images, ISPRS Journal of Photogrammetry and Remote Sensing, 145, 44–59, https://doi.org/10.1016/j.isprsjprs.2018.02.006, 2018.

410 KC, U., Garg, S., Hilton, J., and Aryal, J.: A cloud-based framework for sensitivity analysis of natural hazard models, Environmental Modelling & Software, 134, 104800, https://doi.org/10.1016/j.envsoft.2020.104800, 2020.

Kellermann, P., Schröter, K., Thieken, A. H., Haubrock, S.-N., and Kreibich, H.: The object-specific flood damage database HOWAS 21, Nat. Hazards Earth Syst. Sci., 20, 2503–2519, https://doi.org/10.5194/nhess-20-2503-2020, 2020.

 Knapp, K. R., Kruk, M. C., Levinson, D. H., Diamond, H. J., and Neumann, C. J.: The International Best Track Archivefor
 Climate Stewardship (IBTrACS): Unifying Tropical Cyclone Data, Bull. Amer. Meteor. Soc., 91, 363–376, https://doi.org/10.1175/2009BAMS2755.1, 2010.

Knapp, K. R., Diamond, H. J., Kossin, J. P., Kruk, M. C., and Schreck, C. J.: International Best Track Archive for Climate Stewardship (IBTrACS) Project, Version 4, https://doi.org/10.25921/82TY-9E16, 2018.

Kreibich, H., Thieken, A., Haubrock, S.-N., and Schröter, K.: HOWAS21, the German Flood Damage Database, in:
Geophysical Monograph Series, edited by: Molinari, D., Menoni, S., and Ballio, F., John Wiley & Sons, Inc., Hoboken, NJ, USA, 65–75, https://doi.org/10.1002/9781119217930.ch5, 2017.

Mitchell-Wallace, K. (Ed.): Natural catastrophe risk management and modelling: a practitioner's guide, John Wiley and Sons, Inc, Hoboken, NJ, 2017.

Noacco, V., Sarrazin, F., Pianosi, F., and Wagener, T.: Matlab/R workflows to assess critical choices in Global Sensitivity 425 Analysis using the SAFE toolbox, MethodsX, 6, 2258–2280, https://doi.org/10.1016/j.mex.2019.09.033, 2019.

Phillips, N.: Global value chains:, in: Dictionnaire d'économie politique, Presses de Sciences Po, 247-250, https://doi.org/10.3917/scpo.smith.2018.01.0247, 2018.

 Pinto, J. G., Bellenbaum, N., Karremann, M. K., and Della-Marta, P. M.: Serial clustering of extra tropical cyclones over the North Atlantic and Europe under recent and future climate conditions: CLUSTERING OF EXTRATROPICAL CYCLONES,
 J. Geophys. Res. Atmos., 118, 12,476-12,485, https://doi.org/10.1002/2013JD020564, 2013.

Priestley, M. D. K., Pinto, J. G., Dacre, H. F., and Shaffrey, L. C.: Rossby wave breaking, the upper level jet, and serial clustering of extratropical cyclones in western Europe: WESTERN EUROPE CLUSTERING DYNAMICS, Geophys. Res. Lett., 44, 514–521, https://doi.org/10.1002/2016GL071277, 2017.

Priestley, M. D. K., Dacre, H. F., Shaffrey, L. C., Hodges, K. I., and Pinto, J. G.: The role of serial European windstom
 clustering for extreme seasonal losses as determined from multi-centennial simulations of high-resolution global climatemodel
 data, Nat. Hazards Earth Syst. Sci., 18, 2991–3006, https://doi.org/10.5194/nhess-18-2991-2018, 2018.

Schorlemmer, D., Beutin, T., Cotton, F., Garcia Ospina, N., Hirata, N., Ma, K.-F., Nievas, C., Prehn, K., and Wyss, M.: Global Dynamic Exposure and the OpenBuildingMap - A Big-Data and Crowd-Sourcing Approach to Exposure Modeling, oral, https://doi.org/10.5194/egusphere-egu2020-18920, 2020.

440 Steptoe, H.: The risk of global weather connections. Are at mospheric hazards independent?, MetOffice, 2016.

Steptoe, H., Jones, S. E. O., and Fox, H.: Correlations Between Extreme Atmospheric Hazards and Global Teleconnections: Implications for Multihazard Resilience: ATMOSPHERIC HAZARDS AND GLOBAL TELECONNECTIONS, Rev. Geophys., 56, 50–78, https://doi.org/10.1002/2017RG000567, 2018.

Thompson, M. and Warmink, J. J.: Natural Hazard Modeling and Uncertainty Analysis, in: Geophysical Monograph Series,
 edited by: Riley, K., Webley, P., and Thompson, M., John Wiley & Sons, Inc., Hoboken, NJ, USA, 9–19,
 https://doi.org/10.1002/9781119028116.ch2, 2016.

Tilloy, A., Malamud, B. D., Winter, H., and Joly-Laugel, A.: Evaluating the efficacy of bivariate extreme modelling approaches for multi-hazard scenarios, Nat. Hazards Earth Syst. Sci., 20, 2091–2117, https://doi.org/10.5194/nhess-20-2091-2020, 2020.

Tiwari, S., Wee, H. M., and Daryanto, Y.: Big data analytics in supply chain management between 2010 and 2016: Insights to 450 industries, Computers & Industrial Engineering, 115, 319–330, https://doi.org/10.1016/j.cie.2017.11.017, 2018.

Vitolo, R., Stephenson, D. B., Cook, I. M., and Mitchell-Wallace, K.: Serial clustering of intense European storms, metz, 18, 411-424, https://doi.org/10.1127/0941-2948/2009/0393, 2009.

Ward, P. J., Blauhut, V., Bloemendaal, N., Daniell, J. E., de Ruiter, M. C., Duncan, M. J., Emberson, R., Jenkins, S. F., Kirschbaum, D., Kunz, M., Mohr, S., Muis, S., Riddell, G. A., Schäfer, A., Stanley, T., Veldkamp, T. I. E., and Winsemius, 455 H. C.: Review article: Natural hazard risk assessments at the global scale, Nat. Hazards Earth Syst. Sci., 20, 1069-1096, https://doi.org/10.5194/nhess-20-1069-2020, 2020.

Winter, B., Schneeberger, K., Huttenlau, M., and Stötter, J.: Sources of uncertainty in a probabilistic flood risk model, Nat Hazards, 91, 431-446, https://doi.org/10.1007/s11069-017-3135-5, 2018.

Yu, M., Yang, C., and Li, Y.: Big Data in Natural Disaster Management: A Review, Geosciences, 8, 165, 460 https://doi.org/10.3390/geosciences8050165, 2018.

Zscheischler, J., Martius, O., Westra, S., Bevacqua, E., Raymond, C., Horton, R. M., van den Hurk, B., AghaKouchak, A, Jézéquel, A., Mahecha, M. D., Maraun, D., Ramos, A. M., Ridder, N. N., Thiery, W., and Vignotto, E.: A typology of compound weather and climate events, Nat Rev Earth Environ, 1, 333-347, https://doi.org/10.1038/s43017-020-0060-z, 2020.

465




Figure 1: Loss Modelling Framework composed of 4 components. A simulation engine is used to intersect the exposure information with the catalogue of hazard events and apply the damage ratio characterized with the vulnerability curve, function of hazard and building characteristics. This operation leads to a loss, gross of any financial insurance conditions. Their application is performed in the financial module.