

# Interactions between precipitation, evapotranspiration and soil moisture-based indices to characterize drought with high-resolution remote sensing and land-surface model data.

Jaime Gaona<sup>1,2</sup>, Pere Quintana-Seguí<sup>2</sup>, María José Escorihuela<sup>3</sup>, Aaron Boone<sup>4</sup>, María Carmen Llasat<sup>5</sup>

- 5 <sup>1</sup>Instituto de Investigación en Agrobiotecnología, CIALE, Universidad de Salamanca, Villamayor-Salamanca, 37185, Spain.  
<sup>2</sup>Hydrology and climate change, Observatori de l'Ebre (Universitat Ramon Llull - CSIC), Roquetes, 43520, Catalonia, Spain.  
<sup>3</sup>IsardSAT, Parc Tecnologic Barcelona Activa, Barcelona, 08042, Catalonia, Spain.  
<sup>4</sup>GMME/MOANA (Groupe de Météorologie à Moyenne Echelle), MOdélisation de l'Atmosphère Nuageuse et Analyse CNRM-GAME (URA CNRS & Météo-France), 42, Av. G. Coriolis, 31057, Toulouse Cedex 1, France.  
10 <sup>5</sup>Applied physics department, Faculty of Physics, University of Barcelona, Barcelona, 08028, Catalonia, Spain.

*Correspondence to:* Jaime Gaona (jaimegaona@usal.es)

## Abstract.

The Iberian Peninsula is prone to drought due to the high variability of the Mediterranean climate with severe consequences for drinking water supply, agriculture, hydropower, and ecosystems functioning. In view of the complexity and relevance of droughts  
15 in this region, it is necessary to increase our understanding of the temporal interactions of precipitation, evapotranspiration and soil moisture that originate drought within the Ebro basin, in northeast Spain, as [the](#) study region. Remote sensing and land-surface models provide high spatial and temporal resolution data to characterize evapotranspiration and soil moisture anomalies in detail. The increasing availability of these datasets has [the](#) potential to overcome the lack of in-situ observations of evapotranspiration and soil moisture. In this study, remote sensing data [of](#) evapotranspiration [from MOD16A2ET](#) and soil moisture [data from](#)  
20 [SMOS1km](#) as well as SURFEX-ISBA land-surface model data are used to calculate ~~the weekly values of~~ the EvapoTranspiration Deficit Index (ETDI) and [the](#) Soil Moisture Deficit Index (SMDI) for the period 2010-2017. The study compares the [remote sensing](#) time series of these ETDI and SMDI indices with the ones estimated using the land-surface model SURFEX-~~ISBA~~, including the [Standardized](#) Precipitation Index (SPI) computed at [a](#) weekly scale. The study focuses on the analysis of the [temporal-time](#) lags between the indices to identify the synchronicity and memory of the [feedbacks-anomalies](#) between precipitation, evapotranspiration  
25 and soil moisture ~~to interpret factors involved in drought onset~~. Lag analysis results demonstrate the capabilities of the SPI, ETDI and SMDI drought indices [computed at a weekly scale](#) to inform about the mechanisms of drought propagation [at distinct levels of the land-atmosphere system](#). Relevant [feedbacks](#) both for antecedent and subsequent conditions are identified, with a preminent role of evapotranspiration in the link between rainfall and soil moisture. Both remote sensing and [the](#) land-surface model show [eapable-capability](#) to characterize drought events, with specific advantages and drawbacks of the remote sensing and land-surface  
30 model datasets. Results underline the value of analyzing drought with dedicated indices, preferably at [a](#) weekly scale, to better identify the quick self-intensifying and mitigating mechanisms governing drought, which are relevant for drought monitoring in semi-arid areas.

## 1 Introduction

Drought is a major natural hazard for ~~the~~ societies in semi-arid climates (Van Loon, 2015) which demands increasing levels of  
35 adaptation and resilience measures to guarantee water supply (Watts et al., 2012), particularly in water-stressed environments. Rain-fed agriculture (Tigkas and Tsakiris, 2015), and even the enduring natural vegetation are very exposed to drought, especially under climate change, which has long-lasting implications ~~to-for~~ the local environment (Gudmundsson et al., 2014). Knowing that

complex interactions take place in the land-atmosphere system under drought, the traditional meteorological or hydrologic approach may overlook drought-relevant interactions between evapotranspiration and soil moisture (Teuling et al., 2013).

40

This explains why modern drought monitoring combines evapotranspiration, soil moisture and even vegetation anomalies to track drought status, such as the [Objective Drought Indicator \(OBDI\) integrated in the U.S. Drought Monitor \(Svoboda et al., 2002\)](#) or [the Combined Drought Indicator within the framework of the European Drought Indicator Observatory \(Sepulcre-Cantó et al., 2012\)](#). ~~This approach is on the upward trend, since even parsimonious composite drought indices like the probabilistic precipitation~~ [vegetation index \(PPVI\) \(Monteleone, Bonacorso and Martina et al., 2020\)](#) outperform the capabilities of common indices to [characterize drought. Therefore, composite indices facilitate the characterization of drought from multiple perspectives \(e.g. Meteorological, Hydrological or Agricultural\) but can be impractical to explore the mechanisms of drought](#) ~~due to complex calculations or missing data.~~ Even though long-term anomalies of rainfall, ~~and other meteorological, meteorological, hydrological or vegetation condition variables~~ [evapotranspiration and soil moisture](#) are [currently](#) regularly monitored, [evapotranspiration and soil moisture](#) ~~ones~~ [still face challenging monitoring. Not only the indirect nature of these variables' data but also their limited spatial and temporal availability limit the number of studies adopting them, even despite](#) ~~and their~~ [known to play a role](#) ~~relevance~~ in the recurrence of drought and heat waves (Zampieri et al., 2009; Dasari et al., 2014), ~~often short term anomalies are overlooked.~~ [Provided that especially regarding](#) interactions of evapotranspiration and soil moisture [operate on short time scales \(Teuling et al., 2018\), there is need to address dedicated exploration of their relevance on the evolution of drought at the shortest time scale](#) [available, which for the soil moisture and evapotranspiration data is currently the weekly scale.](#)

45

50

55

Well-known drought indices such as the standardized precipitation index (SPI) (McKee, Doesken and Kleist, 1993) and the Palmer drought severity index (PDSI) (Palmer, 1965), primarily defined ~~at-on~~ [on](#) the monthly scale, can lack detail to identify short-term anomalies of temperature, wind or radiation originating “flash droughts” (Otkin et al., 2013). Rain-fed agriculture and natural vegetation are particularly sensitive to quickly evolving droughts in specific moments of the growing season (Saini and Westgate, 1999), which subsequently generates evapotranspiration and soil moisture anomalies of short and long-term impact (Jimenez et al., 2011). Recently, there is more interest ~~on-in~~ [in](#) using drought indices with high temporal resolution for short-term drought monitoring, such as the SPI and other indices at the weekly scale (Otkin et al., 2015). Indices with this short-term time scale include the weekly-scale evapotranspiration deficit index (ETDI) and the soil moisture deficit index (SMDI) (Narasimhan & Srinivasan, 2005). The ETDI and SMDI indices are variable-specific enabling full characterization of anomalies at specific levels of the atmosphere-surface system. This is especially useful in the Mediterranean climates where not only rainfall anomalies originate drought (Vicente-Serrano et al., 2004).

60

65

This study focuses ~~in-on~~ [in](#) the Ebro basin, which is an important Mediterranean river basin of the [Iberian Peninsula \(IP\)](#). In view of the increase in the frequency of drought events (Sousa et al., 2011) and ~~in~~ [in](#) the number of consecutive dry spells (Turco and Llasat, 2011) identified in the area, we can expect consequences in the long-term environmental state and the balance between water availability and demands. Furthermore, being placed in a semi-arid climate where most of the rainfall evaporates (68%, Table 15, of “Libro blanco del agua” (MMA, 2000)), the Ebro basin represents an example of how important natural water demands are, particularly in the headwaters where runoff decrease due to reforestation (López-Moreno et al., 2014). Rainfed agriculture dominates the rest of [the](#) ~~un-~~ [un-](#)forested areas and represents the other big consumer of water in the basin. However, despite the relevance of rainfed agriculture its analysis is often overshadowed by irrigation, the biggest anthropogenic demand in the basin (Hoerling et al., 2012). Due to the importance of these water demands and others such as hydropower and energy, the Ebro River

75

Basin Agency operates a dense hydrologic monitoring network, but the lack of dedicated soil moisture and evapotranspiration monitoring jeopardizes drought characterization (Seneviratne et al., 2010). Fortunately, the increasing availability of remote sensing (RS) products enables distributed, precise, and frequent monitoring of these coarsely observed variables (Martínez-Fernández et al., 2016).

Space agencies have released multiple ~~remote sensing~~RS products in the last decades facilitating the distributed analysis of drought (AghaKouchak et al., 2015). Optical spectrometry of the atmospheric (rainfall, temperature, water vapor) and surface (vegetation reflectance) variables have often been the basis for distributed characterization of drought indicators. Surface vegetation indices such as the widespread NDVI (Liu and Kogan, 1996) pioneered the application of RS data to assess the impacts of drought, but thereafter the increasing availability of RS data for multiple meteorological variables has increased its usage on drought indices (West et al., 2019). While even common indices like the SPI can now rely on RS data (Sahoo et al., 2015), the many advantages of RS data facilitate integrating multiple data sources into the increasingly operative composite drought indices for weekly drought monitoring (USDN, Svovoda et al., 2002; CDI, Sepulcre-Cantó et al., 2012) even below the weekly scale (Monteleone et al., 2020). Beyond precipitation, temperature and other directly observable meteorological variables, evapotranspiration and soil moisture represent components of the land-atmosphere system which are difficult to measure on the ground, and consequently suitable for the focus of RS. Recent years have seen a rise in the availability of RS-based evapotranspiration databases such as the global dataset included in GLEAM (Miralles et al., 2011; Martens et al., 2017) or the soil moisture global database CCI (Dorigo et al., 2017), that despite their coarse spatial resolution of these global datasets for regional scale analysis, ease the path to processes understanding. Aiming to gain insight into interactions, the availability of datasets focused on such relevant variables of the land-atmosphere facilitates the use of single-variable drought indices such as the SPI, ETDI and SMDI which is particularly advantageous to analyze the mechanisms of interaction during droughts.

100 Fortunately, in parallel to the ~~remote sensing~~RS missions, the development of processing techniques has improved the applicability of RS-derived data products (Wagner et al., 2007). On this basis, there are soil moisture datasets of increasing high resolution available from the combination of passive microwave sensors such as those from SMOS and SMAP missions (Kerr et al., 2010; Entekhabi et al., 2010; respectively) and active microwave sensors such as ASCAT or Sentinel-1 (Bartalis et al., 2007; Hornacek et al., 2012; respectively). This is the case of the high-resolution soil moisture and evapotranspiration products  
105 SMOS1km (Merlin et al., 2013; Molero et al., 2016, Escorihuela and Quintana-Seguí., 2016; Escorihuela et al., 2018) which have been tested in the area and shown to outperform ASCAT and ASMR-E due to its lack of roughness and vegetation effects. SMAP and Sentinel-1 options or similar resolution to SMOS1km but while accurate in the study area (Dari et al., 2021) are of a much shorter series length and consequently not selected. Similarly, high-resolution RS evapotranspiration products such as ~~and the~~ MODIS16A2-ET (Mu et al., 2013) used in this study are currently available. Therefore, it is worth exploring the capabilities and  
110 limitations of high-resolution RS evapotranspiration data for drought monitoring at the regional scale. High-resolution RS data are therefore best suitable for analysis at the basin scale where the resolution of alternative reanalysis or modelled datasets such as ERA5-Land (Muñoz-Sabater et al., 2021) or LISFLOOD (Van der Knijff et al., 2008) GLEAMv3 (Miralles et al., 2011; Martens et al., 2017) lack detail. Both ~~remote sensing~~RS products represent components of the land-atmosphere system which are difficult to measure on the ground, particularly under extreme conditions such as drought (Miralles et al., 2019). To date, relatively few  
115 works have used satellite high-resolution data for drought analysis in the Iberian PeninsulaIP (Vicente-Serrano, 2006; Scaini et al., 2015, Martínez-Fernández et al., 2016; Sánchez et al., 2016; Ribeiro et al., 2019), especially at the spatial and temporal resolution of this study (Pablos et al., 20192017).

Another source of high-temporal and spatial resolution data is land-surface models (LSM). Used in atmospheric models to simulate the interactions between soil, vegetation and the atmosphere, LSMs [represent a suitable alternative to RS to](#) evaluate the surface water and energy balances [at regional to local scales](#). LSMs initiated their development with one-layer models such as the TOPUP (Schultz et al., 1998) or the PROMET (Mauser and Schadlich, 1998) were developed. Avissar and Pielke (1989) inaugurated the mosaic approach, applying just one-layer models to the different fractions of land-use type. One of the mosaic models able to distinguish between soil evaporation and transpiration is the Météo-France developed model SURFEX (Masson et al., 2013), which fed by the atmospheric analysis SAFRAN (Durand et al., 1999) uses the ISBA scheme for natural surfaces (Noilhan and Mahfouf, 1996). SURFEX has been improved to study the continental water cycle in applications such as SIM and SIM2 (Habets et al., 2008; Le Moigne et al., 2020), often in combination with the hydrologic model MODCOU (Ledoux et al., 1989). The modelling chain called SASER (SAFRAN-SURFEX-Eaudyssée-RAPID) used in this study has been applied to Spain before (Barella-Ortiz and Quintana-Seguí., 2019; Quintana-Seguí et al, 2020) and gives the values required for SPI and SMDI. [Despite the limitations of this LSM when applied as an offline model, it has been validated in the area before and successfully reported providing useful evaluations of water resources in the area of study \(Escorihuela and Quintana-Seguí, 2016; Barella-Ortiz and Quintana-Seguí, 2019\) and in the nearby areas like Portugal and France \(Nogueira et al., 2020; Le Moigne et al., 2020\).](#)

This study aims at evaluating the suitability of high-resolution ~~remote sensing~~RS (SMOS1km and MODIS16A2ET) and LSM (SURFEX-ISBA) data generating rainfall (SPI), soil moisture (SMDI) and evapotranspiration (ETDI) drought [\(single-variable\)](#) indices to better understand the mechanisms behind the temporal evolution of drought in semi-arid climates. [The comparison of RS and LSM data is a main aim of the study to detect the factors impacting RS and LSM performance.](#) The study [further](#) evaluates the advantage of the barely explored weekly temporal scale to capture the short-term anomalies of evaporation and soil moisture decisive to drought in semi-arid areas. The study has an agricultural scope focused on drought ~~of~~[in](#) rain-fed environments given its importance on the land-atmospheric feedbacks (Herrera-Estrada et al., 2017) and ~~to~~the regional socioeconomic sustainability.

## 2 Study area

The study area is the Ebro basin, located in the north-east of the Iberian Peninsula (IP). Placed in between Atlantic and Mediterranean climatic influences, the vast area (85534 km<sup>2</sup>) of the basin (Fig. 1a) has a complex topography (Fig. 1c) which defines a wide range of climatic conditions (Fig. 1d) of distinct spatial and temporal patterns of precipitation, evapotranspiration and soil moisture. The northern border has humid cool climates typical of the Atlantic-exposed Cantabric coastline, while the southeast border enjoys a warm Mediterranean climate. The southwest and north-eastern border are dominated by the Iberian and Pyrenees mountains which together with the Cantabric and Mediterranean ranges restrict the oceanic influence on the central part of the basin. Soil types (e.g. gypsum, limestones) intensify the aridity of certain areas of the basin (Fig. 1b). The combination of semi-arid climatic conditions and unfavorable soil types to vegetation development determine extreme regimes of rainfall, soil moisture and evapotranspiration, prone to drought. The basin is densely populated and supplies a wide range of water demands, especially for agriculture and energy. The vast network of irrigated areas, located mostly in the arid central depression, is vulnerable to hydrological drought risks.

### 3 Data

#### 155 3.1 Land-Surface Model Data

SURFEX, the land-surface modelling platform originally developed and currently maintained by Météo-France (Mason et al., 2013; Le Moigne, 2020), has been chosen to perform the LSM model simulation used in this study. Simulation for the ~~Iberian Peninsula~~IP and Balearic Islands developed within the HUMID project have 5km spatial resolution, with the forcing provided by the Iberian application (Quintana-Seguí et al., 2008; 2016; 2017) of the SAFRAN meteorological analysis system (Durand et al., 160 1999). This is a modelling chain whose offline mode operates using the atmospheric forcing of SAFRAN to feed the ~~land-surface model~~LSM SURFEX-ISBA and simulate even the hydrology with Eaudyssée-RAPID (David et al. 2011).

ISBA (Noilhan and Mahouf, 1996) is the SURFEX module in charge of simulating natural surfaces. There are different versions of ISBA, in this study, we have used the diffusion version (ISBA-DIF; Boone, 1999; Decharme et al., 2011), which performs better 165 in the study area than the simpler 3-layer force restore version (Quintana-Seguí et al., 2020). In this version of ISBA, the LAI has a prescribed annual cycle (constant every year), which may limit the ability of the model to reproduce the long-term effects of drought on vegetation. The model simulates the soil column, but it is unable to simulate groundwater, which despite its impact on soil moisture memory is fortunately not very relevant in the Ebro basin. SURFEX-ISBA requires additional physiographic information that is incorporated from the ECOCLIMAP II land cover database (Faroux et al., 2013) which includes topographic, 170 soil and land cover information at high-resolution.

The available SAFRAN forcing data allows us to simulate the period 1979-2017, but the period used for this study is restricted by ~~remote-sensing~~RS data due to the relatively short length of SMOS data (2010-present) compared to the model. To ensure the comparability of RS-based and LSM-based drought indices, the study period is 2010-2017, for which both ~~remote-sensing~~RS and 175 ~~land-surface model~~LSM data ~~is-are~~ available. To ensure that the RS and the LSM soil moisture ~~is-are~~ comparable, we have averaged three first soil layers of the model according to their discretization in the first 5 cm of the soil (1, 3 and 10 cm of depth respectively). The simulation is performed using a regular 5 km resolution grid based on a custom Lambert Conical Conformal projection.

#### 3.2 Remote Sensing Data

##### 3.2.2 Evapotranspiration

180 To evaluate evapotranspiration, barely measured ~~in-on~~ the ground and not directly measurable from space, we adopt a product based on multiple evaporation-related variables observed by MODIS (Moderate Resolution Imaging Spectroradiometer on NASA's Terra satellite): the MOD16A2-~~ET~~ dataset. This is a level 4 product providing 8-day evapotranspiration (ET) and potential evapotranspiration (PET) based on daily meteorological forcing and 8-day ~~remote-sensing~~RS data of vegetation dynamics from MODIS (Mu et al., 2013). The datasets of MOD16A2-~~ET~~ are published in a sinusoidal projection at a resolution of 500 m (Running 185 et al., 2017). In this study, we have re-projected and interpolated all ~~remote-sensing~~RS products to the same 5 km grid that the LSM simulations use. After the re-gridding step, the temporal step of the datasets is rearranged from the original 8-day accumulation period to a more practical 7-day accumulation period which is more suitable for weekly analysis. Values are linearly weighted depending on their contribution to each week-year for the 52 weeks of a year. We also calculate the monthly means of the ~~ET~~evapotranspiration product ~~in-order~~ to evaluate the impact of the time resolution on drought recognition. The formatting of

190 MOD16A2-ET datasets ~~requires~~require the evaluation of the quality control flags (ET\_QC), given that areas of the Pyrenees show missing data. Only the classes classified as good and optimal in the ET\_QC flags are accepted as data for our study.

### 3.2.3 Surface Soil Moisture

~~In view~~Because of the relatively few years of data currently available from the Soil Moisture Active Passive mission SMAP (Entekhabi et al., 2010), the study adopts SMOS data (Kerr et al., 2010), in particular, the [high-resolution SMOS1km](#) dataset (Merlin et al., 2013). This dataset downscales the original coarse resolution SMOS data using the Disaggregation based on Physical And Theoretical scale Change algorithm DISPATCH (Merlin et al., 2012) / C4DIS (Molero et al., 2016) algorithm. The algorithm enables the downscaling of the 40 km resolution of the SMOS soil moisture data available from 2010 into 1 km resolution using two products at 1 km resolution from MODIS, the NDVI and LST, and an elevation map at the same resolution. Precisely because the scale of interest to study relevant interactions to droughts is the weekly scale, the data is primarily used ~~on a~~ weekly scale. [The spatial scale of interest for the study is that of a regular 5 km resolution grid which takes advantage of the high resolution of SMOS1km.](#) The 2010-2017 dataset presents frequent gaps in the mountainous areas of the Pyrenees. In order to fill the gaps, we apply temporal interpolation [pixel by pixel considering a maximum period for temporal interpolation of two weeks.](#) These data are also spatially aggregated and reprojected to the same grid as the LSM model.

## 4 Methods

### 205 4.1 Drought indices

Drought indices allow quantifying several aspects of drought, like the magnitude and duration, and may also focus on particular variables depending on the scope of interest (i.e. precipitation, soil moisture, aridity...etc.). In view of the convenience to combine several indicators to describe the most about the drought evolution and our focus on rain-fed environments, we adopt the use of the Standardized Precipitation Index (~~“SPI”~~) (McKee, ~~Doesken and Kleist et al.~~, 1993), the Evapotranspiration Deficit Index (~~“ETDI”~~) and the Soil Moisture Deficit Index (~~“SMDI”~~) (Narasimhan and Srinivasan, 2005). Using these three indices, the study aims to investigate the interaction between the two main water fluxes (rainfall and evapotranspiration) and the main storage (soil moisture) involved in the water balance of the land-atmosphere system. The aggregation periods of the SPI index inform about the different responding times of rainfall, soil moisture, streamflow, and groundwater anomalies. By evaluating the evolution of these indices along with their interactions, this study aims to characterize drought mechanisms in the Ebro basin. In this study, the indices  
215 have been computed using gridded datasets, thus, generating a time series for each grid point.

#### 4.1.1 SPI

The Standardized Precipitation Index (SPI) (McKee, ~~Doesken and Kleist et al.~~, 1993) is an index of precipitation anomalies, which is calculated by transforming the accumulated precipitation from its original distribution (usually gamma or Pearson Type III) to the normal distribution with zero mean and unit standard deviation. As a result, we obtain a time series that shows, for each time  
220 step, the departure from the expected value in terms of standard deviations. The calculation of the index is usually done based on monthly time series of rainfall, aggregated over multiple accumulation periods, typically at 3, 6, 12 months. However, SPI can also be calculated on a weekly basis, provided that the accumulation periods are at least 4 weeks (1 month). In this way, this study uses the notation SPI<sub>m-i</sub> to denote the SPI at a monthly scale with an accumulation period of *i* months and the SPI<sub>w-i</sub> to denote the weekly SPI with an accumulation period of *i* weeks. Using SAFRAN data, we adopt the non-parametric methodology proposed by

225 Farahmand and Aghakouchak (2015) to calculate the SPI on a monthly and weekly basis using the multi-month accumulation of 1, 3, 6 and 12 months suitable for rainfall, soil moisture, streamflow and groundwater evaluation.

#### 4.1.2 ETDI

230 The second index incorporated [into](#) the analysis is the EvapoTranspiration Deficit Index (*ETDI*) defined by Narasimhan and Srinivasan (2005). The first step for calculating this index implies defining the water stress ratio (*WS*) for each week, which is the difference between potential evapotranspiration (*PET*) and actual evapotranspiration (*AET*) divided by *PET*. Then the Water Stress Anomaly (*WSA*) is computed as follows:

$$WSA_{i,j} = \frac{MWS_j - WS_{i,j}}{MWS_j - \min WS_j} \times 100, \quad \text{if } WS_{i,j} \leq MWS_j$$

$$WSA_{i,j} = \frac{MWS_j - WS_{i,j}}{\max WS_j - MWS_j} \times 100, \quad \text{if } WS_{i,j} > MWS_j \quad (1)$$

235 where *j* denotes the week of the year ( $1 \leq j \leq 52$ ) and *i* denotes the year. *WS<sub>i,j</sub>* is the water stress of the week *j* of the year *i*. *MWS<sub>j</sub>* is the median *WS* for the week *j* of the year, *minWS<sub>j</sub>* corresponds to the minimum and *maxWS<sub>j</sub>* the maximum. This process removes the seasonality of the time series. *WSA* ranges from -100 (maximum water stress) to 100 (minimum water stress). The *WSA* is accumulated over time to define the *ETDI* as in the following equation:

$$ETDI_j = 0.5 ETDI_{j-1} + \frac{WSA_j}{50} \quad (2)$$

240 To define a range between -4 and 4 for the index, *ETDI* of value -4 must correspond to *WSA* of value -100 and *ETDI* of value 4 to *WSA* of value 100. This range adjustment determines the coefficients 0.5 and the divisor 50 of Eq. (2). In this way, *ETDI* becomes a non-seasonal index suitable for comparing time series of diverse climatic characteristics.

245 Monthly values of the *ETDI* are calculated by computing the average of the weekly values of the corresponding month. In this study, we calculate the *ETDI* using [PET-potential](#) and [AET-actual evapotranspiration](#) provided by [MODIS16A2ET](#). We have also calculated it using the [SURFEX-ISBA](#) simulated *AET* and *PET*. By default, [SURFEX-ISBA](#) does not calculate [PET-potential evapotranspiration](#). In order to do so, we have modified the source code to set soil moisture permanently at field capacity and have run a simulation with this modification. The resulting evapotranspiration corresponds to the potential one (~~*PET*~~).

#### 4.1.3 SMDI

250 The third index incorporated [into](#) the analysis is the Soil Moisture Deficit Index (*SMDI*) also defined by Narasimhan and Srinivasan (2005). The sequence to calculate this index follows the same procedure [of-as](#) the *ETDI*. We first calculate a weekly soil moisture deficit as follows:

$$SD_{i,j} = \frac{SW_{i,j} - MSW_j}{MSW_j - \min SW_j} \times 100, \quad \text{if } SW_{i,j} \leq MSW_j$$

$$SD_{i,j} = \frac{SW_{i,j} - MSW_j}{\max SW_j - MSW_j} \times 100, \quad \text{if } SW_{i,j} > MSW_j \quad (3)$$

Then, the time series of *SMDI* is generated accumulating *SD*:

255 
$$SMDI_j = 0.5 SMDI_{j-1} + \frac{SD_j}{50} \quad (4)$$

*SMDI* also ranges between -4 and 4, respectively corresponding to extremely dry and very wet soil moisture conditions. The *SMDI*, similarly to the *ETDI*, becomes a non-seasonal index able to compare time series of diverse climatic characteristics at different soil depths. In this study, we have calculated the *SMDI* using SMOS-1km surface soil moisture data and SURFEX-ISBA simulated surface soil moisture data. In the case of SURFEX-ISBA, we have calculated the weighted averages of the first two layers of the soil, which corresponds to the first 5 cm of the soil.

#### **4.1.4 Temporal consistency of drought indices calculated based on relatively short RS and LSM data series**

Given the relatively short availability of data for the calculation of ETDI and SMDI series, which depend on the maximum, minimum and median values of the available series, we conducted a sensitivity analysis of the indices in reference to the length of the series and the subset of spatial data. Results shown in Table S1 illustrate the relatively low impact of the length of the series thanks to the high spatial resolution of the dataset. The shortening of the series by half or a quarter barely alters the ETDI series compared to the series of full temporal length. The subset of the dataset to a fraction of its spatial resolution increasingly impacts the robustness of ETDI and SMDI series. Therefore, the high-resolution spatial and temporal datasets such as the RS and LSM used for this study seem beneficial to the consistency of drought indices even when data availability remains under a decade long.

## **4.2 Analysis of interaction between the indices**

### **4.2.1 Correlation between the indices**

To evaluate the similarity between the series of the drought indices (SPI, ETDI and SMDI) we use a variant of the procedure applied by Barella-Ortiz and Quintana-Seguí (2019) and Quintana-Seguí et al. (2020), based on Barker et al. (2015). The method consists of computing the r Pearson correlation between each pair of series of these three drought indices (e.g. SPIw-i and ETDIw and SMDIw); where *i* is the accumulation period, which varies from 4 weeks (1 month) to 52 weeks (1 year). We adopt the r Pearson coefficient for this and the following lag-analysis instead of the r Spearman coefficient, more suitable under non-normal distributions, due to the great computational advantage of computing r Pearson compared to the r Spearman. The time consumed with r Pearson is an order of magnitude lower (e.g. weeks) than that required by r Spearman (e.g. months) for processing our RS and LSM datasets. To further support the use of r Pearson despite the concerns of non-normality of SMDI and ETDI distributions, we conducted a similarity test of the correlation series between indices obtained using r Pearson and r Spearman. Results indicated r Pearson and r Spearman correlate generally over  $r = 0.9$  for RS data and moderately lower for LSM data while do not differ significantly in the timing characteristics of correlation series between indices (Fig. S2). Therefore, we can consider r Pearson comparison of drought indices suitable and sufficiently accurate for the approach and focus of the study.

### **4.2.2 Temporal lag-analysis of the indices:**

Following the correlation analysis between the series, we perform a lag-analysis of the correlation of the pairs of drought indices at a weekly scale, introducing lags from -104 weeks to +104 weeks. We compare the ETDIw and SMDIw with the SPIw-i (being the period of accumulation for the SPIw-i= 4, 13, 26 and 52 weeks, equivalent to the SPIm-1, SPIm-3, SPIm-6, SPIm-12 months), as well as the ETDIw ~~in relation to~~with the SMDIw. The purpose of this analysis is to diagnose the reciprocity and memory in the interaction between rainfall and evapotranspiration, and rainfall and surface soil moisture. The relative abundance of positive over

negative lags (and vice versa) provides information about the asymmetry of the interaction between the indices (precedence and delay). Negative lags refer to leading times of ETDI<sub>w</sub> and SMDI<sub>w</sub> in respect of SPI<sub>w-i</sub> (e.g. lag - 104, left side of the time bar) while positive ones (e.g. lag +104) represent lag times when SPI<sub>w-i</sub> precedes ETDI<sub>w</sub> and SMDI<sub>w</sub> (e.g. lag +104, right side of the time bar). The number of consecutive weeks of positive ~~or~~ negative lags with the increasing period of aggregation of the SPI<sub>w-i</sub> can inform about the memory of the interactions. For each time lag, it is indicated the percentage of the basin affected by non-significant / significant correlation values (Grey / coloured scales of bars in Figs. 3-7). Positive /negative correlations indicate a direct / indirect relationship (red /blue bars in Figs. 3-7).

## 5 Results

### 5.1 Correlation between indices: Monthly scale in comparison to the weekly scale

The first two aims of the study are to evaluate the suitability of SPI, ETDI and SMDI indices to characterize the main anomalies in water exchanges of the land-atmosphere system and to evaluate the suitability of adopting the weekly scale for the analysis of drought indices compared to the use of the monthly scale. Regarding the first, results shown in Fig. 2 indicate a general agreement of the SPI, ETDI and SMDI indices (either computed at a monthly or weekly scale) on the major events of dry and wet anomalies of the period 2010-2017. The dry period of 2011-2012 and the wet period from the end of 2012 to 2015 were properly depicted. However, we identified differences, especially in the case of the SMDI. This index ~~tends~~ to show a generally lower variability when calculated with the LSM compared to the other indices. RS results of SMDI differ from the other indices during the start of 2010 due to the uncertainties during the test period of the SMOS mission. The left column of Fig. 2 shows the monthly SPI<sub>m-i</sub> and the monthly averaged ETDI<sub>m</sub> and SMDI<sub>m</sub>. Correlations between SPI<sub>i</sub> and ETDI / SMDI are calculated ~~at~~ on the monthly (Table 1, left columns) and at the weekly scales (right columns) for both RS and LSM data (where  $i=4, 13, 26$  and 52 weeks of aggregation, equivalent to 1, 3, 6 and 12 months). We test the significance of the correlations ( $p\text{-value} < 0.05$ ) between indices for two subsets: the entire period (2010-2017) and a subset of dry periods (~~i.e.~~, when  $SPI < 0, ETDI < 0, SMDI < 0$ ). Then, we compare the observed (RS) and simulated (LSM) estimates of the indices to explore differences between data sources. RS and LSM ETDI<sub>m</sub> and SMDI<sub>m</sub> indices are moderately correlated (barely over 0.5, significant). Table 1 reports a value of  $r=0.58$  (significant) between ETDI<sub>m</sub> RS and SMDI<sub>m</sub> RS which is significantly higher than the  $r=0.32$  between ETDI<sub>m</sub> LSM and SMDI<sub>m</sub> LSM. In general, despite the resemblance of RS and LSM series of ETDI and SMDI shown in Fig. 2, these two series differ. There is a higher agreement between the RS and the LSM estimates of ETDI<sub>m</sub> ( $r = 0.77$ , significant) than between RS and LSM ones of SMDI<sub>m</sub> ( $r = 0.27$ ).

Table 1 also reveals differences in the moderate correlation of the SPI<sub>m-i</sub> with the ETDI and the SMDI of different temporal aggregations ( $i=1,3,6,12$  months of SPI<sub>m-i</sub> accumulation) as well as between their RS and LSM versions. Correlation between SPI<sub>m</sub> and ETDI<sub>m</sub> RS increases with the aggregation period of the SPI<sub>m</sub> ( $r = 0.39, 0.62, 0.72, 0.81$  respectively for SPI<sub>m-1, SPIm-3, SPIm-6, SPIm-12</sub>) while correlations of SPI<sub>m</sub> with ETDI LSM peak at SPI<sub>m-3</sub> ( $r=0.8$ ) and remain high for SPI<sub>m-6</sub> ( $r=0.78$ ) and SPI<sub>m-12</sub> ( $r=0.71$ ). The correlations between SPI<sub>m-i</sub> and SMDI RS show moderate correlation ranging from the SPI<sub>m-1</sub> ( $r=0.42$ ) to the maximum value of the SPI<sub>m-12</sub> ( $r=0.63$ ). SMDI<sub>m</sub> LSM exhibits a decreasing correlation pattern with the increasing aggregation of the SPI<sub>m</sub> from SPI<sub>m-3</sub> to SPI<sub>m-12</sub> ( $r=0.45, 0.31, 0.3, 0.24$  respectively for SPI<sub>m-1, SPIm-3, SPIm-6 and SPIm-12</sub>). Remarkably, correlations of LSM SPI<sub>m</sub> - SMDI<sub>m</sub> are lower than their RS pairs, which suggests data uncertainties in RS and LSM, as reported by Barella-Ortiz and Quintana-Seguí (2019) and by Quintana-Seguí et al. (2020)).

The monthly correlation analysis is additionally conducted for the subsets of dry periods (those ~~of~~with negative signs of the SPI<sub>m</sub>, ETDI<sub>m</sub> and SMDI<sub>m</sub>). Using the dry subset primarily decreases all correlations. Compared to the whole-time series correlations, the RS ETDI<sub>m</sub> – SMDI<sub>m</sub> values decrease a bit while LSM ones increase a bit. Also, the SPI<sub>m</sub>-i – ETDI<sub>m</sub> /SMDI<sub>m</sub> correlations noticeably decline. Despite losing correlation and significance there is still an increase ~~of~~in correlations with the longer aggregation periods as observed in the whole series analysis.

Fig. 2 provides an overview of the effect of adopting the weekly scale (right column) instead of the monthly scale (left column). The ~~re~~ weekly scale substantially improves the temporal resolution of the plots. Subplots of SPI-i (a to d), ETDI (e) and SMDI (f) show how the weekly scale accurately reproduces the magnitude, tendency and duration of the monthly-scale anomalies, while the increase in temporal resolution additionally captures quick changes. Graphically, we noticed that the aggregation period applied to the SPI prevents from increasing the resolution of the weekly compared to the monthly scale, while the weekly scale strongly increases the ETDI and SMDI resolution (Fig. 2 e<sub>2</sub> and f<sub>2</sub>).

The right columns of Table 1 indicate the correlations between indices ~~on~~at weekly scale compared to those ~~on~~at monthly scale, (the ‘w’ sub-index of ETDI<sub>w</sub> / SMDI<sub>w</sub> / SPI<sub>w</sub> denotes the weekly scale). There is an overall decrease ~~of~~in correlations at the weekly scale compared to the monthly scale, accentuated ~~with~~by the increasing period of aggregation (from SPI<sub>w</sub>-3 to SPI<sub>w</sub>-12). Correlations increase a bit compared to the monthly scale at the lowest period of aggregation of SPI<sub>w</sub>-i, especially for the ETDI (from r=0.39 to 0.57 / r=0.51 to 0.68 in SPI<sub>w</sub>-1 - ~~ETDI<sub>w</sub>-SMDI<sub>w</sub>~~RS / LSM). For both RS and LSM data, the weekly scale lowers the correlation values of the SPI<sub>w</sub>-i with the SMDI<sub>w</sub> more than those with the ETDI<sub>w</sub>. Within the dry-period subset, all the correlations decrease too. The weekly scale lowers correlations due to the increase of variability of the weekly time series compared to the monthly ones but enables capturing the increasing complexity of interactions at shorter time scales.

Therefore, since quick shifts (Fig. 2 e<sub>1</sub> and f<sub>1</sub>) are of great interest for drought analysis, we consider the weekly scale as the most convenient for the lag analysis ~~of~~in the next section. This decision benefits comparing drought indices at their highest temporal definition, which for the ETDI and SMDI indices is natively the weekly scale and for the SPI can be easily applied.

## 5.2 Temporal lag analysis:

The analysis of correlations between the temporally lagged time series provides valuable insight ~~on~~into how the indices interact with each other (e.g., in terms of reciprocity) and about the memory periods of one variable ~~into~~ the other (synchronicity). The analysis addresses the characterization of the feedbacks between rainfall, evapotranspiration and soil moisture while also aims to examine whether the land-atmosphere exchange of semi-arid areas under drought events is a quickly evolving or rather inertial system. Plots of the fraction of the area affected by each correlation level (from -1 to 1 in steps of r=0.2) for each lag period are shown in Figures 3 - 7. It is anticipated that the three evaluated indices ~~will~~agree the most in a narrow range around lag = 0 (no lag) and that their correlation fades away progressively for the increasing lag periods beyond the scale of the propagation of drought. Figures 3 to 6 in which the SPI<sub>w</sub>-i takes part in the interaction of interest with ETDI and SMDI illustrate the range of aggregation periods of SPI<sub>w</sub> (i.e. from 4 weeks of aggregation SPI<sub>w</sub>-4 to SPI<sub>w</sub>-52) along which the interpretation of the interactions between drought indices becomes most altered. In general, the upper two subplots of Figs. 3 to 6 identify the admissible aggregation periods to interpret the clusters of interaction (SPI<sub>w</sub>-4 and SPI<sub>w</sub>-13, equivalent to SPI-1 and SPI-3 on a monthly scale) while the bottom two subplots (SPI<sub>w</sub>-26 and SPI<sub>w</sub>-52, equivalent to SPI-6 and SPI-12 at monthly scale) depict the increasingly merged clusters of interaction due to excessively

365 [long aggregation periods of the SPI. This range of aggregation periods aims to open discussion about the optimal time scale to analyze the interacting drought processes.](#)

### 5.2.1 Lag analysis SPI-ETDI

#### RS

370 The lag analysis of SPIw – ETDIw and SPIw – SMDIw shown in Fig. 3 - 6, aims at diagnosing the reciprocity, synchronicity and memory in the interaction between rainfall and evapotranspiration, and rainfall and surface soil moisture. Each subplot of Fig. 3 to 6 shows the correlation between the ETDIw and SMDIw indices with the SPIw-i index calculated for different aggregation periods (-i) at the week scale: the SPIw-4, SPIw-13, SPIw-26 and SPIw-52 which are the equivalents of SPI-1, SPI-3, SPI-6 and SPI-12 at monthly scale. Negative lags refer to leading times of ETDIw and SMDIw in respect of SPIw-i (e.g. e.g., lag - 104, left side of the time bar) while positive ones (e.g. lag +104) represent lag times when SPIw-i precedes ETDIw and SMDIw (e.g., lag +104, right side of the time bar).

In the case of the ETDIw – SPIw-i analysis based on the ~~remote sensing (RS)~~ data, there is a remarkable cluster of positive correlations in the short term (indicated with the tag ‘ST<sub>1</sub>’ (Short-Term) over the subplots of Fig. 3). This ‘ST<sub>1</sub>’ cluster show relevant fractions of the basin affected by significant moderate (0.4-0.6) values of correlation, particularly in the first weeks of the positive range of lags (from lag 0 to +4) (Fig. 3). The cluster lasts more with the increasing period of aggregation (Fig. 3 a) to d)) eventually becoming merged with the mid-term clusters. In view of subplots Fig. 3 a) to d), the ‘ST<sub>1</sub>’ cluster extends from lag -13 to 4 in the case of SPIw-4, from lag -13 to + 13 in the case of SPIw-13, and once merged with ‘MT<sub>1</sub>’ from -26 to +26 in the case of SPIw-26 and from -26 to +39 for the SPIw-52. The mid-term cluster ‘MT<sub>1</sub>’, originally indicating a period of correlation of ETDIw preceding SPIw-i from 4 to 13 weeks of aggregation, displays moderate to low but significant values of correlation (from 0.2 to 0.4) from lag -36 to -13. The merging of the cluster ‘ST<sub>1</sub>’ and ‘MT<sub>1</sub>’ decreases the asymmetry defining the leading role of ETDIw on SPIw-i. The initially asymmetric interaction of ETDIw – SPIw-i that is mostly located in the negative range of lags (‘ST<sub>1</sub>’ and ‘MT<sub>1</sub>’ shown between lag -30 and +10 for the SPIw-4 and SPI-13 cases) propagates and dampens towards the positive range of lags with the increasing aggregation. The dampening eventually shifts the interaction of ETDIw to SPIw-i from preceding to delayed (at 26 and 52 weeks of aggregation, the clusters of significant positive correlations are mostly within the positive range of lags, especially in Fig. 3 d). The loss of asymmetry due to the increasing aggregation translates into an increase of the duration of the cluster as well as of the fraction of the basin significantly affected by correlations. Both effects may indicate the inconvenience of adopting long aggregation periods that alter the interaction magnitude and timing. There is an additional cluster of positive correlations ‘LT<sub>1</sub>’ past the year and half (lags +78 to +104), particularly noticeable at SPIw-13 and -26. Blue bars in Fig. 3 indicate negative correlations for the relationship ETDIw – SPIw-i dominating the long-term between evapotranspiration and rainfall anomalies. There is a couple of clusters around lag +42 (tagged with ‘LT<sub>2</sub>’) and around lag +104 (‘LT<sub>4</sub>’), slightly significant, that similarly to the positive correlations, increased in duration and magnitude with the increase of the aggregation period of the SPI, particularly for SPIw-26 and 52 (Fig. 3c - d) which may indicate their low relevance.

#### LSM

400 Results from the LSM SURFEX-ISBA (Fig. 4) show a less lasting and more concentrated cluster of significant positive correlations around lag 0 (‘ST<sub>1</sub>’ tagged in Fig. 4) than those observed in the RS results (Fig. 3). This result implies there is more synchronicity between SPI and ETDI in LSM data than in the RS data. Furthermore, the LSM provides higher magnitudes of the significant positive correlations of ETDIw – SPIw-i of ST<sub>1</sub> than RS results. Similarly to RS data, the duration of the highly correlated period

'ST<sub>1</sub>' extends with the increasing aggregation of the SPIw-i (Fig. 4 a to d), eventually causing the merge of clusters 'ST<sub>1</sub>' and 'MT<sub>1</sub>'. The initial asymmetry of the positive correlations towards the negative range of lags is due to the cluster 'MT<sub>1</sub>' placed around lag -26 and cluster 'LT<sub>1</sub>' (Fig. 4 a and b). In the SPIw-26 and SPIw-52 cases (Fig. 4 c and d) 'LT<sub>1</sub>' disappears, and the 'ST<sub>1</sub>' merges with 'MT<sub>1</sub>', artificially shifting the initial asymmetry dominating the negative range of lags towards the positive range. Thus, the precedence of ETDIw with SPIw-i prevalent in the period of aggregation of 4 and 13 weeks translates may look as ~~into a~~ precedence of SPIw-i with ETDIw when long aggregation periods such as the ones in SPIw-26 and -52 are applied. The initial asymmetry of the cluster 'ST<sub>1</sub>' – 'MT<sub>1</sub>' is lower in LSM results (Fig. 4) than in RS results (Fig. 3) due to the lower magnitude of 'MT<sub>1</sub>'. The additional cluster of positive correlations 'LT<sub>3</sub>' past the year and a half range of positive lags (lags +78 to +104) in LSM results (Fig. 4a to c) concurs with that of RS results (Fig. 3a to c). LSM results (Fig. 4) show a few more clusters of negative correlations ETDIw – SPIw-i but of lower magnitude than those of RS ones (Fig. 3). These significant clusters merge with the increasing aggregation period 'LT<sub>4</sub>–LT<sub>5</sub>', which agrees with 'LT<sub>2</sub>' shown in RS at the 26- and 52-weeks period of aggregation of SPIw-i. LSM results further include a cluster of significant negative correlations values in the lead time range (cluster 'LT<sub>2</sub>') that is absent in RS results (Fig. 4 vs. Fig. 3). The agreement between LSM and RS results confirms the asymmetrical interaction between ETDIw and SPIw-i. The asymmetry suggests a certain prevalence of the precedence of positive correlations of ETDIw with SPIw-i in the short term (from the month to the seasonal scale) while points to some delayed response of ETDIw to SPIw-i in the negative correlations in the mid to long-term (from seasonal to interannual scale). The LSM results tend to amplify the magnitude and the area affected by positive correlations compared to the RS dataset. The asymmetry of the ETDIw—SPIw-i becomes also more shifted towards the positive range of lags with the increasing aggregation period in LSM than in RS results.

### 5.2.2 Lag analysis SPI-SMDI

#### RS

The interaction of SPIw-i with SMDIw for the RS dataset is not as strong as it was with ETDIw (Fig 5 and 6 vs. Fig. 3 and 4). Both the correlation values and the significant fraction of the basin affected by them are lower than in the case of the ETDIw. Conversely—Similarly to the ~~ETDIw~~ETDIw, the significant positive correlations around lag 0 ('ST<sub>1</sub>') are asymmetrical. viewBecause of this, the SMDIw tends to experience the effect of the preceding conditions of the SPIw-i more-less than influencing those of the SPIw-i. The increasing period of aggregation of the SPIw-13, -26 and -52 widens and lags the short-term influence of the SPIw-i on the SMDIw ('ST<sub>1</sub>') into a short to mid-term influence ('ST<sub>1</sub>-MT<sub>1</sub>'). This widened cluster of positive correlations stays almost entirely in the positive range of lags, which differs from that of the ETDIw where the 'ST<sub>1</sub>-MT<sub>1</sub>' cluster extended both in positive and negative ranges of the lags (Fig. 5 compared to Fig. 3 to 4). The leading range of lags (from lag -104 to lag 0) does not show relevant clusters of significant correlation at all, neither on-in the mid nor in the the long term. Negative correlations of the SMDIw – SPIw-i interaction only occasionally stand up at a small cluster ('LT<sub>1</sub>', Fig. 5 b - d) in the positive range of lags (when SPIw-i precedes SMDIw). The smoothing effect of the aggregation period of SPIw-i increases-increasingly alters the initial bias of interactions towards the leading influence of SMDI on SPIw as well as the delays of this 'LT<sub>1</sub>' cluster but differently from in ETDIw cases, the magnitude of the significant negative correlations decline, most clusters similarly to the case of with ETDIw.

#### LSM

The LSM results for the SMDIw – SPIw-i relationship depicts s strongly dampened patterns of significant correlation between SMDIw and the SPIw-I compared to those of RS SMDIw – SPIw-i or ETDIw – SPIw-i. Only the 'ST<sub>1</sub>' cluster is noticeable around lag 0. The increase of-in the aggregation period does not favor its permanence as a relevant cluster beyond the SPIw-4. The cause can be the generalized low values of both non-significant and significant correlations obtained for these estimates of LSM

SURFEX-ISBA which may indicate the difficulties of the ~~land surface model~~ LSM to describe the response of the surface soil moisture (the SMDI index) to the atmospheric forcing (the SPI index) that we saw in the RS dataset. The periods identified of short-term, mid-term and long-term influence of one SPIw-i in the SMDIw such as 'SP<sub>1</sub>', 'MT<sub>1</sub>', 'LT<sub>1</sub>' cannot be recognized in Fig. 6 compared to Fig. 5. Therefore, the LSM results of the SPIw-i – SMDIw relationship strongly differ from the ones obtained from RS data and is a matter of debate in the discussion section.

### 5.2.3 Lag analysis ETDI-SMDI

#### RS

The remaining interaction in this analysis is the ETDI – SMDI. The results show a less asymmetric relationship between the ETDI and the SMDI compared to the ones between SPIw with ETDI and SMDI. The significant moderate positive correlation values (red bars in Fig. 7a) between lag 0 and +13 and from lag -10 to 0 indicate that the influence of the ETDI on the SMDI lasts longer than the one of the SMDI on the ETDI. The magnitude, though, expresses that SMDI moderately impacts the short-term conditions of the ETDI for about a month in comparison to the sole week the ETDI affects moderately those of SMDI. However, the highest correlations occur for the lag -1 when SMDI precedes ETDI in one week. Negative clusters 'MT<sub>1</sub>', 'MT<sub>2</sub>' at +/- 39 weeks suggest that the interaction between the indices goes beyond the seasonal scale commented above. However, the significance of all these mid to long-term clusters remains low.

#### LSM

The results of ETDI – SMDI interaction based on LSM data show less evident periods of interaction between the indices compared to the RS results. The expected strong correlation around lag 0 is largely diminished. The strongest cluster appears from lag -21 to -39 'MT<sub>1</sub>' (Fig. 7b) when SMDI precedes ETDI. No notable negative clusters can be identified. Apart from the lack of agreement on the symmetry of the interaction between LSM and RS results (Fig. 7b vs. 7a), the notable cluster 'ST<sub>1</sub>' in RS results is less relevant in LSM results. In view of Given the disparity between LSM and RS results, we rise concerns about the accuracy of offline LSM simulations compared to the RS results, addressing them for discussion in the next section.

## 6 Discussion

Results require careful discussion regarding three main aspects: firstly, the effect of adopting the weekly scale for drought indices and analyses, secondly the meaning behind the complex interactions between drought indices and thirdly the comparison of RS and LSM as tools for high-resolution monitoring of drought. All comments refer to the results at-on a weekly scale.

### 6.1 Scales for drought monitoring in semiarid environments

Analysing/Analyzing the differences in correlations between indices at monthly and weekly scales (Figure 2), we support the necessity of adopting the weekly scale to study lags between relevant variables driving drought processes, because the monthly scale preferred for drought assessment from a hydrological perspective may overlook the quick response of the land-atmosphere interactions. The clusters of moderate to high correlation between indices mostly occur within the first month preceding or following an anomaly (Figs. 3-7), particularly in the short to very short-term. Apart from the tendency of high correlations to peak and plunge in the interval of a few weeks, the information about its delay or precedence can only be observed when the weekly scale is adopted and when aggregation periods of the indices remain below the seasonal scale, such as SPIw-13 (equivalent to SPI3). Our results showing soil moisture response to rainfall (-5 to 5 weeks) and evapotranspiration (-10 to 5 weeks) anomalies in

~~the~~ a matter of weeks are consistent with previous works showing soil moisture echoing rainfall anomalies in a range from days to weeks (Scaini et al., 2015; Martínez-Fernández et al., 2016), but also when driven by evapotranspiration (Otkin et al., 2013).  
480 Therefore, in a basin exposed to the high-energy characteristics of semi-arid climates, the weekly scale increases the capability to diagnose quick and often disregarded interactions such as the short-term interactions of ETDI and SMDI on SPI, without losing resolution in the identification of mid to long-term interactions. The need to apply a weekly scale even the SPI index, which is currently barely used below the monthly scale, is recommended not only for interactions such as the rainfall-evapotranspiration anomalies but also for anomalies such as soil moisture which are often assumed to show interactions over the monthly scale. An  
485 additional aspect to bear in mind when shifting to the weekly scale for drought assessment is that the spatial resolution must increase accordingly. We identified that certain inputs of the LSM model (mainly the forcing dataset, which nominally has a resolution of 5 km, but which has a lower real resolution due to the ~~spacing between meteorological stations~~semi-aggregated configuration of the physical input data) may undermine the resolution of LSM results compared to the ones from RS. This is a scale effect reported by Rodriguez-Iturbe et al. (2001), who warned that coarse resolution of the spatial characteristics increases  
490 the temporal scale “at which the processes are effectively correlated”. Therefore, the lag analysis must compile datasets of appropriate spatial and temporal scales for their aims, like the high-resolution RS and LSM data evaluated in this study.

## 6.2 Interpretation of the interactions between drought indices

Adopting specific drought indices for rainfall, evapotranspiration and soil moisture allows us to explore the interactions between variables of different levels of the land-atmosphere system. Fig. 8 a), and graphically 8 b), ~~graphically~~ summarizes the annual  
495 mode of interactions. In the short to mid-term, both the ETDI and SMDI interactions with SPI concur on having moderate significance, with only a few negative and positive low interactions in the mid to long term. Positive correlations around lag=0 of the ETDI and SMDI with SPI indicate direct precedent dependence of the indices, which means changes on ETDI and SMDI correlate positively (negatively) to positive (negative) changes on the SPI. The short-term correlations after rainfalls for both ETDI and SMDI (lagged response of these indices to SPI) have immediate interpretation and ~~was-were~~ reported before in similar Iberian  
500 regions (Martínez-Fernandez et al., 2016). Sustained dry or wet anomalies in both variables ~~favoured~~favoured by a positive correlation between indices are primarily restricted to a length of three seasons, while remaining less probable beyond a year.

-The merging effect of the ‘ST<sub>1</sub> – MT<sub>1</sub>’ clusters with the increase of the aggregation period implies that the reinforcing positive interaction of SPI-ETDI /SMDI can be still be identified easily when indices are aggregated at monthly or even up to the seasonal  
505 scales. ~~This also affects~~ eCorrelations beyond the year scale may suggesting some multi-annual persistence of the positive reinforcing anomalies, nothing infrequent in Mediterranean climates. Nonetheless, given the strong aggregation impact occurring when adopting the SPIw-26 and SPIw-52, the analysis of interactions between rainfall and evapotranspiration and soil moisture may be uncertain when indices are aggregated beyond the seasonal scale in contrast to the suitability of SPI-6 and SPI-12 used for drought monitoring purposes based on single variables. Autocorrelation values of the SPI series are only partly caused by the  
510 aggregation effect since significant autocorrelated values extend always for less than the period of aggregation of SPIw-i (2 weeks on SPIw-1, 3 weeks on SPIw-4, 10 weeks for SPIw-13, 18 on SPIw-26, 35 on SPIw-52, 40 on SPIw-78, 45 on SPIw-104). The partial autocorrelation of both ETDI and SMDI shows mostly two significant week-lags which may imply an AR(2) model (Fig. S3) functioning as a combination of growing and decaying factors, which reflects the waving balance between positive and negative correlations between indices. In consequence, the clusters of interactions identified in Figs 3 to 6 at least partially capture the  
515 interactions between rainfall, evapotranspiration and soil moisture anomalies at short-term temporal scales. This highlights the benefit of assuming that drought evolution, as an expression of the complex exchanges between land-atmosphere variables,

primarily occurs up to seasonal scales. This means the temporal resolution of drought indices must adapt to short time scales such as the weekly scale to detail the interacting processes governing drought. Otherwise, using drought indices at time scales beyond the seasonal scale may neglect the interplay of land-atmosphere processes determining drought progress.

520

The existence of the precedent influence of ETDI and SMDI on SPI (clusters of the negative range of lags) implies some unequal reciprocity (feedback) between evapotranspiration and soil moisture with rainfalls. This precedent influence is weaker than the influence of SPI on subsequent ETDI and SMDI anomalies (lagged response), but still remarkable. It is reasonable that the lagged response of evapotranspiration and soil moisture to rainfall is stronger and more ~~long-long~~-lasting than the precedent influence (Fig. 8 a). Furthermore, the precedent influence period between the ETDI and SPI is stronger and of longer duration than the one of SMDI on SPI. This asymmetry suggests that ETDI, more than SMDI, has a weekly to seasonal precedent influence on rainfall (Fig. 8b). We expected a longer period of positive correlations of SMDI influencing rainfalls, given the multiple reports of soil moisture inducing memory to the near-surface atmosphere (Manning et al., 2018).

525

530

~~One reason that-why~~ the ETDI shows a longer influence on SPI than the SMDI may be that ETDI from MOD16A2~~ET~~ is fed by the whole depth of soil moisture, while SMDI based on SMOS1km is limited to the top 5 ~~centimetres~~centimeters of soil moisture, a very exposed soil level in semi-arid climates. The complexity of soil moisture dynamics, which barely follow a cyclic interaction (Rodriguez-Iturbe et al., 1991), can also explain a weaker relationship between the SMDI and SPI compared to the ETDI. Other reasons ~~to-for~~ this may be in the prevalence of maritime advection as the main contributor to evapotranspiration in the ~~Iberian Peninsula~~IP (Gimeno et al., 2010), compared to the prevalence of local soil moisture recycling common in ~~the-more~~ continental areas of Europe (Bisselink and Dolman, 2008). The advective explanation is supported by the contrast between the few weeks of precedent influence of soil moisture on rainfall we observe in the Ebro basin and the up to 250 days of precedent influence of continental areas prone to soil moisture recycling (Rowntree and Bolton, 1983; Bisselink and Dolman, 2008). Some studies focused on continental climates of relevant summer rainfall have described the implications of the alteration of the recycling due to soil moisture depletion during heatwaves and drought. Eventually, the self-intensification loop may go beyond the interactions described here and which can eventually-alter the atmosphere (Rasmijn et al., 2018; Miralles et al., 2019). In the Mediterranean climate of the Iberian Peninsula characterized by the lack of summer rainfall, soil moisture annually reaches such low levels that we can expect annual summer alterations in the near atmosphere. Differences between areas where soil moisture plays a role, like central Europe, and areas where soil moisture is unable to control the evolution of the system under high-energy conditions, like the Iberian Peninsula, have been reported before in Mediterranean-like Western Australia in comparison to eastern Australia (Herold et al., 2016).

535

540

545

In consequence, our results at the Ebro basin seem compatible with the frequent activation of a reinforcing or self-intensification loop (Brubaker and Entekhabi, 1996), by which the precedent influence of ~~positive~~-(negative)(eventually positive)-anomalies of evapotranspiration ~~increasing~~-(reducing)(increasing)-rainfall cascades into a depletion (rise) of soil moisture that further limits (enhance) a lagged-enhanced-(depleted)-response of evapotranspiration restarting the cycle (Fig. ~~8b~~8c, right column). The weak precedence of soil moisture on rainfall compared to that of evapotranspiration expresses the limited duration ~~power~~-of the reinforcing-control capacity of the soil moisture ~~loop (based on rainfall recycling) to inhibit the self-intensification loop of~~over evapotranspiration in semi-arid climates of Mediterranean type (~~upper-left column~~ of Fig. 8c).

555

Negative correlations when indices differ in sign ( $r < 0$  conditions, Fig. 8a) can be indicative of transitional periods of mid-seasons. The sharp shift from the cluster of short-term positive correlations to the cluster of mid- to long-term negative correlations suggests a limit in the persistence of the self-intensification mechanism. A physical interpretation of the shift may be related to the change in the dominance of [the sequence from the one under high-energy conditions \(Fig. 8c, right column\) to the one under low-energy conditions \(Fig. 8c, left column\)](#). ~~A low-energy inhibiting mechanism was already described by~~ Brubaker and Entekhabi (1996) ~~described an inhibiting feedback mechanism~~ of negative correlations between soil moisture and surface temperature ~~when under~~ [low temperatures is low](#). Given the direct link between evapotranspiration and temperature, the shift from positive to negative interactions beyond the semester, but below the year scale, suggests [that](#) the arrival of winter low-energy conditions terminates the dominance ~~for most of the year~~ of the self-intensifying loop of evapotranspiration.

[In this way, the annual cycle is regulated by the seasonal succession of two sequences: one under the low-energy conditions of winter when evapotranspiration no longer outweighs the inhibiting of soil moisture due to rainfall \(left column of Fig. 8c\), and the other under high-energy conditions driven by evapotranspiration \(right column of Fig. 8c\). While driven by energy variables, the shift between the long period of interactions dominated by evapotranspiration \(right column of Fig. 8c\) and a short period of interactions controlled by rainfall and soil moisture \(lower sequence of Fig. 8c\) may also occur in advance or delayed depending on thresholds of rainfall and soil moisture anomalies.](#) This reason explains [why under high-energy conditions may eventually drought may terminate due to heavy rainfall, while under low-energy conditions drought may persist triggering the onset of the self-intensification loop of evapotranspiration sooner than normal.](#)

~~is~~ [The conceptualization of the interactions illustrated in Fig. 8 aims to in line with reports highlighting the power of evapotranspiration anomalies not only during in hydrological extremes drought generation \(Otkin et al., 2013\) and meteorological extremes \(Seneviratne et al., 2006; Teuling, 2018; Miralles et al., 2019\) but also along the year under the highly variable Mediterranean climate.](#) Another reason supporting the [year-round implications of the](#) dominance of evapotranspiration over soil moisture is that rainfall mostly transfers to evapotranspiration in semi-arid climates (Rodriguez-Iturbe et al., 2001), where the often-underestimated interception (Savenije et al., 2004) further increases evaporation at the expense of soil moisture. [Therefore, the Mediterranean climate likely presents higher thresholds of rainfall, evapotranspiration and soil moisture anomalies different from those triggering hydrometeorological extremes in other areas \(Tramblay et al., 2021\) that determine the initiation of the evapotranspiration-dominated sequence but may not end up in drought. All these aspects together with predictions of comparatively more change of extremes in the Mediterranean area due to climate change \(Samaniego et al., 2018\) recommend assessing changes in the interactions and their balance from the basis of this study.](#)

However, we bear in mind that our results may oversimplify the causality since processes not analyzed in this study may also play a role. [One major factor of uncertainty is due to the non-linear response of vegetation to water stress, particularly in Mediterranean areas of adapted vegetation \(Boulet et al., 2020\). An example of uncertainties due to vegetation comes from the vegetation type whose specific interaction with the boundary-layer of the atmosphere modulates the partitioning of energy governing evapotranspiration \(Lansu et al., 2020\) but similar ones are reported with soil moisture \(Barbeta et al., 2015\). Additional processes factors may be of influence may be, such as, the well-known atmospheric patterns of NAO or WeMO \(Barnston and Livezey, 1987, Conte et al., 1989\) or the oceanic ones like AMO \(Kerr, 2000\). The multiple periods showing neither prevalent positive nor negative correlations between indices indicate a loss of linear interaction. A source of non-linearity is vegetation. Plants can control evapotranspiration and soil moisture in adaptation to water stress in complex manners that depend more on the type of vegetation](#)

(Katul et al., 2012) than ~~in-on~~ the evapotranspiration or soil moisture status. Plant impact both the ETDI and SMDI because interactions of vegetation integrate the status of the atmospheric and the land-surface variables (Peters, ~~Rundquist, & Wilhite, et al.~~, 1991). In the Mediterranean environment of this study, the quick response to drought of rainfed crops and sclerophyll vegetation and the lagged response of other vegetation types (Vicente-Serrano et al., 2019) may obscure interpreting the links between rainfall, evapotranspiration and soil moisture.

### 6.3 The value of remote sensing and land-surface model's estimates

The RS and LSM results of the lag analysis of the ETDI – SPI interactions show consistently comparable results in contrast to the remarkable disagreement between RS and LSM for the SMDI - SPI interaction. Results of SMDI obtained with the LSM show substantially lower correlations than the ones of RS, while also differing in the timing of the clusters of correlation. We expected the opposite, that the LSM, as being simpler than reality, has stronger SPI – ETDI - SMDI correlations than the RS dataset. We assume the implicit accumulation of uncertainties of modelling (Rodriguez-Iturbe et al., 1991), partly inherited from inputs but also from LSM structure, causes the decrease ~~of-in~~ correlations. This is particularly true for soil moisture, a variable integrating exchanges between climate, soil and vegetation (Rodriguez-Iturbe et al., 2001). Secondly, this is an offline simulation, where the atmosphere (SAFRAN) is forcing the land-surface (SURFEX-ISBA), without explicit feedback, because SURFEX-ISBA does not influence back SAFRAN. SAFRAN estimates real conditions by ingesting observations, so the feedback is implicit in results, which may be insufficient to represent reality. Thirdly, the model itself does not consider important processes like the interactive response of vegetation. ISBA has an interactive vegetation module (ISBA-A-gs), but Mediterranean vegetation can be particularly challenging for it. We expect to test the capabilities of interactive modelling vegetation in a follow-up study. Uncertainties of ISBA with vegetation have also roots in the use of ECOCLIMAP2 database, which shows inaccuracies of cover type and LAI. ECOCLIMAP assumes the maximum/minimum LAI occur in June/February in contrast with the early spring and autumn LAI maximums characteristic of the Mediterranean environment (Queguiner et al., 2011). All in all, the differences between LSM and RS datasets are already an important result to improve the LSM and useful insight into the use of offline LSM drought simulations.

Our results positively verify that ~~remote-sensing~~RS represents an effective tool to overcome the problem of sparsely observed soil moisture or evapotranspiration, whose crucial role ~~on-in~~ drought evolution requires ~~high-high~~-resolution data similarly to precipitation (AghaKouchak and Nakhjiri, 2012). Including high-res evapotranspiration products from MODIS (MOD16A2~~ET~~) and soil moisture from SMOS missions (SMOS-1km) together with the distributed rainfall reanalysis data allows dedicated interpretation of the interactions between these two drought-relevant variables and rainfall, and their role in the water balance of the land-surface interface (Dai, 2011). Especially for evapotranspiration, the maps and series of LSM SURFEX-ISBA are comparable ~~of-to~~ those of ~~remote-sensing~~RS, which supports the reliability of LSM despite their limited capability in arid regions (De Kauwe et al., 2015).

The temporal and spatial patterns of the anomalies are overly identified both by ~~remote-sensing~~RS and the LSM model. The RS data seems able to capture a more complex scheme of interactions than the LSM model, despite the intrinsic data issues of the RS sensors and the performance of the algorithms used to generate the products. Conversely, the LSM seems sensitive to uncertainties from input data, especially surface properties, and the offline forcing. The parametrization of the model assumes a semi-distributed approach by sub-basins of the catchment on which each sub-basin is defined based on average values of land over and soil characteristics of the ECOCLIMAP database, which may induce some patchiness of LSM results compared to the RS results. The offline run means that the meteorological data forces the LSM, but the feedbacks are lost beyond the meteorological observations

635 [included as observation in the model. Additional aspects can be, for instance, the impact of groundwater redistributing soil moisture depending on topography, which is underrepresented in the LSM. LSM limitations have been reported in multiple works before and are subject of improvement based on the discussion of these limitations \(Teuling et al., 2006; Samaniego et al., 2018\).](#) Either way, uncertainties are causes of major concern in the lag analysis where they can alter the correlations between indices and obscure the interpretation of the interactions.

640 These aspects together with the complex response of vegetation to semi-arid environments can be critical to ~~improve~~[improving](#) evapotranspiration and soil moisture estimates (Ukkola et al., 2016). Solving these inaccuracies would increase the value of RS and LSM estimates. This study exemplifies the potential of high-resolution RS and LSM products for a wide range of applications, such as drought analysis.

## 645 **7 Conclusions and perspectives**

The analysis of droughts in the Ebro basin using dedicated evapotranspiration and soil moisture drought indices based on high-resolution data from MOD16A2~~ET~~, SMOS1km and the ~~land-surface-model~~[LSM](#) SURFEX-ISBA provides the following insights. The monthly scale commonly adopted for drought evaluation (e.g., SPI-3) may overlook the quick evolution of drought from an agricultural and environmental perspective, especially in the high-energy climates of the Mediterranean basin where the anomalies of rainfall, evapotranspiration and soil moisture can vary in a matter of days. ETDI shows the strongest response at a weekly scale while it remains also influential in the mid-term. SMDI can also quickly evolve with anomalies of evapotranspiration and particularly with lasting anomalies of rainfall. The weekly scale is advantageous to describe trends and shifts in the evolution of the indices and to identify disregarded interactions such as the preceding influence of ETDI on SPI.

655 The ETDI and SMDI indices, together with the SPI adapted to the weekly scale, allow ~~to track~~[tracking](#) the evolution of the anomalies of evapotranspiration, soil moisture and rainfall, as well as their interactions driving water anomalies in the region. There is great consistency between [the](#) time series of ETDI, SMDI and SPI. Lag analysis between these indices clarifies the interactions between anomalies on different levels of the surface-atmosphere system, information that is neglected when using multivariable indices [or indices aggregated beyond the seasonal scale](#). The lag analysis also identifies sequences of interactions defining reinforcing or inhibiting feedbacks. Evapotranspiration dominates the water balance of the Iberian semi-arid climate, especially during high-energy periods. This dominance frequently exceeds the controlling action of rainfall and soil moisture, inducing the reinforcing dry loop. ~~In view~~[Because](#) of the relevance of evapotranspiration, heat waves further ~~fuelling~~[fueling](#) dry events deserve further attention. The weak influence of soil moisture on subsequent evapotranspiration and rainfall limits its capability to control the propagation of the anomalies.

665 ~~Remote sensing~~[RS](#) datasets of MOD16A2~~ET~~ and SMOS-1km accurately estimate the temporal and spatial anomalies in the basin. Evapotranspiration from the ~~land-surface-model~~[LSM](#) SURFEX-ISBA closely resembles the ~~remote-sensing~~[RS](#) one of MOD16A2~~ET~~. Results differ substantially between SMOS1km and SURFEX-ISBA estimates of soil moisture. ~~Remote sensing~~[RS](#) uncertainties arise mainly from data gaps. Land-surface model's estimates can extend the evaluation of soil moisture beyond the surface towards the root zone, but face notable challenges from offline simulation neglecting feedbacks, as well as from the quality of input data that defines surface characteristics. ~~Remote sensing~~[RS](#) outcompetes the ~~land-surface-model~~[LSM](#) ~~on-in~~ the ability to integrate information about challenging processes, such as ~~the~~ vegetation dynamics. Assimilation seems the way forward to

integrate the best aspects of both kinds of data. For as long as ground-based observations remain sparse, [remote sensing](#)RS and [land surface models](#)LSM represent effective tools to assess the water anomalies of the land-atmosphere system and their interaction mechanisms.

## Acknowledgements

Funding: This work was performed within the Project HUMID project [CGL2017-85687-R] which was supported by the Spanish Agency of Research (AEI), in the framework of AEI/FEDER-EU grants.

## References

- 680 AghaKouchak, A., and Nakhjiri, N.: A near real-time satellite-based global drought climate data record, *Environ. Res. Lett.* 7(4), 044037, doi: 10.1088/1748-9326/7/4/044037, 2012.
- AghaKouchak, A., Farahmand, A., Melton, F. S., Teixeira, J., Anderson, M. C., Wardlow, B. D., and Hain, C. R.: Remote sensing of drought: Progress, challenges and opportunities, *Rev. Geophys.*, 53(2), 452-480, doi: 10.1002/2014RG000456, 2015.
- 685 Avissar, R., and Pielke, R. A.: A parameterization of heterogeneous land surfaces for atmospheric numerical models and its impact on regional meteorology, *Mon. Wea. Rev.* 117(10), 2113-2136, doi: 10.1175/1520-0493(1989)117<2113:APOHLS>2.0.CO;2, 1989.
- [Barbeta, A., Mejía-Chang, M., Ogaya, R., Voltas, J., Dawson, T. E., and Peñuelas, J.: The combined effects of a long-term experimental drought and an extreme drought on the use of plant-water sources in a Mediterranean forest. \*Glob. Chang. Biol.\*, 21\(3\), 1213-1225, doi: 10.1111/gcb.12785, 2015.](#)
- 690 Barella-Ortiz, A., and Quintana-Seguí, P. Evaluation of drought representation and propagation in regional climate model simulations across Spain, *Hydrol. Earth Syst. Sci.*, 23(12), 5111-5131, doi: 10.5194/hess-23-5111-2019, 2019.
- Barker, L.J., Hannaford, J., Chiverton, A., and Svensson, C.: From meteorological to hydrological drought using standardized indicators, *Hydrol. Earth Syst. Sci. Disc.*, 12(12):12,827–12,875, doi:10.5194/hessd-12-12827-2015, 2015.
- Barnston, A. G., and Livezey, R. E.: Classification, Seasonality and Persistence of Low-Frequency Atmospheric Circulation Patterns, *Mon. Wea. Rev.*, 115, 1083–1126, doi: 10.1175/1520-0493(1987)115<1083:CSAPOL>2.0.CO;2, 1987.
- 695 Bartalis, Z., Wagner, W., Naeimi, V., Hasenauer, S., Scipal, K., Bonekamp, H., ... and Anderson, C.: Initial soil moisture retrievals from the METOP-A Advanced Scatterometer (ASCAT), *Geophys. Res. Lett.*, 34(20), doi: 10.1029/2007GL031088, 2007.
- Beck, H.E., Zimmermann, N.E., McVicar, T.R., Vergopolan, N., Berg, A., and Wood, E.F.: Present and future Köppen-Geiger climate classification maps at 1-km resolution, *Sci. Data*, 5:180214, doi: 10.1038/sdata.2018.214, 2018.
- 700 Bisselink, B., and Dolman, A. J.: Precipitation recycling: Moisture sources over Europe using ERA-40 data, *J. Hydrometeorol.*, 9(5), 1073-1083, doi: 10.1175/2008JHM962.1, 2008.
- Boone, A., Calvet, J.C., and Noilhan, J.: Inclusion of a Third Soil Layer in a Land Surface Scheme Using the Force–Restore Method, *J. Appl. Meteorol.*, 38(11): 1611–1630, doi: 10.1175/1520-0450, 1999.
- 705 [Boulet, G., Jarlan, L., Olioso, A., and Nieto, H.: Chapter 2: Evapotranspiration in the Mediterranean region. In \*Water resources in the Mediterranean region\* \(pp. 23-49\). Elsevier, doi: 10.1016/B978-0-12-818086-0.00002-9, 2020.](#)
- Brubaker, K. L., and Entekhabi, D.: Analysis of feedback mechanisms in land-atmosphere interaction, *Water Resour. Res.*, 32(5), 1343-1357, doi: 10.1029/96WR00005, 1996.

- Conte, M., Giuffrida, A., and Tedesco, S.: Mediterranean Oscillation: Impact on Precipitation and Hydrology in Italy, In  
710 Conference on Climate and Water, (Vol. 1), 1989.
- Dai, A.: Drought under global warming: a review, *Wiley Interdiscip. Rev. Clim. Change*, 2(1), 45-65, doi: 10.1002/wcc.81, 2011.
- [Dari, J., Quintana-Seguí, P., Escorihuela, M. J., Stefan, V., Brocca, L., and Morbidelli, R.: Detecting and mapping irrigated areas in a Mediterranean environment by using remote sensing soil moisture and a land surface model, \*J. Hydrol.\*, 596, 126129, doi: 10.1016/j.jhydrol.2021.126129, 2021.](#)  
715
- Dasari, H. P., Pozo, I., Ferri-Yáñez, F., and Araújo, M. B.: A regional climate study of heat waves over the Iberian Peninsula, *Atmospheric and Climate Sciences*, 4(05), 841, doi: 10.4236/acs.2014.45074, 2014.
- David, C. H., Habets, F., Maidment, D. R., and Yang, Z. L.: RAPID applied to the SIM-France model, *Hydrol. Process.*, 25(22), 3412-3425, doi: 10.1002/hyp.8070, 2011.
- 720 Decharme, B., Boone, A., Delire, C., and Noilhan, J.: Local evaluation of the Interaction between Soil Biosphere Atmosphere soil multilayer diffusion scheme using four pedotransfer functions, *J. Geophys. Res. Atmos.*, 116(D20), doi: 10.1029/2011JD016002, 2011.
- De Kauwe, M. G., Zhou, S. X., Medlyn, B. E., Pitman, A. J., Wang, Y. P., Duursma, R. A., and Prentice, I. C.: Do land surface models need to include differential plant species responses to drought? Examining model predictions across a mesic-xeric  
725 gradient in Europe, *Biogeosciences*, 12(24), 7503-7518, doi: 10.5194/bg-12-7503-2015, 2015.
- [Dorigo, W., Wagner, W., Albergel, C., Albrecht, F., Balsamo, G., Brocca, L., ... and Lecomte, P.: ESA CCI Soil Moisture for improved Earth system understanding: State-of-the art and future directions. \*Remote Sens. Environ.\*, 203, 185-215, doi: 10.1016/j.rse.2017.07.001, 2017.](#)
- Durand, Y., Giraud, G., Brun, E., Merindol, L. and Martin, E.: A computer-based system simulating snowpack structures as a  
730 tool for regional avalanche forecasting, *J. Glaciol.*, 45, 469-484. ESPERE, cited 2006: Climate encyclopedia—Food and climate, doi: 10.3189/S0022143000001337, 1999.
- Entekhabi, D., Njoku, E. G., O'Neill, P. E., Kellogg, K. H., Crow, W. T., Edelstein, W. N., ... and Kimball, J.: The soil moisture active passive (SMAP) mission, *Proc. IEEE*, 98(5), 704-716, doi: 10.1109/JPROC.2010.2043918, 2010.
- Escorihuela, M. J., and Quintana-Seguí, P.: Comparison of remote sensing and simulated soil moisture datasets in Mediterranean  
735 landscapes, *Remote Sens. Environ.*, 180, 99-114, doi: 10.1016/j.rse.2016.02.046, 2016.
- Escorihuela, M. J., Merlin, O., Stefan, V., Moyano, G., Eweys, O. A., Zribi, M., ... and Ghaout, S.: SMOS based high resolution soil moisture estimates for desert locust preventive management, *Remote Sens. Appl.: Soc. Environ.*, 11, 140-150, doi: 10.1016/j.rsase.2018.06.002, 2018.
- Farahmand, A., and AghaKouchak, A.: A generalized framework for deriving nonparametric standardized drought indicators,  
740 *Adv. Water Resour.*, 76, 140-145, doi: 10.1016/j.advwatres.2014.11.012, 2015.
- Faroux, S., Kaptué Tchuenté, A. T., Roujean, J. L., Masson, V., Martin, E., and Moigne, P. L.: ECOCLIMAP-II/Europe: A twofold database of ecosystems and surface parameters at 1 km resolution based on satellite information for use in land surface, meteorological and climate models, *Geosci. Model. Dev.*, 6(2), 563-582, doi: 10.5194/gmd-6-563-2013, 2013.
- Gimeno, L., Nieto, R., Trigo, R. M., Vicente-Serrano, S. M., and López-Moreno, J. I.: Where does the Iberian Peninsula  
745 moisture come from? An answer based on a Lagrangian approach, *J. Hydrometeorol.*, 11(2), 421-436, doi: 10.1175/2009JHM1182.1, 2010.
- Gudmundsson, L., Rego, F. C., Rocha, M., and Seneviratne, S. I.: Predicting above normal wildfire activity in southern Europe as a function of meteorological drought, *Environ. Res. Lett.*, 9(8), 084008, doi: 10.1088/1748-9326/9/8/084008, 2014.

- Habets, F., Boone, A., Champeaux, J.L., Etchevers, P., Franchistéguy, L., Leblois, E., Ledoux, E., Le Moigne, P., Martin, E.,  
750 Morel, S., 20 Noilhan, J., Quintana-Seguí, P., Rousset-Regimbeau, F., and Viennot, P.: The SAFRAN-ISBA-MODCOU  
hydrometeorological model applied over France, *J. Geophys. Res.*, 113, doi: 10.1029/2007JD008548, 2008.
- [Herold, N., Kala, J., and Alexander, L. V.: The influence of soil moisture deficits on Australian heatwaves. \*Environ. Res. Lett.\*, 11\(6\), 064003, doi: 10.1088/1748-9326/11/6/064003, 2016.](#)
- Herrera-Estrada, J. E., Satoh, Y., and Sheffield, J.: Spatiotemporal dynamics of global drought, *Geophys. Res. Lett.*, 44(5), 2254-  
755 2263, doi:10.1002/2016GL071768, 2017.
- Hoerling, M., Eischeid, J., Perlwitz, J., Quan, X., Zhang, T., and Pegion, P.: On the increased frequency of Mediterranean  
drought, *J. Clim.*, 25(6), 2146-2161, doi: 10.1175/JCLI-D-11-00296.1, 2012.
- Hornacek, M., Wagner, W., Sabel, D., Truong, H. L., Snoeij, P., Hahmann, T., ... and Doubková, M.: Potential for high  
resolution systematic global surface soil moisture retrieval via change detection using Sentinel-1, *IEEE J. Sel. Top. Appl. Earth*  
760 *Obs. Remote Sens.*, 5(4), 1303-1311, doi: 10.1109/JSTARS.2012.2190136, 2012.
- Jiménez, M. A., Jaksic, F. M., Armesto, J. J., Gaxiola, A., Meserve, P. L., Kelt, D. A., and Gutiérrez, J. R.: Extreme climatic  
events change the dynamics and invasibility of semi-arid annual plant communities, *Ecol. Lett.*, 14(12), 1227-1235, doi:  
10.1111/j.1461-0248.2011.01693.x, 2011.
- Katul, G. G., Oren, R., Manzoni, S., Higgins, C., and Parlange, M. B.: Evapotranspiration: a process driving mass transport and  
765 energy exchange in the soil-plant-atmosphere-climate system, *Rev. Geophys.*, 50(3), doi: 10.1029/2011RG000366, 2012.
- Kerr, R. A.: A North Atlantic Climate Pacemaker for the Centuries, *Science*, 288(5473), 1984–1985, doi:  
10.1126/science.288.5473.1984, 2000.
- Kerr, Y. H., Waldteufel, P., Wigneron, J. P., Delwart, S., Cabot, F., Boutin, J., ... and Juglea, S. E.: The SMOS mission: New  
tool for monitoring key elements of the global water cycle, *Proc. IEEE*, 98(5), 666-687, doi: 10.1109/JPROC.2010.2043032,  
770 2010.
- [Lansu, E. M., van Heerwaarden, C. C., Stegehuis, A. I., and Teuling, A. J. : Atmospheric aridity and apparent soil moisture  
drought in European forest during heat waves. \*Geophys. Res. Lett.\*, 47\(6\), e2020GL087091, doi: 10.1029/2020GL087091, 2020.](#)
- Ledoux, E., Girard, G., De Marsily, G., Villeneuve, J. P., and Deschenes, J. : Spatially distributed modeling: conceptual  
approach, coupling surface water and groundwater. In *Unsaturated Flow in Hydrologic Modeling* (pp. 435-454). Springer,  
775 Dordrecht, doi: 10.1007/978-94-009-2352-2\_16, 1989.
- Le Moigne, P., Besson, F., Martin, E., Boé, J., Boone, A., Decharme, B., Etchevers, P., Faroux, S., Habets, F., Lafaysse, M.,  
Leroux, D., and Rousset-Regimbeau, F. : The latest improvements with SURFEX v8.0 of the Safran–Isba–Modcou  
hydrometeorological model for France, *Geosci. Model Dev.*, 13(9), 3925–3946, doi: 10.5194/gmd-13-3925-2020, 2020.
- [Liu, W. T., and Kogan, F. N.: Monitoring regional drought using the vegetation condition index. \*Int. J. Remote Sens.\*, 17\(14\),  
780 2761-2782, doi: 10.1080/01431169608949106, 1996.](#)
- López-Moreno, J. I., Zabalza, J., Vicente-Serrano, S. M., Revuelto, J., Gilaberte, M., Azorin-Molina, C., ... and Tague, C.:  
Impact of climate and land use change on water availability and reservoir management: Scenarios in the Upper Aragón River,  
Spanish Pyrenees, *Sci. Total Environ.*, 493, 1222-1231, doi: 10.1016/j.scitotenv.2013.09.031, 2014.
- Manning, C., Widmann, M., Bevacqua, E., Van Loon, A. F., Maraun, D., and Vrac, M.: Soil moisture drought in Europe: a  
785 compound event of precipitation and potential evapotranspiration on multiple time scales, *J. Hydrometeorol.*, 19(8), 1255-1271,  
doi: 10.1175/JHM-D-18-0017.1, 2018.


- [Martens, B., Miralles, D.G., Lievens, H., van der Schalie, R., de Jeu, R.A.M., Fernández-Prieto, D., Beck, H.E., Dorigo, W.A., and Verhoest, N.E.C.: GLEAM v3: satellite-based land evaporation and root-zone soil moisture, \*Geosci. Model. Dev.\*, 10, 1903–1925, doi: 10.5194/gmd-10-1903-2017, 2017.](#)
- 790 Martínez-Fernández, J., González-Zamora, A., Sánchez, N., Gumuzzio, A., and Herrero-Jiménez, C. M.: Satellite soil moisture for agricultural drought monitoring: Assessment of the SMOS derived Soil Water Deficit Index, *Remote Sens. Environ.*, 177, 277–286, doi: 10.1016/j.rse.2016.02.064, 2016.
- Masson, V., Le Moigne, P., Martin, E., Faroux, S., Alias, A., Alkama, R., ... and Brousseau, P.: The SURFEXv7. 2 land and ocean surface platform for coupled or offline simulation of earth surface variables and fluxes, *Geosci. Model Dev.*, 6, 929–960, doi: 10.5194/gmd-6-929-2013, 2013.
- 795 Mauser, W., and Schadlich, S.: Modelling the Spatial Distribution of Evapotranspiration Using Remote Sensing Data and PROMET, *J. Hydrology*, 213, 250, 267, doi: 10.1016/S0022-1694(98)00228-5, 1998.
- McKee, T.B., Doesken, N.J. and Kleist., J.: The relationship of drought frequency and duration to time scale. In: Proceedings of the Eighth Conference on Applied Climatology, Anaheim, California, 17–22 January 1993, Boston, American Meteorological Society, 179–184, 1993.
- 800 Merlin, O., Rudiger, C., Al Bitar, A., Richaume, P., Walker, J. P., and Kerr, Y. H.: Disaggregation of SMOS soil moisture in Southeastern Australia, *IEEE Trans. Geosci. Remote Sens.*, 50(5), 1556–1571, doi: 10.1109/TGRS.2011.2175000, 2012.
- Merlin, O., Escorihuela, M. J., Mayoral, M. A., Hagolle, O., Al Bitar, A., and Kerr, Y.: Self-calibrated evaporation-based disaggregation of SMOS soil moisture: An evaluation study at 3 km and 100 m resolution in Catalunya, Spain, *Remote Sens. Environ.*, 130, 25–38, doi: 10.1016/j.rse.2012.11.008, 2013.
- 805 [Miralles, D.G., Holmes, T.R.H., de Jeu, R.A.M., Gash, J.H., Meesters, A.G.C.A., Dolman, A.J.: Global land-surface evaporation estimated from satellite-based observations, \*Hydrol. Earth Syst. Sci.\*, 15, 453–469, doi: 10.5194/hess-15-453-2011, 2011.](#)
- Miralles, D. G., Gentile, P., Seneviratne, S. I., and Teuling, A. J. : Land–atmospheric feedbacks during droughts and heatwaves: state of the science and current challenges, *Ann. N. Y. Acad. Sci.*, 1436(1), 19, doi: 10.1111/nyas.13912, 2019.
- 810 MMA: Libro Blanco del Agua en España. Spanish Ministry of Environment (MMA), 2000.
- Molero, B., Merlin, O., Malbêteau, Y., Al Bitar, A., Cabot, F., Stefan, V., ... and Jackson, T. J.: SMOS disaggregated soil moisture product at 1 km resolution: Processor overview and first validation results, *Remote Sens. Environ.*, 180, 361–376, doi: 10.1016/j.rse.2016.02.045, 2016.
- 815 [Monteleone, B., Bonaccorso, B., and Martina, M.: A joint probabilistic index for objective drought identification: the case study of Haiti, \*Nat. Hazards Earth Syst. Sci.\*, 20\(2\), 471–487, doi: 10.5194/nhes-20-471-2020, 2020.](#)
- Mu, Q., Zhao, M., and Running, S. W.: MODIS global terrestrial evapotranspiration (ET) product (NASA MOD16A2/A3). Algorithm Theoretical Basis Document, Collection, 5, 2013.
- 820 [Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., ... and Thépaut, J. N.: ERA5-Land: A state-of-the-art global reanalysis dataset for land applications, \*Earth Syst. Sci. Data\*, 13\(9\), 4349–4383, doi: 10.5194/essd-13-4349-2021, 2021.](#)
- Narasimhan, B., and Srinivasan, R.: Development and evaluation of Soil Moisture Deficit Index (SMDI) and Evapotranspiration Deficit Index (ETDI) for agricultural drought monitoring, *Agric. For. Meteorol.*, 133(1–4), 69–88, doi: 10.1016/j.agrformet.2005.07.012, 2005.
- 825 [Nogueira, M., Albergel, C., Boussetta, S., Johannsen, F., Trigo, I. F., Ermida, S. L., ... and Dutra, E.: Role of vegetation in representing land surface temperature in the CHTESSEL \(CY45R1\) and SURFEX-ISBA \(v8. 1\) land surface models: A case study over Iberia, \*Geosci. Model Dev.\*, 13\(9\), 3975–3993, doi: 10.5194/gmd-13-3975-2020, 2020.](#)

- Noilhan, J., and Mahfouf, J. F.: The ISBA land surface parameterisation scheme, *Glob. Planet. Change.*, 13(1-4), 145-159, doi: 10.1016/0921-8181(95)00043-7, 1996.
- Otkin, J. A., Anderson, M. C., Hain, C., Mladenova, I. E., Basara, J. B., and Svoboda, M.: Examining rapid onset drought  
830 development using the thermal infrared-based evaporative stress index, *J. Hydrometeorol.*, 14(4), 1057-1074, doi: 10.1175/JHM-D-12-0144.1, 2013.
- Otkin, J. A., Anderson, M. C., Hain, C., and Svoboda, M.: Using temporal changes in drought indices to generate probabilistic drought intensification forecasts, *J. Hydrometeorol.*, 16(1), 88-105, doi: 10.1175/JHM-D-14-0064.1, 2015
- Pablos, M., Martínez-Fernández, J., Sánchez, N., and González-Zamora, Á.: Temporal and spatial comparison of agricultural  
835 drought indices from moderate resolution satellite soil moisture data over Northwest Spain, *Remote Sens.*, 9(11), 1168, doi: 10.3390/rs9111168, 2017.
- Palmer, W. C.: Meteorological drought. U.S. Weather Bureau, Res. Pap. No. 45, 58 pp, 1965.
- Peters, A. J., Rundquist, D. C., and Wilhite, D. A.: Satellite detection of the geographic core of the 1988 Nebraska drought, *Agric. For. Meteorol.*, 57(1-3), 35-47, doi: 10.1016/0168-1923(91)90077-4, 1991.
- 840 Queguiner, S., Martin, E., Lafont, S., Calvet, J.-C., Faroux, S., and Quintana-Seguí, P.: Impact of the use of a CO2 responsive land surface model in simulating the effect of climate change on the hydrology of French Mediterranean basins, *Nat. Hazards Earth Syst. Sci.*, 11, 2803–2816, doi: 10.5194/nhess-11-2803-2011, 2011.
- Quintana-Segui, P., Le Moigne, P., Durand, Y., Martin, E., Habets, F., Baillon, M., ... and Morel, S.: Analysis of near-surface atmospheric variables: Validation of the SAFRAN analysis over France, *J Appl. Meteorol. Climatol.*, 47(1), 92-107,  
845 doi:10.1175/2007JAMC1636.1, 2008.
- Quintana-Seguí, P., Peral, M.C., Turco, M., Llasat, M.C., and Martin, E.: Meteorological Analysis Systems in North-East Spain: Validation of SAFRAN and SPAN, *J. Environ. Inform.*, 27(2):116–130, doi:10.3808/jei.201600335, 2016.
- Quintana-Seguí, P., Turco, M., Herrera, S., and Miguez-Macho, G.: Validation of a new SAFRAN-based gridded precipitation product for Spain 25 and comparisons to Spain02 and ERA-Interim, *Hydrol. Earth Syst. Sci.*, 21(4):2187–2201, doi:  
850 10.5194/hess-21-2187- 2017, 2017.
- Quintana-Seguí, P., Barella-Ortiz, A., Regueiro-Sanfiz, S., and Miguez-Macho, G.: The utility of land-surface model simulations to provide drought information in a water management context using global and local forcing datasets, *Water Resour. Manag.*, 34(7), 2135-2156, doi: 10.1007/s11269-018-2160-9, 2020.
- [Rasmijn, L. M., Van der Schrier, G., Bintanja, R., Barkmeijer, J., Sterl, A., and Hazeleger, W.: Future equivalent of 2010 Russian heatwave intensified by weakening soil moisture constraints. \*Nat. Clim. Chang.\*, 8\(5\), 381-385, doi: 10.1038/s41558-018-0114-0, 2018.](#)  
855
- Ribeiro, A. F., Russo, A., Gouveia, C. M., and Páscoa, P.: Modelling drought-related yield losses in Iberia using remote sensing and multiscalar indices, *Theor. Appl. Climatol.*, 136(1-2), 203-220, doi: 10.1007/s00704-018-2478-5, 2019.
- Rodriguez-Iturbe, I., Entekhabi, D., and Bras, R. L.: Nonlinear dynamics of soil moisture at climate scales: 1. Stochastic  
860 analysis, *Water Resour. Res.*, 27(8), 1899-1906, doi: 10.1029/91WR01035, 1991.
- Rodriguez-Iturbe, I., Porporato, A., Laio, F., and Ridolfi, L.: Plants in water-controlled ecosystems: active role in hydrologic processes and response to water stress: I. Scope and general outline, *Adv. Water Resour.*, 24(7), 695-705, doi: 10.1016/S0309-1708(01)00005-7, 2001.
- Rowntree, P. R., and Bolton, J. A.: Simulation of the atmospheric response to soil moisture anomalies over Europe, *Q. J. R. Meteorol. Soc.*, 109(461), 501-526, doi: 10.1002/qj.49710946105, 1983.  
865

- Running, S. W., Mu, Q., Zhao, M., & Moreno, A.: MODIS Global Terrestrial Evapotranspiration (ET) Product (NASA MOD16A2/A3) NASA Earth Observing System MODIS Land Algorithm. NASA: Washington, DC, USA, 2017.
- [Sahoo, A. K., Sheffield, J., Pan, M., and Wood, E. F.: Evaluation of the tropical rainfall measuring mission multi-satellite precipitation analysis \(TMPA\) for assessment of large-scale meteorological drought. \*Remote Sens. Environ.\*, 159, 181-193. doi: 10.1016/j.rse.2014.11.032, 2015.](#)
- 870 Saini, Hargurdeep S., and Westgate, M. E.: Reproductive development in grain crops during drought. *Advances in agronomy*. Vol. 68. Academic Press. 59-96, doi: 10.1016/S0065-2113(08)60843-3, 1999.
- [Samaniego, L., Thober, S., Kumar, R., Wanders, N., Rakovec, O., Pan, M., and Marx, A.: Anthropogenic warming exacerbates European soil moisture droughts. \*Nat. Clim. Change\*, 8\(5\), 421–426. doi:10.1038/s41558-018-0138-5, 2018.](#)
- 875 Sánchez, N., González-Zamora, Á., Piles, M., and Martínez-Fernández, J.: A new Soil Moisture Agricultural Drought Index (SMADI) integrating MODIS and SMOS products: A case of study over the Iberian Peninsula, *Remote Sens.*, 8(4), 287, doi: 10.3390/rs8040287, 2016.
- Savenije, H. H.: The importance of interception and why we should delete the term evapotranspiration from our vocabulary, *Hydrol. Process.*, 18(8), 1507-1511, doi: 10.1002/hyp.5563, 2004.
- 880 Scaini, A., Sánchez, N., Vicente-Serrano, S. M., and Martínez-Fernández, J. : SMOS-derived soil moisture anomalies and drought indices: a comparative analysis using in situ measurements, *Hydrol. Process.*, 29(3), 373-383, doi: 10.1002/hyp.10150, 2015.
- Schultz, K., Franks, S., and Beven, K.: TOPUP—A TOPMODEL based SVAT model to calculate evaporative fluxes between the land surface and the atmosphere, Version 1.1, Program documentation, 1998.
- 885 Seneviratne, S. I., Lüthi, D., Litschi, M., and Schär, C.: Land–atmosphere coupling and climate change in Europe, *Nature*, 443(7108), 205-209, doi: 10.1038/nature05095, 2006.
- Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., ... and Teuling, A. J.: Investigating soil moisture–climate interactions in a changing climate: A review, *Earth Sci. Rev.*, 99(3-4), 125-161, doi: 10.1016/j.earscirev.2010.02.004, [2010](#).
- 890 Sepulcre-Canto, G., Horion, S. M. A. F., Singleton, A., Carrao, H., and Vogt, J.: Development of a Combined Drought Indicator to detect agricultural drought in Europe, *Nat. Hazards Earth Syst. Sci.*, 12(11), doi: 10.5194/nhess-12-3519-2012, 2012.
- Sousa, P. M., Trigo, R. M., Aizpuru, P., Nieto, R., Gimeno, L., and García Herrera, R.: Trends and extremes of drought indices throughout the 20th century in the Mediterranean, *Nat. Hazards Earth Syst. Sci.*, 11(1), 33-51, doi: 10.5194/nhess-11-33-2011.
- [Svoboda, M., LeComte, D., Hayes, M., Heim, R., Gleason, K., Angel, J., ... and Stephens, S.: The drought monitor. \*Bull. Am. Meteorol. Soc.\*, 83\(8\), 1181-1190. doi: 10.1175/1520-0477-83.8.1181, 2002.](#)
- 895 Teuling, A. J., Van Loon, A. F., Seneviratne, S. I., Lehner, I., Aubinet, M., Heinesch, B., ... and Spank, U.: Evapotranspiration amplifies European summer drought, *Geophys. Res. Lett.*, 40(10), 2071-2075, doi: 10.1002/grl.50495, 2013.
- [Teuling, A. J. A.: hot future for European droughts. \*Nature. Climate Change\*, 8\(5\), 364-365. doi: 10.1038/s41558-018-0154-5, 2018.](#)
- 900 Tigkas, D., and Tsakiris, G.: Early estimation of drought impacts on rainfed wheat yield in Mediterranean climate, *Environ. Process.*, 2(1), 97-114, doi: 10.1007/s40710-014-0052-4, 2015.
- [Tramblay, Y., Koutroulis, A., Samaniego, L., Vicente-Serrano, S. M., Volaire, F., Boone, A., and Polcher, J.: Challenges for drought assessment in the Mediterranean region under future climate scenarios. \*Earth Sci. Rev.\*, 103348. doi:10.1016/j.earscirev.2020.1033, 2021.](#)

- 905 Turco, M., and Llasat, M. C.: Trends in indices of daily precipitation extremes in Catalonia (NE Spain), 1951–2003, *Nat. Hazards Earth Syst. Sci.*, 11(12), 3213–3226, doi: 10.5194/nhess-11-3213-2011, 2011.
- Ukkola, A. M., Pitman, A. J., Decker, M., De Kauwe, M. G., Abramowitz, G., Kala, J., and Wang, Y. P.: Modelling evapotranspiration during precipitation deficits: identifying critical processes in a land surface model, *Hydrol. Earth Syst. Sci.*, 20(6), 2403–2419, doi: 10.5194/hess-20-2403-2016, 2016.
- 910 [Van Der Knijff, J. M., Younis, J., and De Roo, A. P. J.: LISFLOOD: a GIS-based distributed model for river basin scale water balance and flood simulation. \*Int. J. Geogr. Inf. Sci.\*, 24\(2\), 189–212, doi: 10.1080/13658810802549154, 2008.](#)
- Van Loon, Anne F.: Hydrological drought explained. *Wiley Interdiscip. Rev. Water*, 2.4, 2015: 359–392, doi: 10.1002/wat2.1085, 2015.
- Vicente-Serrano, S. M., González-Hidalgo, J. C., de Luis, M., and Raventós, J.: Drought patterns in the Mediterranean area: the Valencia region (eastern Spain), *Clim. Res.*, 26(1), 5–15, doi: 10.3354/cr026005, 2004.
- 915 Vicente-Serrano, S. M.: Evaluating the Impact of Drought Using Remote Sensing in a Mediterranean, Semi-arid Region, *Nat. Hazards*, 40(1), 173–208, doi:10.1007/s11069-006-0009-7,-,2006.
- [Vicente-Serrano, S. M., Azorin-Molina, C., Peña-Gallardo, M., Tomas-Burguera, M., Domínguez-Castro, F., Martín-Hernández, N., ... and García, M.: A high-resolution spatial assessment of the impacts of drought variability on vegetation activity in Spain from 1981 to 2015. \*Nat. Hazards Earth Syst. Sci.\*, 19\(6\), 1189–1213, doi: 10.5194/nhess-19-1189-2019, 2019.](#)
- 920 Wagner, W., Blöschl, G., Pampaloni, P., Calvet, J. C., Bizzarri, B., Wigneron, J. P., and Kerr, Y.: Operational readiness of microwave remote sensing of soil moisture for hydrologic applications, *Hydrol. Res.*, 38(1), 1–20, doi: 10.2166/nh.2007.029, 2007.
- Watts, G., Christerson, B. von, Hannaford, J., and Lonsdale, K.: Testing the resilience of water supply systems to long droughts, *J. Hydrol.*, 414–415, 255–267, doi: 10.1016/j.jhydrol.2011.10.038, 2012.
- 925 [West, H., Quinn, N., and Horswell, M.: Remote sensing for drought monitoring & impact assessment: Progress, past challenges and future opportunities. \*Remote Sensing of Environment\*, 232, 111291, doi: 10.1016/j.rse.2019.111291, 2019.](#)
- Zampieri, M., D'Andrea, F., Vautard, R., Ciais, P., de Noblet-Ducoudré, N., and Yiou, P. : Hot European summers and the role of soil moisture in the propagation of Mediterranean drought, *J. Clim.*, 22(18), 4747–4758, doi: 10.1175/2009JCLI2568.1, 2009.

930



935 **Figure 1: a) Ebro basin location in the Iberian Peninsula (OSM OpenTopoMap). b) Land cover of the basin describes a contrasted basin between the forested areas of the mountains and the steppes of the central depression (© ESRI satellite online maps, World Imagery, 2022). c) Altitudinal range of the basin (IGN ES/FR MDT25, CC-BY 4.0). D) Climatic classes present in the basin according to the climatic classification of Köppen-Geiger (from data of Beck et al. (2018)).**

Pairs of indices	r-Pearson correlation coefficient of the monthly time-series (ETDI <sub>m</sub> , SMDI <sub>m</sub> , SPI <sub>m</sub> -i)		r-Pearson correlation coefficient of the weekly time-series (ETDI <sub>w</sub> , SMDI <sub>w</sub> , SPI <sub>w</sub> -i)	
	2010-2017	Dry periods	2010-2017	Dry periods
ETDI <sub>RS</sub> and ETDI <sub>LSM</sub>	<b>0.77</b>	<b>0.60</b>	<b>0.55</b>	<b>0.34</b>
SMDI <sub>RS</sub> and SMDI <sub>LSM</sub>	<b>0.27</b>	0.13	<b>0.19</b>	0.12
ETDI <sub>RS</sub> and SMDI <sub>RS</sub>	<b>0.58</b>	<b>0.50</b>	<b>0.50</b>	<b>0.46</b>
ETDI <sub>LSM</sub> and SMDI <sub>LSM</sub>	<b>0.32</b>	<b>0.43</b>	<b>0.21</b>	0.18
SPI <sub>m</sub> -1/SPI <sub>w</sub> -4 and ETDI <sub>RS</sub>	<b>0.39</b>	0.23	<b>0.57</b>	<b>0.30</b>
SPI <sub>m</sub> -1/SPI <sub>w</sub> -4 and ETDI <sub>LSM</sub>	<b>0.51</b>	<b>0.37</b>	<b>0.68</b>	<b>0.49</b>
SPI <sub>m</sub> -1/SPI <sub>w</sub> -4 and SMDI <sub>RS</sub>	<b>0.42</b>	0.52	<b>0.57</b>	<b>0.57</b>
SPI <sub>m</sub> -1/SPI <sub>w</sub> -4 and SMDI <sub>LSM</sub>	<b>0.45</b>	<b>0.59</b>	<b>0.33</b>	<b>0.25</b>
SPI <sub>m</sub> -3/SPI <sub>w</sub> -13 and ETDI <sub>RS</sub>	<b>0.62</b>	<b>0.42</b>	<b>0.63</b>	<b>0.32</b>
SPI <sub>m</sub> -3/SPI <sub>w</sub> -13 and ETDI <sub>LSM</sub>	<b>0.8</b>	<b>0.67</b>	<b>0.74</b>	<b>0.47</b>
SPI <sub>m</sub> -3/SPI <sub>w</sub> -13 and SMDI <sub>RS</sub>	<b>0.61</b>	<b>0.41</b>	<b>0.51</b>	<b>0.36</b>
SPI <sub>m</sub> -3/SPI <sub>w</sub> -13 and SMDI <sub>LSM</sub>	<b>0.41</b>	0.32	<b>0.21</b>	<b>0.29</b>
SPI <sub>m</sub> -6/SPI <sub>w</sub> -26 and ETDI <sub>RS</sub>	<b>0.72</b>	<b>0.45</b>	<b>0.57</b>	<b>0.21</b>
SPI <sub>m</sub> -6/SPI <sub>w</sub> -26 and ETDI <sub>LSM</sub>	<b>0.78</b>	<b>0.39</b>	<b>0.60</b>	<b>0.30</b>
SPI <sub>m</sub> -6/SPI <sub>w</sub> -26 and SMDI <sub>RS</sub>	<b>0.58</b>	0.19	<b>0.40</b>	<b>0.40</b>
SPI <sub>m</sub> -6/SPI <sub>w</sub> -26 and SMDI <sub>LSM</sub>	<b>0.30</b>	0.17	<b>0.16</b>	0.02
SPI <sub>m</sub> -12/SPI <sub>w</sub> -52 and ETDI <sub>RS</sub>	<b>0.81</b>	<b>0.75</b>	<b>0.48</b>	<b>0.31</b>
SPI <sub>m</sub> -12/SPI <sub>w</sub> -52 and ETDI <sub>LSM</sub>	<b>0.71</b>	<b>0.53</b>	<b>0.45</b>	<b>0.16</b>
SPI <sub>m</sub> -12/SPI <sub>w</sub> -52 and SMDI <sub>RS</sub>	<b>0.63</b>	<b>0.40</b>	<b>0.43</b>	<b>0.39</b>
SPI <sub>m</sub> -12/SPI <sub>w</sub> -52 and SMDI <sub>LSM</sub>	<b>0.24</b>	0.22	<b>0.15</b>	-0.03

\* Dry periods considered when SPI<0, ETDI<0, SMDI<0

940 **Table 1: Significant (in bold) correlation coefficients considering p-values=0.05 for pairs of indices at monthly and weekly scale, for the period 2010-2017 of the SPI i, ETDI and SMDI series, including results for the subset of dry periods in the series.**

2010-2017 - Monthly scale

		ETDI		SMDI	
		RS	LSM	RS	LSM
ETDI	RS		<b>0,77</b>	<b>0,58</b>	
	LSM	<b>0,77</b>			<b>0,32</b>
SMDI	RS	<b>0,58</b>			<b>0,27</b>
	LSM		<b>0,32</b>	<b>0,27</b>	
SPI	SPI <sub>m-1</sub>	<b>0,39</b>	<b>0,51</b>	<b>0,42</b>	<b>0,45</b>
	SPI <sub>m-3</sub>	<b>0,62</b>	<b>0,80</b>	<b>0,61</b>	<b>0,41</b>
	SPI <sub>m-6</sub>	<b>0,72</b>	<b>0,78</b>	<b>0,58</b>	<b>0,30</b>
	SPI <sub>m-12</sub>	<b>0,81</b>	<b>0,71</b>	<b>0,63</b>	<b>0,24</b>

2010-2017 - Monthly scale - Dry periods

		ETDI		SMDI	
		RS	LSM	RS	LSM
ETDI	RS		<b>0,60</b>	<b>0,50</b>	
	LSM	<b>0,60</b>			<b>0,43</b>
SMDI	RS	<b>0,50</b>			0,13
	LSM		<b>0,43</b>	0,13	
SPI	SPI <sub>m-1</sub>	0,23	<b>0,37</b>	0,52	<b>0,59</b>
	SPI <sub>m-3</sub>	<b>0,42</b>	<b>0,67</b>	<b>0,41</b>	0,32
	SPI <sub>m-6</sub>	<b>0,45</b>	<b>0,39</b>	0,19	0,17
	SPI <sub>m-12</sub>	<b>0,75</b>	<b>0,53</b>	<b>0,40</b>	0,22

2010-2017 - Weekly scale


		ETDI		SMDI	
		RS	LSM	RS	LSM
ETDI	RS		<b>0,55</b>	<b>0,50</b>	
	LSM	<b>0,55</b>			<b>0,21</b>
SMDI	RS	<b>0,50</b>			<b>0,19</b>
	LSM		<b>0,21</b>	<b>0,19</b>	
SPI	SPI <sub>w-4</sub>	<b>0,57</b>	<b>0,68</b>	<b>0,57</b>	<b>0,33</b>
	SPI <sub>w-13</sub>	<b>0,63</b>	<b>0,74</b>	<b>0,51</b>	<b>0,21</b>
	SPI <sub>w-26</sub>	<b>0,57</b>	<b>0,60</b>	<b>0,40</b>	<b>0,16</b>
	SPI <sub>w-52</sub>	<b>0,48</b>	<b>0,45</b>	<b>0,43</b>	<b>0,15</b>

2010-2017 - Weekly scale - Dry periods


		ETDI		SMDI	
		RS	LSM	RS	LSM
ETDI	RS		<b>0,34</b>	<b>0,46</b>	
	LSM	<b>0,34</b>			0,18
SMDI	RS	<b>0,46</b>			0,12
	LSM		0,18	0,12	
SPI	SPI <sub>w-4</sub>	<b>0,30</b>	<b>0,49</b>	<b>0,57</b>	<b>0,25</b>
	SPI <sub>w-13</sub>	<b>0,32</b>	<b>0,47</b>	<b>0,36</b>	<b>0,29</b>
	SPI <sub>w-26</sub>	<b>0,21</b>	<b>0,30</b>	<b>0,40</b>	0,02
	SPI <sub>w-52</sub>	<b>0,31</b>	<b>0,16</b>	<b>0,39</b>	-0,03

\* Dry periods considered when SPI<0, ETDI<0, SMDI<0

**Table 1: Matrices of significant (in bold) correlation coefficients considering p-values=0.05 for pairs of indices at monthly and weekly scale for the period 2010-2017 of the SPI I, ETDI and SMDI series, and the corresponding dry period subsets. Higher correlation values show a more intense red color. Since the correlations of interest refer to comparing the same type of data sources for different indices, dark grey cells identify unsuitable combinations for the analysis such as comparing RS results from one index with LSM results from another index.**




950 **Figure 2:** Time series of SPI, ETDI and SMDI indices at monthly scale (left column of subplots identified with sub-index 1) and weekly scale (right column of subplots identified with sub-index 2). a) to d) correspond to SPI<sub>m</sub>-1, SPI<sub>m</sub>-3, SPI<sub>m</sub>-6 and SPI<sub>m</sub>-12, e) and f) display the temporal evolution of the ETDI and SMDI. These ones show two rows corresponding to the series based on RS and LSM data. Red /blue color represents dry /wet periods. Periods of drought occur when SPI<0, ETDI>0, SMDI<0.




955

**Figure 3:** Lag plots of SPI-1, SPI-3, SPI-6 and SPI-12 (expressed as SPIw-4, 13, 26, and 52 weeks respectively) with remote sensing (RS) ETDIw index at weekly scale for the period 2010-2017. Lags are calculated for the -104 leading and +104 lagged time steps of ETDIw in reference to the SPIw-i. The height of the bars of the plot indicates the area of the basin affected by non-significant (grey scale) or significant (coloured scale) correlations. The saturation of the colored scale indicates the magnitude of the significant positive (red) or negative (blue) Pearson correlation coefficient of SPIw-i and ETDIw.

960



**Figure 4: Lag plots of SPI-1, SPI-3, SPI-6 and SPI-12 (expressed as SPIw-4, 13, 26, and 52 weeks respectively) with land-surface model (LSM) ETDIw index at weekly scale for the period 2010-2017. Lags are calculated for the -104 leading and +104 lagged time steps of ETDIw in reference to the SPIw-i. The height of the bars of the plot indicates the area of the basin affected by non-significant (grey-scale) or significant (coloured scale) correlations. The saturation of the colored scale indicates the magnitude of the significant positive (red) or negative (blue) Pearson correlation coefficient of SPIw-i and ETDIw.**



970 **Figure 5:** Lag plots of SPI-1, SPI-3, SPI-6 and SPI-12 (expressed as SPIw-4, 13, 26, and 52 weeks respectively) with remote sensing (RS) SMDIw index at weekly scale for the period 2010-2017. Lags are calculated for the -104 leading and +104 lagged time steps of SMDIw in reference to the SPIw-i. The height of the bars of the plot indicates the area of the basin affected by non-significant (grey-scale) or significant (coloured) correlations. The saturation of the colored scale indicates the magnitude of the significant positive (red) or negative (blue) Pearson correlation coefficient of SPIw-i and SMDIw.

975





Figure 6: Lag plots of SPI-1, SPI-3, SPI-6 and SPI-12 (expressed as SPIw-4, 13, 26, and 52 weeks respectively) with land-surface model (LSM) SMDIw index at the weekly scale for the period 2010-2017. Lags are calculated for the -104 leading and +104 lagged time steps of SMDIw in reference to the SPIw-i. The height of the bars of the plot indicates the area of the basin affected by non-significant (grey scale) or significant (coloured) correlations. The saturation of the colored scale indicates the magnitude of the significant positive (red) or negative (blue) Pearson correlation coefficient of SPIw-i and SMDIw.

980



985 **Figure 7:** Lag plots of ETDIw – SMDIw at weekly time scale in the period 2010-2017 for a) RS data and b) LSM data. Lags are calculated for the -104 leading and for the +104 lagged time steps of SMDIw in reference to the [ETDIwSPIw-4](#). The five levels of red and blue indicate the positive (red) or negative (blue) Pearson correlation coefficient between the ETDIw and the SMDIw for each time step (lead or lag time step). The height of the bars of the plot indicates the area of the basin affected by that level-level of  $r$  Pearson values.



ones (negative correlations of blue arrows in b)). (b) Summary of the annual magnitude and timing of the interactions. Red arrows represent positive correlations while blue arrows the negative ones. Arrow width represents the magnitude of the correlation. Arrow direction determines the direction of the interaction. The timeline ~~in-at~~ the bottom indicates the scale of the interactions ranging from short-term to long-term. (c) Sequences of prevalent reinforcing (upper) and inhibiting (lower) conditions alternating during the annual cycle based on the scheme of interactions described in a) and b). The upper sequence is the self-intensifying loop driven by evapotranspiration under high-energy conditions. The lower sequence displays the inhibiting role of rainfall ~~+~~ soil moisture under low-energy conditions.

995

1000