

Model Comparisons for Predicting Grassland Fire Occurrence Probability in Inner Mongolia Autonomous Region, China

Chang Chang^{1,2}, Yu Chang^{1,*}, Zaiping Xiong¹, Xiaoying Ping^{1,2}, Heng Zhang³, Meng Guo⁴, Yuanman Hu^{1,5}

5 ¹CAS Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110016, China

²University of Chinese Academy of Sciences, Beijing 100049, China

³ College of Forestry, Inner Mongolia Agricultural University, Hohhot 010019, China

⁴School of Geographical Sciences, Northeast Normal University, Changchun 130024, China

10 ⁵E'erguna Wetland Ecosystem National Research Station, Inner Mongolia, 022250, China

Correspondence to: Yu Chang (changyu@iae.ac.cn)

Abstract. Grassland fires threaten grassland ecosystem, human life and economic development greatly. However, there are limited researches focusing on grassland fire prediction, thereby it is necessary to find a better method to predict probability of grassland fire occurrence. Here we selected 16 environmental variables that may have impacts on fire occurrence, then built regression models of grassland fire probability based on historical fire points and variables in Inner Mongolia by three methods to identify the grassland fire drivers and to predict fire probability. The three methods were global logistic regression, geographically weighted logistic regression and random forest. The results showed that random forest model had the best predictive effect. The influence of 9 variables selected by geographically weighted logistic regression model on grassland fire was unbalanced spatially. The three models all showed that meteorological factors were of great importance to grassland fire occurrence. In Inner Mongolia, grassland fires occurred in different areas had various responses to the influencing drivers, and the areas with different natural and geographical condition had different fire prevention periods. Thus, the grassland fire management strategy based on local conditions should be advocated.

1 Introduction

Grassland ecosystems are precious natural resources and have crucial ecological functions, such as maintaining water, regulating local climate condition, providing food and herbs (Steiner et al., 2020; Jiang et al., 2018). Grassland ecosystems are known as an important green ecological safeguard (Pan et al., 2018). Fire is a common disturbance factor in grassland ecosystems (Thomson et al., 2020; Bond and Keeley, 2005). In long term evolutionary process, grassland ecosystems have established a harmonious and balanced relationship with fire. Fire has become an integrated part of grassland ecosystems (Lamont and He, 2017). Fire promotes grassland regeneration, succession, species evolution, and maintains biodiversity in ecosystems (Chandra and Bhardwaj, 2015). However, in the context of global climate change along with human activities, fire frequency is increasing and fire seasons are lengthening (Jolly et al., 2015). Fire has become a threat to biodiversity and



ecosystem stability in the last decades (Verbesselt et al., 2006). Grassland fire is also a sudden and highly destructive disaster that can not only alter the structure, function, pattern and processes of the landscape, but also affect the carbon cycle in grassland ecosystems (Yu and Zhuang, 2019). Besides, it poses threats to herdsman's lives, infrastructures, and valuable grassland resources (Podur et al., 2003). Prevention of grassland fires has become the main task of local governments to protect herdsman's lives and infrastructures. Accurate prediction of grassland fire occurrences could help to locate high fire risk areas for fire prevention and fighting.

Prediction of grassland fire occurrence needs to properly choose independent variables. With human population increasing and development of social economy, factors affecting grassland fires are becoming more complicated (Alexandre et al., 2015). Fire occurrence is the consequence of a combination of numerous anthropogenic and natural factors. Among natural factors, meteorological factors are commonly treated as the fire drivers (Oliveira et al., 2012), while vegetation condition or fuels load has important impact on grassland fire burning and spread. Some scholars added variables such as Spring NDVI, Autumn NDVI (Wu et al., 2020a), vegetation condition index (VCI) (Zapata-Rios et al., 2021), percentage of grassland, shrubland, forest area (Pavlek et al., 2017) that can reflect the vegetation condition or fuels load into the analysis of fire occurrence pattern. In addition, although topography factors show a relatively weaker contribution on fire occurrence pattern, scholars still preferred to select few variables such as topographic roughness index, elevation, slope, aspect index, and surface curvature in fire occurrence prediction (Su et al., 2021). Regarding for the anthropogenic factors, distance to nearest road and settlement is often used as proxy for the intensity of human activities (Liu et al., 2012) for the closer to road and settlement, the more human activities. Besides choosing variables, appropriate models are also needed to predict grassland fire occurrences.

Over the last decades, a lot of prediction models were developed to explore the relationship between fire occurrence and independent variables (Miranda et al., 2012; Wu et al., 2014). Global logistic regression (GLR) model is the most widely used conventional method on predicting probability of wildfire occurrence (Guo et al., 2016a). It assumes that the interaction between fire occurrence and its drivers is constant and stable in space. However, due to the environmental heterogeneity in different spatial locations, the relative importance of factors affecting fire occurrence may be different inevitably. This phenomenon is called spatial non-stationarity (Fotheringham, 2002). To solve this problem, geographically weighted logistic regression (GWLR) model was adopted to predict fire occurrence considering the different roles of variables across various spatial locations. GWLR model can deal with the various relationships between dependent variable and independent variable across geographical locations (Liang et al., 2017). Recently, to improve prediction accuracy, a variety of machine learning methods, including neural networks, support vector machines and random forests (Rodrigues and De La Riva, 2014; De Vasconcelos et al., 2001; Phelps and Woolford, 2021b) also have been used in fire occurrence prediction, among which Random Forest (RF) is deemed to be a flexible method to assess complex interactions among variables for it can automatically select the important variables, regardless of how many variables are input at the beginning, and can overcome the problem of over-fitting (Guo et al., 2016a; Oliveira et al., 2012). Although GWLR overcomes the shortcomings of GLR, and new machine

learning methods were proposed, model comparisons are commonly taken to determine the best-fitted model in fire prediction (Mohajane et al., 2021; Phelps and Woolford, 2021a). For examples, Guo (2016) and Oliveira (2012) both compared the GLR model to RF model in fire predictions, and found that RF model performs better than GLR model. Liang had compared the RF model and GWLR model, then got the conclusion that GWLR model had the better performance in forest fire forecasting Fujian province (Liang et al., 2017). Phelps and Woolford employed the logistic regression, bagged classification trees, random forest, neural network to predict the fire occurrence, however, they found all the models were unsuitable for fire occurrence prediction in their study area (Phelps and Woolford, 2021a).

China possesses nearly 3.92×10^6 km² area of grasslands, accounting for 12% of the world's grasslands and 41.7% of the national land area. Grassland is the main land cover type in Inner Mongolia Autonomous Region (Liu et al., 2018). Meanwhile, the Inner Mongolia grassland is the main component of the temperate grassland of Eurasia, where grassland fire is extremely active (Le Page et al., 2008). Till now, plentiful scholars always paid more attention to forest fire occurrence (Arnan et al., 2013; Matin et al., 2017; Renard et al., 2012; Wotton et al., 2010) and less to grassland fires. Nevertheless, the driving factors and spatial patterns of grassland fires in Inner Mongolia are currently unclear.

In this study, we aim to (1) explore the temporal and spatial distribution patterns of historical grassland fires from 2000 to 2018 in Inner Mongolia to get a whole view of fire points distribution; (2) predict grassland fire occurrence probability based on fire points and environmental variables by three models to identify the drivers of grassland fire occurrence, and to find the best fitting model by comparing the performances of the three models; (3) apply the three models to check whether they can predict probability of grassland fire occurrence in Inner Mongolia in 2014 accurately and put forward some suggestions on grassland fire management. Our results will facilitate formulating region-specific strategies for grassland resource protection and fire forecast, warning, as well as management. The paper is organized as follows. The second part introduces the general situation of study area, data source, data processing and study methods. In the third part, we present the study results including temporal and spatial distribution patterns of fire occurrences in study area, three regression models and their evaluation as well as applying the three models. The fourth part discusses the results and their implications for grassland fire management. The fifth part concludes the content of the study.

2 Method

2.1 Study area

Inner Mongolia autonomous region locates in the north of China (37°24'—53°23'N, 97°12'—126°04'E), and bordered by Mongolia and Russia. The distance between the east and the west is about 2400 km, and the span between the north and the south is about 1700 km. It has a long and narrow shape extending from the northeast to the southwest with total area of

118.3×10⁴km² (Fig.1a). There are Inner Mongolia plateau, Greater Khingan ge, Hetao Plain in study area. According to the Seventh National Census in China in 2020, the resident population of the district totalled 24.05 million.

100 Inner Mongolia belongs to temperate continental monsoon climate with the annual average temperature ranging from 0 to 8°C and annual precipitation from 50 to 450mm (Jia et al., 2020). The climate is characterized by large temperature difference, long sunshine duration, limited and imbalance precipitation. From northeast to southwest, it comprises temperate humid zone, semi-humid zone, semi-arid zone, arid zone and extreme arid zone (Li and Feng, 2019). The grassland vegetation in Inner Mongolia has a distinct zonal distribution pattern from northeast to southwest as well. Three main types of grassland are

105 temperate meadow grassland, temperate typical grassland, and temperate desert grassland (Zhou et al., 2016) (Fig.1b).




Figure 1: Geographic location of the study area. (a)The relative position and elevation of Inner Mongolia, (b)fire ignition points and vegetation type map, and (c) administrative map on the city level. (All the base map layers except ignition points were from © Institute of Geographic Sciences and Natural Resources Research, CAS).

110 2.2 Data source and pre-processing

2.2.1 Dependent variable

We used the historical grassland fire occurrence records as the dependent variable. These fire records came from Inner Mongolia fire management department that contained detailed information of each grassland fire in Inner Mongolia, including geographical location, fire starting time, fire size, fire cause, casualties and so on. There were 1049 fires recorded from 2000 to 2018, of which 685 fires had geographical coordinate information, thus the remains without geographical information were removed. The geographical distribution patterns of fire occurrences were shown in Fig.1b. Based on the 685 valid fire occurrence points, the non-fire occurrence points were created that had the 1.5 times number of fire points (Catry et al., 2009) by ArcGIS10.6. Therefore, the total sampling points were 1711. Then an attribute value of the points was coded 0 for the non-fire points, and 1 for fire points. This attribute was the dependent variable, which requires for coding in a binary format in logistic regression model (Wheeler, 2007).

2.2.2 Explanatory variables

Explanatory variables used to predict fire occurrence probability were divided into 4 categories, meteorological, anthropogenic activity, vegetation and topography. All the explanatory variables selected were as follows: mean annual temperature and precipitation, daily average wind speed, daily average temperature, daily average specific humidity, daily cumulative precipitation; distance to nearest settlement, distance to nearest road, distance to nearest river, distance to nearest railway, population density; Normalized Difference Vegetation Index (NDVI), global vegetation moisture index (GVMI); elevation, aspect and slope. The data source, abbreviation and unit of all the variables were listed in Table1.

Meteorological factors

Meteorology plays an important role on fire occurrence. Mean annual temperature and precipitation were selected as climate factors because they influence fuel moisture content which is a crucial factor for fire occurrences. In addition, they are traditional indicators for the degree of climate change (Scholze et al., 2006). Daily average temperature, wind speed, specific humidity and cumulative precipitation influence the moisture content of fuels more directly (Shmuel et al., 2022). Meteorological factors in this study came from China Meteorological Forcing Data (1979-2018) (Yang, 2018). NetCDF is the original format of the data, which can be disposed by ArcGIS. Mean annual temperature and precipitation, daily mean wind speed, daily mean temperature, daily mean specific humidity and daily cumulative precipitation, were extracted from this dataset. The value of each variable was extracted by the locations and date of the sampling points. In addition, daily meteorological data of non-fire points was extracted by a random date without fire occurrence in the period of 1990 to 2018.

Anthropogenic factors

Human activities are closely related to the human-caused fire occurrence (Vilar et al., 2010). The distance to human infrastructure, such as road, railway and settlement is usually used to represent the intensity of human activity (Liu et al., 2012;

Chang et al., 2013). In Inner Mongolia, herdsman prefer to feed water for livestock by rivers, so the closer to the river, the greater the intensity of human activity. Based on this, we selected the distance to nearest settlement, road, railway and river as the anthropogenic factor. Digital maps (1:250,000) of railway, road, river, and resident points were collected from National Catalogue Service for Geographic Information. We calculated the Euclidean distances from each fire location to the nearest railway, road, river, and settlement. These distances were acted as proxies of intensity of human activities. Then the values of these distances were extracted by the locations of the fire records. To some extent, population density showed strong correlation with human activity intensity. Population density data of counties in the whole study area was drawn into raster layers and was extracted by the locations of the sampling points.

150 *Vegetation factors*

NDVI can be used to indicate the gross of live and dead fuels above the surface (Purevdorj et al., 1998). Normalized Difference Vegetation Index (NDVI) has the similarity with FVC, so it can be replaced by FVC under some circumstance. Global Vegetation Moisture Index (GVMI) can provide direct information on vegetation water content at canopy level (Ceccato et al., 2002a) thus reflect the moisture content of live fuels. Therefore, two index, NDVI and GVMI were selected to represent fuel load and fuel moisture conditions respectively. We calculated the average NDVI value of the years from 2000 to 2018, creating a NDVI layer. GVMI is based on combination of near-infrared and short-wave infrared bands of the electromagnetic spectrum to estimate vegetation moisture content (Ceccato et al. 2002). Band 2 and band 6 of product MOD09A1 of MODIS surface reflectance was used. The method to calculate the GVMI can be found in Ceccato's research (Ceccato et al., 2002b).

Topographic factors

160 The topographic factors affect vegetation distribution, composition and flammability (Syphard et al., 2008). The higher the elevation, the lower the temperature. Different aspect will receive imbalance radiation from the sun. Hence, the vegetation type changes with elevation and aspect. A study found that human-caused fires occurred more frequently in gentle slopes (Conedera et al., 2011). Elevation data came from Geospatial Data Cloud. The 90m spatial resolution digital elevation model (DEM) was adopted then the processes of spatial correction mosaic, clip, and resample on DEM were indispensable. By the 3D Analyst tool of ArcGIS10.6, raster layers of slope and aspect can be obtained. Aspect needed to be exponentiated, which was convenient for analysis. The aspect was converted into aspect index according to specific method provided by Su (Su et al., 2021).

170 Explanatory variables could be extracted from the corresponding map layers, the value of each explanatory variable was extracted as an attribute of the sampling points to form the completed sampling dataset collection, then we randomly chose 60% from fire points and non-fire points respectively as training dataset, and the remained 40% as test dataset.

Table1 Explanatory variables and their sources

Factors	Variables	Abbreviation	Data source	Units	Resolution
	Mean annual temperature	Temp		K	0.1°


Meteorological factors	Mean annual precipitation	Prec	China Meteorological Forcing Data (1979-2018) (Yang, 2018)	m/s	0.1 °
	Daily average specific humidity	Humi_dy		kg/kg	0.1 °
	Daily cumulative precipitation	Prec_dy		mm/hr	0.1 °
	Daily average temperature	Temp_dy		K	0.1 °
	Daily average wind speed	Wind_dy		m/s	0.1 °
Anthropogenic factors	Distance to nearest settlement	D_resp	National Catalogue Service for Geographic Information (https://www.webmap.cn)	km	500m
	Distance to nearest road	D_road		km	500m
	Distance to nearest river	D_river		km	500m
	Distance to nearest railway	D_rail		km	500m
	Population density	P_density	National Bureau of Statistics of China (http://www.stats.gov.cn)	per/km ²	500m
Vegetation factors	Global vegetation moisture index	GVMI	Level-1 and Atmosphere Archive & Distribution System Distributed Active Archive Center (https://ladsweb.modaps.eosdis.nasa.gov/)	-	500m
	Normalized Difference Vegetation Index	NDVI	Resource and Environment Science and Data Center (Xu, 2018)	-	1000m
Topographic factors	elevation	Elev	Geospatial Data Cloud (http://www.gscloud.cn)	Meter	500m
	aspect index	Aspect		-	500m
	slope	Slope		degree	500m






2.3 Data Analysis methods

175 2.3.1 Multicollinearity diagnosis between explanatory variables.


If there is strong multicollinearity among explanatory variables, the regression fitting may produce biased parameter estimation, leading excessive standard error of regression coefficient, thus the model will be unreliable (Wheeler, 2007; Chang et al., 2013). Variance inflation factor (VIF) is a common method to evaluate multicollinearity among the variables. It is stated that if $VIF > 10$, there is severe collinearity; $5 < VIF < 10$, there is moderate collinearity; $2 < VIF < 5$, there is mild collinearity (Krebs et al., 2012). Therefore, we did a multicollinearity diagnosis for our sampling dataset by VIF. The VIF test was manipulated by car package of R studio 4.0.3. In this study, the VIF value of variables Temp, Prec, Humi_dy, Prec_dy, Temp_dy, Wind_dy, D_resp, D_road, D_river, D_rail, P_density, GVMI, NDVI, Elev, Aspect, Slope were 1.024, 5.292, 1.8434, 1.253, 1.613, 1.140, 1.396, 1.430, 1.507, 1.352, 1.189, 1.103, 5.381, 1.800, 1.032, 1.490 respectively. This indicated that there was moderate

collinearity for Prec variable and NDVI variable with their VIF greater than 5. In consideration of this, we deleted Prec variable
185 in the global logistic regression and geographically weighted logistic regression to ensure the model reliability. 

2.3.2 Trend analysis of grassland fires

In order to get a concrete and overall distribution of all the historical fire occurrence points, we need a tool to present them
clearly. Trend analysis is an analytical tool in ArcGIS geographic statistics module, which can analyse the spatial distribution
trend of data (Mou, 2012). In order to find the spatial distribution patterns of grassland fire occurrences, the Inner Mongolia
190 map was divided into 6064 grids of 20k k by the create fishnet tool, then the fire points in each grid were calculated. Trend 
analysis in ArcGIS was used to get a three-dimension overview of fire points in geographical space. Additionally, to find the
characteristics of time change in a year, sample quantity statistics was done at the half-month interval according to the fire
occurrence date from 2000 to 2018. 

2.3.3 Modelling methods

195 Regression methods have been widely applied in fire occurrence modelling, such as linear regression (Oliveira et al., 2012) or
logistic regression (Phelps and Woolford, 2021b; Chang et al., 2013). These models are often built with the goal of using the
fewest predictors to explain the greatest variability in the dependent variable (Graham, 2003). Random forest model has 
showed its strong predictive ability in fire occurrence probability (Guo et al., 2016b; Oliveira et al., 2012). Therefore, we
selected the logistic regression and random forest model to build our fire occurrence models. Geographically weighted logistic
200 regression shows its ability on dealing with spatial heterogeneity, so we picked it. The following is an introduction to the three
models.

Global logistic regression (GLR)

Generalized Linear Models are widespread methods in plenty of research fields and logistic regression is one of the most
popular approaches in occurrence modelling in fire science (Bar Massada et al., 2013). When predicting a binary variable
205 through a series of continuous or categorical predictor variables, logistic regression is a useful tool. LGR assumes that the
samples are independent and have a constant relationship between the dependent variable and the independent variables in the
entire study area (Zhang et al., 2013). The global logistic regression was performed by AER package in R studio4.0.3.


Geographically weighted logistic regression (GWLR)

210 GWLR firstly assumes that the model structure is spatially non-stationary which means the relationship between the binary
dependent variable and the continuous independent variables changes with geographic locations. Then, it tests whether the
spatial non-stationary relationship of this hypothesis is significant. LGWR shares similarity with its global counterpart,
producing the β regression coefficients and significance tests. It is worth noting that rather than a single parameter, a collection
of parameters for each point or polygon can be obtained. Therefore, such model often outperforms global regression models
(Fotheringham, 2002). A conventional GWR is described by the following expression:

215

$$y_i = \sum_k \beta_k(a_i, b_i) x_{k,i} + \varepsilon_i \quad (1)$$

where y_i , $x_{k,i}$ and ε_i are, dependent variable, k th explanatory variable, and the Gaussian error at location i respectively; (a_i, b_i) is the X/ Y coordinate of the i th location; and coefficients $\beta(a_i, b_i)$ are varying depending on the location.

 We used the GWR4 software freely accessible to build GWLR model and chose Adaptive bi-square as kernel type, golden section search as the bandwidth selection method. Generally, the fixed kernel specifies an equal distance threshold for each regression point, while the adaptive kernel specifies the number of neighbours to be considered for each regression point. Therefore, fixed kernels approach should be suitable in a circumstance where the points are regularly distributed in space and the adaptive approach is more appropriate for spatially clustered points (Nunes et al., 2016). According to the spatial distribution pattern of our fire occurrence data, Adaptive approach was selected to build model. Akaike information criterion (AIC) approach was used to assess the goodness-of-fit for each model.

225 *Random forest (RF)*

Random forest model is a kind of supervised machine learning algorithms based on decision trees. RF model gathers a large number of classification trees to improve the prediction accuracy of the model. It is unnecessary to set the function form in advance, and also can overcome the complex interactions between covariables to get a high classification accuracy (Gao et al., 2020). It can handle a large number of independent variables and pick out important ones. Besides, it is not affected by the multicollinearity between variables and can evaluate the complex interactions between variables more flexibly (Ma et al., 2020). Random forest has the advantages of having higher accuracy than individual decision trees and low sensitive to parameter adjustment than other machine learning models (Su et al., 2020). However, random forest model also has the limitation that it can't calculate the specific regression coefficient and confidence interval. We built random forest model by randomForest package in R studio4.0.3.

235 **2.3.4 Model evaluation methods**

To evaluate the goodness of fit of the three models we chose, four statistical measures, akaikie information criterion (AIC), area under curve (AUC), mean absolute error (MAE) and R^2 , were selected. AIC is based on the concept of entropy, which can weigh the complexity of the estimated model and the goodness of model fitting data. It is an index obtained in the process of model fitting. The receiver operating characteristic (ROC) curve was obtained by plotting sensitivity versus specificity for various probability thresholds. The area under the curve (AUC) is also often used to evaluate model performance (Jiménez-Valverde, 2012). MAE is the average value of absolute error, which can better reflect the actual situation of predicted value error. MAE is defined in the following expression:

$$MAE = \frac{1}{n} \sum_{i=1}^n |\hat{y}_i - y_i| \quad (2)$$

R^2 can reflect the fitting degree of the regression line to the observed value, which can be calculated by the following equation:

$$R^2 = 1 - \frac{\sum_{i=1}^n (\hat{y}_i - y_i)^2}{\sum_{i=1}^n (\hat{y}_i - \bar{y}_i)^2} \quad (3)$$

where n is the number of samples, \hat{y}_i is fire occurrence probability predicted by the model, y_i is a binary value representing whether or not a fire exists, \bar{y}_i is the arithmetic mean of binary values.

3 Results

3.1 Temporal and spatial distribution patterns of fire occurrences

250 From a spatial perspective, the fire points gathered in the east and central parts of Inner Mongolia, and the east part had more fire points than the central, while the west part had limited fire points in longitude direction. The fire points distribution was more balanced in latitudinal direction, except for the northernmost part where no fire occurred due to the lack of grassland (Fig.2).

255 From a temporal perspective, there were two peaks of fire occurrences during a year. The first one appeared in the first half of April, and the second appeared in the first half of October (Fig.3). The number of fires during the first fire prevention period was significantly higher than that during the second fire prevention period. According to the *Regulations on Fire Prevention of Forests and Grasslands in the Inner Mongolia Autonomous Region*, there are two fire prevention periods, March 15th to June 15th, and September 15th to November 15th. We found that the periods when the number of fires was greater than 20
260 coincided with the fire prevention periods regulated in the region.





Figure 2. The spatial distribution of grassland fires in Inner Mongolia from 2000 to 2018. (The height of the orange line represents number of fires; the grey point is the position of the observation point and the projection on the XY plane. The green point is the projection of the observations on the XZ plane; the blue point is the projection of the observations onto the YZ plane)



265

Figure 3. Changes of the number of grassland fires in Inner Mongolia at the half-month interval. The grey dotted line is the reference line for 20 fires.

3.2 Model fitting and evaluation

Based on GLR model, only 9 variables had significant coefficients ($p < 0.05$) among the 16 explanatory variables, namely Temp, Humi_dy, Wind_dy, D_river, D_resp, D_rail, GVMI, NDVI and Slope that were regarded as grassland fire drivers. Among the drivers, only NDVI and Wind_dy were positively correlated with fire occurrences, and the rest of drivers were negatively correlated with fire occurrences (Table 2).


270

Table 2 Parameter estimation of explanatory variables by the GLR model

Variable	Abbreviation	Coefficient	Standard error	P-value
Intercept	Intercept	38.543	10.218	<0.0001
Mean annual temperature	Temp	-0.138	0.036	<0.0001
Distance to nearest river	D_river	-0.009	<0.0001	<0.0001
Distance to nearest settlement	D_resp	-0.142	<0.0001	<0.0001
Distance to nearest rail	D_rail	-0.004	<0.0001	0.024
Global vegetation moisture index	GVMI	-7.523	1.650	<0.0001
Normalized Difference Vegetation Index	NDVI	2.599	0.560	<0.0001
Daily average specific humidity	Humi_dy	-857.033	81.818	<0.0001
Daily average wind speed	Wind_dy	0.217	0.050	<0.0001

Slope	Slope	-0.108	0.024	<0.0001
-------	-------	--------	-------	---------

275 The drivers selected by the GWLR model were consistent with the GLR model. The GWLR model spatialized the coefficient of each of the selected explanatory variables (Fig.4). These coefficients demonstrated large spatial variations (Table 3).



280 Figure 4. The spatial distribution of significant areas of the estimated coefficient for each variable selected by GWLR model. These maps were drawn according the t-value for each pixel. The pixels with the corresponding t-value between -1.96 and 1.96 (insignificant area) were not displayed on the graph. Those pixels with the t-value greater than 1.96 or less than -1.96 (significant area) were displayed on the graph. The negative coefficient values were coloured in cool colours, and the positive coefficient values were coloured in warm colours.

Table 3 Coefficient statistics of explanatory variables selected by the GWLR model

Variable	Mean	Min	1Q	Median	3Q	Max
Intercept	30.55973	8.352316	14.07005	28.78026	40.13325	109.0789
Humi_dy	-711.947	-1407.05	-1115.85	-1037.32	-26.1174	262.0729
Wind_dy	0.204153	-0.37813	-0.21256	0.123083	0.534187	1.327401
NDVI	1.294488	-1.53458	0.749144	1.351105	1.899675	4.123084

GVMI	-7.48481	-9.86252	-8.10707	-7.24877	-6.92158	-3.26509
Temp	-0.10685	-0.38292	-0.14078	-0.09896	-0.0535	-0.02163
Slope	-0.06249	-0.10811	-0.0675	-0.0622	-0.05487	-0.03551
D_rail	-0.00752	-0.0189	-0.01085	-0.00803	-0.00173	-0.00097
D_resp	-0.11175	-0.25096	-0.14876	-0.12328	-0.05028	0.016634
D_river	-0.00949	-0.02133	-0.0166	-0.00881	-0.00415	0.002383

285 Among the meteorological factors, the significant area of Humi_dy was distributed in the eastern and central area of the study area, accounting for more than 64.3% of the total area. The positively significant area of Wind_dy located in the northernmost and westernmost area, including Hulunbuir and almost the entire Alxa, the negatively significant areas were concentrated in the middle. Temp was significant in some parts of the east, showing a negative correlation. Among the vegetation factors, GVMI was more significant in the eastern region with small areas distributing in the central and western regions. NDVI


290 positively significant correlated with fire occurrence in a small area in the west. As for the topographic factors, only Slope was negatively correlated with significant area accounting for only 15% of the study area, scattered in the eastern and central regions. Among anthropic factors, the significant area of D_resp was found in Hulunbuir and the western region, and had a greater impact on fire occurrences in the west. D_river was significantly negatively correlated in the western Inner Mongolia and the west of Hulunbuir. The significant area of D_rail resided in southern Inner Mongolia.

295

The RF model picked all 16 explanatory variables into the model during the fitting process. In order to understand the importance of each variable in the model, the Mean Decrease Accuracy of IncNodePurity was used to calculate the importance of each variable, then the importance ranking of variables was shown in Fig.5. In general, the importance of each explanatory variable was quite different. The importance of the two daily-scale meteorological variables, Humi_dy and Temp_dy, was


300 much greater than other variables. In the top 6 of importance ranking, there were five meteorological variables, indicating the meteorological conditions were the main cause of the grassland fire occurrences. The importance of NDVI variable which could reflect the load of fuels, was relatively high in accordance with expectation. Among the three topographical variables, Slope and Aspect were of lower importance, indicating that topographical conditions have less impact on the occurrences of grassland fires in Inner Mongolia.





305

Figure 5. The importance ranking of the explanatory variables in RF model

Table 4 listed several statistical measures to evaluate the performance of GLR, GWLR and RF models. The RF model as a machine learning algorithm cannot obtain the AIC value, so only the GLR and GWLR models were compared according to AIC. The AIC value of GWLR model was smaller than the GLR model, which meant GWLR model had a higher goodness of fit. The AUC value of GLR, GWLR and RF model was 0.841, 0.909 and 0.944; MAE value was 0.148, 0.111 and 0.053; R² was 0.363, 0.537, and 0.779 respectively. By comparisons we believed that RF model performed better than both GWLR model and GLR model, and could more accurately predict grassland fire occurrences in the region. 

310

Table 4 Comparisons of the fitting goodness of three models

Model	Akaike information criterion (AIC)	Area under Curve (AUC)	Cut-off value	Mean Absolute Error (MAE)	R ²
GLR	1018.496	0.841	0.475	0.148	0.363
GWLR	807.330	0.909	0.471	0.111	0.537
RF	–	0.944	0.500	0.053	0.779

3.3 Model application

315 According to RF model, daily meteorological conditions were of high importance compared to other explanatory variables. The vast territory of Inner Mongolia and geographical features lead to great spatial variations in meteorological conditions, which would result in greater differences in the probability of fire occurrence in east and west region. Therefore, we randomly chose two points, one in the eastern and the other in the western area. Point 1 was located in the Hulunbuir grassland in northeastern Inner Mongolia, and Point 2 was located in the Ordos Plateau, both of which were areas with a high historical fire

320 points density. Based on the 19-year fire records obtained, we found that 2014 was the year with the highest number of grassland fires, with a total of 111 occurrences. For this reason, the year of 2014 was taken an example. We used the three models to estimate their probability of fire occurrence with daily meteorological variables in 2014 and other explanatory variables.

325 Fig.6a showed the estimated probability of fire occurrence by the three models at Point 1, and Fig.6c at Point 2 on each day in 2014. We found that the estimated probability of fire occurrence by GLR and GWLR were higher than RF at both locations. During none fire prevention periods, the probability by GLR and GWLR were also much higher while lower by RF, indicating that RF model could reasonably capture the fire prevention periods. In addition, we also found that the probability of fire occurrence estimated by each model at Point 1 was higher than Point 2. This may be due to higher fuel loads at Point 1 than

330 Point 2.




Figure 6. The fire occurrence probability of two typical points in Inner Mongolia in 2014. (a) and (c) are the daily fire occurrence probability of two typical points in 2014. The grey area is the fire prevention periods, and the grey lines is the value of 0.5; (b) is the kernel density level of historical grassland fire occurrences.

4.1 Temporal and spatial distribution patterns of historical grassland fires

Historical fires were densely distributed in eastern and central Inner Mongolia in space, because typical grassland and meadow grassland were widely distributed here (Fig.2c), which was an area with sufficient fuel. Besides, there were dense traffic routes, and the intensity of human activities is higher than that in western Inner Mongolia. In the northern part of Hulunbuir, the main vegetation type was boreal deciduous coniferous forest (Fig.2c), so grassland fires rarely occurred. In temporal, historical fires was concentrated in spring and autumn (Fig.3), which is closely related to the local climate characteristics. The area was dominated by temperate continental monsoon climate (Xue et al., 2021). In spring, the temperature rose rapidly, the wind speed was high, and the air humidity was low, leading to frequent fires. In autumn and winter, air was dry, increasing the number of fires. However, because of the high latitude, snow covered dead fuel in winter, reducing the probability of fire occurrence (Zhao et al., 2009).

4.2 Model comparison in goodness of fitting and predictive ability

Model comparisons have become a hot topic in choosing most suitable prediction models in the last decades (De Vasconcelos et al., 2001; Phelps and Woolford, 2021a; Guo et al., 2016b; Phelps and Woolford, 2021b). In our study, the fitting goodness of GLR model was not competitive with other two models (GWLR and RF) (Table 4). Our results were consistent with previous fire prediction studies (Guo et al., 2016b; Phelps and Woolford, 2021a; Su et al., 2021) which demonstrated that GWLR and RF performs better than GLR in predicting fire occurrences. In addition, the GLR model improperly predicted high probability of fire occurrences during winter at both locations (Fig.6a and c). Temporal distribution of historical fire occurrences showed only fewer fires occurred in winter (Fig.3), suggesting that the prediction ability of GLR model was not ideal in our study area. This may be caused by other crucial environmental variables we did not consider in our study, such as water deficit (Syphard et al., 2018), drought (Rodrigues et al., 2018) etc. needing further to explore in future studies.

Regarding to the GWLR model, it is an improvement of the GLR model by adding a spatial weight matrix to reduce model residuals (Monjarás-Vega et al., 2020), and therefore the goodness of fitting was better than GLR (Table 4). However, the GWLR model also erroneously predicted high probability of fire occurrences during winter and summer at both locations (Fig.6a and c), which was on the contrary to the historical fire statistics (Fig.3). We speculated that GWLR model focused excessively on more significant meteorological variables and ignored the effects of other variables on fire occurrence.

As for the RF model, it had the best goodness of fitting among the three models (Table 4). In addition, the predictions for the two locations generally captured the seasonality of fire probability in our study area, which was higher during spring and autumn, and lower in winter and summer (Shabbir et al., 2020). The predictions were also consistent with the fire prevention periods (Fig.6). Therefore, RF model could be used to predict grassland fire occurrences in our study area.

4.3 Factors affecting grassland fire occurrences

Many factors may impact fire occurrences, including vegetation conditions, meteorological/climatic factors, topography and human activities (Muller et al., 2020; Parajuli et al., 2020; Sousa et al., 2021). In our study, meteorological/climatic factors (daily average specific humidity, daily average wind speed and mean annual temperature) showed higher importance as revealed by the GWLR models. Our results were in line with previous studies (Chang et al., 2013; Guo et al., 2017). Daily average specific humidity had a negative relationship to fire occurrences (Table 2 and Table 3) as with the increase of air humidity, the fuel moisture content also increases, which makes the fuel hard to ignite (Masinda et al., 2021). However, the importance of daily average specific humidity had great spatial variations. More importance was observed in the eastern region (Fig.4a) since the eastern region was near the sea, the spatial difference of daily average humidity was more obvious.

Generally speaking, the higher the wind speed, the more water vapor produced by plant transpiration will be taken away, and the more sufficient oxygen will be brought to the combustion of fuels (Masinda et al., 2020), thereby increasing the probability of fire occurrences. Therefore, in this study, daily average wind speed was positively related to fire occurrences (Table2). However, this positive correlation was not spatially uniform with some parts of the central area showed a negative correlation (Fig.4b). We speculated that this may be due to the characteristic of temperate continental monsoon climate. When the wind blows from sea to continental, and brings humid airflow, thereby reducing the probability of fire occurrences.

It is **generally believed** that temperature positively correlated with fire occurrences (Syphard et al., 2019). On the contrary, negative relation was showed in our study (Table 2 and Table 3) because the climate characteristics of rain and heat synchronization, and fuel moisture is high and uneasy to be ignite in hot weather. In addition, this negative relation was also spatially non-stationary. In some parts of eastern Inner Mongolia, significant difference was observed (Fig.4c) due to the greater differences of natural geographical features and cultural landscapes in our study area (Rodrigues et al., 2018; Su et al., 2021).

Regarding to anthropogenic factors, the three significant human-related variables were all negatively correlated with fire occurrence as expected (Table 2 and Table 3). Our results were consistent with previous researches (Liu et al., 2012). Historical fires records showed that majority of fires were human-originated. Human activities are more common near roads, settlements and rivers resulting in more fire occurrences (Chang et al., 2013). Therefore, the negative relations of human factors with fire occurrences were observed. In addition, these relations were also spatially heterogeneous (Fig. 4g, h, i) due to the variations in spatial distribution of roads, rivers and settlements.

As for vegetation conditions, fuel moisture and fuel loads have crucial impacts on fire occurrences (Zhang, 2014; Wu et al., 2020b; Naderpour et al., 2021). We found that the global vegetation moisture index (GVMI) was negatively correlated with

400 fire occurrence (Table 2 and Table 3), because with the increase of fuel moisture, the fuel is hard to ignite (Su et al., 2021). In addition, significant coefficients of GVMI were located in the eastern region (Fig.4d). This could be explained by the climate constrains hypothesis (Krawchuk and Moritz, 2011) as the fuel in the eastern region is abundant and climate conditions determine fire occurrences. Our results also showed that NDVI was positively correlated with fire occurrences (Table 2 and Table 3) for sufficient fuels is one of the burning conditions (Guo et al., 2017). However, spatially, only a small part of the western Inner Mongolia had significant NDVI coefficients (Fig.4e). This may support the fuel constrains hypothesis (Krawchuk and Moritz, 2011). Vegetation in the western Inner Mongolia is sparse. Once vegetation coverage increases, it would significantly increase the fire occurrence probability.

410 Finally, topographic factors also affect fire occurrence patterns (Su et al., 2019; Naderpour et al., 2021). In this study, we found that the slope and fire occurrence showed a negative correlation in some parts (Table 2 and Table 3). Our results were consistent with the results obtained by Zhang (Zhang, 2014). We believed this was due to the fact that grasslands of Inner Mongolia were mainly distributed on the flat plateau lacking of steep slopes, resulting in a negative correlation between grassland fire occurrences and Slope.

4.4 Implications for grassland fire management

415 From a spatial perspective, our results also showed spatial non-stationarities for various variables in predicting fire occurrences. For examples, the influence of Humi_dy, Wind_dy, GVMI and D_resp on fire occurrences was significant in the eastern Inner Mongolia, and NDVI, D_resp and D_river in the western Inner Mongolia (Fig.4). The spatial non-stationarity of variables may have implications for accurate mapping fire risk zones to fire management based on different environmental factors for various regions, instead of the traditional mapping methods which regard a variable having equal importance across a studied area.

420 Furthermore, our results showed that RF model performed best in our study area, which should be used to map fire occurrence probability. The spatial pattern of the probability of fire occurrence is of crucial importance in fire management (Sağlam et al., 2008). It could be applied to allocate priority areas for grassland fuel treatments (Wei et al., 2008). From a temporal perspective, fire prevention period in spring and autumn should be adjusted according to local conditions. For example, Point 1 was taken from Hulunbuir region, which has high latitude and insufficient heat. According to RF model prediction, the period of high fire probability in autumn started and ended earlier, and the period of high fire probability in spring was shorter (Fig. 6a).

425 Therefore, the fire prevention period should be adjusted accordingly.

In our study, we found that majority of grassland fires were human-originated according to the historical fire records, and the peaks of fire occurrences were consistent with the fire prevention periods in Inner Mongolia. Therefore, strict measures should be taken to prevent fires from occurring, such as forbidding any wildland fire uses during fire prevention periods, building fire monitor systems to timely detect fires so as to extinguish them at the earliest time. In addition, with the increasing intensity of human activities, a large area of grassland was occupied by cultivated land, and the increasing number of agricultural straw

burnings would bring more dangers for grassland. In the circumstance of global warming, the drivers of grassland fire would become more complex. Grassland management strategies must pay more attention to the changes of these drivers and their relationship with grassland fires to ensure effective fire prevention.

5 Conclusions

This study analysed the temporal and spatial characteristics and driving factors of grassland fires in Inner Mongolia Autonomous Region from 2000 to 2018, and compared the prediction accuracy of global logistic regression, geographically weighted logistic regression and random forest models. The spatial distribution of historical fires in Inner Mongolia grassland was more in the east than in the central than in the west, and the fire high-incidence season was generally same as fire prevention period regulated for Inner Mongolia forest and grassland. The fitting goodness of random forest model was the highest, and geographically weighted logistic regression models also had reference significance to some extent. The influence of 9 drivers selected by geographically weighted logistic regression model on grassland fire was spatially unbalanced. The random forest and geographically weighted logistic regression both showed that meteorological factors were of great importance to grassland fire. Inner Mongolia had a vast territory, and different areas had different sensitivities to different drivers. Areas with different hydrothermal condition had different fire prevention periods, thus the grassland fire management strategy based on local conditions should be advocated.

Competing interests. The contact author has declared that neither they nor their co-authors have any competing interests.

Acknowledgments. This research was funded by the National Key Research and Development Program of China Strategic International Cooperation in Science and Technology Innovation Program (2018YFE0207800) and the National Natural Science Foundation of China (grant no. 31971483). We thank the anonymous reviewers for their constructive suggestions to improve the manuscripts.

References

- Alexandre, P. M., Mockrin, M. H., Stewart, S. I., Hammer, R. B., and Radeloff, V. C.: Rebuilding and new housing development after wildfire, *International Journal of Wildland Fire*, 24, 138-149, 10.1071/wf13197, 2015.
- Arnan, X., Quevedo, L., and Rodrigo, A.: Forest fire occurrence increases the distribution of a scarce forest type in the Mediterranean Basin, *Acta Oecologica*, 46, 39-47, <http://dx.doi.org/10.1016/j.actao.2012.10.005>, 2013.
- Bar Massada, A., Syphard, A. D., Stewart, S. I., and Radeloff, V. C.: Wildfire ignition-distribution modelling: a comparative study in the Huron-Manistee National Forest, Michigan, USA, *International Journal of Wildland Fire*, 22, 174-183, 10.1071/wf11178, 2013.

- Bond, W. J. and Keeley, J. E.: Fire as a global 'herbivore': the ecology and evolution of flammable ecosystems, *Trends Ecol Evol*, 20, 387-394, 10.1016/j.tree.2005.04.025, 2005.
- 465 Catry, F. X., Rego, F. C., Bação, F. L., and Moreira, F.: Modeling and mapping wildfire ignition risk in Portugal, *International Journal of Wildland Fire*, 18, 921-931, doi:10.1071/WF07123, 2009.
- Ceccato, P., Flasse, S., and Gregoire, J. M.: Designing a spectral index to estimate vegetation water content from remote sensing data - Part 2. Validation and applications, *Remote Sensing of Environment*, 82, 198-207, 10.1016/s0034-4257(02)00036-6, 2002a.
- 470 Ceccato, P., Gobron, N., Flasse, S., Pinty, B., and Tarantola, S.: Designing a spectral index to estimate vegetation water content from remote sensing data: Part 1 - Theoretical approach, *Remote Sensing of Environment*, 82, 188-197, 10.1016/s0034-4257(02)00037-8, 2002b.
- Chandra, K. K. and Bhardwaj, A. K.: Incidence of forest fire in India and its effect on terrestrial ecosystem dynamics, nutrient and microbial status of soil, *Int. J. Agric. For.*, 5, 69-78, 2015.
- 475 Chang, Y., Zhu, Z., Bu, R., Chen, H., Feng, Y., Li, Y., Hu, Y., and Wang, Z.: Predicting fire occurrence patterns with logistic regression in Heilongjiang Province, China, *Landscape Ecology*, 28, 1989-2004, 10.1007/s10980-013-9935-4, 2013.
- Conedera, M., Torriani, D., Neff, C., Ricotta, C., Bajocco, S., and Pezzatti, G. B.: Using Monte Carlo simulations to estimate relative fire ignition danger in a low-to-medium fire-prone region, *Forest Ecology and Management*, 261, 2179-2187, 10.1016/j.foreco.2010.08.013, 2011.
- 480 de Vasconcelos, M. J. P., Silva, S., Tome, M., Alvim, M., and Pereira, J.: Spatial prediction of fire ignition probabilities: comparing logistic regression and neural networks, *Photogrammetric engineering and remote sensing*, 67, 73-81, 2001.
- Fotheringham, A. S., Brunson, C., Charlton, M.E: *Geographically weighted regression: the analysis of spatially varying relationships*, John Wiley & Sons Ltd, Chichester, UK2002.
- Gao, C., Lin, H.-L., Hu, H.-Q., and Song, H.: A review of models of forest fire occurrence prediction in China, *Ying yong sheng tai xue bao = The journal of applied ecology*, 31, 3227-3240, 10.13287/j.1001-9332.202009.014, 2020.
- 485 Graham, M. H.: Confronting multicollinearity in ecological multiple regression, *Ecology*, 84, 2809-2815, 10.1890/02-3114, 2003.
- Guo, F., Selvalakshmi, S., Lin, F., Wang, G., Wang, W., Su, Z., and Liu, A.: Geospatial information on geographical and human factors improved anthropogenic fire occurrence modeling in the Chinese boreal forest, *Canadian Journal of Forest Research*, 46, 582-594, 10.1139/cjfr-2015-0373, 2016a.
- 490 Guo, F., Su, Z., Wang, G., Sun, L., Tigabu, M., Yang, X., and Hu, H.: Understanding fire drivers and relative impacts in different Chinese forest ecosystems, *Sci Total Environ*, 605-606, 411-425, 10.1016/j.scitotenv.2017.06.219, 2017.
- Guo, F., Wang, G., Su, Z., Liang, H., Wang, W., Lin, F., and Liu, A.: What drives forest fire in Fujian, China? Evidence from logistic regression and Random Forests, *International Journal of Wildland Fire*, 25, 10.1071/wf15121, 2016b.
- 495 Jia, Y., Cui, X., Liu, Y., Liu, Y., Xu, C., Li, T., Ran, Q., and Wang, Y.: Drought vulnerability assessment in Inner Mongolia, *Acta Ecologica Sinica*, 40, 9070-9082, 2020.

- Jiang, L., Yu, S., Wulantuya, and Duwala: Summary of Grassland Fire Research, *Acta Agrestia Sinica*, 26, 791-803, 2018.
- Jiménez-Valverde, A.: Insights into the area under the receiver operating characteristic curve (AUC) as a discrimination measure in species distribution modelling, *Global Ecology and Biogeography*, 21, 498-507, 10.1111/j.1466-8238.2011.00683.x, 2012.
- 500
- Jolly, W. M., Cochrane, M. A., Freeborn, P. H., Holden, Z. A., Brown, T. J., Williamson, G. J., and Bowman, D. M. J. S.: Climate-induced variations in global wildfire danger from 1979 to 2013, *Nature Communications*, 6, 10.1038/ncomms8537, 2015.
- Krawchuk, M. A. and Moritz, M. A.: Constraints on global fire activity vary across a resource gradient, *Ecology*, 92, 121-132, doi:10.1890/09-1843.1, 2011.
- 505
- Krebs, P., Koutsias, N., and Conedera, M.: Modelling the eco-cultural niche of giant chestnut trees: new insights into land use history in southern Switzerland through distribution analysis of a living heritage, *Journal of Historical Geography*, 38, 372-386, 10.1016/j.jhg.2012.01.018, 2012.
- Lamont, B. B. and He, T.: Fire-Proneness as a Prerequisite for the Evolution of Fire-Adapted Traits, *Trends in Plant Science*, 22, 278-288, 10.1016/j.tplants.2016.11.004, 2017.
- 510
- Le Page, Y., Pereira, J. M. C., Trigo, R., da Camara, C., Oom, D., and Mota, B.: Global fire activity patterns (1996-2006) and climatic influence: an analysis using the World Fire Atlas, *Atmospheric Chemistry and Physics*, 8, 1911-1924, 10.5194/acp-8-1911-2008, 2008.
- Li, J. and Feng, C.: Ecosystem service values and ecological improvement based on land use change: a case study of the Inner Mongolia Autonomous Region, *Acta Ecologica Sinica*, 39, 4741-4750, 2019.
- 515
- Liang, H., Wang, W., Guo, F., Lin, F., and Lin, Y.: Comparing the application of logistic and geographically weighted logistic regression models for Fujian forest fire forecasting, *Acta Ecologica Sinica*, 37, 4128-4141, 2017.
- Liu, M., Dries, L., Heijman, W., Huang, J., Zhu, X., Hu, Y., and Chen, H.: The Impact of Ecological Construction Programs on Grassland Conservation in Inner Mongolia, China, *Land Degradation & Development*, 29, 326-336, 10.1002/ldr.2692, 2018.
- 520
- Liu, Z., Yang, J., Chang, Y., Weisberg, P. J., and He, H. S.: Spatial patterns and drivers of fire occurrence and its future trend under climate change in a boreal forest of Northeast China, *Global Change Biology*, 18, 2041-2056, 10.1111/j.1365-2486.2012.02649.x, 2012.
- Ma, W., Feng, Z., Cheng, Z., and Wang, F.: Study on driving factors and distribution pattern of forest fires in Shanxi province, *Journal of Central South University of Forestry & Technology*, 40, 57-69, 2020.
- 525
- Masinda, M. M., Sun, L., Wang, G., and Hu, T.: Moisture content thresholds for ignition and rate of fire spread for various dead fuels in northeast forest ecosystems of China, *Journal of Forestry Research*, 10.1007/s11676-020-01162-2, 2020.
- Masinda, M. M., Li, F., Liu, Q., Sun, L., and Hu, T.: Prediction model of moisture content of dead fine fuel in forest plantations on Maoer Mountain, Northeast China, *Journal of Forestry Research*, 10.1007/s11676-020-01280-x, 2021.

- Matin, M. A., Chitale, V. S., Murthy, M. S. R., Uddin, K., Bajracharya, B., and Pradhan, S.: Understanding forest fire patterns and risk in Nepal using remote sensing, geographic information system and historical fire data, *International Journal of Wildland Fire*, 26, 276-286, <https://doi.org/10.1071/WF16056>, 2017.
- Miranda, B. R., Sturtevant, B. R., Stewart, S. I., and Hammer, R. B.: Spatial and temporal drivers of wildfire occurrence in the context of rural development in northern Wisconsin, USA, *International Journal of Wildland Fire*, 21, 141-154, <http://dx.doi.org/10.1071/WF10133>, 2012.
- Mohajane, M., Costache, R., Karimi, F., Pham, Q. B., Essahlaoui, A., Nguyen, H., Laneve, G., and Oudija, F.: Application of remote sensing and machine learning algorithms for forest fire mapping in a Mediterranean area, *Ecological Indicators*, 129, 17, 10.1016/j.ecolind.2021.107869, 2021.
- Monjarás-Vega, N. A., Briones-Herrera, C. I., Vega-Nieva, D. J., Calleros-Flores, E., Corral-Rivas, J. J., López-Serrano, P. M., Pompa-García, M., Rodríguez-Trejo, D. A., Carrillo-Parra, A., González-Cabán, A., Alvarado-Celestino, E., and Jolly, W. M.: Predicting forest fire kernel density at multiple scales with geographically weighted regression in Mexico, *Science of The Total Environment*, 718, 137313, <https://doi.org/10.1016/j.scitotenv.2020.137313>, 2020.
- Mou, N., Liu, W., Wang, H., Dai, H.: *ArcGIS 10 Tutorial: From Beginner to Master*, Sinomap Press, Beijing, China (in Chinese)2012.
- Muller, M. M., Vila-Villardell, L., and Vacik, H.: Towards an integrated forest fire danger assessment system for the European Alps, *Ecological Informatics*, 60, 10.1016/j.ecoinf.2020.101151, 2020.
- Naderpour, M., Rizeei, H. M., and Ramezani, F.: Forest Fire Risk Prediction: A Spatial Deep Neural Network-Based Framework, *Remote Sensing*, 13, 24, 10.3390/rs13132513, 2021.
- Nunes, A. N., Lourenço, L., and Meira, A. C. C.: Exploring spatial patterns and drivers of forest fires in Portugal (1980–2014), *Science of The Total Environment*, 573, 1190-1202, <http://dx.doi.org/10.1016/j.scitotenv.2016.03.121>, 2016.
- Oliveira, S., Oehler, F., San-Miguel-Ayanz, J., Camia, A., and Pereira, J. M. C.: Modeling spatial patterns of fire occurrence in Mediterranean Europe using Multiple Regression and Random Forest, *Forest Ecology and Management*, 275, 117-129, 10.1016/j.foreco.2012.03.003, 2012.
- Pan, Q., Xue, J., Tao, J., Xu, M., and Zhang, W.: Current status of grassland degradation and measures for grassland restoration in northern China, *Chinese Science Bulletin*, 63, 1642-1650, 2018.
- Parajuli, A., Gautam, A. P., Sharma, S. P., Bhujel, K. B., Sharma, G., Thapa, P. B., Bist, B. S., and Poudel, S.: Forest fire risk mapping using GIS and remote sensing in two major landscapes of Nepal, *Geomatics, Natural Hazards and Risk*, 11, 2569-2586, 10.1080/19475705.2020.1853251, 2020.
- Pavlek, K., Biscevic, F., Furcic, P., Grdan, A., Gugic, V., Malesic, N., Moharic, P., Vragovic, V., Fuerst-Bjelis, B., and Cvitanovic, M.: Spatial patterns and drivers of fire occurrence in a Mediterranean environment: a case study of southern Croatia, *Geogr. Tidsskr.*, 117, 22-35, 10.1080/00167223.2016.1266272, 2017.
- Phelps, N. and Woolford, D. G.: Guidelines for effective evaluation and comparison of wildland fire occurrence prediction models, *International Journal of Wildland Fire*, 30, 10.1071/wf20134, 2021a.

- Phelps, N. and Woolford, D. G.: Comparing calibrated statistical and machine learning methods for wildland fire occurrence prediction: a case study of human-caused fires in Lac La Biche, Alberta, Canada, *International Journal of Wildland Fire*, 10.1071/wf20139, 2021b.
- Podur, J., Martell, D. L., and Csillag, F.: Spatial patterns of lightning-caused forest fires in Ontario, 1976-1998, *Ecological Modelling*, 164, 1-20, 2003.
- Renard, Q., Péliissier, R., Ramesh, B. R., and Kodandapani, N.: Environmental susceptibility model for predicting forest fire occurrence in the Western Ghats of India, *International Journal of Wildland Fire*, 21, 368-379, <http://dx.doi.org/10.1071/WF10109>, 2012.
- Rodrigues, M. and de la Riva, J.: An insight into machine-learning algorithms to model human-caused wildfire occurrence, *Environmental Modelling & Software*, 57, 192-201, <http://dx.doi.org/10.1016/j.envsoft.2014.03.003>, 2014.
- Rodrigues, M., Jiménez-Ruano, A., Peña-Angulo, D., and de la Riva, J.: A comprehensive spatial-temporal analysis of driving factors of human-caused wildfires in Spain using Geographically Weighted Logistic Regression, *Journal of Environmental Management*, 225, 177-192, <https://doi.org/10.1016/j.jenvman.2018.07.098>, 2018.
- Sağlam, B., Bilgili, E., Durmaz, B. D., Kadioğulları, A. İ., and Küçük, Ö.: Spatio-Temporal Analysis of Forest Fire Risk and Danger Using LANDSAT Imagery, *Sensors*, 8, 3970-3987, DOI: 10.3390/s8063970, 2008.
- Scholze, M., Knorr, W., Arnell, N. W., and Prentice, I. C.: A climate-change risk analysis for world ecosystems, *Proceedings of the National Academy of Sciences of the United States of America*, 103, 13116-13120, 10.1073/pnas.0601816103, 2006.
- Shabbir, A. H., Zhang, J., Groninger, J. W., van Etten, E. J. B., Sarkodie, S. A., Lutz, J. A., and Valencia, C.: Seasonal weather and climate prediction over area burned in grasslands of northeast China, *Sci Rep*, 10, 19961, 10.1038/s41598-020-76191-2, 2020.
- Shmuel, A., Ziv, Y., and Heifetz, E.: Machine-Learning-based evaluation of the time-lagged effect of meteorological factors on 10-hour dead fuel moisture content, *Forest Ecology and Management*, 505, 10.1016/j.foreco.2021.119897, 2022.
- Sousa, D., Cruz-Jesus, F., Sousa, A., and Painho, M.: A multivariate approach to assess the structural determinants of large wildfires: evidence from a Mediterranean country, *International Journal of Wildland Fire*, 30, 10.1071/wf20119, 2021.
- Steiner, J. L., Wetter, J., Robertson, S., Teet, S., Wang, J., Wu, X., Zhou, Y., Brown, D., and Xiao, X.: Grassland Wildfires in the Southern Great Plains: Monitoring Ecological Impacts and Recovery, *Remote Sensing*, 12, 10.3390/rs12040619, 2020.
- Su, H., Shen, W., Wang, J., Ali, A., and Li, M.: Machine learning and geostatistical approaches for estimating aboveground biomass in Chinese subtropical forests, *Forest Ecosystems*, 7, 10.1186/s40663-020-00276-7, 2020.
- Su, Z., Zheng, L., Luo, S., Tigabu, M., and Guo, F.: Modeling wildfire drivers in Chinese tropical forest ecosystems using global logistic regression and geographically weighted logistic regression, *Natural Hazards*, 10.1007/s11069-021-04733-6, 2021.
- Su, Z. W., Tigabu, M., Cao, Q. Q., Wang, G. Y., Hu, H. Q., and Guo, F. T.: Comparative analysis of spatial variation in forest fire drivers between boreal and subtropical ecosystems in China, *Forest Ecology and Management*, 454, 10.1016/j.foreco.2019.117669, 2019.

- Syphard, A. D., Sheehan, T., Rustigian-Romsos, H., and Ferschweiler, K.: Mapping future fire probability under climate change: Does vegetation matter?, *PLoS One*, 13, e0201680, 10.1371/journal.pone.0201680, 2018.
- 600 Syphard, A. D., Rustigian-Romsos, H., Mann, M., Conlisk, E., Moritz, M. A., and Ackerly, D.: The relative influence of climate and housing development on current and projected future fire patterns and structure loss across three California landscapes, *Global Environmental Change*, 56, 41-55, 10.1016/j.gloenvcha.2019.03.007, 2019.
- Syphard, A. D., Radeloff, V. C., Keuler, N. S., Taylor, R. S., Hawbaker, T. J., Stewart, S. I., and Clayton, M. K.: Predicting spatial patterns of fire on a southern California landscape, *International Journal of Wildland Fire*, 17, 602-613, 10.1071/wf07087, 2008.
- 605 Thomson, D. M., Bonapart, A. D., King, R. A., Schultz, E. L., Startin, C. R., and Ward, D.: Long-term monitoring of a highly invaded annual grassland community through drought, before and after an unintentional fire, *Journal of Vegetation Science*, 31, 307-318, 10.1111/jvs.12833, 2020.
- Verbesselt, J., Somers, B., van Aardt, J., Jonckheere, I., and Coppin, P.: Monitoring herbaceous biomass and water content with SPOT VEGETATION time-series to improve fire risk assessment in savanna ecosystems, *Remote Sensing of Environment*, 101, 399-414, 10.1016/j.rse.2006.01.005, 2006.
- 610 Vilar, L., Woolford, D. G., Martell, D. L., and Pilar Martin, M.: A model for predicting human-caused wildfire occurrence in the region of Madrid, Spain, *International Journal of Wildland Fire*, 19, 325-337, 10.1071/wf09030, 2010.
- Wei, Y., Rideout, D., and Kirsch, A.: An optimization model for locating fuel treatments across a landscape to reduce expected fire losses, *Canadian Journal of Forest Research*, 38, 868-877, 2008.
- 615 Wheeler, D. C.: Diagnostic tools and a remedial method for collinearity in geographically weighted regression, *Environment and Planning a-Economy and Space*, 39, 2464-2481, 10.1068/a38325, 2007.
- Wotton, B. M., Nock, C. A., and Flannigan, M. D.: Forest fire occurrence and climate change in Canada, *International Journal of Wildland Fire*, 19, 253-271, doi:10.1071/WF09002, 2010.
- 620 Wu, Z., He, H. S., Yang, J., Liu, Z., and Liang, Y.: Relative effects of climatic and local factors on fire occurrence in boreal forest landscapes of northeastern China, *Science of The Total Environment*, 493, 472-480, <http://dx.doi.org/10.1016/j.scitotenv.2014.06.011>, 2014.
- Wu, Z., He, H. S., Keane, R. E., Zhu, Z., Wang, Y., and Shan, Y.: Current and future patterns of forest fire occurrence in China, *International Journal of Wildland Fire*, 29, 104-119, <https://doi.org/10.1071/WF19039>, 2020a.
- 625 Wu, Z. W., He, H. S., Keane, R. E., Zhu, Z. L., Wang, Y. Q., and Shan, Y. L.: Current and future patterns of forest fire occurrence in China, *International Journal of Wildland Fire*, 29, 104-119, 10.1071/wf19039, 2020b.
- XU, X.: China monthly Vegetation index (NDVI) spatial distribution dataset, Data Registration and Publishing System of Resource and Environmental Science Data Center, Chinese Academy of Sciences, 10.12078/2018060602, 2018.
- Xue, Z. C., Kappas, M., and Wyss, D.: Spatio-Temporal Grassland Development in Inner Mongolia after Implementation of the First Comprehensive Nation-Wide Grassland Conservation Program, *Land*, 10, 10.3390/land10010038, 2021.

- 630 Yang, K.: China Meteorological Forcing Data (1979-2018), Big Data System for Pan-Third Pole, 10.11888/AtmosphericPhysics.tpe.249369.file, 2018.
- Yu, T. and Zhuang, Q.: Quantifying global N₂O emissions from natural ecosystem soils using trait-based biogeochemistry models, *Biogeosciences*, 16, 207-222, 10.5194/bg-16-207-2019, 2019.
- Zapata-Rios, X., Lopez-Fabara, C., Navarrete, A., Torres-Paguay, S., and Flores, M.: Spatiotemporal patterns of burned areas, 635 fire drivers, and fire probability across the equatorial Andes, *Journal of Mountain Science*, 18, 952-972, 10.1007/s11629-020-6402-y, 2021.
- Zhang, H.: Spatial analysis of fire-influencing factors in Henan Province, *Progress in Geography*, 33, 958-968, 2014.
- Zhang, H., Han, X., and Dai, S.: Fire Occurrence Probability Mapping of Northeast China With Binary Logistic Regression Model, *Ieee Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 6, 121-127, 640 10.1109/jstars.2012.2236680, 2013.
- Zhao, F., Shu, L., Di, X., Tian, X., and Wang, M.: Changes in the Occurring Date of Forest Fires in the Inner Mongolia Daxing'anling Forest Region Under Global Warming, *Scientia Silvae Sinicae*, 45, 166-172, 2009.
- Zhou, H., Wang, Y., and Zhou, G.: Temporal and spatial dynamics of grassland fires in Inner Mongolia, *Acta Prataculturae Sinica*, 25, 16-25, 2016.

645