

Predictive Understanding of Socioeconomic Flood Impact in Data-Scarce Regions Based on Channel Properties and Storm Characteristics: Application in High Mountain Asia (HMA)

Mariam Khanam¹, Giulia Sofia¹, Wilmalis Rodriguez¹, Efthymios I. Nikolopoulos², Binghao Lu³,
5 Dongjin Song³ and Emmanouil N. Anagnostou¹

¹Civil & Environmental Engineering, University of Connecticut, Storrs, 06269, USA

²Civil & Environmental Engineering, Rutgers University, Piscataway, 08854, USA

³Computer Science and Engineering, University of Connecticut, Storrs, 06269, USA

10 *Correspondence to:* Mariam Khanam (mariam.khanam@uconn.edu), Giulia Sofia (giulia.sofia@uconn.edu)

Abstract. High Mountain Asia (HMA) faces heightened vulnerability to natural disasters due to its extreme conditions and the escalating impacts of climate change. Understanding the long-term response of this landscape to hydroclimatic fluctuations is imperative, given the profound effects these changes have on millions of people annually. Heavy rains, and monsoon seasons, bring forth floods and debris flows, resulting in significant damage to crops, infrastructure, and communities, causing
15 widespread human impacts. Despite efforts to estimate flood risk locally, traditional techniques often fall short due to the scarcity of high-quality, consistent data, especially in ungauged basins. To overcome this challenge, we propose a novel approach: a geomorphologically guided machine learning (ML) method for mapping flood effects across HMA. Central to our methodology is the Lifeyear Index (LYI), a systematic measure that quantifies both the financial and human losses incurred by disasters, specifically for this study fluvial and pluvial flooding. Our model was trained using a dataset comprising over
20 6000 flood events spanning from 1980 to 2020, along with their corresponding five-year and ten-year LYI. Key predictors included: (1) five-year rainfall concentrations derived from ERA5 daily data, (2) a geomorphic classifier based on hydraulic scaling functions derived from high-resolution digital elevation models (DEM), and (3) population density. Results demonstrate the model's effectiveness in identifying flood susceptibility hotspots on a national scale and delineating their evolution from 1980 to 2020. Moreover, the study underscores the severity of hydroclimatic extremes across the entire HMA
25 region. Importantly, the proposed framework is versatile and can be adapted to generate various pluvial and fluvial flood vulnerability and risk maps in ungauged regions.

1 Introduction

High Mountain Asia (HMA) presents complex terrain characterized by dynamic hydrological and geomorphological processes. Over recent years, the region has been significantly affected by climate change, notably witnessing accelerated
30 glacial melts (Shrestha and Aryal, 2011; Byers et al., 2022) and shifts in precipitation patterns and intensity (Haag et al., 2019;

Kirschbaum et al., 2020). These environmental changes, compounded by anthropogenic influences such as landscape alterations, have escalated the region's susceptibility to flooding (Byers et al., 2022; Pervin et al., 2020; Shrestha et al., 2010; G. Zheng et al., 2021), with consequent increasing threats to lives, agriculture, and critical infrastructure (Fischer et al., 2022; Pervin et al., 2020; Rentschler et al., 2022; Sharma et al., 2019; Torti, 2012). The direct impacts caused by the flood are only
35 part of the picture; the enduring socioeconomic repercussions further compound the crisis. These include loss of livelihoods, the urgent need for rehabilitation efforts, and the psychological toll exacted on affected communities.

Flood disasters are generally associated with hydroclimatic extremes. The variability of precipitation patterns over time, space, and intensity is indeed crucial to their occurrence, but changes in catchment characteristics can also alter flood magnitude and frequency. The complex geomorphology and orographic characteristics in the HMA region cause significant
40 spatiotemporal heterogeneity of precipitation patterns and extremes (Haag et al., 2019). Furthermore, the geomorphic structure of basins in HMA can influence the flood characteristics more than land cover does (Marston et al., 1996). Many floods in HMA carry huge amounts of sediment and water that adversely affect downstream areas where most population resides, and can remain in the landscape for years afterward (Kafle et al., 2017; Simonovic et al., 2022).

Changes in river morphology and channel shifting resulting from sediment variability are recognized causes of flood
45 risk (Blench, 1969; Criss & Shock, 2001; Lane et al., 2007; Neuhold et al., 2009; Pinter et al., 2008; Slater et al., 2015; Stover & Montgomery, 2001). Several researchers have highlighted how the morphometric characteristics of watersheds provide useful insights into their hydrologic response to rainfall (Borga et al., 2008) since their morphometric characteristics are a crucial influence on flash flood intensity. In HMA, however, these control mechanisms are difficult to model at a large scale.

Accurate evaluation of the socioeconomic impacts of natural disasters is paramount to mitigate the sufferings of the
50 affected people and rehabilitation (Cavallo & Noy, 2010; Meyer et al., 2013; Noy, 2015, 2016a). To date, available studies (Diehl et al., 2021; Mohanty & Simonovic, 2022; Pangali Sharma et al., 2019; Pervin et al., 2020; Piacentini et al., 2020; Yang & Tsai, 2000) have primarily concentrated on vulnerability mapping and risk analysis, employing case studies and descriptive event-based methodologies at a local level. Scaling up the analysis over the entire HMA region is indeed a difficult task, as it requires collecting data from several countries and multiple sources, and this poses challenges due scarcity of ground
55 observations covering consistent timeframes homogeneously (Barandun et al., 2020; Dollan et al., 2024; Miles et al., 2021). Especially in the context of the impact of floods using socioeconomic data, the analysis involves examining the number of fatalities, injured and people otherwise affected, as well as the financial damage that natural disasters cause, and this information is generally collected at the local scale based on reported events. Significant disasters are documented in global databases like The International Disaster Database (EMDAT, www.emdat.be) or, as an example for HMA and this study, the
60 Nepal Disaster Risk Reduction Portal (<http://drrportal.gov.np/>). However, these databases typically operate at a global or national level resolution, potentially overlooking minor disasters. For example, EMDAT only considers events with at least one of the following criteria: 1) 10 fatalities; 2) 100 affected people; 3) a declaration of state of emergency; 4) a call for international assistance. Additionally, those databases utilized to support insurance may prioritize countries with existing or potential insurance coverage (World Bank, 2012).

65 The integration of geomorphic properties, population data, and rainfall characteristics for assessing socioeconomic flood impact is seldom explored comprehensively on a large scale. For HMA, this is primarily due to the inherent challenges associated with conducting on-site surveys in rugged and often inaccessible terrain. However, leveraging remote sensing data has emerged as a valuable approach for delving deeper into these dynamics and effectively quantifying flood impacts. Modern global datasets, featuring improved resolution and coverage, further enhance the utility of remote sensing in this regard (Diehl et al., 2021; Jongejan & Maaskant, 2015; Mosavi et al., 2018; Bentivoglio et al., 2022; Mazzoleni et al., 2022; Hawker et al., 2018; Kirschbaum et al., 2020; Mohanty and Simonovic, 2022; Pangali Sharma et al., 2019; Sanyal and Lu, 2004; Yang and Tsai, 2000; Zheng et al., 2018).

Furthermore, machine learning (ML) techniques have emerged as increasingly popular tools in advanced prediction systems over the past two decades. They offer more cost-effective solutions with performance that can be aggregated, surpassing the complexity and time demands associated with simulating the complex development of flood processes. Recent research (Bentivoglio et al., 2022; Deroliya et al., 2022; Mosavi et al., 2018) has showcased encouraging advancements by integrating machine learning (ML) techniques with global datasets. This contemporary approach to mapping flood vulnerability notably streamlines the computational processes associated with data-intensive simulations, enhancing flood risk management strategies. However, ML systems rely on existing data for learning. Insufficient or incomplete data coverage can hinder effective learning, leading to suboptimal performance when deployed in real-world scenarios. Therefore, ensuring robust data enrichment, encompassing both quantity and quality, is imperative.

In this study, we introduce a streamlined methodology for preliminary flood vulnerability assessment on a large scale, leveraging available global datasets. Specifically, we introduce a flood-risk assessment model designed to quantify spatially distributed socioeconomic susceptibility in flood-prone regions. We utilize this model to augment disaster understanding by integrating remotely sensed data, including climate variables and high-resolution terrain information. Finally, we apply this model in the High Mountain Asia (HMA) regions to analyze changes in socioeconomic flood impacts spanning from 1980 to 2020.

2 Materials and Methods

2.1 Study Area

90 HMA, otherwise known as the Hindu Kush-Himalayan region, comprises Nepal, Pakistan, Bangladesh, Bhutan, India, Afghanistan, Kazakhstan, Kyrgyzstan, Tajikistan, Uzbekistan, Mongolia, China, and part of many other countries in Asia. HMA is home to some of the world's highest mountain systems, including the Himalayas and the Hindu Kush. This rugged terrain has a highly variable climate ranging from tropical to subpolar, essentially controlled by altitude. Around 1.5 billion people (<https://nsidc.org/data/highmountainasia>) dwelling in the region are at risk of natural disasters (such as heavy rainfall, flooding (pluvial/ fluvial/ flash), earthquakes, avalanches, and landslides) due to the topographic characteristics, changing climate patterns, and high population density. Some of the world's largest rivers and deltas, such as the Indus and the Ganges

are located in this region. In the summertime (June to September), monsoon rains bring a vast amount of water (Kayastha & Kayastha, 2019) to the rivers and valleys in the southern part of HMA (Northern India, Nepal, Bangladesh, and Pakistan). Kirschbaum et al., 2020 have projected that the greatest increase in very high intensities of precipitation (>20 mm/day) will occur during the monsoon season with the enormous amount of rain causing all types of devastating floods (Talchabhadel et al., 2018). Referring to the data reported for example in EMDAT, among all other hydroclimatic disasters in HMA from 1980 to 2020, floods affected the most people (53% among all other hydroclimatic disasters) and caused the highest total damage (56% among all other hydroclimatic disasters). Bangladesh, Nepal, Pakistan, and parts of India were hotspots with the highest casualties (source: EMDAT).

This study considers approximately 6,000 watersheds across HMA as the main target area (Figure 1): the watersheds were selected to be consistent with the HMA domain and all the datasets produced throughout the different phases of the NASA-funded HiMAT project (<https://himat.org/>). The analysis initially centered on training and testing a machine-learning model specifically for Nepal. To achieve this, we collected fine-resolution topographic data along with district-scale socioeconomic information pertaining to population characteristics and documented flood impacts for this region. Subsequently, leveraging the insights gained from this initial phase, we extended the application of the trained model to predict socioeconomic impacts across all watersheds in HMA.




Figure 1: Study area- watersheds across High Mountain Asia (HMA), with highlighted the training domain (Nepal) and the overall rainfall variability across the region. The watershed displayed in black represents the 6000 watersheds that were used in the study. The watershed were selected to be consistent with the HMA domain and all the datasets produced throughout the different phases of the NASA-funded HiMAT project (<https://himat.org/>)

2.2 Methods

Figure 2 illustrates the conceptual framework guiding this study. We employed machine learning (ML) analysis, utilizing climatic and geomorphologic variables, to forecast the socioeconomic impact of extreme fluvial and pluvial flood events spanning from 1980 to 2020 across High Mountain Asia (HMA). To capture the link between flooding and climatic and geomorphologic processes, the model considers as predictors a climatic index derived from ERA5 rainfall, and a geomorphological index, the Flood Geomorphic Potential -FGP- that characterizes the flood-proneness of the landscape, together with population data. A notable advantage of the proposed approach lies in its reliance on automatic techniques leveraging globally available datasets, thereby facilitating its applicability across diverse geographical regions to forecast socioeconomic flood impacts. The framework also benefits from leveraging geomorphologically-driven information, to have an improved characterization of the different aspects of the underlying physical processes shaping the landscape and possibly impacting flood characteristics. By incorporating such domain knowledge into the ML model, the framework can better generalize across different regions and conditions, improving robustness and reliability for risk mapping in diverse environments and facilitating informed decision-making for flood management and mitigation strategies.

To represent exposure and socioeconomic impacts, we introduced, respectively, a variable for population and “Lifeyears Index” (LYI) (Noy, 2014, 2016a, 2016b), a unit of measurement used to describe a disaster’s impact in terms of the total years of life lost (see section 2.3.1 for details). To predict the LYI, we applied XGBoosting (eXtreme Gradient Boosting) (Chen et al., 2018; Chen & Guestrin, 2016). The predictor and response variables of the ML framework are described in the subsections below.




Figure 2: Conceptual framework. Considered predictors are Flood Geomorphic Potential (FGP), Rainfall, and Population. The predicted value is the socioeconomic impact, characterized as the Lifyears Index (LYI) (Noy, 2016a; Noy, 2015). Readers should refer to the following sections for an explanation of the predictors and predicted values.

140 The analysis follows a multistep approach, beginning with data at both watershed and district scales. Initially, the focus was on the district scale, as socioeconomic data for Nepal, selected as the primary training ground, were readily available at this level through the Nepal Disaster Risk Reduction Portal (<http://drrportal.gov.np/>). For this region, furthermore, there is a comprehensive coverage of high-resolution (8-meter) Digital Elevation Models (DEMs) from prior High Mountain Asia (HMA) work (High Mountain Asia 8-meter DEMs Derived from Along-track Optical Imagery, 10.5067/0MCWJJH5ABYO).
 145 Subsequently, all the information is aggregated at the watershed scale, as phenomena such as fluvial and pluvial flooding occur at this level, necessitating a dataset tailored to this scale.

 To transfer the demographic information from the district to the watershed scale, we performed a weighted spatial join between the watersheds and districts. For each watershed, we attributed the statistical characteristics of the intersecting districts, with weights based on the overlapping areas. Generally, the districts in Nepal are smaller in extent compared to the
 150 various watersheds.

2.3 Datasets

2.3.1. Socioeconomic Flood Impacts

 The research focused on predicting the socioeconomic impact of floods. Measured economic loss and tangible
 155 damages were analyzed by considering the Lifyears Index (LYI) (Noy, 2014, 2016a, 2016b). This index is presented by Noy, 2016 as “Lifyears lost” and it is a variation of the WHO Disability Adjusted Life Years (DALYs) lost due to diseases and injuries (WHO, 2014). We calculated LYI for Nepal by using damage statistics and demographic information collected from different data portals in Nepal.

160 The Index is described by Equation 1 and the parameters used in the equation are described in Table 1:

$$LYI = M(A_{exp} - A_{med}) + e * T * N + (1 - c)Y/PCGDP \tag{1}$$

Table 1: Parameters used to calculate LYI

Variable	Description	References
M	Mortality (number of deaths due to disaster	Nepal Disaster Risk Reduction Portal (http://drrportal.gov.np/)
	Average life expectancy at birth (by year)	
A _{exp}		WHO (https://data.who.int/countries/524)
A _{med}	Median age (by year)	WHO (https://data.who.int/countries/524)

e	Welfare reduction weight associated with being exposed to a disaster	set to $e = 0.054$ according to Noy, (2016a), based on Mathers et al., 2013
T	Time taken by the affected person to get back to normal	Noy, (2016a)
N	Number of affected people	Nepal Disaster Risk Reduction Portal (http://drrportal.gov.np/)
c	Percent of time not used in work-related activities (.75)	Noy, (2016a)
Y	Y = Financial damage (value of destroyed/damaged infrastructure)	Nepal Disaster Risk Reduction Portal (http://drrportal.gov.np/)
PCGDP	Income per capita (by year)	The World Bank (https://data.worldbank.org/country/Nepal)

In this study, we classified Lifyears Index (LYI) values into three distinct categories: Low for cases where $\log(\text{LYI}) < 2$; Medium for values falling between 2 and 3; and High for $\log(\text{LYI}) > 3$. This classification scheme indicates that a watershed or district is deemed to be at high risk if the average LYI exceeds 1000 years, while Medium risk spans LYI values ranging from 100 to 1000 years, and Low risk encompasses LYI values less than 100 years. For instance, if the calculated LYI is 100 years, it implies that the estimated impact of the given disaster equates to a potential loss of 100 years of life per 100,000 people.

The cumulative LYI for Nepal (Figure 3) can provide an idea of how the cumulated flood impact has been increasing in a country with time. It also highlights how the index itself captures major disasters, such as those occurring in 1981 (ICIMOD, 2011; Kiran S et al., 2008), 1993 (Nepal - Floods and Landslides, 1993), in 1996 (Nepal - Floods Situation Report No. 1, 26 July 1996), and in the monsoon seasons in 2003 and 2014 (Nepal Annual Report, 2003.; Nepal: Landslides and Floods - Aug 2014). The most changes can be noticed in the LYI for the years 1981, 1993, and 2014, the cumulative step change for these years from the previous year are subsequently 9999, 82865, and 976238 years.




Figure 3: Cumulative lifeyears lost over the years in Nepal. Highlighted years represent jumps in the cumulative value, mostly related to well-known disasters): 1981 (ICIMOD, 2011; Kiran S et al., 2008), 1993 (Nepal - Floods and Landslides, 1993), 1996 (Nepal - Floods Situation Report No. 1, 26 July 1996), and in the monsoon seasons in 2003 and 2014 (Nepal Annual Report, 2003.; Nepal: Landslides and Floods - Aug 2014).

2.3.2. Floodplain Mapping

The identification of areas with the potential to be inundated is fundamental to preserving and protecting human lives and property while safely supporting economic activities. Hence, we applied a large-scale floodplain delineation algorithm to identify such areas at the basin scale across the HMA. Many researchers (e.g., Dingle et al., 2020; Lindersson et al., 2021; Piacentini et al., 2020) have used DEM-derived geomorphic index as a high-resolution flood mapping tool. We opted for considering a variation of the Samela et al., (2017) Geomorphic Flood Index (GFI), thereby described as Flood Geomorphic Potential (FGP).

The index is calculated as the logarithm of the ratio between the bankfull elevations (estimated using a hydraulic scaling function, or HSF, based on contributing area) in the element of the river network closest to the point under examination and the elevation difference between these two points (Figure 4). The index was improved over a main aspect: the automatic identification of the HSF directly from terrain data, applying the technique of (Sofia, et al., 2017b; Sofia et al., 2015) to retrieve the bankfull location automatically through the landscape. This has the advantage of allowing for full automation of the mapping starting purely from terrain data.

For this analysis, we trained the model considering FGP derived from the unique 8-meter Digital Elevation Models (DEMs) for Nepal that are available at the NASA National Snow and Ice Data Center Distributed Active Archive Center (NSIDC DAAC) (Shean, 2017c, 2017b, 2017a). While Nepal is entirely covered by the 8m DEM, extending the model to the whole HMA region is complicated by the gaps in the input satellite strip resulting from limited coverage, clouds, or failed stereo correlation. For this reason, we also considered the 30m DEM by Copernicus (European Space Agency, Sinergise. Copernicus Global Digital Elevation Model, 2021), a digital surface model (DSM) that represents the surface of the Earth, including buildings, infrastructure, and vegetation. Importantly, this DSM is derived from World DEM, an edited DSM in which the flattening of water bodies and the consistent flow of rivers have been included. Shore- and coastlines, special features such as airports, and implausible terrain structures have also been edited.

We identified flood-prone areas by grouping them into six classes by their FGP index. For each watershed, we then considered the areas covered by the classes with FGP greater than 4, which, when compared to published data, proved to correspond realistically with areas subject to floods of about 100-year depth. Figure 4b compares the Flood Geomorphic Potential (FGP) automatic classes derived for select rivers in Nepal, with baseline inundation scenarios evaluated using standard inundation depths associated with critical flood events and their return periods provided in the work of Delalay et al. (2018). This visual comparison serves to highlight the efficacy of flood inundation mapping facilitated by the FGP.

It's worth noting that the FGP methodology has been previously published and applied in various contexts (Samela et al., 2017). While testing the quality of the FGP lies beyond the scope of this work, its effectiveness for flood mapping has been well-established in previous studies (Manfreda et al., 2011, 2014; Manfreda & Samela, 2019; Samela et al., 2016, 2018), which have demonstrated the utility of the methodology, particularly in ungauged conditions, for preliminary identification of

215 flooded areas in regions where conducting expensive and time-consuming hydrologic-hydraulic simulations may not be feasible.




Figure 4: a. Flood Geomorphic Potential (FGP) (modified from Samela et al. 2017); b. FGP automatic classes compared to baseline inundation depth scenarios (aerial imagery © Google Earth 2015).

220 2.3.3. Rainfall Characteristic

The climatology in HMA is highly variable (Dollan et al. 2024). Summer monsoons drive precipitation in the Ganges-Brahmaputra basins and the Tibetan Plateau (Bookhagen and Burbank, 2010; Shamsudduha and Panda, 2019); synoptic storms dominate winter precipitation impacting areas in the northwestern Karakorum mountains (Winiger et al., 2005; Barlow et al., 2005). Overall, as well, variations in elevation gradients contribute to diverse microclimates, exemplified by Nepal's swift
225 transition from high mountains to lowlands (Kansakar et al., 2004; Karki et al., 2016). Winter precipitation in the area is

primarily influenced by the westerly weather system, with western disturbances originating in the Mid-Atlantic or Mediterranean Sea and traversing through northwest India to western Nepal after passing over Afghanistan and Pakistan (Kansakar et al., 2004; Hamal et al., 2020). In Nepal, which was used as the training site for the model, regional climate variations exist, mostly driven by changes in elevation, with an overall homogeneity in trends (aside from a few hotspots) and regional statistics of precipitation, in line with the variability of HMA, as highlighted by the recent study by (Khanal et al., 2023).

For this work, for the main rainfall driver of the model, we focused on daily climate concentration. As climate concentration values are mostly related to the temporal variability of the rainfall, not to the total amount or the average yearly and seasonal statistics, using this index allows to capture well various climates globally (Monjo and Martin-Vide, 2016a). The variability of climate concentration, furthermore, has been proven to be highly linked to pluvial/fluvial flooding impacts in various regions of the world, including for example Italy (both in mountainous landscapes and floodplains (Sofia et al., 2019), the US (Saki et al., 2023) [over a variety of physiographic regions], or China (Du et al., 2023). Different authors have adopted different methods to determine the temporal concentration of precipitation, and the Concentration Index (CI) (Equation 2) is one of the most used parameters (Caloiero et al., 2019; Martin-Vide, 2004; Monjo, 2016; Sangüesa et al., 2018; Serrano-Notivoli et al., 2018).

This index was proposed by Martin-Vide (2004) originally to explore the contribution of the days with major rainfall to the total amount within a certain time range. The benefit of this index is that it can describe the temporal variability of rainfall at daily, annual, and seasonal scales using a single metric, as well as spatial variability at pixel or watershed scale. In the present study, we computed CI (Martin-Vide, 2004) using the ERA5 hourly rainfall data from 1980 to 2019. The source of rainfall data was selected as various works for HMA highlighted its effectiveness in capturing extreme events quite accurately compared to other products (Maggioni & Massari, 2018; Maina et al., 2023; Dollan et al. 2024).

We identified storm events from this dataset primarily based on the criterion of rainfall of more than 0.5 mm, and we separated events when rainfall was below this threshold for more than 12 hours. Furthermore, we calculated CI using the cumulative amount of rainfall (y) and the cumulative frequency of the event duration (x) (Figure 5) for the selected events. The method (similar to Cortesi et al., 2012 and Monjo & Martin-Vide, 2016a) eventually aggregates the amount of precipitation that falls during each event into increasing categories and determines the relative contribution (as a percentage) of the progressively accumulated precipitation, y, as a function of the accumulated percentage of the durations of the events (x). The concentration index is then calculated as the ratio of the area between the line of equality (y=x) and the fitted curve (S), and the total area under the line of equality (A+S) (Figure 5, equation 2). The index is defined by the relationship between the accumulated percentage of time, and the accumulated rainfall.

$$CI = \frac{S}{S+A} \quad (2)$$




Figure 5: Example of line of equality, and empirical curve for the rainfall concentration calculation. The concentration index is equal to the area between the line of equality and the fitted curve (S) divided by the total area below the line of equality (S+A)

2.3.4. Exposure (Population)

As all the parameters of the LYI are not always readily available at the watershed scale (as highlighted by most published literature, that considered LYI at the country scale), we added population counts as one of the predictors to train the model. For Nepal, we selected the data from the country's national census (<https://censusnepal.cbs.gov.np/Home/Index/EN>) and aggregated it at the watershed scale by using the previously mentioned weighted join. To extend the model to the whole HMA, we computed the population for each watershed across the region from the Gridded Population of the World (GPW), v4 | SEDAC, 2024). This dataset provides spatially explicit estimates of population density for the years 2000, 2005, 2010, 2015, and 2020, based on counts consistent with national censuses and population registers, as raster data to facilitate data integration. We used a simple linear regression to retrieve data for the missing years.

2.4. Machine Learning Model

XGBoosting is primarily used to solve classification problems. To generate the results, the XGBoost algorithm uses an ensemble of boosted trees. An ensemble is a collection of predictors that together can give a final prediction while reducing errors significantly. In this case, the predictors were climatic variables, geomorphologic variables, and exposure. Boosted algorithms are those in which each successive model attempts to correct the errors of its predecessor (similar to adaptive

learning). The basic XGBoost algorithm can be understood as an ensemble of boosted trees. The idea behind such an ensemble is that multiple trees are built in sequence, each tree built on the previous one's prediction. And each successive tree built considers the errors of the previous trees. This means that when we take an average of all the trees at the end, we get a final tree that is better than any individual tree within the model. We applied the XGBoosting model to the geomorphologic, climatic, and exposure variables to predict classes of LYI in different basins in Nepal and HMA.

2.4.1 Validation of the System at the HMA Scale

We conducted thorough testing and validation of our model within Nepal, comparing the predicted value of LYI to the calculated Lifeyears Index (LYI) data from tabular values specific to the region. Upon extending the model's applicability to the entire High Mountain Asia (HMA) region, we rigorously assessed the quality of our results by comparing the predicted social impact with that reported in established flood databases covering the region. To verify our findings, we compared the predictions at the HMA level with flood events reported in the Dartmouth Flood Observatory's (DFO) Global Active Archive of Large Flood Events, 1985–Present. This comprehensive database compiles information on major floods sourced from diverse channels such as news reports, governmental records, ground observations, and remote sensing data. Notably, the DFO dataset encompasses various flood types, including lowland floods and mountainous river floods characterized as fluvial and pluvial floods.

The dataset provides point locations, representing the centroids of affected areas during floods. While acknowledging that flood centroids may oversimplify the complexities driving flood events, we utilized this dataset to showcase our model's capability to target high-risk locations historically impacted by floods within the specified timeframe. Identifying high-risk areas with recorded flood occurrences centered around these locations underscores the robustness of the model beyond the confines of its training and validation site in Nepal.

Meteorological and climatological severity reported in the DFO cannot directly capture the social impact of the floods; furthermore, they refer to events that may span multiple watersheds. To overcome these limitations, we compared our predicted results to the DFO data by evaluating a proxy of the social severity reported for each flood—the numbers of “Deaths” and “Displaced” that resulted—rather than the flood's meteorological characteristics. We then considered classes of DFO “social severity” of 10^n . Here n is the severity level declared in the DFO database. Next, we calculated the marginal probability that events with different severity in the DFO happened in watersheds with different predicted LYIs. Finally, we calculated the conditional probability, as in the probability of an event of DFO severity of some kind occurring over watersheds where our LYI prediction was of a certain type. This conditional probability could provide us with information on how our system performed for different time frames—for example, the probability of a watershed's being classified as high impact by our model might be only 10% of the total, but if most of DFO events with great severity (i.e., >1000 Deaths+displaced) did happen in those watersheds, then our system correctly identified the risk there.

3.1. Variability of the Predictors

The topographical characteristics of an area can influence the local climate and population distribution. Figure 6 shows an example of how climate concentration and population vary in HMA, as compared to watersheds that have areas of high FGP of greater or lesser extent. The figure reports the average for the time frame 1980–2020 for CI and population, while the FGP is a static value for the time frame (since it is based on a unique DEM dataset), and it represents the overall geomorphic characteristics of Nepal.

From this analysis, we can see how the variability of CI is complex. If expectedly, the variability of the index is related to atmospheric characteristics (Sangüesa et al., 2018), the index varies also due to geographical factors influencing climate (Tuladhar et al., 2020). In their study based on Nepal, Karki et al., 2017 highlighted the difference in the spatial pattern of high-intensity storm events from that of annual and monsoon events. The rapid rate at which physical processes (e.g., convection) take place regulates the high temporal concentration of precipitation in the regions where the sea surface and ground are highly affected by warmer temperatures (Monjo & Martin-Vide, 2016b). On the other hand, the low temporal concentration of rainfall is characterized as a normal pattern caused by cyclical weather events (Monjo & Martin-Vide, 2016). Watersheds with lesser floodplain extents (that is, less areas with high FGP) are related to higher and steeper mountains, with complex orography. Research has shown that low areas in Nepal are susceptible to receiving high-intensity storm events even though they have fewer wet days (Karki et al., 2017). The authors of the same study also observed that the low-intensity events (annual and monsoonal precipitation) were mostly predominant over Nepal’s western middle mountains and central high mountains. In another study, however, Subba et al., 2019 stated that the frequency of extreme events had decreased significantly over the past two decades in the eastern part of Nepal. For our case, areas having the larger physical potential to flood (high FGP), appear to be areas showing the largest variation in CI, with values ranging from low (0.2) as well as very high (0.75), indicating a potential compound effect of highly torrential rains (CI=0.7) in locations where much of the landscape is potentially floodable (FGP high) and most population reside. Readers should consider that higher FGP values do not imply locations having wider channels, but rather they indicate how the landscape is potentially more flood-prone than as highlighted by (Samela et al., 2017; Manfreda & Samela, 2019; Samela et al., 2016, 2018).

Much of the population of Nepal tends to be concentrated in areas with higher FGP, as is typical for mountainous areas, where population and economic activities are mostly located in the river valleys. Globally, the floodplains of rivers are preferred living spaces for the population and provide favorable locations for economic development. These areas are commonly exposed to floods, however, an increasing population, together with the changes in storminess, mean that the risks from flooding are expected to be higher.




Figure 6: Average variability of the CI (top) and population (bottom) compared to FGP from 1980-2020


3.2. Variable Importance and Model Performance

Based on the methodology described in section 2.4, in this section, we present a variable importance comparison (Figure 7) based on the F score. The initial variable importance indicated that population (Pop) was the most important variable, which was consistent with our expectation in the sense that the socioeconomic impact depends largely on the exposure. The climate variable (CI) happened to be the next important variable, showing the significance of the region's climate on the socioeconomic impact of flood occurrences.

The precision, recall, and F1 score are metrics used to evaluate the performance of a classification model. Precision is the fraction of true positives among the predicted positives. Recall is the fraction of true positives among the actual positives. The F1 score is the harmonic mean of precision and recall.

The evaluation metrics reveal in Table 2 that the model performs best in the High class, with the highest precision, recall, and F1 score. The Medium class also demonstrates relatively high performance across these metrics. However, the Low class exhibits the lowest performance, suggesting that the model may face challenges in accurately distinguishing between the Low and Medium classes or may demonstrate a bias toward predicting the Medium and High classes. These findings provide valuable insights into the strengths and limitations of the classification model and can guide future efforts to improve its

performance. Overall, considering that the model aims to target substantial risk areas, a higher rate of predicting impacts is acceptable, compared to an underestimation of the risk.



355 **Figure 7: Feature Importance**

Table 2: Performance metrics of a classification model

	precision	recall	f1-score
Low	0.54	0.57	0.56
Medium	0.64	0.63	0.64
High	0.73	0.71	0.72

3.3. Predicted versus Observed Flood Impact in Nepal

360 Comparing predicted Lifeyears Index (LYI) flood impacts with observed data showed good correspondence between high-risk areas identified by the ML method and historical flood locations in Nepal. This suggests that the proposed approach effectively delineates flood risk on a national scale. Figure 8 illustrates this comparison, showcasing observed (empirically evaluated) and ML-predicted LYI values at both watershed (upper row) and district (lower row) levels.

365 The 'observed' LYI values were empirically calculated from observational data (Table 1) and categorized into three groups: 'low', 'medium', or 'high', with basins/districts labeled as 'high' for LYI values exceeding 1000 years, 'medium' between 100 and 1000 years, and 'low' below 10 years. The 'predicted' values represent the outputs from the machine learning model.

In Nepal, we achieved an overall training accuracy of 97% and a test accuracy of 63%. Notably, training the model at the watershed level yielded higher accuracy compared to the district level. This is attributed to watersheds being hydrologic units that integrate geomorphological and climatic properties, thus providing a more accurate representation of flood dynamics compared to administrative district boundaries.

370

At the watershed level, nearly all year ranges exhibited a 100% match with observed impacts. In instances where the model's accuracy fell below 100% (e.g., 1985–90 and 1990–95), the LYI values in the affected watersheds were low, indicating that the predictors considered were more indicative of major flooding events.

375 The superior accuracy achieved at the watershed level underscores the value of implementing the model at this scale when scaling up the system.




Figure 8: Comparison of prediction with actual socioeconomic impact for watersheds and districts in Nepal. Basin/districts are marked as “high” for LYI over 1000 years. Medium is between 100 and 1000, and low is less than 10. Numbers in parentheses represent accuracy.

380

3.4. Prediction of Socioeconomic Impact of Heavy Rainfall over HMA

We applied the trained model for the watersheds in HMA to five-year intervals from 1980 to 2020. As an example, Figure 9(c, d) shows the predicted basin-averaged LYIs (Low-Med-High) for the watersheds in HMA for two different timelines. The yellow circles highlight the changes in flood impact over the decades. One must consider that most HMA has low population density (blue color in Fig. 9b), and as expected the proposed model predicts low flood socioeconomic impacts for these regions. Hotspots of high impacts (Red colors in Figures 9c and d) are present, where population exposure is higher.

385




Figure 9: a) Mean Monthly Rainfall (yellow contours) overlaid on Elevation; b) Population density 2020; c,d) Example of predicted basin-averaged flood impact for HMA (left, 1985–90; right, 2015–20). Yellow circles denote the changes in flood impact between the two timelines.

Summarizing the results presented in Table 3, we can say that, for the years shown, we predicted almost 57% of watersheds (marginal) having LYIs between 1 and 100 years (Low), 35.9% for LYIs between 100 and 1000 years (Med), and only 6% for LYIs greater than 1000 years (High). For the entire period, most of the time we predicted LYIs of 1 to 100 years, for which we captured events of DFO severity around 2 (10^2 Deaths+displaced) (conditional = 28.6%). This suggests that most “Low” class DFO events did happen in the watersheds within the lowest predicted LYI range. Readers must consider that “Low” in this case means the flood impact can range from 1 to 100 years lost, and a DFO value of 2 means total deaths and displaced is on the order of 10^2 people. The events with a DFO value of 4 happened mostly in watersheds with predicted LYIs ranging both between 1 and 100 years and between 100 and 1000 years. The events with DFO 6 and 8 happened mostly in ranges greater than 1000 years and between 100 and 1000 years.

Table 3: LYI compared to DFO flood damage.

DFO	LYI	NO	Prop	Marginal Probability	Conditional Probability
-----	-----	----	------	----------------------	-------------------------

2	1–100yr	54	16.6	58.2	28.6
2	100–1000yr	26	8.0	35.7	22.4
2	>1000year	5	1.5	6.2	25.0
4	1–100yr	92	28.3	58.2	48.7
4	100–1000yr	45	13.8	35.7	38.8
4	>1000year	5	1.5	6.2	25.0
6	1–100yr	42	12.9	58.2	22.2
6	100–1000yr	44	13.5	35.7	37.9
6	>1000year	8	2.5	6.2	40.0
8	1–100yr	1	0.3	58.2	0.5
8	100–1000yr	1	0.3	35.7	0.9
8	>1000year	2	0.6	6.2	10.0

We further investigated how our predicted LYI behaved when it was related to the total population (Table 4), evaluating, as suggested by (Noy, 2014), the LYI per capita (that is, the number of lifeyears lost per 100k people). As Table 4 shows, we correctly predicted over the years almost 64% of watersheds (marginal) have LYI/100k people less than 1 year (10^0), 24.3% at 10year/100k people (10^1), 11% at 100year/100k people, and 0.6% at 1000years/100k people. We noticed that LYI/100k people reached, at most, 6000 for Nepal (at the country scale) and the study by Noy. 2016a also reported similar values for Nepal in 1987. (Noy, 2016a) reported actual LYI data in the range of LYI > 1000/100k people in South Asia and stated that the higher number of damages in East and South Asia is likely due to wide-scale flooding. This gave assurance of the consistency of our prediction with the actual data available. When looking at LYI/100k people, we found that, for the whole timeframe, most of the floods that registered in the DFO with low severity (DFO = 10^2 Deaths+displaced) happened in watersheds for which the predicted LYIs were between 1 and 100 years (conditional = 29.8%). This confirmed once again that, in most cases, “low”-risk events did happen in the watersheds having the lowest predicted range (similar to the findings presented in Table 3). As before, while the probability of a watershed’s being labeled as high risk (LYI>1000year/1000k people) by our system was only 6%, the probability of these watersheds having experienced events recorded by the DFO as having a great impact (DFO severity > 6, meaning over 1 million people) rose to 40% and 10%.

Table 4: LYI/100k compared to DFO flood damage.

DFO	LYI	NO	Prop	Marginal Probability	Conditional Probability
0	0	13	3.8	65.0	5.9
0	1	1	0.3	23.5	1.3
0	2	1	0.3	10.9	2.7

2	0	62	18.2	65.0	28.1
2	1	13	3.8	23.5	16.3
2	2	9	2.6	10.9	24.3
2	3	1	0.3	0.6	50.0
4	0	97	28.5	65.0	43.9
4	1	34	10.0	23.5	42.5
4	2	10	2.9	10.9	27.0
4	3	1	0.3	0.6	50.0
6	0	47	13.8	65.0	21.3
6	1	32	9.4	23.5	40.0
6	2	15	4.4	10.9	40.5
8	0	2	0.6	65.0	0.9
8	2	2	0.6	10.9	5.4

Figure 10 shows the LYI per 100k people (LYI/100k) evaluated for different time frames for all the locations reported in the DFO database to compare the DFO severity with our predictions. Overall, the DFO and predicted results were quite consistent instead of some minor variability for some scattered areas. When we compared the changes over time, we noticed an increase in vulnerability. As the plot makes evident, the largest changes took place in 1990–95 and 2010–15; the two concentrated areas were Nepal and China. As Figure 3 shows, two big jumps occurred during these timelines for Nepal because of extreme storm-induced flood events. In Figure 3 we have discussed the predominant events that occurred in these timelines. Regarding China, as of June 2010, more than 29 million people had been affected by flooding, with up to 2.37 million evacuated and 195,000 homes destroyed (China: Floods Information Bulletin N° 1 GLIDE N°, 2010).





Figure 10: Comparison of DFO and LYI/ 100K people for all the timelines

3.6. Change in Socioeconomic Impact over Time

Figure 11a presents our maps of the watersheds where flood impacts increased over time. Furthermore, Figure 11b shows our evaluation of the percentage changes in the number of watersheds between timelines, focusing on three different changes: low to medium (LtoM); medium to high (MtoH); and low to high (LtoH). Some watersheds have not changed, and some have decreased impact. For the sake of highlighting potential increases in flood impacts, we focused on those locations where risk increased over time, from low to medium, or medium to high. The largest changes were from LtoM for all the timelines, which represented a notable change in vulnerability. Several watersheds showed higher flood impacts (from low to medium, medium to high, and low to high) in recent years as compared to 1985–90. Again, we observed the largest changes for 1990–95 and 2010–15, which was consistent with Figure 10. The exposure changed significantly, along with the intensity of the events; hence, the risk of flooding was heightened in these areas.

Impact changes from Low to High were next, according to the number of watersheds changed for all the timelines. It was obvious that more changes would happen overall, but the comparison of the 1990–95 and 1995–2000 timelines demonstrated that heightened flood impact occurred in a considerable number of watersheds within a brief period. For many watersheds, the risk was heightened by a population boom during the overall period.

a



b




Figure 11: Flood impact change in HMA over time

4. Conclusions

445 High Mountain Asia (HMA) presents a multifaceted landscape characterized by rugged terrain, diverse climates, rich vegetation, and substantial population exposure to natural disasters. Given its susceptibility to natural disasters, effective management is imperative for the region's long-term sustainability. Addressing the considerable threat posed by flooding demands a comprehensive strategy involving disaster risk reduction, sustainable land use practices, and climate change mitigation.

450 In this study, we introduced a simplified approach to identify vulnerability hotspots within the HMA region, focusing on intense rainfall events. To map the socioeconomic flood vulnerability, we employed a remotely sensed data-driven model integrating geomorphological and climate variability factors. This adaptable framework can be tailored to various regions, provided that similar terrain and climate datasets are available, accommodating adjustments to flood drivers such as climate and geomorphology, as well as population dynamics. The resulting predictions offer valuable insights into vulnerabilities
455 across HMA watersheds, facilitating proactive flood management planning.

The novelty of our study lies in the efficiency and versatility of the proposed predictive model. Requiring only a small number of variables, our model accurately forecasts the socioeconomic impact of pluvial and fluvial flooding events. In densely populated, possibly ungaged regions with rapidly changing climates, such a model serves as a valuable decision-making tool for stakeholders. The efficacy of the framework, as demonstrated in Nepal, underscores its potential applicability across
460 regions with similar climatic and morphological characteristics.

With advancing technology, we can now predict the drivers of impending extreme events, enabling proactive measures to mitigate their impact. Stakeholders could leverage our model to forecast vulnerability to future flood events with precision, enhancing hazard assessment, decision-making, planning, and mitigation efforts.

However, while this study demonstrates the promise of accurate flood impact prediction, the use of static Flood
465 Geomorphic Potential (FGP) maps presents limitations. Flooding alters channel morphology and downstream topography, impacting future flood dynamics. Therefore, dynamic flood topographies are essential for robust hazard assessment. Although high-resolution topographical data post-extreme events can enhance prediction accuracy, the availability of such data is constrained by acquisition frequency. Hence, efforts to improve data availability post-disaster are crucial for enhancing the reliability of predictive models.

470

Competing interests: The contact author has declared that none of the authors has any competing interests.

Acknowledgment: This work was supported by the NASA High Mountain Asia program (grant #80NSSC20K1300).

Data Availability: The FGP products are available at NASA National Snow and Ice Data Center (NSIDC) (https://nsidc.org/data/hma2_fgp/versions/1)

- Anon: China: Floods Information bulletin n° 1 GLIDE n°, 2010.
- Barlow, M., Wheeler, M., Lyon, B., and Cullen, H.: Modulation of daily precipitation over Southwest Asia by the Madden-Julian oscillation, *Mon Weather Rev*, 133, 3579–3594, <https://doi.org/10.1175/MWR3026.1>, 2005.
- Bentivoglio, R., Isufi, E., Jonkman, S. N., and Taormina, R.: Deep learning methods for flood mapping: a review of existing
 480 applications and future research directions, *Hydrol Earth Syst Sci*, 26, 4345–4378, [https://doi.org/10.5194/HESS-26-4345-](https://doi.org/10.5194/HESS-26-4345-2022)
 2022, 2022.
- Blench 1906-1993, T.: *Mobile-bed fluviology : a regime theory treatment of rivers for engineers and hydrologists*, Edmonton (Ca.) : University of Alberta press, 1969.
- Bookhagen, B. and Burbank, D. W.: Toward a complete Himalayan hydrological budget: Spatiotemporal distribution of
 485 snowmelt and rainfall and their impact on river discharge, *J Geophys Res Earth Surf*, 115,
<https://doi.org/10.1029/2009JF001426>, 2010.
- Borga, M., Gaume, E., Creutin, J. D., and Marchi, L.: Surveying flash floods: Gauging the ungauged extremes, *Hydrol Process*, 22, 3883–3885, <https://doi.org/10.1002/HYP.7111>, 2008.
- Byers, A. C., Shugar, D. H., Chand, M. B., Portocarrero, C., Shrestha, M., Rounce, D. R., and Watanabe, T.: Three Recent and
 490 Lesser-Known Glacier-Related Flood Mechanisms in High Mountain Environments, *Mt Res Dev*, 42, A12–A22,
<https://doi.org/10.1659/MRD-JOURNAL-D-21-00045.1>, 2022.
- Caloiero, T., Coscarelli, R., and Gaudio, R.: Spatial and temporal variability of daily precipitation concentration in the Sardinia region (Italy), *International Journal of Climatology*, 39, 5006–5021, <https://doi.org/10.1002/JOC.6123>, 2019.
- Cavallo, E. and Noy, I.: *The Economics of Natural Disasters A Survey*, 2010.
- 495 Chen, T. and Guestrin, C.: XGBoost: A Scalable Tree Boosting System, *Proceedings of the ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, 13-17-August-2016, 785–794,
<https://doi.org/10.1145/2939672.2939785>, 2016.
- Chen, T., He, T., and Benesty, M.: XGBoost : eXtreme Gradient Boosting, R package version 0.71-2, 1–4, 2018.
- Criss, R. E. and Shock, E. L.: Flood enhancement through flood control, *Geology*, 29, 875, [https://doi.org/10.1130/0091-](https://doi.org/10.1130/0091-7613(2001)029<0875:FETFC>2.0.CO;2)
 500 7613(2001)029<0875:FETFC>2.0.CO;2, 2001.
- Deroliya, P., Ghosh, M., Mohanty, M. P., Ghosh, S., Rao, K. H. V. D., and Karmakar, S.: A novel flood risk mapping approach with machine learning considering geomorphic and socio-economic vulnerability dimensions, *Science of the Total Environment*, 851, <https://doi.org/10.1016/j.scitotenv.2022.158002>, 2022.
- Diehl, R. M., Gourevitch, J. D., Drago, S., and Wemple, B. C.: Improving flood hazard datasets using a low-complexity, probabilistic floodplain mapping approach, *PLoS One*, 16, <https://doi.org/10.1371/journal.pone.0248683>, 2021.
- 505 Dingle, E. H., Creed, M. J., Sinclair, H. D., Gautam, D., Gourmelen, N., Borthwick, A. G. L., and Attal, M.: Dynamic flood topographies in the Terai region of Nepal, *Earth Surf Process Landf*, 45, 3092–3102, <https://doi.org/10.1002/esp.4953>, 2020a.

- Dingle, E. H., Creed, M. J., Sinclair, H. D., Gautam, D., Gourmelen, N., Borthwick, A. G. L., and Attal, M.: Dynamic flood topographies in the Terai region of Nepal, *Earth Surf Process Landf*, 45, 3092–3102, <https://doi.org/10.1002/esp.4953>, 2020b.
- 510 European Space Agency, Sinergise. Copernicus Global Digital Elevation Model: <https://doi.org/10.5069/G9028PQB>, last access: 20 March 2023.
- Fischer, M., Brettin, J., Roessner, S., Walz, A., Fort, M., and Korup, O.: Rare flood scenarios for a rapidly growing high-mountain city: Pokhara, Nepal, *Natural Hazards and Earth System Sciences*, 22, 3105–3123, <https://doi.org/10.5194/NHESS-22-3105-2022>, 2022.
- 515 Global Active Archive of Large Flood Events, Dartmouth Flood Observatory, University of Colorado, USA.: <https://floodobservatory.colorado.edu/Archives/index.html>, last access: 20 March 2023.
- Gridded Population of the World (GPW), v4 | SEDAC: <https://sedac.ciesin.columbia.edu/data/collection/gpw-v4/citations>, last access: 30 April 2024.
- Haag, I., Jones, P. D., and Samimi, C.: Central Asia’s Changing Climate: How Temperature and Precipitation Have Changed across Time, Space, and Altitude, *Climate 2019*, Vol. 7, Page 123, 7, 123, <https://doi.org/10.3390/CLI7100123>, 2019.
- 520 Hamal, K., Sharma, S., Baniya, B., Khadka, N., and Zhou, X.: Inter-Annual Variability of Winter Precipitation Over Nepal Coupled With Ocean-Atmospheric Patterns During 1987–2015, *Front Earth Sci (Lausanne)*, 8, <https://doi.org/10.3389/FEART.2020.00161>, 2020.
- Hawker, L., Rougier, J., Neal, J., Bates, P., Archer, L., and Yamazaki, D.: Implications of Simulating Global Digital Elevation Models for Flood Inundation Studies, *Water Resour Res*, 54, 7910–7928, <https://doi.org/10.1029/2018WR023279>, 2018.
- 525 High Mountain Asia 8-meter DEM Mosaics Derived from Optical Imagery, Version 1 | National Snow and Ice Data Center: https://nsidc.org/data/hma_dem8m_mos/versions/1, last access: 20 March 2023.
- High Mountain Asia 8-meter DEMs Derived from Along-track Optical Imagery, Version 1 | National Snow and Ice Data Center: https://nsidc.org/data/hma_dem8m_at/versions/1, last access: 20 March 2023.
- 530 High Mountain Asia 8-meter DEMs Derived from Cross-track Optical Imagery, Version 1 | National Snow and Ice Data Center: https://nsidc.org/data/hma_dem8m_ct/versions/1, last access: 20 March 2023.
- ICIMOD: Glacial Lakes and Glacial Lake Outburst Floods in Nepal, 2011.
- Jongejan, R. B. and Maaskant, B.: Quantifying flood risks in the Netherlands, *Risk Analysis*, 35, 252–264, <https://doi.org/10.1111/risa.12285>, 2015.
- 535 Kafle, K. R., Khanal, S. N., and Dahal, R. K.: Consequences of Koshi flood 2008 in terms of sedimentation characteristics and agricultural practices, *Geoenvironmental Disasters*, 4, 1–13, <https://doi.org/10.1186/S40677-017-0069-X/FIGURES/15>, 2017.
- Kansakar, S. R., Hannah, D. M., Gerrard, J., and Rees, G.: Spatial pattern in the precipitation regime in Nepal, *International Journal of Climatology*, 24, 1645–1659, <https://doi.org/10.1002/JOC.1098>, 2004.
- Karki, R., Talchabhadel, R., Aalto, J., and Baidya, S. K.: New climatic classification of Nepal, *Theor Appl Climatol*, 125, 799–808, <https://doi.org/10.1007/S00704-015-1549-0>, 2016.
- 540

- Karki, R., ul Hasson, S., Schickhoff, U., Scholten, T., and Böhner, J.: Rising Precipitation Extremes across Nepal, *Climate* 2017, Vol. 5, Page 4, 5, 4, <https://doi.org/10.3390/CLI5010004>, 2017.
- Kayastha, R. B. and Kayastha, R.: Glacio-Hydrological Degree-Day Model (GDM) Useful for the Himalayan River Basins, *Himalayan Weather and Climate and their Impact on the Environment*, 379–398, https://doi.org/10.1007/978-3-030-29684-1_19, 2019.
- Khanal, S., Tiwari, S., Lutz, A. F., Hurk, B. V.D., and Immerzeel, W. W.: Historical Climate Trends over High Mountain Asia Derived from ERA5 Reanalysis Data, *J Appl Meteorol Climatol*, 62, 263–288, <https://doi.org/10.1175/JAMC-D-21-0045.1>, 2023.
- Kiran S, Yogacharya, and Gautam, D. K.: Floods in Nepal: Genesis, Magnitude, Frequency and Consequences, in: *Proc. of the International Conference on Hydrology and Climate Change in Mountainous Areas*, <https://doi.org/10.13140/RG.2.1.2376.8489>, 2008.
- Kirschbaum, D., Kapnick, S. B., Stanley, T., and Pascale, S.: Changes in Extreme Precipitation and Landslides Over High Mountain Asia, *Geophys Res Lett*, 47, <https://doi.org/10.1029/2019GL085347>, 2020.
- Lane, S. N., Tayefi, V., Reid, S. C., Yu, D., and Hardy, R. J.: Interactions between sediment delivery, channel change, climate change and flood risk in a temperate upland environment, *Earth Surf Process Landf*, 32, 429–446, <https://doi.org/10.1002/esp.1404>, 2007.
- Lindersson, S., Brandimarte, L., Märd, J., and Di Baldassarre, G.: Global riverine flood risk - How do hydrogeomorphic floodplain maps compare to flood hazard maps?, *Natural Hazards and Earth System Sciences*, 21, 2921–2948, <https://doi.org/10.5194/nhess-21-2921-2021>, 2021.
- Marston, R., Kleinman, J., and Miller, M.: Geomorphic and forest cover controls on monsoon flooding, central Nepal Himalaya, *Mt Res Dev*, 16, 257–264, <https://doi.org/10.2307/3673948>, 1996.
- Martin-Vide, J.: Spatial distribution of a daily precipitation concentration index in peninsular Spain, *International Journal of Climatology*, 24, 959–971, <https://doi.org/10.1002/JOC.1030>, 2004.
- Mathers, C., Stevens, G., Ho, J., Ma Fat, D., Retno Mahanani, W., Andreev, K., Bassani, D., Black, B., Boerma, T., Boucher, P., Bray, F., Burton, T., Campbell, H., Chou, D., Cibulskis, R., Cousens, S., Ferlay, J., Gacic-Dobo, M., Garfield, R., Gemmill, A., Gerland, P., Ghys, P., Glaziou, P., Gu, D., Hill, K., Iaych, K., Inoue, M., Jakob, R., Jamison, D., Jha, P., Johnson, H., Lawn, J., Li, N., Liu, L., Lozano, R., Mahi, M., Murray, C., Newman, L., Oestergaard, M., Parkin, M., Peden, M., Pelletier, F., Rehm, J., Rudan, I., Say, L., Simons, E., Sismanidis, C., Spoorenberg, T., Stanecki, K., Stover, J., Strebel, P., Suzuki, E., Toroyan, T., Vos, T., Wardlaw, T., White, R., Wilmoth, J., and You, D.: WHO methods and data sources for global burden of disease estimates 2000–2011, 2013.
- Mazzoleni, M., Dottori, F., Cloke, H. L., and Di Baldassarre, G.: Deciphering human influence on annual maximum flood extent at the global level, *Commun Earth Environ*, 3, <https://doi.org/10.1038/s43247-022-00598-0>, 2022.
- Meyer, V., Becker, N., Markantonis, V., Schwarze, R., Van Den Bergh, J. C. J. M., Bouwer, L. M., Bubeck, P., Ciavola, P., Genovese, E., Green, C., Hallegatte, S., Kreibich, H., Lequeux, Q., Logar, I., Papyrakis, E., Pfurtscheller, C., Poussin, J.,

- 575 Przulski, V., Thieken, A. H., and Viavattene, C.: Review article: Assessing the costs of natural hazards-state of the art and knowledge gaps, <https://doi.org/10.5194/nhess-13-1351-2013>, 2013.
- Mohanty, M. P. and Simonovic, S. P.: A Comprehensive Approach for Floodplain Mapping through Identification of Hazard Using Publicly Available Data Sets over Canada, *Water (Switzerland)*, 14, <https://doi.org/10.3390/w14142280>, 2022.
- Monjo, R. and Martin-Vide, J.: Daily precipitation concentration around the world according to several indices, *International Journal of Climatology*, 36, 3828–3838, <https://doi.org/10.1002/JOC.4596>, 2016a.
- 580 Monjo, R. and Martin-Vide, J.: Daily precipitation concentration around the world according to several indices, *International Journal of Climatology*, 36, 3828–3838, <https://doi.org/10.1002/JOC.4596>, 2016b.
- Monjo, R.: Measure of rainfall time structure using the dimensionless n-index, *Clim Res*, 67, 71–86, <https://doi.org/10.3354/cr01359>, 2016.
- 585 Mosavi, A., Ozturk, P., and Chau, K. W.: Flood prediction using machine learning models: Literature review, <https://doi.org/10.3390/w10111536>, 27 October 2018.
- Nepal - Floods and Landslides Jul 1993 UN DHA Situation Reports 1-8 - Nepal | ReliefWeb: <https://reliefweb.int/report/nepal/nepal-floods-and-landslides-jul-1993-un-dha-situation-reports-1-8>, last access: 6 February 2023.
- 590 Nepal - Floods Situation Report No. 1, 26 July 1996 - Nepal | ReliefWeb: <https://reliefweb.int/report/nepal/nepal-floods-situation-report-no-1-26-july-1996>, last access: 6 February 2023.
- Nepal Appeal No. 01.55/2003 Annual Report - Nepal | ReliefWeb: <https://reliefweb.int/report/nepal/nepal-appeal-no-01552003-annual-report>, last access: 6 February 2023.
- Nepal: Landslides and Floods - Aug 2014 | ReliefWeb: <https://reliefweb.int/disaster/ls-2014-000103-npl>, last access: 6
- 595 February 2023.
- Neuhold, C., Stanzel, P., and Nachtnebel, H. P.: Incorporating river morphological changes to flood risk assessment: Uncertainties, methodology and application, *Natural Hazards and Earth System Science*, 9, 789–799, <https://doi.org/10.5194/nhess-9-789-2009>, 2009.
- Noy, I.: A Global Comprehensive Measure of the Impact of Natural Hazards and Disasters, *Glob Policy*, 7, 56–65, <https://doi.org/10.1111/1758-5899.12272>, 2016a.
- 600 Noy, I.: Comparing the direct human impact of natural disasters for two cases in 2011: The Christchurch earthquake and the Bangkok flood, *International Journal of Disaster Risk Reduction*, 13, 61–65, <https://doi.org/10.1016/j.ijdr.2015.03.009>, 2015.
- Noy, I.: Natural disasters in the Pacific Island Countries: new measurements of impacts, *Natural Hazards*, 84, 7–18, <https://doi.org/10.1007/s11069-015-1957-6>, 2016b.
- 605 Noy, I.: Prepared for the 2015 Global Assessment Report on Disaster Risk Reduction A NON-MONETARY GLOBAL MEASURE OF THE DIRECT IMPACT OF NATURAL DISASTERS, 2014.
- Pangali Sharma, T. P., Zhang, J., Koju, U. A., Zhang, S., Bai, Y., and Suwal, M. K.: Review of flood disaster studies in Nepal: A remote sensing perspective, <https://doi.org/10.1016/j.ijdr.2018.11.022>, 1 March 2019.

- Pervin, I. A., Rahman, S. M. M., Nepal, M., Haque, A. K. E., Karim, H., and Dhakal, G.: Adapting to urban flooding: A case of two cities in South Asia, *Water Policy*, 22, 162–188, <https://doi.org/10.2166/wp.2019.174>, 2020.
- Piacentini, T., Carabella, C., Boccabella, F., Ferrante, S., Gregori, C., Mancinelli, V., Pacione, A., Pagliani, T., and Miccadei, E.: Geomorphology-Based analysis of flood critical areas in small hilly catchments for civil protection purposes and earlywarning systems: The case of the feltrino stream and the Lanciano Urban Area (Abruzzo, Central Italy), *Water (Switzerland)*, 12, <https://doi.org/10.3390/w12082228>, 2020.
- Pinter, N., Jemberie, A. A., Remo, J. W. F., Heine, R. A., and Ickes, B. S.: Flood trends and river engineering on the Mississippi River system, *Geophys. Res. Lett*, 35, 23404, <https://doi.org/10.1029/2008GL035987>, 2008.
- Rentschler, J., Salhab, M., and Jafino, B. A.: Flood exposure and poverty in 188 countries, *Nat Commun*, 13, <https://doi.org/10.1038/s41467-022-30727-4>, 2022.
- Saki, S. A., Sofia, G., and Anagnostou, E. N.: Characterizing CONUS-wide spatio-temporal changes in daily precipitation, flow, and variability of extremes, *J Hydrol (Amst)*, 626, 130336, <https://doi.org/10.1016/J.JHYDROL.2023.130336>, 2023.
- Samela, C., Troy, T. J., and Manfreda, S.: Geomorphic classifiers for flood-prone areas delineation for data-scarce environments, *Adv Water Resour*, 102, 13–28, <https://doi.org/10.1016/j.advwatres.2017.01.007>, 2017.
- Sangüesa, C., Pizarro, R., Ibañez, A., Pino, J., Rivera, D., García-Chevesich, P., and Ingram, B.: Spatial and temporal analysis of rainfall concentration using the Gini Index and PCI, *Water (Switzerland)*, 10, <https://doi.org/10.3390/w10020112>, 2018.
- Sanyal, J. and Lu, X. X.: Application of Remote Sensing in Flood Management with Special Reference to Monsoon Asia: A Review, *Natural Hazards*, 283–301 pp., 2004.
- Serrano-Notivoli, R., Martín-Vide, J., Saz, M. A., Longares, L. A., Beguería, S., Sarricolea, P., Meseguer-Ruiz, O., and de Luis, M.: Spatio-temporal variability of daily precipitation concentration in Spain based on a high-resolution gridded data set, *International Journal of Climatology*, 38, e518–e530, <https://doi.org/10.1002/JOC.5387>, 2018.
- Shamsudduha, M. and Panda, D. K.: Spatio-temporal changes in terrestrial water storage in the Himalayan river basins and risks to water security in the region: A review, *International Journal of Disaster Risk Reduction*, 35, <https://doi.org/10.1016/J.IJDRR.2019.101068>, 2019.
- Sharma, P. p. T., Zhang, J., Koju, A. U., Zhang, S., Bai, Y., and Suwal, K. M.: International Journal of Disaster Risk Reduction Review of flood disaster studies in Nepal : A remote sensing perspective Til Prasad Pangali Sharma, *International Journal of Disaster Risk Reduction*, 34, 18–27, 2019.
- Shrestha, A. B. and Aryal, R.: Climate change in Nepal and its impact on Himalayan glaciers, *Reg Environ Change*, 11, 65–77, <https://doi.org/10.1007/s10113-010-0174-9>, 2011.
- Shrestha, A. B., Eriksson, M., Mool, P., Ghimire, P., Mishra, B., and Khanal, N. R.: Glacial lake outburst flood risk assessment of Sun Koshi basin, Nepal, *Geomatics, Natural Hazards and Risk*, 1, 157–169, <https://doi.org/10.1080/19475701003668968>, 2010.

- Simonovic, S. P., Karmakar, S., Cheng, Z., Kansal, M. L., and Singh, S.: Flood Management Issues in Hilly Regions of Uttarakhand (India) under Changing Climatic Conditions, *Water* 2022, Vol. 14, Page 1879, 14, 1879, <https://doi.org/10.3390/W14121879>, 2022.
- Slater, L. J., Khouakhi, A., and Wilby, R. L.: River channel conveyance capacity adjusts to modes of climate variability, *Sci Rep*, 9, 1–10, <https://doi.org/10.1038/s41598-019-48782-1>, 2019.
- Slater, L. J., Singer, M. B., and Kirchner, J. W.: Hydrologic versus geomorphic drivers of trends in flood hazard, *Geophys Res Lett*, 42, 370–376, <https://doi.org/10.1002/2014GL062482>, 2015a.
- Slater, L. J., Singer, M. B., and Kirchner, J. W.: Hydrologic versus geomorphic drivers of trends in flood hazard, *Geophys Res Lett*, 42, 370–376, <https://doi.org/10.1002/2014GL062482>, 2015b.
- Sofia, G. and Nikolopoulos, E. I.: Floods and rivers: a circular causality perspective, *Sci Rep*, 10, <https://doi.org/10.1038/s41598-020-61533-x>, 2020.
- Sofia, G., Di Stefano, C., Ferro, V., and Tarolli, P.: Morphological Similarity of Channels: From Linear Erosional Features (Rill, Gully) to Alpine Rivers, *Land Degrad Dev*, 28, 1717–1728, <https://doi.org/10.1002/ldr.2703>, 2017b.
- Sofia, G., Di Stefano, C., Ferro, V., and Tarolli, P.: Morphological Similarity of Channels: From Linear Erosional Features (Rill, Gully) to Alpine Rivers, *Land Degrad Dev*, 28, 1717–1728, <https://doi.org/10.1002/ldr.2703>, 2017c.
- Sofia, G., Ragazzi, F., Giandon, P., Dalla Fontana, G., and Tarolli, P.: On the linkage between runoff generation, land drainage, soil properties, and temporal patterns of precipitation in agricultural floodplains, *Adv Water Resour*, 124, 120–138, <https://doi.org/10.1016/j.advwatres.2018.12.003>, 2019.
- Sofia, G., Roder, G., Dalla Fontana, G., and Tarolli, P.: Flood dynamics in urbanised landscapes: 100 years of climate and humans' interaction, *Sci Rep*, 7, 1–12, <https://doi.org/10.1038/srep40527>, 2017a.
- Sofia, G., Tarolli, P., Cazorzi, F., and Dalla Fontana, G.: Downstream hydraulic geometry relationships: Gathering reference reach-scale width values from LiDAR, *Geomorphology*, 250, 236–248, <https://doi.org/10.1016/j.geomorph.2015.09.002>, 2015.
- Stover, S. C. and Montgomery, D. R.: Channel change and flooding, Skokomish River, Washington, *J Hydrol (Amst)*, 243, 272–286, [https://doi.org/10.1016/S0022-1694\(00\)00421-2](https://doi.org/10.1016/S0022-1694(00)00421-2), 2001.
- Subba, S., Ma, Y., and Ma, W.: Spatial and Temporal Analysis of Precipitation Extremities of Eastern Nepal in the Last Two Decades (1997–2016), *Journal of Geophysical Research: Atmospheres*, 124, 7523–7539, <https://doi.org/10.1029/2019JD030639>, 2019.
- Talchabhadel, R., Karki, R., Thapa, B. R., Maharjan, M., and Parajuli, B.: Spatio-temporal variability of extreme precipitation in Nepal, *International Journal of Climatology*, 38, 4296–4313, <https://doi.org/10.1002/JOC.5669>, 2018.
- Torti, J. M. I.: Floods in Southeast Asia: A health priority, *J Glob Health*, 2, <https://doi.org/10.7189/jogh.02.020304>, 2012.
- Tuladhar, D., Dewan, A., Kuhn, M., and Corner, R. J.: Spatio-temporal rainfall variability in the Himalayan mountain catchment of the Bagmati River in Nepal., *Theor Appl Climatol*, 139, 599–615, <https://doi.org/10.1007/S00704-019-02985-8>, 2020.

675 WHO: GlobalDALYmethodsJapan_2011_WHO, World Health Organization, 4, 2014.

Winiger, M., Gumpert, M., and Yamout, H.: Karakorum-Hindukush-western Himalaya: Assessing high-altitude water resources, *Hydrol Process*, 19, 2329–2338, <https://doi.org/10.1002/HYP.5887>, 2005.

World Bank, 2012. Weather and Climate Services in Europe and Central Asia: A Regional Review.

Yang, C. R. and Tsai, C. T.: Development of a GIS-based flood information system for floodplain modeling and damage
680 calculation, *J Am Water Resour Assoc*, 36, 567–577, <https://doi.org/10.1111/j.1752-1688.2000.tb04287.x>, 2000.

Zheng, G., Allen, S. K., Bao, A., Ballesteros-Cánovas, J. A., Huss, M., Zhang, G., Li, J., Yuan, Y., Jiang, L., Yu, T., Chen, W., and Stoffel, M.: Increasing risk of glacial lake outburst floods from future Third Pole deglaciation, *Nat Clim Chang*, 11, 411–417, <https://doi.org/10.1038/s41558-021-01028-3>, 2021.

Zheng, X., Maidment, D. R., Tarboton, D. G., Liu, Y. Y., and Passalacqua, P.: GeoFlood: Large-Scale Flood Inundation
685 Mapping Based on High-Resolution Terrain Analysis, *Water Resour Res*, 54, 10,013-10,033, <https://doi.org/10.1029/2018WR023457>, 2018.