



1 **Spring water anomalies before two consecutive earthquakes (Mw 7.7 and Mw 7.6)**
2 **in Kahramanmaraş (Türkiye) on 6 February 2023**

3

4 Sedat İnan^{1,*}, Hasan Çetin², Nurettin Yakupoğlu¹

5

6 ¹Department of Geological Engineering, Istanbul Technical University, Ayazağa,
7 Istanbul, Türkiye, 34467

8 ²Department of Geology, Çukurova University, Adana, Türkiye, 01330

9 *Corresponding Author: sedatinan@itu.edu.tr

10 **ABSTRACT**

11 Understanding earthquake phenomena is always challenging. Search for reliable
12 precursors of earthquakes are important but requires systematic and long-time monitoring
13 employing multi-disciplinary techniques. In search of possible precursors, we obtained
14 commercially bottled spring waters dated before and after the earthquakes of 6 February
15 2023. Hydrogeochemical precursors have been detected in commercially bottled natural
16 spring waters (Ayran Spring and Bahçepınar Spring) which are at a distance of about 100
17 km and 175 km from the epicenters of the Mw 7.7 and Mw 7.6 Kahramanmaraş (Türkiye)
18 Earthquakes of 6 February 2023, respectively. The available water samples cover the
19 period from March 2022 to March 2023. The pre-earthquake anomaly is characterized by
20 an increase in electrical conductivity and major ions (Ca^{2+} , Mg^{2+} , K^+ , Na^+ , Cl^- , and SO_4^{2-}
21) compared to the background for Ayran Spring water samples. The pre-earthquake
22 anomaly lasted for at least six months. The anomaly in major ions sharply declined and
23 the ion content approached the background values about two weeks after the
24 earthquakes. Although only 6.5 kilometers away from the Ayran Spring, the bottled water
25 samples of the Bahçepınar Spring did not show any anomalies in electrical conductivity;
26 therefore, the samples were not analyzed for ion content. Bahçepınar water is collected
27 from shallow boreholes dug into alluvial deposits which we believe are decoupled from
28 the basement rocks and this may be the reason for the lack of abnormal water chemistry
29 prior to the earthquakes. This attests to the fact that sampling locations are very important
30 in the detection of possible earthquake precursors. Results on the Ayran spring water
31 samples indicate that spring water chemical anomalies of discrete samples may provide



32 valuable information on pre-earthquake crustal deformation. Monitoring of spring waters,
33 along with other monitoring techniques in a multidisciplinary network, and for a sufficiently
34 long time, could potentially enable obtaining reliable proxy indicators of pre-earthquake
35 **crustal deformation.**

36

37 **Keywords:** geochemical anomalies, spring water, earthquake precursors,
38 Kahramanmaraş earthquakes, East Anatolian Fault Zone (EAFZ), Türkiye

39

40 1. Introduction

41 Two devastating earthquakes (Mw 7.7 and Mw 7.6) struck the Kahramanmaraş area in
42 Southern Turkey on 6 February 2023; the earthquakes occurred about 9 hours apart. The
43 earthquakes caused devastation **claiming more than 50,000 deaths**; leaving behind
44 thousands injured and/or homeless. Earthquakes of destructive magnitudes (e.g., M>7)
45 are naturally expected to occur at plate boundary settings (Figure 1) and Kahramanmaras
46 province is at the junction of the East Anatolian Fault System (EAFS) and the Dead Sea
47 Fault System (DAFS). However, the reason why such natural events turn into disasters
48 is mainly due to a lack of preparedness. Where buildings are not built to be sufficiently
49 earthquake-resistant, monitoring of crustal deformation and searching for reliable pre-
50 earthquake signals become more important. This is obviously a big challenge for earth
51 scientists to overcome. Although there is still a long way to go on this front, the scientific
52 literature is full of scattered but promising and encouraging cases.

53 Earthquakes are complex natural phenomena and their predictions have been long
54 viewed as difficult, if not impossible (e.g., Geller et al., 1997). Geochemical observations
55 to identify earthquake precursors were initiated in the late 1960s (Rikitake, 1979; Wakita
56 1996). Reviewing twenty years of relevant data Turcotte (1991) concluded that large
57 earthquakes are not preceded by reliable seismic precursors. Moreover, Geller et al.
58 (1997) claimed that earthquakes can never be predicted. However, for the last few
59 decades, there have been numerous reports of **ground-based anomalies** preceding major
60 earthquakes. (including but not limited to Rikitake, 1979; Dobrovolsky et al., 1979;
61 Birchard and Libby, 1978; Hauksson, 1981; Wakita et al., 1988; Sultankhodhaev, 1984;



62 Thomas et al., 1986; Rikitake, 1987; Etiope et al., 1997; Bella et al., 1998; Virk and Singh,
63 1993; King et al., 1995; Planinic et al., 2004; Claesson et al., 2004; Hartmann and Levy,
64 2006; Papadopoulos et al., 2006; Uyeda et al., 2008; İnan et al., 2008; İnan et al., 2010;
65 İnan et al., 2012a,b,c; Skelton et al., 2014 and 2019; Barberio et al., 2017; Ouzounov et
66 al., 2021; Gori and Barberio, 2022; Xiang and Peng, 2023). Compiling a review of claimed
67 precursors, Cicerone et al. (2009) conducted a survey of published scientific literature on
68 earthquake precursors and concluded that precursory anomalies seem to be recorded
69 where there is modern instrumentation. İnan et al. (2010 and 2012a) provided hints to
70 select monitoring sites. Recently, Conti et al. (2021) have provided a short review of
71 ground-based observations before earthquakes

72 **Hydro-geochemical anomalies observed nearby seismic events** are generally interpreted
73 to be related to the alteration of the groundwater circulating system by the changes in the
74 crustal stress/strain before earthquakes and mixing of different aquifers (e.g., Scholz et
75 al., 1973; Nur, 1974; Sibson et al., 1975; Sugisaki et al., 1996; Tsunogai and Wakita,
76 1995; Toutain et al., 1997; Claesson et al., 2004; Pérez et al., 2008; İnan et al., 2010;
77 Grant et al., 2011; İnan et al., 2012c; Doglioni et al., 2014; Ingebritsen and Manga, 2014;
78 Skelton et al., 2014 and 2019; Barberio et al., 2017; Gori and Barberio, 2022; Xiang and
79 Peng, 2023). However, another different approach based on “stress-activated positive
80 hole currents” has been suggested to play a role in the development of physicochemical
81 pre-earthquake stress indicators (Freund, 2011; Paudel et al., 2018)


82 As suggested by Nur (1974) and later by Rikitake (1987) precursory phenomena may
83 have a common physical basis which Scholz et al. (1973) called the “Dilatation and water
84 diffusion (DWD) model”. Roeloffs (1996) noted that with respect to earthquake hydrology,
85 mechanical and fluid-dynamic effects can be modeled using poroelasticity. More recently,
86 the DWD model has been explained further (e.g., Doglioni et al., 2014; Wang and Manga,
87 2021). However, other authors have proposed a fundamentally different approach
88 (Freund et al., 2006; Freund, 2008; Freund, 2011; Paudel et al., 2018) to study and
89 evaluate physicochemical pre-earthquake stress indicators. Until the mechanism
90 controlling pre-earthquake processes is fully understood, it is worth noting that the
91 success of any pre-earthquake stress indicators may be compromised by the ever-
92 present crustal heterogeneity, anisotropy, and/or crustal blocks (Areshidze et al., 1992;



93 Tansi et al., 2005; Sol et al., 2007; İnan et al., 2012a; Yu et al., 2023). Microplate and/or
94 block boundaries are obstacles to pre-earthquake strain to transfer from one block to the
95 other (İnan et al., 2012a; Yu et al., 2023).

96

97 A multi-disciplinary earthquake observation network (GPS, seismology, soil radon, and
98 spring water monitoring stations) was established in Kahramanmaraş and surrounding
99 provinces along the fault zones (Adana, Hatay, Malatya, Elazığ, Bingöl) in 2007 under the
100 scope of the TURDEP Project (İnan et al. 2007). In the Kahramanmaraş area, due to its
101 quiescence, also borehole tilt monitoring stations were established. Continuous
102 monitoring was continued until the middle of 2012 and valuable multi-disciplinary data
103 were collected. However, throughout these five years, no earthquake of significant
104 magnitude (e.g. $M > 6$) occurred to test the usefulness of the monitoring network, the
105 project was terminated by the funding organization due mainly to a lack of vision. As a
106 result, the earth science community was caught unprepared when two consecutive
107 devastating earthquakes struck the area on 6 February 2023. No ground (except GPS
108 and seismology) monitoring station data were available to detect possible pre-earthquake
109 anomalies. However, following the Mw 7.7 and Mw 7.6 Kahramanmaraş earthquakes, we
110 searched for bottled spring waters to analyze in search of possible pre-earthquake
111 anomalies. This proved difficult as the water supply to the large community affected by
112 the earthquakes was quite limited and businesses providing bottled spring waters were
113 also mostly shut down. Finally, we were able to obtain commercially bottled water
114 samples (dated before and after the earthquake) from the Ayran and Bahçepinar springs
115 which are located within about a 6.5-kilometer distance in the Osmaniye Province. The
116 spring waters are about 100 kilometers and 175 kilometers from the epicenter of the first
117 (Mw 7.7) and the second (Mw 7.6) earthquakes, respectively (Figure 1B). In this study,
118 we conducted electrical conductivity (Ec) measurements on bottled waters, and based on
119 the Ec results, we selected samples for analysis of major ions in water in search of pre-
120 earthquake anomalies. The spring water samples cover the range from March 2022 to
121 March 2023.



122

123 **Figure 1. A)** Neotectonics map of the Türkiye and surroundings (compiled from Sengör
124 and Yılmaz, 1981; Sengör et al., 1985; Hancock and Barka, 1987; Şaroğlu et al., 1992;
125 Barka and Reilinger, 1997; Bozkurt, 2001). **B)** Active fault map of the region affected by
126 the February 2023 Earthquakes (Perinçek and Çemen, 1990; Şaroğlu et al., 1992a; Cetin
127 et al., 2003). Red stars show the epicenters of the Mw 7.7 and 7.6 Kahramanmaraş
128 Earthquakes of 6 February 2023, and Mw 6.4 Hatay Earthquake of 20 February 2023.
129 Filled red circles show the locations of the earthquakes (Mw>4.5) that occurred in the
130 area (circle area with a radius of 150 km from the location of the water spring) between
131 September 2022 and 5 February 2023. Details of the earthquakes are given in Table 1.
132 TTJ is Türkoğlu Triple Junction. Beach balls are fault plane solutions of earthquakes and
133 were obtained from the Bogazici University Kandilli Observatory and Earthquake
134 Research Institute (KOERI) of Turkey; www.koeri.edu.tr



135 2. Active tectonics of the Kahramanmaraş region

136 Kahramanmaraş region takes place in the suture zone formed by the collision between
137 Arabian and Anatolian plates (Figure 1A). After this collision, very important strike-slip
138 fault zones were developed in the Anatolian plate due to the continuous northward
139 movement of the Arabian plate and the resulting westward movement or escape of the
140 Anatolian plate along two major fault zones, the North Anatolian Fault Zone (NAFZ) and
141 the East Anatolian Fault Zone (EAFZ) (Ketin, 1948; McKenzie, 1972; Dewey and Şengör,
142 1979; Şengör and Yılmaz, 1981; Hempton, 1982; Şengör et al., 1985).

143 The East Anatolian Fault Zone (EAFZ) is approximately 550 km long, northeast-
144 southwest trending, sinistral strike-slip fault (Figure 1A). It was first described by Allen
145 (1969) and mapped by Arpat and Şaroğlu (1972). The EAFZ starts from Karlıova Triple
146 Junction in the northeast, and it runs in the southwest direction, passes near the east-
147 southeast of Kahramanmaraş, and joins another triple junction at Türkoğlu (TTJ in Figure
148 1B). The EAFZ then continues to the Hatay in the south direction to merge into the Dead
149 Sea Fault Zone (DSFZ) (Allen, 1969; Arpat and Şaroğlu, 1972; Dewey and Sengör, 1979;
150 Rotstein, 1984; Şengör et al., 1985; Kelling et al., 1987; Şaroğlu et al., 1992a and 1992b;
151 Cetin et al., 2003; Yönlü et al., 2017). There are different interpretations, however, for the
152 remainder of the fault zone after Türkoğlu Triple Junction (marked as TTJ in Figure 1B).
153 Some studies extend the fault zone southwesterly to the Mediterranean Sea (McKenzie,
154 1972; Dewey et al., 1973; Jackson and McKenzie, 1984; Barka and Kadinsky-Cade,
155 1988; Karig and Kozlu, 1990; Kempler and Garfunkel, 1991; Westaway and Arger, 1996),
156 joining it with the Cyprian Arc along which the convergence is taking place between the
157 African and Anatolian plates (McKenzie, 1976; Dewey and Şengör, 1979). Others think
158 that the fault zone ends around the TTJ (Lovelock, 1984; Chorowicz et al., 1994).
159 According to Muehlberger and Gordon (1987), the EAFZ becomes the northern branch
160 of the DSFZ

161 The seismicity of the study area is controlled by a complex interaction of the African,
162 Arabian, and Eurasian plates (McKenzie, 1972). The seismicity of the EAFZ has been
163 minimal for most of the last 100 years (Ambraseys, 1989). Historical earthquake records



164 show that Kahramanmaraş and its surroundings were affected by **the** two major
165 earthquakes in AD 1114 and AD 1513 (Soysal et al., 1981; Ambraseys, 1989). There had
166 been a long quiescence of more than 500 years in the Kahramanmaraş area before the
167 Mw 7.7 and Mw 7.6 earthquakes struck on 6 February 2023. About one year before these
168 earthquakes occurred, the area had been seismically quiet as suggested by only a few
169 $M > 4.5$ earthquakes occurring in a circular area with a radius of 150 km; taking the Ayran
170 spring water as the center (Figure 1B and Table 1). The fault plane solutions (FPS) for
171 earthquakes #3, #4, and #5 suggest mainly normal faulting, whereas, for others
172 (earthquakes #1, #2, and #6), FPS suggest movement on dominantly left lateral strike-
173 slip faults (Figure 1B) as expected for left-lateral strike-slip nature of the EAFZ.

174 **Table 1.** Earthquakes' time, magnitude, and locations as received from www.koeri.edu.tr.
175 Earthquakes #1, #2, and #3 are the earthquakes of February 2023. Earthquakes #4, #5,
176 and #6 are those that have occurred in the circular area (with a radius of 150 km from the
177 Ayran spring water location) between September 1st, 2022 and 5 February 2023. The
178 locations of these earthquakes are given on the map (Figure 1B).

Earthquake #	Magnitude (Mw)	Date	Time (GMT)	Latitude	longitude
1	7.7	06.02.2023	01:17	37.1757	37.0850
2	7.6	06.02.2023	10:24	38.0818	37.1773
3	6.4	20.02.2023	17:04	36.0713	36.1012
4	4.6	12.01.2023	20:40	35.5712	36.6723
5	4.9	18.12.2022	18:13	36.3978	36.4455
6	5.0	11.10.2022	15:48	37.3025	36.2403

179

180 3. Samples and methods

181

182 3.1. Spring water samples

183 The spring water samples were received in commercial polyethylene bottles and brought
184 to Istanbul Technical University Laboratory for electrical conductivity measurements and
185 major ion analyses. Some of the samples had been bottled up to several months before
186 the analyses. However, this does not create any concern because much longer storage
187 in this kind of bottle has been reported to be appropriate in terms of keeping reliable
188 concentrations (Tsunogai and Wakita, 1995; İnan et al., 2012c).



189 The spring water samples cover the range from March 2022 to March 2023. It is worth
190 noting that the oldest sample predating the earthquakes was AYR 1 (dated 8 March 2022)
191 from the Ayran Spring. Other bottled water samples we could obtain from both springs
192 were dated between September 2022 and March 2023. In fact, we could not obtain any
193 samples dated between 8 March and 14 September 2022. The samples from September,
194 October, and November 2022 are limited but from December 2022 to January 2023,
195 available samples are several per month (Table 2).

196 **3.2. Spring water analysis**

197 We first screened the bottled water samples by conducting electrical conductivity (Ec)
198 measurements, and based on the results, we selected samples for analysis of major ions.
199 Samples of the AYR spring water were analyzed by ion chromatography as discussed by
200 Zeyrek et al. (2010). Briefly, the samples were filtered at 0.45 μm and split into two
201 portions before analysis using an ion chromatography instrument (Dionex ICS 1000).
202 Sodium carbonate and methane-sulfonic acid were used as eluents for anion and cation
203 analyses, respectively. For calibration, DIONEX Certified Reference Standards were
204 used. Deionized water with a resistance better than 18.2 Megaohm was used for the
205 preparation of all eluents. Repeated measurements ascertained that the analytical
206 uncertainties for all anions and cations were below 5%. Electrical conductivity (Ec)
207 measurements for the bottled Bahçepinar (BPN) spring water samples and both Ec and
208 Ion analysis results for the bottled Ayran spring waters are listed in Table 2.

209 **3.3. Statistical analysis of the data**

210 For the statistical treatment of the data on major ion contents of the water samples, we
211 calculated the weighted average (weighted compared to the analytical error for each
212 point) and computed the 2σ external error ($2 \times \alpha_e$) from the following equation

213

$$\alpha_e = \frac{\sum_{i=1}^n (x_i - \bar{x})^2 / \sigma_i^2}{(n-1) \sum_{i=1}^n 1 / \sigma_i^2}$$

214



215 where x is the average and σ the analytical error on each measured point. The 2σ external
216 error (α_e) considers the general variability of all datasets and the analytical error on each
217 point; thus, we obtained the total error envelope for the samples that we consider
218 representing background (from 15 February to 31 March 2023; see Table 2 and Figure
219 3B).

220

221 **3.4. Relation between earthquake magnitude, distance, and precursory duration**

222 Slightly different relations between earthquake magnitude, duration of a precursory
223 anomaly, and the distance of the monitoring site to the earthquake epicenter have been
224 proposed. Dobrovolsky et al. (1979) proposed a theoretical relation ($D = 10^{0.43M}$) between
225 earthquake magnitude and maximum epicentral distance at which geochemical
226 anomalies may be observed. This relation assumes a homogenous and isotropic crust.
227 Where M is the earthquake magnitude and D is the distance in kilometers to the
228 earthquake epicenter. Rikitake (1987) noted a slightly different relation ($\log T = a + b \cdot M$;
229 where a and b are constants, T is the duration of anomaly and M is the magnitude of an
230 earthquake). Moreover, Sultankhodhaev (1984) also reported a relation, between
231 earthquake magnitude, the distance of the monitoring site to the earthquake epicenter,
232 and duration of precursory anomaly ($\log(DT) = 0.63 \cdot M - b$; where D is the distance in
233 km, T is the duration of a precursory anomaly in days, and M is earthquake magnitude; b
234 is a constant taken as 0.15. All of **these three relations** provide a helpful initial idea about
235 what to expect of precursory anomalies in terms of duration and distance to the
236 earthquake epicenter. Inan et al. (2008 and 2010) verified Dobrovolsky et al.'s (1979)
237 relation for medium-size earthquakes ($M < 5.3$). Accordingly, for an earthquake of
238 magnitude 4.5, the maximum distance for detection of possible geochemical anomalies
239 in the Ayran Spring water will be about 100 km. For contingency, we took a 150 km radius
240 and listed in Table 1 the earthquakes with $M > 4.5$ occurring between September 2022
241 and 5 February 2023 in order to compare with the water geochemical data we obtained
242 in this study.

243




244

245

4. Results and Discussion

246

247 The bottled water samples from the Bahcepinar (BPN) did not show any meaningful (e.g.,
248 significant) variations in electrical conductivity (Ec) values; varying in a narrow range
249 between 220 and 230 microsiemens/cm (Table 2). Therefore, these samples were not
250 analyzed for major ions content because change (increase/decrease) of major ions
251 contents is expected to result in Ec variation (İnan et al., 2010; İnan et al., 2012c).
252 However, the bottled water samples from the Ayran (AYR) spring showed major variations
253 in the Ec values; varying in range between 50 and 200 microsiemens/cm. Therefore, the
254 AYR samples were analyzed for major ions. Possible reasons for not detecting any
255 anomaly in the Ec measurements of the BNP spring water samples have been
256 investigated. The investigation suggests that the reason may be the geological
257 environment of the springs. The AYR spring water emanates from Middle-Upper
258 Ordovician age metamorphic rocks (Kardere Formation) made up of quartzite,
259 metasandstone, metasiltstone, and metashale (Usta et al., 2015 and 2017), whereas the
260 BPN spring water is collected from shallow boreholes dug into valley-filling Quaternary
261 age alluvial deposits that are underlain by ophiolite (Figure 2). The alluvial deposits reach
262 a thickness of about two hundred meters and the water reservoir within the alluvium
263 deposit is fed by precipitation and a nearby Bekdemir stream flowing towards the alluvial
264 deposit. It is interesting that the streams disappear to the south; suggesting that the
265 stream (creek) water is captured by the alluvial deposit. Since the BPN water is collected
266 from shallow boreholes (less than 100 meters) dug into alluvial deposits, we believe that
267 the alluvial deposits are decoupled from the basement rocks (which undergo pre-
268 earthquake stress) and this may be the reason for the lack of anomaly in water chemistry
269 prior to the earthquakes. This testifies to the importance of adequate geological
270 knowledge of the area before sampling discrete geochemical samples (water or soil gas)
271 and/or continuous monitoring in search of pre-earthquake signals (İnan et al., 2008; İnan
272 et al., 2010).



273

274 **Figure 2.** Locations and local geology of the water springs. (Modified from Usta et al.,
 275 2015 and 2017). The Ayrancı spring water emanates from a fault in the Metamorphic
 276 Kardere Formation (blue triangle shown at the lower right in the map) whereas the
 277 Bahçe spring water is obtained from the Quaternary Alluvium (blue triangle shown
 278 at the upper left in the map).

279

280 Variations of major ions in the AYR spring water samples are significant. It is clear that
 281 pre-earthquake anomalies exceed the α_e (Figures 3B and Table 2). Before any
 282 interpretation, we need to make sure that geochemical time series are not affected by
 283 meteorological conditions. In this context, meteorological data have been obtained from



284 the Osmaniye Meteorology Station (located about 32 km SW of the AYR spring) and the
 285 daily average air temperature and rainfall are shown in Figure 3c.

286 **Table 2.** Ec and major ion analysis results for the Ayran (AYR) and the Ec analysis results
 287 for the Bahçepınar (BPN) bottled waters. The data for Ca^{2+} , Mg^{2+} , K^+ , Na^+ , Cl^- , SO_4^{2-} for
 288 the AYR samples are plotted in Figure 3B. Standard deviation (2σ) has been computed
 289 considering cations/anions contents of samples dated from 15 February to 31 March
 290 2023; the period which is considered to nearly represent background concentrations of
 291 the water. These samples are marked in bold fonts.

Sample ID	Date	Cl^-	SO_4^{2-}	Na^+	K^+	Mg^{+2}	Ca^{+2}	AYR Ec	Date	BPN Ec
AYR 1	08.03.2022	2.99	8.34	4.34	0.39	3.22	6.92	50	19.09.2022	220
AYR 2	14.09.2022	7.37	13.08	12.10	1.08	7.73	17.54	150	07.11.2022	230
AYR 3	06.10.2022	9.73	14.79	15.08	1.34	9.20	20.10	180	12.12.2022	230
AYR 4	03.11.2022	9.99	15.52	15.66	1.39	9.50	20.72	170	19.12.2022	220
AYR 5	13.12.2022	7.45	13.43	11.93	1.05	7.59	16.58	150	30.12.2022	220
AYR 6	26.12.2022	11.06	17.35	16.49	1.56	10.19	22.19	190	08.01.2023	220
AYR 7	29.12.2022	11.08	17.20	16.84	1.50	10.20	22.33	180	20.01.2023	220
AYR 8	30.12.2022	10.97	17.29	16.78	1.50	10.17	22.28	190	24.01.2023	220
AYR 9	03.01.2023	10.62	17.23	16.26	1.45	10.06	23.04	170	28.01.2023	220
AYR 10	06.01.2023	11.12	17.56	16.91	1.49	10.29	22.51	190	04.02.2023	220
AYR 11	11.01.2023	11.41	17.96	16.90	1.50	10.43	22.81	190	11.02.2023	220
AYR 12	12.01.2023	11.60	18.21	17.22	1.53	10.50	22.99	200	17.02.2023	220
AYR 13	27.01.2023	9.83	16.20	14.24	1.25	8.89	19.35	160	18.02.2023	230
AYR 14	31.01.2023	11.04	17.62	15.81	1.39	9.87	21.58	180	02.03.2023	230
AYR 15	01.02.2023	11.43	17.85	16.21	1.43	10.04	21.97	190	13.03.2023	220
AYR 16	10.02.2023	9.09	15.59	13.29	1.16	8.46	18.33	180	22.03.2023	230
AYR 17	12.02.2023	6.00	12.47	9.36	0.79	6.29	13.51	120		
AYR 18	13.02.2023	4.25	10.69	6.96	0.56	4.95	10.38	90		
AYR 19	15.02.2023	3.54	10.30	6.28	0.50	4.79	9.65	80		
AYR 20	16.02.2023	3.56	13.64	7.51	0.67	6.91	12.60	110		
AYR 21	28.02.2023	3.29	10.54	5.79	0.47	4.60	9.23	80		
AYR 22	02.03.2023	3.26	10.08	5.48	0.44	4.20	8.59	70		
AYR 23	11.03.2023	3.36	9.85	5.49	0.43	4.21	8.62	70		
AYR 24	13.03.2023	3.28	9.91	5.47	0.44	4.22	8.68	70		
AYR 25	20.03.2023	3.28	9.96	5.36	0.43	4.22	8.73	70		
AYR 26	24.03.2023	3.20	10.02	5.35	0.42	4.14	8.45	70		
AYR 27	31.03.2023	3.31	10.13	5.40	0.43	4.16	8.47	70		
	<i>mean</i>	0.37	1.13	0.60	0.05	0.46	0.94	7.78		
	Σ	0.33	1.19	0.79	0.08	0.88	1.34	13.64		
	2σ	0.65	2.38	1.57	0.17	1.76	2.68	27.28		
	<i>mean +1 σ</i>	3.64	11.32	6.19	0.51	5.04	9.81	83.64		
	<i>mean +2 σ</i>	2.99	8.94	4.62	0.35	3.28	7.13	56.36		

292

293 Air temperature gradually decreases from about 30°C in September 2022 to less than
 294 10°C in February 2023 (Figure 3C). Daily rainfall is noticeably present in November 2022
 295 and March 2023. Normally, variations in air temperature are not expected to affect the
 296 chemical contents of the spring water (İnan et al., 2010 and 2012) but the effect of rainfall



297 on soil radon concentration **is dominant** (Inan et al., 2008, 2010, 2012b, Seyis et al.,
298 2022). All earthquakes listed in Table 1 are plotted on the meteorology time series in
299 Figure 3C and this shows that major and heavy rainfall took place right after the
300 devastating earthquakes of 6 February 2023. Based on the relatively low EC and low
301 major ion contents of the AYR spring water (Table 2) that is bottled and commercially
302 distributed, it can be said that this water is of shallow origin (Di Luccio et al., 2018). A
303 comparison of the geochemical time series and significant variations shown in Figure 3B
304 and the daily average rainfall data shown in Figure 3C reveals no correlations. Inan et al.
305 (2010 and 2012) compared meteorological time series with hydrogeochemical time series
306 and noted that meteorological conditions do not seem to play a role in water's major ion
307 contents. In this study, we compare rainfall data and geochemical time series (Figure 3)
308 and, as there is no correlation, we conclude that the increase of major ion contents
309 observed in AYR spring waters are not related to atmospheric variations (e.g., rainfall).
310 Therefore, it is safe to conclude that the chemical changes recorded in the spring water
311 must be related to crustal deformation associated with earthquake stress buildup.

312 As shown in Figure 3B, changes in the concentration of the major ionic species dissolved
313 in the AYR spring water were observed. Positive anomalies are recorded in the Ca^{2+} ,
314 Mg^{2+} , K^+ , Na^+ , Cl^- , and SO_4^{2-} contents (mg/l) before the 6 February Mw 7.7 and 7.6
315 Kahramanmaraş Earthquakes (Figure 3b; Table 2). These positive anomalies (increase
316 in dissolved ion content) started as early as September 2022; suggesting a pre-
317 earthquake anomaly of nearly six months. Considering Sultankhodhaev's (1984) relation
318 ($\log(DT) = 0.63 * M - b$) between earthquake magnitude, precursory anomaly duration,
319 and the distance of the earthquake epicenter to the monitoring site, such a long duration
320 (six months) of a precursory anomaly we report in this study is very likely because the
321 magnitudes of the 7.7 and 7.6 devastating earthquakes are sufficiently big to cause such
322 a long precursory anomaly at a location about 100 km from the epicenter. Considering
323 the relation proposed by Sultankhodhaev (1984), such a magnitude of the earthquake
324 theoretically **should lead to months-long of precursory anomaly** in the geochemical
325 parameters at locations hundreds of kilometers far from the epicentral area.



326 In regard to changes in the dissolved ions in the AYR spring water, the following changes
327 are **imminent**. The Ca^{2+} and Na^+ content increase (for the period between September
328 2022 to 15 February 2023) above the background by about 14 (mg/l) and 10 (mg/l),
329 respectively, and reach up to 22 (mg/l) and 16 (mg/l), respectively. This increase started
330 about six months before the 6 February earthquakes (EQ # 1 and EQ #2). Since we could
331 not obtain samples between 8 March 2022 and 14 September 2022, the anomaly could
332 have possibly started even earlier (any time between March and August 2022); so the
333 positive anomaly (e.g., increase) in the major ions started at least six months before the
334 6 February 2023 earthquakes. The Mg^{2+} content also increased from about 4 (mg/l) to 10
335 (mg/l) in the period September 2022 to 15 February 2023. Similar major increases were
336 also detected in Cl^- , and SO_4^{2-} contents. Water samples are relatively poor in K^+ content
337 therefore the increase, due to the scale of the graph, is not very obvious in Figure 3B.
338 However, the values given in Table 2 clearly indicate about four times an increase in the
339 K^+ content compared to the background concentrations (post-seismic samples collected
340 between February 15 and 31 March 2023).


341 **The pre-earthquake anomaly** in the AYR water samples is characterized by an increase
342 of up to 400% for the Ec and also major ions; namely Ca^{2+} , Mg^{2+} , K^+ , Na^+ , Cl^- , and SO_4^{2-}
343 before the 6 February 2023 Mw 7.7 and Mw 7.6 earthquakes (Figure 3B). Post-
344 earthquake samples show decreasing trends in all major ions. Analyses results of the
345 post-earthquake dated samples show that the spring water has had chemical stability
346 since the Middle February-Early March 2023; just two to three weeks after the
347 earthquakes (Figure 3B). We have also obtained a chemical analysis report on AYR water
348 submitted with the business license application of the company dated 29 August 2012.
349 The chemical analysis data of the samples collected more than 10 years ago include
350 values only for Na^+ , Cl^- , and SO_4^{2-} as 3.86, 3.12, and 8.37 mg/l, respectively. These
351 values are very close to the analysis result of the AYR water sample dated 8 March 2022
352 (AYR 1 which is the oldest sample in our data set) and the AYR water samples collected
353 after 15 February (Table 2); confirming that these samples represent background values
354 for the AYR spring water.



355 Immediately after the earthquake, the values started to decrease suggesting a reversible
356 chemical change (Figure 3B; Table 2). It is worth mentioning that the broad positive
357 anomaly detected in the AYR water chemistry (Figure 3B) that lasted for about six months
358 before the Mw 7.7 and Mw 7.6 earthquakes shows some transient decreases (about
359 Middle December 2022 and toward the end of January 2023). Following each transient
360 decrease, an increase in ion contents is observed and the broad positive anomaly
361 (starting from September 2022) is sustained until the date of the major earthquakes of 6
362 February 2023. The observations of sudden decrease and rebound in the major ion
363 contents of the water samples (taking place in Mid December 2022 and end of January
364 2023) may suggest sudden and short-lived crustal stress release related to smaller
365 earthquakes (e.g., EQ # 4 and EQ # 5). Soon after the major earthquakes (EQ # 1 and
366 EQ # 2), the major ion contents of the water samples show a sharp decline; almost
367 approaching the background values as early as 15 February 2023. One single positive
368 anomaly after the major earthquakes (EQ #1 and EQ #2) is detected in the sample dated
369 16 February 2023. The further increase of the ion contents of this sample seems to
370 suggest a short-term stress buildup prior to EQ # 3 (Mw 6.4) that occurred about 120 km
371 to the south of the Ayran Spring water location (Figure 1B). Considering Dobrovolsky et
372 al.'s (1979) theoretical relation ($R= 10^{0.43 \cdot M}$), an increase in major ions contents of the
373 Ayran Spring water is very likely to take place due to an earthquake of magnitude 6.4
374 occurring in 120 km distance.

375


376



377

378

379 **Figure 3.** Time variation graphs of Ec for the Ayran (AYR) and the Bahçepinar (BPN)
 380 bottled waters (A) and major ions for the AYR bottled waters (B). All data are listed in
 381 Table 2.



382

383 **Figure 3. Cont.** Daily average air temperature and rainfall at the Osmaniye meteorology
384 station (37°07'N, 36°25'E; 32 km SW of the Ayran spring) between 1 September 2022
385 and 30 March 2023 (C).

386 ([https://www.meteoblue.com/tr/hava/historyclimate/weatherarchive/
387 osmaniye_turkiye_303195](https://www.meteoblue.com/tr/hava/historyclimate/weatherarchive/osmaniye_turkiye_303195)). EQ1 through EQ6 are the earthquakes listed in Table 1.

388

389 We have shown and discussed the reliable precursory anomalies in the major ions of the
390 bottled AYR spring water prior to the Mw 7.7 and Mw 7.6 earthquakes that occurred in
391 the Kahramanmaraş region on 6 February 2023. However, the process(es) leading to the
392 build-up of geochemical anomalies related to the earthquake cannot be inferred with
393 certainty. However, some inferences based on previous observations can be made. For
394 instance, Sibson (1992) suggested that extensive hydro-fracture dilatancy might develop
395 prior to failure leading to the earthquake. Development of fractures probably enhances
396 water circulation and mixing of different reservoirs leading to pre-earthquake anomalies
397 (Italiano et al., 2004; Federico et al., 2008; İnan et al., 2010; İnan et al., 2012c; Skelton
398 et al., 2014; Ingebritsen and Manga, 2014; Doglioni et al., 2014; Barberio et al., 2017;
399 Skelton et al., 2019; Wang and Manga, 2021; Gori and Barberio, 2022;). Although the
400 process(es) responsible for chemical anomalies detected in the Ayran spring waters prior
401 to the 6 February 2023 earthquakes cannot be suggested with any certainty at this stage,
402 two immediate mechanisms emerge: 1) a simple increase in fluid flow in the surrounding
403 of the future epicenter and selective dissolution of some K–Mg–Ca-rich rocks (e.g.,



404 Federico et al., 2008); or 2) “electro-corrosion” whereby the dissolution of rocks is
405 accelerated by the flow of stress-activated positive hole currents (Balk et al., 2009;
406 Freund, 2011; Paudel et al., 2018). Following the second mechanism, the increased
407 content of major ions in water could be related to the oxidation of water to hydrogen
408 peroxide at the rock-water interface (Balk et al., 2009; Paudel et al., 2018). Freund (2011)
409 suggested that with the positive hole current flowing, the “corrosion” of the rock is
410 accelerated releasing into the water major cations and anions. Further work to be
411 conducted in this area may enable us to suggest the process(es) responsible for the pre-
412 earthquake geochemical anomalies we have discussed in the AYR spring water.

413

414 5. Conclusions

415

416 Hydrogeochemical precursors have been detected in commercially bottled water samples
417 of natural springs (Ayran Spring and Bahçepinar Spring) emanating from a location about
418 100 km distance from the epicenter of the Mw 7.7 Kahramanmaraş Earthquake of 6
419 February 2023. The pre-earthquake anomaly is characterized by an increase in Ca^{2+} ,
420 Mg^{2+} , K^+ , Na^+ , Cl^- , and SO_4^{2-} content in the bottled water samples of the Ayran spring.
421 Samples that are dated after the earthquakes (covering about two months after the
422 earthquake) show decreasing trends in all major ions. About three weeks after the
423 earthquake, the major ion contents of the spring water attained stability. At least six
424 months of pre-earthquake anomaly (increase) in the major ions content of the Ayran
425 spring water is imminent. It is worth noting that the Bahçepinar Spring water samples did
426 not show any anomalies in electrical conductivity therefore the samples were not
427 analyzed for ion content. Bahçepinar water is collected from shallow boreholes dug into
428 alluvial deposits which, we believe, are decoupled from the basement rocks and this may
429 be the reason for the lack of any significant change in the water chemistry prior to the
430 earthquakes. Here, we remind that geological knowledge of the investigated area and the
431 sampling site have paramount importance in sampling discrete samples for geochemical
432 analysis and/or conducting continuous monitoring. The results of this study suggest that
433 spring water chemical anomalies may be monitored as proxy indicators of pre-earthquake



434 crustal deformation. The physical mechanisms of the observed precursors are yet
435 impossible to explain with certainty at this stage. In order to be able to suggest the
436 mechanism(s) leading to the reported pre-earthquake geochemical anomalies, more work
437 needs to be conducted; especially multi-disciplinary (seismological, geodetical,
438 geochemical) and continuous earthquake monitoring networks must be established and
439 run for a sufficiently long time.

440

441 **Acknowledgements**

442 We appreciate all the technical help we have received on ion chromatography analyses
443 from Ms. Sevde Korkut at the Istanbul Technical University MEM-TEK laboratory. We
444 thank Mr. Asen Sabuncu (Istanbul Technical University) and Dr. Teach. Assist. Emre
445 Pınarcı (Çukurova University) for help in drafting the figures. We also thank Assoc. Prof
446 Dr. Tülay İnan for help in conducting electrical conductivity measurements of the bottled
447 water samples. This work has been partially supported by Istanbul Technical University
448 Scientific Research Fund (ITU BAPSIS) Project # 44774.

449 **Authors contributions**

450 S.I. and H.C. conceived the project; H.C. collected the samples; N.Y. coordinated
451 laboratory analysis, compiled seismic events, and prepared the figures; S.I. was the
452 primary interpreter of the data. S.I. and H.C. were writers of the manuscript with
453 contributions from N.Y.

454 **References**

- 455 Allen, C.R.: Active faulting in northern Turkey, Division of Geological Sciences, California
456 Institute of Technology, Contribution No. 1577. 32 pp, 1969.
- 457 Ambraseys, N.N.: Temporary seismic quiescence: SE Turkey. *Geophys. J. Int.*, 96, 311–
458 331, <https://doi.org/10.1111/j.1365-246X.1989.tb04453.x>, 1989.
- 459 Areshidze, G., Bella, F., Biagi, P. F., Caputo, M., Della Monica, G., Ermini, A.,
460 Manjgaladze, P., Melikazdze, G., Sgrigna, V. and Zilpimani, D.: No preseismic evidence
461 from hydrogeochemical parameters on the occasion of April 29, 1991 Georgian



- 462 Earthquake, *Caucasus, Tectonophysics*, 213, 353–358, [https://doi.org/10.1016/0040-1951\(92\)90463-Gö](https://doi.org/10.1016/0040-1951(92)90463-Gö) 1992.
- 463
- 464 Arpat, E., Şaroğlu, F.: Dogu Anadolu fayı ile ilgili bazı gözlem ve düşünceler (Some
465 observations and thoughts on the East Anatolian fault). *Bulletin of the General Directorate
466 of Mineral Research and Exploration of Turkey (MTA)* 73, 44–50, 1972.
- 467 Balk, M., Bose, M., Ertem, G., Rogoff, D. A., Rothschild, L. J. and Freund, F.T.: Oxidation
468 of water to hydrogen peroxide at the rock-water interface due to stress-activated electric
469 currents in rocks, *Earth Planet Sc. Lett.*, 283, 87–92.
470 <https://doi.org/10.1016/j.epsl.2009.03.044>, 2009.
- 471 Barberio, M.D., Barbieri, M., Billi, A., Doglioni, C., Petitta, M.: Hydrogeochemical changes
472 before and during the 2016 Amatrice-Norcia seismic sequence (central Italy). *Sci. Rep-
473 UK*, 7, 11735. <https://doi.org/10.1038/s41598-017-11990-8>, 2017.
- 474 Barka, A. A. and Reilinger, R.: Active tectonics of the Eastern Mediterranean region
475 deduced from GPS, neotectonic and seismicity data, *Ann. Geophys-Italy*, 40, 587-610,
476 <http://hdl.handle.net/2122/1520>, 1997.
- 477 Barka, A.A., Kadinsky-Cade, K.: Strike-slip fault geometry in Turkey and its influence on
478 earthquake activity, *Tectonics*, 7, 663–684, <https://doi.org/10.1029/TC007i003p00663>,
479 1988.
- 480 Bella, F., Biagi, P.F., Caputo, M., Cozzi, Della Monica, G., Ermini, A., Gordeez, E.I.,
481 Khatkevich, Y.M., Martinelli, G., Plastino, W., Scandone, R., Sgrigna, V., Zilpimiani, D.:
482 Hydrogeochemical anomalies in Kamchatka (Russia), *Phys. Chem. Earth.*, 23, 921-925,
483 [https://doi.org/10.1016/S0079-1946\(98\)00120-7](https://doi.org/10.1016/S0079-1946(98)00120-7), 1998.
- 484 Birchard, G.F., Libby, W.F.: Earthquake associated radon anomalies possible
485 mechanisms, *Eos Trans. AGU* 59, 4, 329, 1978.
- 486 Bozkurt, E.: Neotectonics of Turkey—a synthesis, *Geodin. Acta*, 14, 3–30,
487 <https://doi.org/10.1080/09853111.2001.11432432>, 2001.
- 488 Cetin, H., Guneyli, H., Mayer, L.: Paleoseismology of the Palu–Lake Hazar segment of
489 the East Anatolian Fault Zone, Turkey, *Tectonophysics* 374, 163–197,
490 <http://dx.doi.org/10.1016/j.tecto.2003.08.003>, 2003.
- 491 Chorowicz, J., Luxey, P., Lyberis, N., Carvalho, J., Parrot, J.F., Yurur, T. and Gundogdu,
492 N.: The Maras Triple Junction (southern Turkey) based on digital elevation model and
493 satellite imagery interpretation, *J. Geophys. Res.*, 99, B10, 20225–20242,
494 <https://doi.org/10.1029/94JB00321>, 1994.
- 495 Cicerone, R.D., Ebel, J.E., Britton, J.: A systematic compilation of earthquake precursors,
496 *Tectonophysics*, 476, 371-396. <https://doi.org/10.1016/j.tecto.2009.06.008>, 2009.



- 497 Claesson, L., Skelton, A., Graham, C., Dietl, C., Mörth, M., Torssander, P., Kockum, I.:
498 Hydrogeochemical changes before and after a major earthquake, *Geology*, 32, 641–644,
499 <https://doi.org/10.1130/G20542.1>, 2004.
- 500 Conti, L., Picozza, P., Sotgiu, A.: A critical review of ground-based observations of
501 earthquake precursors, *Front. Earth Sci.*, 9, <https://doi.org/10.3389/feart.2021.676766>,
502 2021.
- 503 Dewey, J.F. and Şengör, A.M.C.: Aegean and surrounding regions: complex multi-plate
504 and continuum tectonics in a convergent zone, *Geol. Soc. Am. Bull.*, 90, 84–92,
505 [https://doi.org/10.1130/0016-
506 7606%281979%2990%3C84%3AAASRCM%3E2.0.CO%3B2](https://doi.org/10.1130/0016-7606%281979%2990%3C84%3AAASRCM%3E2.0.CO%3B2), 1979.
- 507 Dewey, J.F., Pitman, W.C., Ryan, W.B.F., Bonnin, J.: Plate tectonics and the evolution of
508 the Alpine system. *Geol. Soc. Am. Bull.*, 84, 3137–3180, [https://doi.org/10.1130/0016-
509 7606\(1973\)84%3C3137:PTATEO%3E2.0.CO;2](https://doi.org/10.1130/0016-7606(1973)84%3C3137:PTATEO%3E2.0.CO;2), 1973.
- 510 Di Luccio, F., Chiodini, G., Caliro, S., Cardellini, C., Convertito, V., Pino, N.A., Tolomei,
511 C., Ventura, G.: Seismic signature of active intrusions in mountain chains. *Science*
512 *Advances* 4, e1701825. <https://doi.org/10.1126/sciadv.1701825>, 2018.
- 513 Dobrovolsky, I.P., Zubkov, S.I. and Miachkin, V.I.: Estimation of the size of earthquake
514 preparation zones, *Pure and Applied Geophysics* 117, 1025–1044,
515 <https://doi.org/10.1007/BF00876083>, 1979.
- 516 Doglioni, C., Barba, S., Carminati, E., Riguzzi, F.: Fault on–off versus coseismic fluids
517 reaction, *Geosci. Front.*, 5 (6), 767–780, <https://doi.org/10.1016/j.gsf.2013.08.004>, 2014.
- 518 Etiope, G., Calcara, M. and Quattrocchi, F.: Seismogeochemical algorithms for
519 earthquake prediction: a review, *Ann. Geophys-Italy*, 40(6), 1483–1492,
520 <http://hdl.handle.net/2122/1524>, 1997.
- 521 Federico C., Pizzino, L., Cinti, D., De Gregorio, S., Favara, R., Galli, G., Giudice, G.,
522 Guerrieri, S., Quattrocchi, F. and Voltattorni, N.: Inverse and forward modeling of
523 groundwater circulation in a seismically active area (Monferrato, Piedmont, NW Italy):
524 insights into stress-induced variations in water chemistry, *Chem. Geol.*, 248, 14–39,
525 <http://dx.doi.org/10.1016/j.chemgeo.2007.10.007>, 2008.
- 526 Freund, F. T.: Pre-earthquake signals: Underlying physical processes. *J. Asian Earth Sci.*,
527 41, 383–400, <https://doi.org/10.1016/j.jseaes.2010.03.009>, 2011.
- 528 Freund, F.T.: Earthquake probabilities and pre-earthquake signals, *Current Science*, 94,
529 311–313, <https://www.jstor.org/stable/24100338>, 2008.
- 530 Freund, F.T., Takeuchi, A., Lau, B.W.S.: Electric currents streaming out of stressed
531 igneous rocks – A step towards understanding pre-earthquake low frequency EM



- 532 emissions, *Phys. Chem. Earth*, 31, 389-396, <https://doi.org/10.1016/j.pce.2006.02.027>,
533 2006.
- 534 Gori, F. and Barberio, M.D.: Hydrogeochemical changes before and during the 2019
535 Benevento seismic swarm in central-southern Italy, *J. Hydrol.*, 604,
536 <https://doi.org/10.1016/j.jhydrol.2021.127250>, 2022.
- 537 Grant, R. A., Halliday, T., Balderer, W. P., Leuenberger, F., Newcomer, M., Cyr, G. and
538 Freund, F. T.: Groundwater chemistry changes before major earthquakes and possible
539 effects on animals, *Int. J. Environ. Res. PU*, 1936–1956,
540 <https://doi.org/10.3390%2Fijerph8061936>, 2011.
- 541 Geller, R. J., Jackson, D. D., Kagan, Y. Y., and Mulargia, F.: Earthquake cannot be
542 predicted, *Science*, 275, 1616–1617, <https://doi.org/10.1126/science.275.5306.1616>,
543 1997.
- 544 Hancock, P. L. and Barka, A. A.: Kinematic indicators on active normal faults in western
545 Turkey. *J. Struct Geol.*, 9, 415-430, [https://doi.org/10.1016/0191-8141%2887%2990142-
546 8](https://doi.org/10.1016/0191-8141%2887%2990142-8), 1987.
- 547 Hartmann, J. and Levy, J. K.: The influence of seismotectonic on precursory changes in
548 groundwater composition for the 1995 Kobe earthquake, Japan. *Hydrology Journal*, 14,
549 1307–1318, <https://doi.org/10.1007/s10040-006-0030-7>, 2006.
- 550 Hempton, M.R.: The North Anatolian fault and complexities of continental escape. *J. Struct*
551 *Geol.*, 4, 502 – 504, [https://doi.org/10.1016/0191-8141\(82\)90041-4](https://doi.org/10.1016/0191-8141(82)90041-4), 1982.
- 552 Hauksson, E.: Radon content of groundwater as an earthquake precursor: Evaluation of
553 worldwide data and physical basis, *J. Geophys. Res.* 86, B10, 9397-9410,
554 <https://doi.org/10.1029/JB086iB10p09397>, 1981.
- 555 İnan, S., Ergintav, S., Saatçılar, S., Tüzel, B. and İravul, Y.: Turkey makes major
556 investments in earthquake research. *EOS Transactions* 88, 333–334,
557 <https://doi.org/10.1029/2007EO340002>, 2007.
- 558 İnan, S., Akgül, T., Seyis, C., Saatçılar, R., Baykut, S., Ergintav, S. and Bas, M.:
559 Geochemical monitoring in the Marmara Region (NW Turkey): A search for precursors of
560 seismic activity, *J. Geophys. Res.* 113, B03401, <https://doi.org/10.1029/2007JB005206>,
561 2008.
- 562 İnan, S., Ertekin, K., Seyis, C., Şimşek, S., Kulak, F., Dikbaş, A., Tan, O., Ergintav, S.,
563 Çakmak, R., Yörük, A., Çergel, M., Yakan, H., Karakuş, H., Saatçılar, R., Akçiğ, Z., İravul,
564 Y. and Tüzel, B. (2010) Multi-disciplinary earthquake researches in Western Turkey: Hints
565 to select sites to study geochemical transients associated to seismicity. *Acta Geophys.*,
566 58, 767–813, <https://doi.org/10.2478/s11600-010-0016-7>, 2010.
- 567 İnan, S., Pabuçcu, Z., Kulak, F., Ergintav, S., Tatar, O., Altunel, E., Akyüz, S., Tan, O.,
568 Seyis, C., Çakmak, R., Saatçılar, R. and Eyidogan, H.: Microplate boundaries hindering ~



- 569 pre-earthquake strain transfer (Western Turkey): inferences from continuous
570 geochemical monitoring, *J. Asian Earth Sci.*, 48, 56-71,
571 <http://dx.doi.org/10.1016/j.jseaes.2011.12.016>, 2012a.
- 572 İnan, S., Kop, A., Çetin, H., Kulak, F., Pabuççu, Z., Seyis, C., Ergintav, S., Tan, O.,
573 Saatçılar, S. and Bodur, M. N.: Seasonal variations in soil radon emanation: Long term
574 continuous monitoring in light of seismicity, *Nat. Hazards*, 62, 575-591,
575 <https://doi.org/10.1007/s11069-012-0096-6>, 2012b.
- 576 İnan, S., Balderer, W.P., Leuenberger-West, F., Yakan, H., Özvan, A., Freund, F.T.:
577 Springwater chemical anomalies prior to the Mw = 7.2 Van Earthquake
578 (Turkey). *Geochem J.*, 46, e11-e16, <https://doi.org/10.2343/geochemj.1.0159>, 2012c.
- 579
580 Ingebritsen, S.E., Manga, M.: Earthquakes: hydrogeochemical precursors. *Nat. Geosci.*,
581 7, 697–698. <https://doi.org/10.1038/ngeo2261>, 2014.
- 582 Italiano, F., Martinelli, G. and Rizzo, A.: Geochemical evidence of seismogenic-induced
583 anomalies in the dissolved gases of thermal waters: A case study of Umbria (Central
584 Apennines, Italy) both during and after the 1997–1998 seismic swarm, *Geochem.*
585 *Geophys. Geosy.*, 5, Q11001, <https://doi.org/10.1029/2004GC000720>, 2004.
- 586 Jackson, J. and McKenzie, D.P.: Active tectonics of the Alpine–Himalayan Belt between
587 western Turkey and Pakistan, *Geophys. J. Roy. Astr. S.*, 77, 185–264,
588 <https://doi.org/10.1111/j.1365-246X.1984.tb01931.x>, 1984.
- 589 Karig, D.E. and Kozlu, H.: Late Palaeogene – Neogene evolution of the triple junction
590 region near Maras, south-central Turkey, *J. Geol. Soc. London*, 147, 1023–1034,
591 <https://doi.org/10.1144/gsjgs.147.6.1023>, 1990.
- 592 Kelling, G., Gökçen, S.L., Floyd, P.A. and Gökçen, N.: Neogene tectonics and plate
593 convergence in the Eastern Mediterranean: new data from southern Turkey, *Geology*, 15,
594 425–429, [https://doi.org/10.1130/0091-7613\(1987\)15%3C425:NTAPCI%3E2.0.CO;2](https://doi.org/10.1130/0091-7613(1987)15%3C425:NTAPCI%3E2.0.CO;2)
595 1987.
- 596 Kempler, D. and Garfunkel, Z.: Northeast Mediterranean triple junction from a plate
597 kinematics point of view, *Bulletin of the Technical University of Istanbul, Special Issue* 44,
598 425–454, 1991.
- 599 Ketin, I.: Über die tektonisch-mechanischen Folgerungen aus den grossen anatolischen
600 Erdbeben des letzten Dezenniums, *Geol. Rundsch.*, 36, 77–83,
601 <https://doi.org/10.1007/BF01791916>, 1948.
- 602 King, J. Y., Koizumi, Y. and Kitagawa, Y.: Hydrogeochemical anomalies and the 1995
603 Kobe Earthquake, *Science*, 269, 38–39, <https://doi.org/10.1126/science.269.5220.38>,
604 1995.



- 605 Lovelock, P.E.R.: A review of the tectonics of the northern Middle East region. *Geol. Mag.*,
606 121, 577–587, Online ISSN: 1469-5081, 1984.
- 607 Nur, A.: Matsushiro, Japan, earthquake swarm: Confirmation of the dilatancy-fluid
608 diffusion model, *Geology*, 2, 217–221, [https://doi.org/10.1130/0091-
609 7613\(1974\)2%3C217:MJESCO%3E2.0.CO;2](https://doi.org/10.1130/0091-7613(1974)2%3C217:MJESCO%3E2.0.CO;2), 1974.
- 610 McKenzie, D.P.: Active tectonics of the Mediterranean Region, *Geophys. J. Int.*, 30, 109–
611 185, <https://doi.org/10.1111/j.1365-246X.1972.tb02351.x>, 1972.
- 612 McKenzie, D.P.: The East Anatolian fault: a major structure in eastern Turkey. *Earth
613 Planet Sc. Lett.*, 29, 189–193, [https://doi.org/10.1016/0012-821X\(76\)90038-8](https://doi.org/10.1016/0012-821X(76)90038-8), 1976.
- 614 Muehlberger, W.B. and Gordon, M.B.: Observations on the complexity of the East
615 Anatolian fault, Turkey. *J. Struct Geol.*, 9, 899–903, [http://dx.doi.org/10.1016/0191-
616 8141\(87\)90091-5](http://dx.doi.org/10.1016/0191-8141(87)90091-5), 1987.
- 617 Ouzounov, D., Pulinets, S., Davidenko, D., Rozhnoi, A., Solovieva, M., Fedun, V.,
618 Dwivedi, B. N., Rybin, A., Kafatos, M., and Taylor, P.: Transient effects in atmosphere
619 and ionosphere preceding the 2015 M7.8 and M7.3 Gorkha-Nepal earthquakes, *Front.
620 Earth Sci.*, 9, <https://doi.org/10.3389/feart.2021.757358>, 2021.
- 621 Papadopoulos, G. A., Latoussakis, I., Daskalaki, E., Diakogianni, G., Fokaefs, A., Kolligri,
622 M., Liadopoulou, K., Orfanogiannaki, K., and Pirentis, A.: The East Aegean Sea strong
623 earthquake sequence of October–November 2005: lessons learned for earthquake
624 prediction from foreshocks, *Nat. Hazards Earth Syst. Sci.*, 6, 895–901,
625 <https://doi.org/10.5194/nhess-6-895-2006> , 2006.
- 626 Paudel, S.R., Banjara, S.P., Wagle, A., Freund, F.T.: Earthquake chemical precursors in
627 groundwater: a review, *J. Seismol.*, 22, 1293–1314, [https://doi.org/10.1007/s10950-018-
628 9739-8](https://doi.org/10.1007/s10950-018-9739-8), 2018.
- 629 Pérez, N. M., Hernández, P. A., Igarashi, G., Trujillo, I., Nakai, S., Sumino, H. and Wakita,
630 H.: Searching and detecting earthquake geochemical precursors in CO₂-rich
631 groundwaters from Galicia, Spain, *Geochem J.*, 42, 75–83,
632 <http://dx.doi.org/10.2343/geochemj.42.75> , 2008.
- 633 Perinçek, D. and Çemen, I.: The structural relationship between the East Anatolian and
634 Dead Sea fault zones in southeastern Turkey, *Tectonophysics*, 172, 331–340,
635 [http://dx.doi.org/10.1016/0040-1951\(90\)90039-B](http://dx.doi.org/10.1016/0040-1951(90)90039-B), 1990.
- 636 Rikitake, T.: Earthquake Precursors in Japan: Precursor Time and Detectability,
637 *Tectonophysics*, 136, 265–282, [https://doi.org/10.1016/0040-1951\(87\)90029-1](https://doi.org/10.1016/0040-1951(87)90029-1) , 1987.
- 638 Rikitake, T.: Classification of earthquake precursors, *Tectonophysics*, 54, 293-309,
639 [https://doi.org/10.1016/0040-1951\(79\)90372-X](https://doi.org/10.1016/0040-1951(79)90372-X), 1979.



- 640 Roeloffs, E.: Poroelastic techniques in the study of earthquake-related hydrologic
641 phenomena, *Adv. Geophys.*, 37, 135–195, [https://doi.org/10.1016/S0065-](https://doi.org/10.1016/S0065-2687%2808%2960270-8)
642 [2687%2808%2960270-8](https://doi.org/10.1016/S0065-2687%2808%2960270-8), 1996.
- 643 Rotstein, Y.: Counterclockwise rotation of the Anatolian block, *Tectonophysics*, 108, 71–
644 91, [https://doi.org/10.1016/0040-1951\(84\)90155-0](https://doi.org/10.1016/0040-1951(84)90155-0), 1984.
- 645 Şaroğlu, F., Emre, O., Kuscu, I.: Active Fault map of Turkey (1:2 000 000 scale), General
646 Directorate of Mineral Research and Exploration (MTA), Ankara, Turkey, 1992a
- 647 Şaroğlu, F., Emre, O., Kuscu, I.: The East Anatolian fault zone of Turkey. *Annales*
648 *Tectonicae: International Journal of Structural Geology and Tectonics*, 6, 99–125 (Special
649 Issue-Supplement to Volume VI), 1992b.
- 650 Scholz, R., Sykes, L. R. and Aggarwal, Y. P.: Earthquake prediction: A physical basis,
651 *Science*, 181, 803–810, <https://www.science.org/doi/10.1126/science.181.4102.803>,
652 1973.
- 653 Sengör, A. M. C. and Yılmaz, Y.: Tethyan evolution of Turkey: A plate tectonic approach,
654 *Tectonophysics*, 75, 181– 241, [http://dx.doi.org/10.1016/0040-1951\(81\)90275-4](http://dx.doi.org/10.1016/0040-1951(81)90275-4), 1981.
- 655 Sengör, A. M. C., Görür, N. and Şaroğlu, F.: Strike-slip faulting and related basin
656 formation in zones of tectonic escape: Turkey as a case study, in: *Strike-slip Deformation,*
657 *Basin Formation, and Sedimentation*, edited by: Biddle, K. T. and Christie-Blick, N., *Soc.*
658 *Econ. Pa*, 37, 227–264, 1985.
- 659 Seyis, C., İnan, S., Yalçın, M.N.: Major factors affecting soil radon emanation, *Nat.*
660 *Hazards*, 114, 2139–2162, <http://link.springer.com/10.1007/s11069-022-05464-y>,
661 2022.
662 .
- 663 Sibson, R. H.: Implications of fault-valve behavior for rupture nucleation and recurrence,
664 *Tectonophysics*, 211, 283–293, [https://doi.org/10.1016/0040-1951\(92\)90065-E](https://doi.org/10.1016/0040-1951(92)90065-E), 1992.
- 665 Sibson, R.H., Moore, J.M.M., Rankin, A.H.: Seismic pumping—a hydrothermal fluid
666 transport mechanism, *J. Geol. Soc. London*, 131, 653–659,
667 <https://doi.org/10.1144/gsjgs.131.6.0653>, 1975.
- 668 Skelton, A., Liljedahl-Claesson, L., Wästeby, N., Andrén, M., Stockmann, G., Sturkell, E.,
669 et al.: Hydrochemical changes before and after earthquakes based on long-term
670 measurements of multiple parameters at two sites in northern Iceland—a review, *J.*
671 *Geophys. Res-Sol EA.*, 124, 2702–2720. <https://doi.org/10.1029/2018JB016757>, 2019.
- 672 Skelton, A., Andrén, M., Kristmannsdóttir, H., Stockmann, G., Mörth, C-M.,
673 Sveinbjörnsdóttir, A., Jónsson, S., Sturkell, E., Guðrúnardóttir, H.R., Hjartarson, H.,
674 Siegmund, H., Kockum, I.: Changes in groundwater chemistry before two consecutive
675 earthquakes in Iceland. *Nat. Geosci.*, 7, 752-756. <http://dx.doi.org/10.1038/NGEO2250>,
676 2014.



- 677 Sol, S., Meltzer, A., Bürgmann, R., van der Hilst, R. D., King, R., Chen, Z., Koons, P. O.,
678 Lev, E., Liu, Y. P., Zeitler, P. K., Zhang, X., Zhang, J. and Zurek, B.: Geodynamics of the
679 southern Tibetan Plateau from seismic anisotropy and geodesy, *Geology*, 35, 563–566,
680 <https://doi.org/10.1130/G23408A.1>, 2007.
- 681 Soysal, H., Sipahioğlu, S., Kolçak, D. and Altınok, Y.: Historical earthquake catalogue of
682 Turkey and surrounding area (2100 B.C. – 1900 A.D.), The Scientific and Technological
683 Research Council of Turkey Technical Report, TBAG-341, 1981.
- 684 Sugisaki, R., Ito, T., Nagamine, K. and Kawabe, I.: Gas geochemical changes at mineral
685 springs associated with the 1995 southern Hyogo earthquake (M = 7.2), Japan, *Earth
686 Planet Sc. Lett.*, 139, 239–249, [https://doi.org/10.1016/0012-821X\(96\)00007-6](https://doi.org/10.1016/0012-821X(96)00007-6), 1996.
- 687 Sultankhodhaev, G. A.: International symposium on earthquake prediction, UNESCO,
688 Paris 1979, 181-191, 1984.
- 689 Tansi, C., Tallarico, A., Iovine, G., Folino Gallo, M. and Falcone, G.: Interpretation of
690 radon anomalies in seismotectonic and tectonic-gravitational settings: the southeastern
691 Crati graben (Northern Calabria, Italy), *Tectonophysics*, 396, 181–193,
692 <https://doi.org/10.1016/j.tecto.2004.11.008>, 2005.
- 693 Thomas, D.M., Cuff, K.E., Cox, M.E.: The association between ground gas radon
694 variations and geologic activity in Hawaii, *J. Geophys. Res-Sol EA*, 91, 12186-12198,
695 [https://ui.adsabs.harvard.edu/link_gateway/1986JGR....9112186T/doi:10.1029/JB091iB
696 12p12186](https://ui.adsabs.harvard.edu/link_gateway/1986JGR....9112186T/doi:10.1029/JB091iB12p12186). 1986.
- 697 Toutain, J. P., Munoz, M., Poitrasson, F. and Lienard, F.: Springwater chloride ion
698 anomaly prior to a ML = 5.2 Pyrenean earthquake, *Earth Planet Sc. Lett.*, 149, 113–119.
699 [https://doi.org/10.1016/S0012-821X\(97\)00066-6](https://doi.org/10.1016/S0012-821X(97)00066-6), 1997.
- 700 Tsunogai, U. and Wakita, H.: Precursory chemical changes in groundwater: Kobe
701 earthquake, Japan, *Science*, 269, 61-63, <https://doi.org/10.1126/science.269.5220.61>,
702 1995.
- 703 Turcotte, W. L.: Earthquake prediction, *Annu. Rev. Earth Pl. Sc.*, 19, 263 – 281,
704 <https://doi.org/10.1146/annurev.ea.19.050191.001403>, 1991.
- 705 Usta, D., Ateş, Ş., Beyazpınar, M., Kanar, F., Yıldız, H., Uçar, L., İsmail Akça, İ., Tufan,
706 E. and Örtlek, A. T.: New data on the stratigraphy of the central and northern Amonous
707 mountains (Osmaniye - Gaziantep - K. Maraş). *Bulletin of the Turkish Association of
708 Petroleum Geologists*, 27, 57-98, 2015.
- 709 Usta, D., Ateş, Ş., Kanar, F., Beyazpınar, M., Uçar, L., Yıldız, H., Tufan, E., Akça, İ.,
710 Örtlek, A. T.: Doğu Toroslar'ın jeolojisi ve jeodinamik evrimi projesi (K.Maras, Osmaniye,
711 Gaziantep, Adana, Hatay). General Directorate of Minerals and Exploration of Turkey
712 (MTA) unpublished Report No 13568, 494 pages, Ankara (in Turkish), 2017.



- 713 Uyeda, S., Nagao, T. and Kamogawa, M.: Short-term earthquake prediction: Current
714 status of seismo-electromagnetics, *Tectonophysics*, 470, 205–213.
715 <https://doi.org/10.1016/j.tecto.2008.07.019>, 2008.
- 716 Wakita, H.: Geochemical challenge to earthquake prediction, *P. Natl. Acad. Sci. USA*, 93,
717 9, 3781-3786, DOI: 10.1073/pnas.93.9.3781, 1996.
- 718 Wakita, H., Nakamura, Y. and Sano, Y.: Short-term and intermediate-term geochemical
719 precursors, *Pure and Applied Geophysics*, 125, 267–278,
720 <https://doi.org/10.1007/BF00878999>, 1988.
- 721 Wang, C-Y. and Manga, M.: *Water and Earthquakes. Lecture Notes in Earth System*
722 *Sciences*. Springer, 307 pp. <https://doi.org/10.1007/978-3-030-64308-9>. 2021.
- 723 Westaway, R. and Arger, J.: The Gölbasi basin, southeastern Turkey: a complex
724 discontinuity in a major strike-slip fault zone, *J. Geol. Soc. London*, 153, 729 – 744,
725 <https://doi.org/10.1144/gsjgs.153.5.0729>, 1996.
- 726 Xiang, Y. and Peng, S.: Hydrochemical and stable isotopes ($\delta^2\text{H}$ and $\delta^{18}\text{O}$) changes
727 of groundwater from a spring induced by local earthquakes, Northwest China. *Frontiers*
728 *in Earth Sciences*, 09 January 2023. *Solid Earth Geophysics*.
729 <https://doi.org/10.3389/feart.2022.1100068>, 2023.
- 730 Virk, H.S., and Singh, B.: Radon anomalies in soil gas and groundwater as earthquake
731 precursor phenomena, *Tectonophysics*, 227, 215-224, [https://doi.org/10.1016/0040-](https://doi.org/10.1016/0040-1951(93)90096-3)
732 [1951\(93\)90096-3](https://doi.org/10.1016/0040-1951(93)90096-3), 1993.
- 733 Yönlü, Ö., Altunel, E. and Karabacak, V.: Geological and geomorphological evidence for
734 the southwestern extension of the East Anatolian Fault Zone, Turkey. *Earth Planet Sc.*
735 *Lett.*, 469, 1–14, <http://dx.doi.org/10.1016/j.epsl.2017.03.034>, 2017.
- 736 Yu, H., Liu, L., Ma, Y., Yan, R., Liu, J., Ma, Y., Li, Z., Zhang, X., Zhao, J., Yu, C.: Observed
737 hydrological changes associated with active tectonic blocks before three consecutive
738 earthquakes in Qinghai, China *Scientific Reports* 13, 8988.
739 <https://doi.org/10.1038/s41598-023-36274-2>, 2023.
- 740 Zeyrek, M., Ertekin, K., Kacmaz, S., Seyis, C., İnan, S.: An ion chromatography method
741 for the determination of major anions in geothermal water samples, *Geostand. Geoanal.*
742 *Res.*, 34, 67-77, <http://dx.doi.org/10.1111/j.1751-908X.2009.00020.x>, 2010.
743