

1 **Spring water anomalies before two consecutive earthquakes (Mw 7.7 and Mw 7.6)**
2 **in Kahramanmaraş (Türkiye) on 6 February 2023**

3
4 Sedat İnan^{1,*}, Hasan Çetin², Nurettin Yakupoğlu¹

5
6 ¹Department of Geological Engineering, Istanbul Technical University, Ayazağa,
7 Istanbul, Türkiye, 34467

8 ²Department of Geology, Çukurova University, Adana, Türkiye, 01330

9 *Corresponding Author: sedatinan@itu.edu.tr

10 **ABSTRACT**

11 Understanding earthquake phenomena is always challenging. Search for reliable
12 precursors of earthquakes are important but requires systematic and long-time monitoring
13 employing multi-disciplinary techniques. In search of possible precursors, we obtained
14 commercially bottled spring waters dated before and after the earthquakes of 6 February
15 2023. Hydrogeochemical precursors have been detected in commercially bottled natural
16 spring waters (Ayran Spring and Bahçepınar Spring) which are at a distance of about 100
17 km and 175 km from the epicenters of the Mw 7.7 and Mw 7.6 Kahramanmaraş (Türkiye)
18 Earthquakes of 6 February 2023, respectively. The available water samples cover the
19 period from March 2022 to March 2023. The pre-earthquake anomaly is characterized by
20 an increase in electrical conductivity and major ions (Ca^{2+} , Mg^{2+} , K^+ , Na^+ , Cl^- , and SO_4^{2-}
21) compared to the background for Ayran Spring water samples. The pre-earthquake
22 anomaly lasted for at least six months. The anomaly in major ions sharply declined and
23 the ion content approached the background values about two weeks after the
24 earthquakes. Although only 6.5 kilometers away from the Ayran Spring, the bottled water
25 samples of the Bahçepınar Spring did not show any anomalies in electrical conductivity;
26 therefore, the samples were not analyzed for ion content. Bahçepınar water is collected
27 from shallow boreholes dug into alluvial deposits which we believe are decoupled from
28 the basement rocks and this may be the reason for the lack of abnormal water chemistry
29 prior to the earthquakes. This attests to the fact that sampling locations are very important
30 in the detection of possible earthquake precursors. Results of the Ayran spring water
31 samples indicate that spring water chemical anomalies of discrete samples may provide

32 valuable information on pre-earthquake crustal deformation. Monitoring of spring waters,
33 along with other monitoring techniques in a multidisciplinary network, and for a sufficiently
34 long time, could potentially enable obtaining reliable proxy indicators of pre-earthquake
35 crustal deformation.

36

37 **Keywords:** geochemical anomalies, spring water, earthquake precursors,
38 Kahramanmaraş earthquakes, East Anatolian Fault Zone (EAFZ), Türkiye

39

40 **1. Introduction**

41 Two devastating earthquakes (Mw 7.7 and Mw 7.6) struck the Kahramanmaraş area in
42 Southern Turkey on 6 February 2023; the earthquakes occurred about 9 hours apart. The
43 earthquakes caused devastation causing more than 50,000 deaths; leaving behind
44 thousands injured and/or homeless. Earthquakes of destructive magnitudes (e.g., $M > 7$)
45 are naturally expected to occur at plate boundary settings (Figure 1) and Kahramanmaras
46 province is at the junction of the East Anatolian Fault System (EAFS) and the Dead Sea
47 Fault System (DAFS). However, the reason why such natural events turn into disasters
48 is mainly due to a lack of preparedness. Where buildings are not built to be sufficiently
49 earthquake-resistant, monitoring of crustal deformation and searching for reliable pre-
50 earthquake signals become more important. This is obviously a big challenge for earth
51 scientists to overcome. Although there is still a long way to go on this front, the scientific
52 literature is full of scattered but promising and encouraging cases.

53 Earthquakes are complex natural phenomena and their predictions have been long
54 viewed as difficult, if not impossible (e.g., Geller et al., 1997). Geochemical observations
55 to identify earthquake precursors were initiated in the late 1960s (Rikitake, 1979; Wakita
56 1996). Reviewing twenty years of relevant data Turcotte (1991) concluded that large
57 earthquakes are not preceded by reliable seismic precursors. Moreover, Geller et al.
58 (1997) claimed that earthquakes can never be predicted. However, for the last few
59 decades, there have been numerous reports of ground-based anomalies (e.g., changes
60 in soil gas, water chemistry and ground water level, rock deformation detected by
61 tiltmeters, electrical and electromagnetic fields) preceding major earthquakes. (including

62 but not limited to Rikitake, 1979; Dobrovolsky et al., 1979; Birchard and Libby, 1978;
63 Hauksson, 1981; Wakita et al., 1988; Sultankhodhaev, 1984; Thomas et al., 1986;
64 Rikitake, 1987; Martinelli, and Ferrari, 1991; Etiope et al., 1997; Vallianatos and Tzanis,
65 1998; Bella et al., 1998; Virk and Singh, 1993; King et al., 1995; Planinic et al, 2004;
66 Claesson et al., 2004; Vallianatos et al., 2004; Hartmann and Levy, 2006; Papadopoulos
67 et al., 2006; Uyeda et al., 2008; İnan et al., 2008; Walia et al., 2009; İnan et al., 2010;
68 İnan et al, 2012a,b,c; Skelton et al., 2014 and 2019; Barberio et al., 2017; He and Singh
69 2019; Ouzounov et al., 2021; Gori and Barberio, 2022; Xiang and Peng, 2023). Compiling
70 a review of claimed precursors, Cicerone et al. (2009) conducted a survey of published
71 scientific literature on earthquake precursors and concluded that precursory anomalies
72 seem to be recorded where there is modern instrumentation. İnan et al. (2010 and 2012a)
73 provided hints to select monitoring sites. Recently, Conti et al. (2021) have provided a
74 short review of ground-based observations before earthquakes

75 Hydro-geochemical anomalies observed prior to seismic events are generally interpreted
76 to be related to the alteration of the groundwater circulating system by the changes in the
77 crustal stress/strain before earthquakes and mixing of different aquifers (e.g., Scholz et
78 al., 1973; Nur, 1974; Sibson et al., 1975; Sugisaki et al., 1996; Tsunogai and Wakita,
79 1995; Toutain et al., 1997; Claesson et al., 2004; Pérez et al., 2008; İnan et al., 2010;
80 Grant et al., 2011; İnan et al., 2012c; Doglioni et al., 2014; Ingebritsen and Manga, 2014;
81 Skelton et al., 2014 and 2019; Barberio et al., 2017; Gori and Barberio, 2022; Xiang and
82 Peng, 2023). However, another different approach based on “stress-activated positive
83 hole currents” has been suggested to play a role in the development of physicochemical
84 pre-earthquake stress indicators (Freund, 2011; Paudel et al., 2018)


85 As suggested by Nur (1974) and later by Rikitake (1987) precursory phenomena may
86 have a common physical basis which Scholz et al. (1973) called the “Dilatation and water
87 diffusion (DWD) model”. Roeloffs (1996) noted that with respect to earthquake hydrology,
88 mechanical and fluid-dynamic effects can be modeled using poroelasticity. More recently,
89 the DWD model has been explained further (e.g., Doglioni et al., 2014; Wang and Manga,
90 2021). However, other authors have proposed a fundamentally different approach
91 (Freund et al., 2006; Freund, 2008; Freund, 2011; Paudel et al., 2018) to study and
92 evaluate physicochemical pre-earthquake stress indicators. Until the mechanism

93 controlling pre-earthquake processes is fully understood, it is worth noting that the
94 success of any pre-earthquake stress indicators may be compromised by the ever-
95 present crustal heterogeneity, anisotropy, and/or crustal blocks (Areshidze et al., 1992;
96 Tansi et al., 2005; Sol et al., 2007; İnan et al., 2012a; Yu et al., 2023). Microplate and/or
97 block boundaries are obstacles to pre-earthquake strain to transfer from one block to the
98 other (İnan et al., 2012a; Yu et al., 2023).

99

100 A multi-disciplinary earthquake observation network (GPS, seismology, soil radon, and
101 spring water monitoring stations) was established in Kahramanmaraş and surrounding
102 provinces along the fault zones (Adana, Hatay, Malatya, Elazığ, Bingöl) in 2007 under the
103 scope of the TURDEP Project (İnan et al. 2007). In the Kahramanmaraş area, due to
104 seismic gap interpretations, also borehole tilt monitoring stations were established.
105 Continuous monitoring was continued until the middle of 2012 and valuable multi-
106 disciplinary data were collected. However, throughout these five years, no earthquake of
107 significant magnitude (e.g. $M > 6$) occurred to test the usefulness of the monitoring
108 network, the project was terminated by the funding organization due mainly to a lack of
109 vision. As a result, the earth science community was caught unprepared when two
110 consecutive devastating earthquakes struck the area on 6 February 2023. No ground
111 (except GPS and seismology) monitoring station data were available to detect possible
112 pre-earthquake anomalies. However, following the Mw 7.7 and Mw 7.6 Kahramanmaraş
113 earthquakes, we searched for bottled spring waters to analyze in search of possible pre-
114 earthquake anomalies. This proved difficult as the water supply to the large community
115 affected by the earthquakes was quite limited and businesses providing bottled spring
116 waters were also mostly shut down. Finally, we were able to obtain commercially bottled
117 water samples (dated before and after the earthquake) from the Ayran and Bahçepınar
118 springs which are located within about a 6.5-kilometer distance in the Osmaniye Province.
119 The spring waters are about 100 kilometers and 175 kilometers from the epicenter of the
120 first (Mw 7.7) and the second (Mw 7.6) earthquakes, respectively (Figure 1B). In this
121 study, we conducted electrical conductivity (E_c) measurements on bottled waters, and
122 based on the E_c results, we selected samples for analysis of major ions in water in search

123 of pre-earthquake anomalies. The spring water samples cover the range from March 2022
 124 to March 2023.



125
 126 **Figure 1. A)** Neotectonics map of the Türkiye and surroundings (compiled from Sengör
 127 and Yılmaz, 1981; Sengör et al., 1985; Hancock and Barka, 1987; Şaroğlu et al., 1992;
 128 Barka and Reilinger, 1997; Bozkurt, 2001). **B)** Active fault map of the region affected by
 129 the February 2023 Earthquakes (Perinçek and Çemen, 1990; Şaroğlu et al., 1992a; Cetin
 130 et al., 2003). Red stars show the epicenters of the Mw 7.7 and 7.6 Kahramanmaraş
 131 Earthquakes of 6 February 2023, and Mw 6.4 Hatay Earthquake of 20 February 2023.
 132 Filled red circles show the locations of the earthquakes (Mw>4.5) that occurred in the
 133 area (circle area with a radius of 150 km from the location of the water spring) between
 134 September 2022 and 5 February 2023. Details of the earthquakes are given in Table 1.

135 TTJ is Türkoğlu Triple Junction. Beach balls are fault plane solutions of earthquakes and
136 were obtained from the Bogazici University Kandilli Observatory and Earthquake
137 Research Institute (KOERI) of Turkey; www.koeri.edu.tr

138 **2. Active tectonics of the Kahramanmaraş region**

139 Kahramanmaraş region is located in the suture zone formed by the collision between
140 Arabian and Anatolian plates during Late Miocene-Pliocene (Figure 1A). After this
141 collision, very important strike-slip fault zones were developed in the Anatolian plate due
142 to the continuous northward movement of the Arabian plate and the resulting westward
143 movement or escape of the Anatolian plate along two major fault zones, the North
144 Anatolian Fault Zone (NAFZ) and the East Anatolian Fault Zone (EAFZ) (Ketin, 1948;
145 McKenzie, 1972; Dewey and Şengör, 1979; Şengör and Yılmaz, 1981; Hempton, 1982;
146 Şengör et al., 1985).

147 The East Anatolian Fault Zone (EAFZ) is approximately 550 km long, northeast-
148 southwest trending, sinistral strike-slip fault (Figure 1A). It was first described by Allen
149 (1969) and mapped by Arpat and Şaroğlu (1972). The EAFZ starts from Karlıova Triple
150 Junction in the northeast, and it runs in the southwest direction, passes near the east-
151 southeast of Kahramanmaraş, and joins another triple junction at Türkoğlu (TTJ in Figure
152 1B). The EAFZ then continues to the Hatay in the south direction to merge into the Dead
153 Sea Fault Zone (DSFZ) (Allen, 1969; Arpat and Şaroğlu, 1972; Dewey and Şengör, 1979;
154 Rotstein, 1984; Şengör et al., 1985; Kelling et al., 1987; Şaroğlu et al., 1992a and 1992b;
155 Cetin et al., 2003; Yönlü et al., 2017). There are different interpretations, however, for the
156 remainder of the fault zone after Türkoğlu Triple Junction (marked as TTJ in Figure 1B).
157 Some studies extend the fault zone southwesterly to the Mediterranean Sea (McKenzie,
158 1972; Dewey et al., 1973; Jackson and McKenzie, 1984; Barka and Kadinsky-Cade,
159 1988; Karig and Kozlu, 1990; Kempler and Garfunkel, 1991; Westaway and Arger, 1996),
160 joining it with the Cyprian Arc along which the convergence is taking place between the
161 African and Anatolian plates (McKenzie, 1976; Dewey and Şengör, 1979). Others think
162 that the fault zone ends around the TTJ (Lovelock, 1984; Chorowicz et al., 1994).
163 According to Muehlberger and Gordon (1987), the EAFZ becomes the northern branch
164 of the DSFZ.

165 The seismicity of the study area is controlled by a complex interaction of the African,
 166 Arabian, and Eurasian plates (McKenzie, 1972). The seismicity of the EAFZ has been
 167 minimal for most of the last 100 years (Ambraseys, 1989). Historical earthquake records
 168 show that Kahramanmaraş and its surroundings were affected by two major earthquakes
 169 in AD 1114 and AD 1513 (Soysal et al., 1981; Ambraseys, 1989). There had been a long
 170 quiescence of more than 500 years in the Kahramanmaraş area before the Mw 7.7 and
 171 Mw 7.6 earthquakes struck on 6 February 2023. About one year before these
 172 earthquakes occurred, the area had been seismically quiet as suggested by only a few
 173 M>4.5 earthquakes occurring in a circular area with a radius of 150 km; taking the Ayran
 174 spring water as the center (Figure 1B and Table 1). The fault plane solutions (FPS) for
 175 earthquakes #3, #4, and #5 suggest mainly normal faulting, whereas, for others
 176 (earthquakes #1, #2, and #6), FPS suggest movement on dominantly left lateral strike-
 177 slip faults (Figure 1B) as expected for left-lateral strike-slip nature of the EAFZ.

178 **Table 1.** Earthquakes' time, magnitude, and locations as received from www.koeri.edu.tr.
 179 Earthquakes #1, #2, and #3 are the earthquakes of February 2023. Earthquakes #4, #5,
 180 and #6 are those that have occurred in the circular area (with a radius of 150 km from the
 181 Ayran spring water location) between September 1st, 2022 and 5 February 2023. The
 182 locations of these earthquakes are given on the map (Figure 1B).

Earthquake #	Magnitude (Mw)	Date	Time (GMT)	Latitude	longitude
1	7.7	06.02.2023	01:17	37.1757	37.0850
2	7.6	06.02.2023	10:24	38.0818	37.1773
3	6.4	20.02.2023	17:04	36.0713	36.1012
4	4.6	12.01.2023	20:40	35.5712	36.6723
5	4.9	18.12.2022	18:13	36.3978	36.4455
6	5.0	11.10.2022	15:48	37.3025	36.2403

183

184 3. Samples and methods

185

186 3.1. Spring water samples

187 The spring water samples were received in commercial polyethylene bottles and brought
 188 to Istanbul Technical University Laboratory for electrical conductivity measurements and
 189 major ion analyses. Some of the samples had been bottled up to several months before
 190 the analyses. However, this does not create any concern because much longer storage

191 in this kind of bottle has been reported to be appropriate in terms of keeping reliable
192 concentrations (Tsunogai and Wakita, 1995; İnan et al., 2012c; Rapti et al., 2023).

193 The spring water samples cover the range from March 2022 to March 2023. It is worth
194 noting that the oldest sample predating the earthquakes was AYR 1 (dated 8 March 2022)
195 from the Ayran Spring. Other bottled water samples we could obtain from both springs
196 were dated between September 2022 and March 2023. In fact, we could not obtain any
197 samples dated between 8 March and 14 September 2022. The samples from September,
198 October, and November 2022 are limited but from December 2022 to January 2023,
199 available samples are several per month (Table 2).

200 **3.2. Spring water analysis**

201 We first screened the bottled water samples by conducting electrical conductivity (Ec)
202 measurements, and based on the results, we selected samples for analysis of major ions.

203 Samples of the AYR spring water were analyzed by ion chromatography as discussed by
204 Zeyrek et al. (2010). Briefly, the samples were filtered at 0.45 μm and split into two
205 portions before analysis using an ion chromatography instrument (Dionex ICS 1000).
206 Sodium carbonate and methane-sulfonic acid were used as eluents for anion and cation
207 analyses, respectively. For calibration, DIONEX Certified Reference Standards were
208 used. Deionized water with a resistance better than 18.2 Megaohm was used for the
209 preparation of all eluents. Repeated measurements ascertained that the analytical
210 uncertainties for all anions and cations were below 5%. Electrical conductivity (Ec)
211 measurements for the bottled Bahçepinar (BPN) spring water samples and both Ec and
212 Ion analysis results for the bottled Ayran spring waters are listed in Table 2.

213 **3.3. Statistical analysis of the data**

214 For the statistical treatment of the data on major ion contents of the water samples, we
215 calculated the weighted average (weighted compared to the analytical error for each
216 point) and computed the 2σ external error ($2 \times \alpha e$) from the following equation

217

$$\alpha_e = \frac{\sum_{i=1}^n (x_i - \bar{x})^2 / \sigma_i^2}{(n-1) \sum_{i=1}^n 1 / \sigma_i^2}$$

218

219 where \bar{x} is the average and σ the analytical error on each measured point. The 2σ external
 220 error (α_e) considers the general variability of all datasets and the analytical error on each
 221 point; thus, we obtained the total error envelope for the samples that we consider
 222 representing background (from 15 February to 31 March 2023; see Table 2 and Figure
 223 3B).

224

225 **3.4.Relation between earthquake magnitude, distance, and precursory duration**

226 Slightly different relations between earthquake magnitude, duration of a precursory
 227 anomaly, and the distance of the monitoring site to the earthquake epicenter have been
 228 proposed. Dobrovolsky et al. (1979) proposed a theoretical relation between earthquake
 229 magnitude and maximum epicentral distance at which geochemical anomalies may be
 230 observed.

$$231 \quad D = 10^{0.43 * M} \quad (1)$$

232 Relation 1 assumes a homogenous and isotropic crust. Where M is the earthquake
 233 magnitude and D is the distance in kilometers to the earthquake epicenter (Dobrovolsky
 234 et al., 1979). Rikitake (1987) noted a slightly different relation;

$$235 \quad \log T = a + b * M \quad (2)$$

236 where a and b are constants, T is the duration of anomaly and M is the magnitude of an
 237 earthquake (Rikitake, 1987). Moreover, Sultankhodhaev (1984) also reported a relation,
 238 between earthquake magnitude, the distance of the monitoring site to the earthquake
 239 epicenter, and duration of precursory anomaly

$$240 \quad \log (DT) = 0.63 * M - b \quad (3)$$

241 where D is the distance in km, T is the duration of a precursory anomaly in days, and M
 242 is earthquake magnitude; b is a constant taken as 0.15 Sultankhodhaev (1984).

243 All of these three relations provide a helpful initial idea about what to expect of precursory
244 anomalies in terms of duration and distance to the earthquake epicenter. [The closer the](#)
245 [epicenter of an earthquake of a given magnitude, the longer duration of anomaly at](#)
246 [monitoring site.](#)

247 Obviously, we should keep in mind that the relations proposed by Dobrovolsky et al.
248 (1979), Sultankhodhaev (1984), and Rikitake (1987) assume homogenous and isotropic
249 crust where pre-earthquake stress and resultant strain propagates in all directions. In
250 fact, we know that this assumption is not totally correct as microplates and/or block
251 boundaries hinder stress transfer (e.g., Inan et al. 2012b). This issue should be seriously
252 considered and care should be exercised.

253 Inan et al. (2008 and 2010) verified Dobrovolsky et al.'s (1979) relation for medium-size
254 earthquakes ($M < 5.3$). Accordingly, for an earthquake of magnitude 4.5, the maximum
255 distance for detection of possible geochemical anomalies in the Ayran Spring water will
256 be about 100 km. For contingency, we took a 150 km radius and listed in Table 1 the
257 earthquakes with $M > 4.5$ occurring between September 2022 and 5 February 2023 in
258 order to compare with the water geochemical data we obtained in this study.


259

260 **4. Results and Discussion**

261

262 The bottled water samples from the Bahcepinar (BPN) Spring did not show any
263 meaningful (e.g., significant) variations in electrical conductivity (Ec) values; varying in a
264 narrow range between 220 and 230 microsiemens/cm (Table 2). Therefore, these
265 samples were not analyzed for major ions content because change (increase/decrease)
266 of major ions contents is expected to result in Ec variation (Inan et al., 2010; Inan et al.,
267 2012c). However, the bottled water samples from the Ayran (AYR) spring showed major
268 variations in the Ec values; varying in range between 50 and 200 microsiemens/cm.
269 Therefore, the AYR samples were analyzed for major ions. Possible reasons for not
270 detecting any anomaly in the Ec measurements of the BNP spring water samples have
271 been investigated. The investigation suggests that the reason may be the geological

272 environment of the springs. The AYR spring water emanates from Middle-Upper
273 Ordovician age metamorphic rocks (Kardere Formation) made up of quartzite,
274 metasediments, metasiltstone, and metashale (Usta et al., 2015 and 2017), whereas the
275 BPN spring water is collected from shallow boreholes dug into valley-filling Quaternary
276 age alluvial deposits that are underlain by ophiolite (Figure 2). The alluvial deposits reach
277 a thickness of about two hundred meters and the water reservoir within the alluvium
278 deposit is fed by precipitation and a nearby Bekdemir stream flowing towards the alluvial
279 deposit. It is interesting that the streams disappear to the south; suggesting that the
280 stream (creek) water is captured by the alluvial deposit. Since the BPN water is collected
281 from shallow boreholes (less than 100 meters) dug into alluvial deposits, we believe that
282 the alluvial deposits are decoupled from the basement rocks (which undergo pre-
283 earthquake stress) and this may be the reason for the lack of anomaly in water chemistry
284 prior to the earthquakes. This has testified again to the importance of adequate geological
285 knowledge of the area before sampling discrete geochemical samples (water or soil gas)
286 and/or continuous monitoring in search of pre-earthquake signals (İnan et al., 2008; İnan
287 et al., 2010).



288

289 **Figure 2.** Locations and local geology of the water springs. (Modified from Usta et al.,
 290 2015 and 2017). The Ayran spring water emanates from a fault in the Metamorphic
 291 Kardere Formation (blue triangle shown at the lower right in the map) whereas the
 292 Bahçepınar spring water is obtained from the Quaternary Alluvium (blue triangle shown
 293 at the upper left in the map).

294

295 Variations of major ions in the AYR spring water samples were significant. It is clear that
 296 pre-earthquake anomalies have exceeded the α_e (Figures 3B and Table 2). Before any
 297 interpretation, we needed to make sure that geochemical time series are not affected by
 298 meteorological conditions. In this context, meteorological data were obtained from the

299 Osmaniye Meteorology Station (located about 32 km SW of the AYR spring) and the daily
 300 average air temperature and rainfall are shown in Figure 3c.

301 **Table 2.** Ec and major ion analysis results for the Ayran (AYR) and the Ec analysis results
 302 for the Bahçepınar (BPN) bottled waters. The data for Ca²⁺, Mg²⁺, K⁺, Na⁺, Cl⁻, SO₄²⁻ for
 303 the AYR samples are plotted in Figure 3B. Standard deviation (2σ) has been computed
 304 considering cations/anions contents of samples dated from 15 February to 31 March
 305 2023; the period which is considered to nearly represent background concentrations of
 306 the water. These samples are marked in bold fonts.

Sample ID	Date	Cl ⁻	SO ₄ ²⁻	Na ⁺	K ⁺	Mg ²⁺	Ca ²⁺	AYR Ec	Date	BPN Ec
AYR 1	08.03.2022	2.99	8.34	4.34	0.39	3.22	6.92	50	19.09.2022	220
AYR 2	14.09.2022	7.37	13.08	12.10	1.08	7.73	17.54	150	07.11.2022	230
AYR 3	06.10.2022	9.73	14.79	15.08	1.34	9.20	20.10	180	12.12.2022	230
AYR 4	03.11.2022	9.99	15.52	15.66	1.39	9.50	20.72	170	19.12.2022	220
AYR 5	13.12.2022	7.45	13.43	11.93	1.05	7.59	16.58	150	30.12.2022	220
AYR 6	26.12.2022	11.06	17.35	16.49	1.56	10.19	22.19	190	08.01.2023	220
AYR 7	29.12.2022	11.08	17.20	16.84	1.50	10.20	22.33	180	20.01.2023	220
AYR 8	30.12.2022	10.97	17.29	16.78	1.50	10.17	22.28	190	24.01.2023	220
AYR 9	03.01.2023	10.62	17.23	16.26	1.45	10.06	23.04	170	28.01.2023	220
AYR 10	06.01.2023	11.12	17.56	16.91	1.49	10.29	22.51	190	04.02.2023	220
AYR 11	11.01.2023	11.41	17.96	16.90	1.50	10.43	22.81	190	11.02.2023	220
AYR 12	12.01.2023	11.60	18.21	17.22	1.53	10.50	22.99	200	17.02.2023	220
AYR 13	27.01.2023	9.83	16.20	14.24	1.25	8.89	19.35	160	18.02.2023	230
AYR 14	31.01.2023	11.04	17.62	15.81	1.39	9.87	21.58	180	02.03.2023	230
AYR 15	01.02.2023	11.43	17.85	16.21	1.43	10.04	21.97	190	13.03.2023	220
AYR 16	10.02.2023	9.09	15.59	13.29	1.16	8.46	18.33	180	22.03.2023	230
AYR 17	12.02.2023	6.00	12.47	9.36	0.79	6.29	13.51	120		
AYR 18	13.02.2023	4.25	10.69	6.96	0.56	4.95	10.38	90		
AYR 19	15.02.2023	3.54	10.30	6.28	0.50	4.79	9.65	80		
AYR 20	16.02.2023	3.56	13.64	7.51	0.67	6.91	12.60	110		
AYR 21	28.02.2023	3.29	10.54	5.79	0.47	4.60	9.23	80		
AYR 22	02.03.2023	3.26	10.08	5.48	0.44	4.20	8.59	70		
AYR 23	11.03.2023	3.36	9.85	5.49	0.43	4.21	8.62	70		
AYR 24	13.03.2023	3.28	9.91	5.47	0.44	4.22	8.68	70		
AYR 25	20.03.2023	3.28	9.96	5.36	0.43	4.22	8.73	70		
AYR 26	24.03.2023	3.20	10.02	5.35	0.42	4.14	8.45	70		
AYR 27	31.03.2023	3.31	10.13	5.40	0.43	4.16	8.47	70		
	<i>mean</i>	0.37	1.13	0.60	0.05	0.46	0.94	7.78		
	Σ	0.33	1.19	0.79	0.08	0.88	1.34	13.64		
	2σ	0.65	2.38	1.57	0.17	1.76	2.68	27.28		
	<i>mean +1 σ</i>	3.64	11.32	6.19	0.51	5.04	9.81	83.64		
	<i>mean +2 σ</i>	2.99	8.94	4.62	0.35	3.28	7.13	56.36		

307
 308 Air temperature gradually decreases from about 30°C in September 2022 to less than
 309 10°C in February 2023 (Figure 3C). Daily rainfall is noticeably present in November 2022
 310 and March 2023. Normally, variations in air temperature are not expected to affect the
 311 chemical contents of the spring water (İnan et al., 2010 and 2012) but the effect of rainfall

312 on soil radon concentration is dominant (Inan et al., 2008, 2010, 2012b, Seyis et al.,
313 2022). All earthquakes listed in Table 1 are plotted on the meteorology time series in
314 Figure 3C and this shows that major and heavy rainfall took place right after the
315 devastating earthquakes of 6 February 2023. Based on the relatively low EC and low
316 major ion contents of the AYR spring water (Table 2) that is bottled and commercially
317 distributed, it can be said that this water is of shallow origin (Di Luccio et al., 2018). A
318 comparison of the geochemical time series and significant variations shown in Figure 3B
319 and the daily average rainfall data shown in Figure 3C reveals no correlations. Inan et al.
320 (2010 and 2012) compared meteorological time series with hydrogeochemical time series
321 and noted that meteorological conditions do not seem to play a role in water's major ion
322 contents. In this study, we compare rainfall data and geochemical time series (Figure 3)
323 and, as there is no correlation, we conclude that the increase of major ion contents
324 observed in AYR spring waters are not related to atmospheric variations (e.g., rainfall).
325 Therefore, it is safe to conclude that the chemical changes recorded in the spring water
326 must be related to crustal deformation associated with earthquake stress buildup.

327 As shown in Figure 3B, changes in the concentration of the major ionic species dissolved
328 in the AYR spring water were observed. Positive anomalies are recorded in the Ca^{2+} ,
329 Mg^{2+} , K^+ , Na^+ , Cl^- , and SO_4^{2-} contents (mg/l) before the 6 February Mw 7.7 and 7.6
330 Kahramanmaraş Earthquakes (Figure 3b; Table 2). These positive anomalies (increase
331 in dissolved ion content) started as early as September 2022; suggesting a pre-
332 earthquake anomaly of nearly six months. Considering Sultankhodhaev's (1984) relation
333 ($\log(\text{DT}) = 0.63 * M - b$) between earthquake magnitude, precursory anomaly duration,
334 and the distance of the earthquake epicenter to the monitoring site, such a long duration
335 (six months) of a precursory anomaly we report in this study is very likely because the
336 magnitudes of the 7.7 and 7.6 devastating earthquakes are sufficiently big to cause such
337 a long precursory anomaly at a location about 100 km from the epicenter. Considering
338 the relation proposed by Sultankhodhaev (1984), such a magnitude of the earthquake
339 theoretically should lead to months-long of precursory anomaly in the geochemical
340 parameters at locations hundreds of kilometers far from the epicentral area. For instance,
341 in Western Turkey, Inan et al. (2010) reported 32 days of chemical anomaly at a water
342 monitoring site located 32 kilometers from the epicenter of a M 4.2 earthquake.


343 In regard to changes in the dissolved ions in the AYR spring water, the following changes
344 are prominent. The Ca^{2+} and Na^+ content increase (for the period between September
345 2022 to 15 February 2023) above the background by about 14 (mg/l) and 10 (mg/l),
346 respectively, and reach up to 22 (mg/l) and 16 (mg/l), respectively. This increase started
347 about six months before the 6 February earthquakes (EQ # 1 and EQ #2). Since we could
348 not obtain samples between 8 March 2022 and 14 September 2022, the anomaly could
349 have possibly started even earlier (any time between March and August 2022); so the
350 positive anomaly (e.g., increase) in the major ions started at least six months before the
351 6 February 2023 earthquakes. The Mg^{2+} content also increased from about 4 (mg/l) to 10
352 (mg/l) in the period September 2022 to 15 February 2023. Similar major increases were
353 also detected in Cl^- , and SO_4^{2-} contents. Water samples are relatively poor in K^+ content
354 therefore the increase, due to the scale of the graph, is not very obvious in Figure 3B.
355 However, the values given in Table 2 clearly indicate about four times an increase in the
356 K^+ content compared to the background concentrations (post-seismic samples collected
357 between February 15 and 31 March 2023).

358 The pre-earthquake anomaly in the AYR water samples is characterized by an increase
359 of up to 400% for the Ec and also major ions; namely Ca^{2+} , Mg^{2+} , K^+ , Na^+ , Cl^- , and SO_4^{2-}
360 before the 6 February 2023 Mw 7.7 and Mw 7.6 earthquakes (Figure 3B). It is worth noting
361 that the Ec values and major ion concentrations for the Ayran spring waters show similar
362 trends. Post-earthquake samples show decreasing trends in all major ions. Analyses
363 results of the post-earthquake dated samples show that the spring water has had
364 chemical stability since the Middle February-Early March 2023; just two to three weeks
365 after the earthquakes (Figure 3B). We have also obtained a chemical analysis report on
366 AYR water submitted with the business license application of the company dated 29
367 August 2012. The chemical analysis data of the samples collected more than 10 years
368 ago include values only for Na^+ , Cl^- , and SO_4^{2-} as 3.86, 3.12, and 8.37 mg/l, respectively.
369 These values are very close to the analysis result of the AYR water sample dated 8 March
370 2022 (AYR 1 which is the oldest sample in our data set) and the AYR water samples
371 collected after 15 February (Table 2); confirming that these samples represent
372 background values for the AYR spring water.


373 Immediately after the earthquake, the values started to decrease suggesting a reversible
374 chemical change (Figure 3B; Table 2). It is worth mentioning that the broad positive
375 anomaly detected in the AYR water chemistry (Figure 3B) that lasted for about six months
376 before the Mw 7.7 and Mw 7.6 earthquakes shows some transient decreases (about
377 Middle December 2022 and toward the end of January 2023). Following each transient
378 decrease, an increase in ion contents is observed and the broad positive anomaly
379 (starting from September 2022) is sustained until the date of the major earthquakes of 6
380 February 2023. The observations of sudden decrease and rebound in the major ion
381 contents of the water samples (taking place in Mid December 2022 and end of January
382 2023) may suggest sudden and short-lived crustal stress release related to smaller
383 earthquakes (e.g., EQ # 4 and EQ # 5). Soon after the major earthquakes (EQ # 1 and
384 EQ # 2), the major ion contents of the water samples show a sharp decline; almost
385 approaching the background values as early as 15 February 2023. One single positive
386 anomaly after the major earthquakes (EQ #1 and EQ #2) is detected in the sample dated
387 16 February 2023. The further increase of the ion contents of this sample seems to
388 suggest a short-term stress buildup prior to EQ # 3 (Mw 6.4) that occurred about 120 km
389 to the south of the Ayran Spring water location (Figure 1B). Considering Dobrovolsky et
390 al.'s (1979) theoretical relation ($R= 10^{0.43*M}$), an increase in major ions contents of the
391 Ayran Spring water is very likely to take place due to an earthquake of magnitude 6.4
392 occurring in 120 km distance.

393

394




395



396

397 **Figure 3.** Time variation graphs of Ec for the Ayran (AYR) and the Bahçepinar (BPN)
 398 bottled waters (A) and major ions for the AYR bottled waters (B). All data are listed in
 399 Table 2.



400

401 **Figure 3. Cont.** Daily average air temperature and rainfall at the Osmaniye meteorology
 402 station (37°07'N, 36°25'E; 32 km SW of the Ayran spring) between 1 September 2022
 403 and 30 March 2023 (C).

404 ([https://www.meteoblue.com/tr/hava/historyclimate/weatherarchive/
 405 osmaniye_turkiye_303195](https://www.meteoblue.com/tr/hava/historyclimate/weatherarchive/osmaniye_turkiye_303195)). EQ1 through EQ6 are the earthquakes listed in Table 1.

406

407 We have shown and discussed the reliable precursory anomalies in the major ions of the
 408 bottled AYR spring water prior to the Mw 7.7 and Mw 7.6 earthquakes that occurred in
 409 the Kahramanmaraş region on 6 February 2023. However, the process(es) leading to the
 410 build-up of geochemical anomalies related to the earthquake cannot be inferred with
 411 certainty. However, some inferences based on previous observations can be made. For
 412 instance, Sibson (1992) suggested that extensive hydro-fracture dilatancy might develop
 413 prior to failure leading to the earthquake. Development of fractures probably enhances
 414 water circulation and mixing of different reservoirs leading to pre-earthquake anomalies
 415 (Italiano et al., 2004; Federico et al., 2008; İnan et al., 2010; İnan et al., 2012c; Skelton
 416 et al., 2014; Ingebritsen and Manga, 2014; Doglioni et al., 2014; Barberio et al., 2017;
 417 Skelton et al., 2019; Wang and Manga, 2021; Gori and Barberio, 2022;). Although the
 418 process(es) responsible for chemical anomalies detected in the Ayran spring waters prior
 419 to the 6 February 2023 earthquakes cannot be suggested with any certainty at this stage,
 420 two immediate mechanisms emerge: 1) a simple increase in fluid flow in the surrounding
 421 of the future epicenter and selective dissolution of some K–Mg–Ca-rich rocks (Federico

422 et al., 2008); or 2) “electro-corrosion” whereby the dissolution of rocks is accelerated by
423 the flow of stress-activated positive hole currents (Balk et al., 2009; Freund, 2011; Paudel
424 et al., 2018). Following the second mechanism, the increased content of major ions in
425 water could be related to the oxidation of water to hydrogen peroxide at the rock-water
426 interface (Balk et al., 2009; Paudel et al., 2018). Freund (2011) suggested that with the
427 positive hole current flowing, the “corrosion” of the rock is accelerated releasing into the
428 water major cations and anions. Further work to be conducted in this area may enable us
429 to suggest the process(es) responsible for the pre-earthquake geochemical anomalies
430 we have discussed in the AYR spring water.

431

432 **5. Conclusions**

433

434 Hydrogeochemical precursors have been detected in commercially bottled water samples
435 of natural springs (Ayran Spring and Bahçepinar Spring) emanating from a location about
436 100 km distance from the epicenter of the Mw 7.7 Kahramanmaraş Earthquake of 6
437 February 2023. The pre-earthquake anomaly is characterized by an increase in Ca^{2+} ,
438 Mg^{2+} , K^+ , Na^+ , Cl^- , and SO_4^{2-} content in the bottled water samples of the Ayran spring.
439 Samples that are dated after the earthquakes (covering about two months after the
440 earthquake) show decreasing trends in all major ions. About three weeks after the
441 earthquake, the major ion contents of the spring water attained stability. At least six
442 months of pre-earthquake anomaly (increase) in the major ions content of the Ayran
443 spring water is prominent. It is worth noting that the Bahçepinar Spring water samples did
444 not show any anomalies in electrical conductivity therefore the samples were not
445 analyzed for ion content. Bahçepinar water is collected from shallow boreholes dug into
446 alluvial deposits which, we believe, are decoupled from the basement rocks and this may
447 be the reason for the lack of any significant change in the water chemistry prior to the
448 earthquakes. Here, we remind that geological knowledge of the investigated area and the
449 sampling site have paramount importance in sampling discrete samples for geochemical
450 analysis and/or conducting continuous monitoring. The results of this study suggest that
451 spring water chemical anomalies may be monitored as proxy indicators of pre-earthquake

452 crustal deformation. The physical mechanisms of the observed precursors are yet
453 impossible to explain with certainty at this stage. In order to be able to suggest the
454 mechanism(s) leading to the reported pre-earthquake geochemical anomalies, more work
455 needs to be conducted; especially multi-disciplinary (seismological, geodetical,
456 geochemical) and continuous earthquake monitoring networks must be established and
457 run for a sufficiently long time.

458

459 **Acknowledgements**

460 We appreciate all the technical help we have received on ion chromatography analyses
461 from Ms. Sevde Korkut at the Istanbul Technical University MEM-TEK laboratory. We
462 thank Mr. Asen Sabuncu (Istanbul Technical University) and Dr. Teach. Assist. Emre
463 Pınarcı (Çukurova University) for help in drafting the figures. We also thank Assoc. Prof
464 Dr. Tülay İnan for help in conducting electrical conductivity measurements of the bottled
465 water samples. We thank Prof. Dr. Filippos Vallianatos for editorial handling. We also
466 appreciate constructive reviews by Dr. Giovanni Martinelli, Dr. Vivek Valia, and three
467 anonymous reviewers that helped us improve the manuscript. This work has been
468 partially supported by Istanbul Technical University Scientific Research Fund (ITU
469 BAPSIS) Project # 44774.

470 **Authors contributions**

471 S.I. and H.C. conceived the project; H.C. collected the samples; N.Y. coordinated
472 laboratory analysis, compiled seismic events, and prepared the figures; S.I. was the
473 primary interpreter of the data. S.I. and H.C. were writers of the manuscript with
474 contributions from N.Y.

475 **References**

476 Allen, C.R.: Active faulting in northern Turkey, Division of Geological Sciences, California
477 Institute of Technology, Contribution No. 1577. 32 pp, 1969.

478 Ambraseys, N.N.: Temporary seismic quiescence: SE Turkey. *Geophys. J. Int.*, 96, 311–
479 331, <https://doi.org/10.1111/j.1365-246X.1989.tb04453.x>, 1989.

480 Areshidze, G., Bella, F., Biagi, P. F., Caputo, M., Della Monica, G., Ermini, A.,
481 Manjgaladze, P., Melikazdze, G., Sgrigna, V. and Zilpimani, D.: No preseismic evidence
482 from hydrogeochemical parameters on the occasion of April 29, 1991 Georgian
483 Earthquake, Caucasus, Tectonophysics, 213, 353–358, [https://doi.org/10.1016/0040-](https://doi.org/10.1016/0040-1951(92)90463-Gö)
484 [1951\(92\)90463-Gö](https://doi.org/10.1016/0040-1951(92)90463-Gö) 1992.

485 Arpat, E., Şaroğlu, F.: Dogu Anadolu fayı ile ilgili bazı gözlem ve düşünceler (Some
486 observations and thoughts on the East Anatolian fault). Bulletin of the General Directorate
487 of Mineral Research and Exploration of Turkey (MTA) 73, 44–50, 1972.

488 Balk, M., Bose, M., Ertem, G., Rogoff, D. A., Rothschild, L. J. and Freund, F.T.: Oxidation
489 of water to hydrogen peroxide at the rock-water interface due to stress-activated electric
490 currents in rocks, Earth Planet Sc. Lett., 283, 87–92.
491 <https://doi.org/10.1016/j.epsl.2009.03.044>, 2009.

492 Barberio, M.D., Barbieri, M., Billi, A., Doglioni, C., Petitta, M.: Hydrogeochemical changes
493 before and during the 2016 Amatrice-Norcia seismic sequence (central Italy). Sci. Rep-
494 UK, 7, 11735. <https://doi.org/10.1038/s41598-017-11990-8>, 2017.

495 Barka, A. A. and Reilinger, R.: Active tectonics of the Eastern Mediterranean region
496 deduced from GPS, neotectonic and seismicity data, Ann. Geophys-Italy, 40, 587-610,
497 <http://hdl.handle.net/2122/1520>, 1997.

498 Barka, A.A., Kadinsky-Cade, K.: Strike-slip fault geometry in Turkey and its influence on
499 earthquake activity, Tectonics, 7, 663–684, <https://doi.org/10.1029/TC007i003p00663>,
500 1988.

501 Bella, F., Biagi, P.F., Caputo, M., Cozzi, Della Monica, G., Ermini, A., Gordeez, E.I.,
502 Khatkevich, Y.M., Martinelli, G., Plastino, W., Scandone, R., Sgrigna, V., Zilpimiani, D.:
503 Hydrogeochemical anomalies in Kamchatka (Russia), Phys. Chem. Earth., 23, 921-925,
504 [https://doi.org/10.1016/S0079-1946\(98\)00120-7](https://doi.org/10.1016/S0079-1946(98)00120-7), 1998.

505 Birchard, G.F., Libby, W.F.: Earthquake associated radon anomalies possible
506 mechanisms, Eos Trans. AGU 59, 4, 329, 1978.

507 Bozkurt, E.: Neotectonics of Turkey—a synthesis, Geodin. Acta, 14, 3–30,
508 <https://doi.org/10.1080/09853111.2001.11432432>, 2001.

509 Cetin, H., Guneyli, H., Mayer, L.: Paleoseismology of the Palu–Lake Hazar segment of
510 the East Anatolian Fault Zone, Turkey, Tectonophysics 374, 163–197,
511 <http://dx.doi.org/10.1016/j.tecto.2003.08.003>, 2003.

512 Chorowicz, J., Luxey, P., Lyberis, N., Carvalho, J., Parrot, J.F., Yurur, T. and Gundogdu,
513 N.: The Maras Triple Junction (southern Turkey) based on digital elevation model and
514 satellite imagery interpretation, J. Geophys. Res., 99, B10, 20225–20242,
515 <https://doi.org/10.1029/94JB00321>, 1994.

516 Cicerone, R.D., Ebel, J.E., Britton, J.: A systematic compilation of earthquake precursors,
517 Tectonophysics, 476, 371-396. <https://doi.org/10.1016/j.tecto.2009.06.008>, 2009.

518 Claesson, L., Skelton, A., Graham, C., Dietl, C., Mörth, M., Torssander, P., Kockum, I.:
519 Hydrogeochemical changes before and after a major earthquake, Geology, 32, 641-644,
520 <https://doi.org/10.1130/G20542.1>, 2004.

521 Conti, L., Picozza, P., Sotgiu, A.: A critical review of ground-based observations of
522 earthquake precursors, Front. Earth Sci., 9, <https://doi.org/10.3389/feart.2021.676766>,
523 2021.

524 Dewey, J.F. and Şengör, A.M.C.: Aegean and surrounding regions: complex multi-plate
525 and continuum tectonics in a convergent zone, Geol. Soc. Am. Bull., 90, 84–92,
526 [https://doi.org/10.1130/0016-
527 7606%281979%2990%3C84%3AAASRCM%3E2.0.CO%3B2](https://doi.org/10.1130/0016-7606%281979%2990%3C84%3AAASRCM%3E2.0.CO%3B2), 1979.

528 Dewey, J.F., Pitman, W.C., Ryan, W.B.F., Bonnin, J.: Plate tectonics and the evolution of
529 the Alpine system. Geol. Soc. Am. Bull., 84, 3137–3180, [https://doi.org/10.1130/0016-
530 7606\(1973\)84%3C3137:PTATEO%3E2.0.CO;2](https://doi.org/10.1130/0016-7606(1973)84%3C3137:PTATEO%3E2.0.CO;2), 1973.

531 Di Luccio, F., Chiodini, G., Caliro, S., Cardellini, C., Convertito, V., Pino, N.A., Tolomei,
532 C., Ventura, G.: Seismic signature of active intrusions in mountain chains. Science
533 Advances 4, e1701825. <https://doi.org/10.1126/sciadv.1701825>, 2018.

534 Dobrovolsky, I.P., Zubkov, S.I. and Miachkin, V.I.: Estimation of the size of earthquake
535 preparation zones, Pure and Applied Geophysics 117, 1025-1044,
536 <https://doi.org/10.1007/BF00876083>, 1979.

537 Doglioni, C., Barba, S., Carminati, E., Riguzzi, F.: Fault on–off versus coseismic fluids
538 reaction, Geosci. Front., 5 (6), 767–780, <https://doi.org/10.1016/j.gsf.2013.08.004>, 2014.

539 Etioppe, G., Calcara, M. and Quattrocchi, F.: Seismogeochemical algorithms for
540 earthquake prediction: a review, Ann. Geophys-Italy, 40(6), 1483–1492,
541 <http://hdl.handle.net/2122/1524>, 1997.

542 Federico C., Pizzino, L., Cinti, D., De Gregorio, S., Favara, R., Galli, G., Giudice, G.,
543 Guerrieri, S., Quattrocchi, F. and Voltattorni, N.: Inverse and forward modeling of
544 groundwater circulation in a seismically active area (Monferrato, Piedmont, NW Italy):
545 insights into stress-induced variations in water chemistry, Chem. Geol., 248, 14–39,
546 <http://dx.doi.org/10.1016/j.chemgeo.2007.10.007>, 2008.

547 Freund, F. T.: Pre-earthquake signals: Underlying physical processes. J. Asian Earth Sci.,
548 41, 383–400, <https://doi.org/10.1016/j.jseaes.2010.03.009>, 2011.

549 Freund, F.T.: Earthquake probabilities and pre-earthquake signals, Current Science, 94,
550 311-313, <https://www.jstor.org/stable/24100338>, 2008.

551 Freund, F.T., Takeuchi, A., Lau, B.W.S.: Electric currents streaming out of stressed
552 igneous rocks – A step towards understanding pre-earthquake low frequency EM
553 emissions, *Phys. Chem. Earth*, 31, 389-396, <https://doi.org/10.1016/j.pce.2006.02.027>,
554 2006.

555 Gori, F. and Barberio, M.D.: Hydrogeochemical changes before and during the 2019
556 Benevento seismic swarm in central-southern Italy, *J. Hydrol.*, 604,
557 <https://doi.org/10.1016/j.jhydrol.2021.127250>, 2022.

558 Grant, R. A., Halliday, T., Balderer, W. P., Leuenberger, F., Newcomer, M., Cyr, G. and
559 Freund, F. T.: Groundwater chemistry changes before major earthquakes and possible
560 effects on animals, *Int. J. Environ. Res. PU*, 1936–1956,
561 <https://doi.org/10.3390%2Fijerph8061936>, 2011.

562 Geller, R. J., Jackson, D. D., Kagan, Y. Y., and Mulargia, F.: Earthquake cannot be
563 predicted, *Science*, 275, 1616–1617, <https://doi.org/10.1126/science.275.5306.1616>,
564 1997.

565 Hancock, P. L. and Barka, A. A.: Kinematic indicators on active normal faults in western
566 Turkey. *J. Struct Geol.*, 9, 415-430, [https://doi.org/10.1016/0191-8141%2887%2990142-
567 8](https://doi.org/10.1016/0191-8141%2887%2990142-8), 1987.

568 Hartmann, J. and Levy, J. K.: The influence of seismotectonic on precursory changes in
569 groundwater composition for the 1995 Kobe earthquake, Japan. *Hydrology Journal*, 14,
570 1307–1318, <https://doi.org/10.1007/s10040-006-0030-7>, 2006.

571 Hempton, M.R.: The North Anatolian fault and complexities of continental escape. *J. Struct*
572 *Geol.*, 4, 502 – 504, [https://doi.org/10.1016/0191-8141\(82\)90041-4](https://doi.org/10.1016/0191-8141(82)90041-4), 1982.

573 Hauksson, E.: Radon content of groundwater as an earthquake precursor: Evaluation of
574 worldwide data and physical basis, *J. Geophys. Res.* 86, B10, 9397-9410,
575 <https://doi.org/10.1029/JB086iB10p09397>, 1981.

576 He, A. and Singh, R.P.: Groundwater level response to the Wenchuan earthquake of
577 May 2008, *Geomatics, Natural Hazards and Risk* 10, 336-352,
578 <https://doi.org/10.1080/19475705.2018.1523236> . 2019.

579 İnan, S., Ergintav, S., Saatçılar, S., Tüzel, B. and İravul, Y.: Turkey makes major
580 investments in earthquake research. *EOS Transactions* 88, 333–334,
581 <https://doi.org/10.1029/2007EO340002>, 2007.

582 İnan, S., Akgül, T., Seyis, C., Saatçılar, R., Baykut, S., Ergintav, S. and Bas, M.:
583 Geochemical monitoring in the Marmara Region (NW Turkey): A search for precursors of
584 seismic activity, *J. Geophys. Res.* 113, B03401, <https://doi.org/10.1029/2007JB005206>,
585 2008.

586 İnan, S., Ertekin, K., Seyis, C., Şimşek, S., Kulak, F., Dikbaş, A., Tan, O., Ergintav, S.,
587 Çakmak, R., Yörük, A., Çergel, M., Yakan, H., Karakuş, H., Saatçılar, R., Akçiğ, Z., İravul,
588 Y. and Tüzel, B. (2010) Multi-disciplinary earthquake researches in Western Turkey: Hints
589 to select sites to study geochemical transients associated to seismicity. *Acta Geophys.*,
590 58, 767–813, <https://doi.org/10.2478/s11600-010-0016-7>, 2010.

591 İnan, S., Pabuçcu, Z., Kulak, F., Ergintav, S., Tatar, O., Altunel, E., Akyüz, S., Tan, O.,
592 Seyis, C., Çakmak, R., Saatçılar, R. and Eyidogan, H.: Microplate boundaries hindering
593 pre-earthquake strain transfer (Western Turkey): inferences from continuous
594 geochemical monitoring, *J. Asian Earth Sci.*, 48, 56-71,
595 <http://dx.doi.org/10.1016/j.jseaes.2011.12.016>, 2012a.

596 İnan, S., Kop, A., Çetin, H., Kulak, F., Pabuçcu, Z., Seyis, C., Ergintav, S., Tan, O.,
597 Saatçılar, S. and Bodur, M. N.: Seasonal variations in soil radon emanation: Long term
598 continuous monitoring in light of seismicity, *Nat. Hazards*, 62, 575-591,
599 <https://doi.org/10.1007/s11069-012-0096-6>, 2012b.

600 İnan, S., Balderer, W.P., Leuenberger-West, F., Yakan, H., Özvan, A., Freund, F.T.:
601 Springwater chemical anomalies prior to the Mw = 7.2 Van Earthquake
602 (Turkey). *Geochem J.*, 46, e11-e16, <https://doi.org/10.2343/geochemj.1.0159>, 2012c.

603

604 Ingebritsen, S.E., Manga, M.: Earthquakes: hydrogeochemical precursors. *Nat. Geosci.*,
605 7, 697–698. <https://doi.org/10.1038/ngeo2261>, 2014.

606 Italiano, F., Martinelli, G. and Rizzo, A.: Geochemical evidence of seismogenic-induced
607 anomalies in the dissolved gases of thermal waters: A case study of Umbria (Central
608 Apennines, Italy) both during and after the 1997–1998 seismic swarm, *Geochem.*
609 *Geophys. Geosy.*, 5, Q11001, <https://doi.org/10.1029/2004GC000720>, 2004.

610 Jackson, J. and McKenzie, D.P.: Active tectonics of the Alpine–Himalayan Belt between
611 western Turkey and Pakistan, *Geophys. J. Roy. Astr. Soc.*, 77, 185–264,
612 <https://doi.org/10.1111/j.1365-246X.1984.tb01931.x>, 1984.

613 Karig, D.E. and Kozlu, H.: Late Palaeogene – Neogene evolution of the triple junction
614 region near Maras, south-central Turkey, *J. Geol. Soc. London*, 147, 1023–1034,
615 <https://doi.org/10.1144/gsjgs.147.6.1023>, 1990.

616 Kelling, G., Gökçen, S.L., Floyd, P.A. and Gökçen, N.: Neogene tectonics and plate
617 convergence in the Eastern Mediterranean: new data from southern Turkey, *Geology*, 15,
618 425–429, [https://doi.org/10.1130/0091-7613\(1987\)15%3C425:NTAPCI%3E2.0.CO;2](https://doi.org/10.1130/0091-7613(1987)15%3C425:NTAPCI%3E2.0.CO;2)
619 1987.

620 Kempler, D. and Garfunkel, Z.: Northeast Mediterranean triple junction from a plate
621 kinematics point of view, *Bulletin of the Technical University of Istanbul, Special Issue 44*,
622 425–454, 1991.

- 623 Ketin, I.: Über die tektonisch-mechanischen Folgerungen aus den grossen anatolischen
624 Erdbeben des letzten Dezenniums, Geol. Rundsch, 36, 77–83,
625 <https://doi.org/10.1007/BF01791916>, 1948.
- 626 King, J. Y., Koizumi, Y. and Kitagawa, Y.: Hydrogeochemical anomalies and the 1995
627 Kobe Earthquake, Science, 269, 38–39, <https://doi.org/10.1126/science.269.5220.38>,
628 1995.
- 629 Lovelock, P.E.R.: A review of the tectonics of the northern Middle East region. Geol. Mag.,
630 121, 577–587, Online ISSN: 1469-5081, 1984.
- 631 Martinelli, G. and Ferrari, G.: Earthquake forerunners in a selected area of Northern Italy:
632 recent developments in automatic geochemical monitoring. Tectonophysics 193, 397-
633 410. [https://doi.org/10.1016/0040-1951\(91\)90348-v](https://doi.org/10.1016/0040-1951(91)90348-v) , 1991.
- 634 McKenzie, D.P.: Active tectonics of the Mediterranean Region, Geophys. J. Int., 30, 109–
635 185, <https://doi.org/10.1111/j.1365-246X.1972.tb02351.x>, 1972.
- 636 McKenzie, D.P.: The East Anatolian fault: a major structure in eastern Turkey. Earth
637 Planet Sc. Lett., 29, 189–193, [https://doi.org/10.1016/0012-821X\(76\)90038-8](https://doi.org/10.1016/0012-821X(76)90038-8), 1976.
- 638 Muehlberger, W.B. and Gordon, M.B.: Observations on the complexity of the East
639 Anatolian fault, Turkey. J. Struct Geol., 9, 899–903, [http://dx.doi.org/10.1016/0191-8141\(87\)90091-5](http://dx.doi.org/10.1016/0191-8141(87)90091-5), 1987.
- 641 Nur, A.: Matsushiro, Japan, earthquake swarm: Confirmation of the dilatancy-fluid
642 diffusion model, Geology, 2, 217–221, [https://doi.org/10.1130/0091-7613\(1974\)2%3C217:MJESCO%3E2.0.CO;2](https://doi.org/10.1130/0091-7613(1974)2%3C217:MJESCO%3E2.0.CO;2), 1974.
- 644 Ouzounov, D., Pulinets, S., Davidenko, D., Rozhnoi, A., Solovieva, M., Fedun, V.,
645 Dwivedi, B. N., Rybin, A., Kafatos, M., and Taylor, P.: Transient effects in atmosphere
646 and ionosphere preceding the 2015 M7.8 and M7.3 Gorkha-Nepal earthquakes, Front.
647 Earth Sci., 9, <https://doi.org/10.3389/feart.2021.757358>, 2021.
- 648 Papadopoulos, G. A., Latoussakis, I., Daskalaki, E., Diakogianni, G., Fokaefs, A., Kolligri,
649 M., Liadopoulou, K., Orfanogiannaki, K., and Pirentis, A.: The East Aegean Sea strong
650 earthquake sequence of October–November 2005: lessons learned for earthquake
651 prediction from foreshocks, Nat. Hazards Earth Syst. Sci., 6, 895–901,
652 <https://doi.org/10.5194/nhess-6-895-2006> , 2006.
- 653 Paudel, S.R., Banjara, S.P., Wagle, A., Freund, F.T.: Earthquake chemical precursors in
654 groundwater: a review, J. Seismol., 22, 1293–1314, <https://doi.org/10.1007/s10950-018-9739-8>, 2018.
- 656 Pérez, N. M., Hernández, P. A., Igarashi, G., Trujillo, I., Nakai, S., Sumino, H. and Wakita,
657 H.: Searching and detecting earthquake geochemical precursors in CO₂-rich

658 groundwaters from Galicia, Spain, *Geochem J.*, 42, 75–83,
659 <http://dx.doi.org/10.2343/geochemj.42.75> , 2008.

660 Perinçek, D. and Çemen, I.: The structural relationship between the East Anatolian and
661 Dead Sea fault zones in southeastern Turkey, *Tectonophysics*, 172, 331–340,
662 [http://dx.doi.org/10.1016/0040-1951\(90\)90039-B](http://dx.doi.org/10.1016/0040-1951(90)90039-B), 1990.

663 Rapti, D., Martinelli, G., Zheng, G., Vincenzi, C.: Bottled Mineral Waters as
664 Unconventional Sampling in Hydro-Geological Research. *Water* 15, 3466.
665 <https://doi.org/10.3390/w15193466>. 2023

666 Rikitake, T.: Earthquake Precursors in Japan: Precursor Time and Detectability,
667 *Tectonophysics*, 136, 265–282, [https://doi.org/10.1016/0040-1951\(87\)90029-1](https://doi.org/10.1016/0040-1951(87)90029-1) , 1987.

668 Rikitake, T.: Classification of earthquake precursors, *Tectonophysics*, 54, 293-309,
669 [https://doi.org/10.1016/0040-1951\(79\)90372-X](https://doi.org/10.1016/0040-1951(79)90372-X), 1979.

670 Roeloffs, E.: Poroelastic techniques in the study of earthquake-related hydrologic
671 phenomena, *Adv. Geophys.*, 37, 135–195, [https://doi.org/10.1016/S0065-](https://doi.org/10.1016/S0065-2687%2808%2960270-8)
672 [2687%2808%2960270-8](https://doi.org/10.1016/S0065-2687%2808%2960270-8), 1996.

673 Rotstein, Y.: Counterclockwise rotation of the Anatolian block, *Tectonophysics*, 108, 71–
674 91, [https://doi.org/10.1016/0040-1951\(84\)90155-0](https://doi.org/10.1016/0040-1951(84)90155-0), 1984.

675 Şaroğlu, F., Emre, O., Kuscu, I.: Active Fault map of Turkey (1:2 000 000 scale), General
676 Directorate of Mineral Research and Exploration (MTA), Ankara, Turkey, 1992a

677 Şaroğlu, F., Emre, O., Kuscu, I.: The East Anatolian fault zone of Turkey. *Annales*
678 *Tectonicae: International Journal of Structural Geology and Tectonics*, 6, 99–125 (Special
679 Issue-Supplement to Volume VI), 1992b.

680 Scholz, R., Sykes, L. R. and Aggarwal, Y. P.: Earthquake prediction: A physical basis,
681 *Science*, 181, 803–810, <https://www.science.org/doi/10.1126/science.181.4102.803>,
682 1973.

683 Sengör, A. M. C. and Yılmaz, Y.: Tethyan evolution of Turkey: A plate tectonic approach,
684 *Tectonophysics*, 75, 181– 241, [http://dx.doi.org/10.1016/0040-1951\(81\)90275-4](http://dx.doi.org/10.1016/0040-1951(81)90275-4), 1981.

685 Sengör, A. M. C., Görür, N. and Şaroğlu, F.: Strike-slip faulting and related basin
686 formation in zones of tectonic escape: Turkey as a case study, in: *Strike-slip Deformation,*
687 *Basin Formation, and Sedimentation*, edited by: Biddle, K. T. and Christie-Blick, N., *Soc.*
688 *Econ. Pa.*, 37, 227–264, 1985.

689 Seyis, C., İnan, S., Yalçın, M.N.: Major factors affecting soil radon emanation, *Nat.*
690 *Hazards*, 114, 2139–2162, <http://link.springer.com/10.1007/s11069-022-05464-y>,
691 2022.
692 .

693 Sibson, R. H.: Implications of fault-valve behavior for rupture nucleation and recurrence,
694 *Tectonophysics*, 211, 283–293, [https://doi.org/10.1016/0040-1951\(92\)90065-E](https://doi.org/10.1016/0040-1951(92)90065-E), 1992.

695 Sibson, R.H., Moore, J.M.M., Rankin, A.H.: Seismic pumping—a hydrothermal fluid
696 transport mechanism, *J. Geol. Soc. London*, 131, 653–659,
697 <https://doi.org/10.1144/gsjgs.131.6.0653>, 1975.

698 Skelton, A., Liljedahl-Claesson, L., Wästeby, N., Andrén, M., Stockmann, G., Sturkell, E.,
699 et al.: Hydrochemical changes before and after earthquakes based on long-term
700 measurements of multiple parameters at two sites in northern Iceland—a review, *J.*
701 *Geophys. Res-Sol EA.*, 124, 2702–2720. <https://doi.org/10.1029/2018JB016757>, 2019.

702 Skelton, A., Andrén, M., Kristmannsdóttir, H., Stockmann, G., Mörth, C-M.,
703 Sveinbjörnsdóttir, A., Jónsson, S., Sturkell, E., Guðrúnardóttir, H.R., Hjartarson, H.,
704 Siegmund, H., Kockum, I.: Changes in groundwater chemistry before two consecutive
705 earthquakes in Iceland. *Nat. Geosci.*, 7, 752-756. <http://dx.doi.org/10.1038/NGEO2250>,
706 2014.

707 Sol, S., Meltzer, A., Bürgmann, R., van der Hilst, R. D., King, R., Chen, Z., Koons, P. O.,
708 Lev, E., Liu, Y. P., Zeitler, P. K., Zhang, X., Zhang, J. and Zurek, B.: Geodynamics of the
709 southern Tibetan Plateau from seismic anisotropy and geodesy, *Geology*, 35, 563–566,
710 <https://doi.org/10.1130/G23408A.1>, 2007.

711 Soysal, H., Sipahioğlu, S., Kolçak, D. and Altınok, Y.: Historical earthquake catalogue of
712 Turkey and surrounding area (2100 B.C. – 1900 A.D.), The Scientific and Technological
713 Research Council of Turkey Technical Report, TBAG-341, 1981.

714 Sugisaki, R., Ito, T., Nagamine, K. and Kawabe, I.: Gas geochemical changes at mineral
715 springs associated with the 1995 southern Hyogo earthquake (M = 7.2), Japan, *Earth*
716 *Planet Sc. Lett.*, 139, 239–249, [https://doi.org/10.1016/0012-821X\(96\)00007-6](https://doi.org/10.1016/0012-821X(96)00007-6), 1996.

717 Sultankhodhaev, G. A.: International symposium on earthquake prediction, UNESCO,
718 Paris 1979, 181-191, 1984.

719 Tansi, C., Tallarico, A., Iovine, G., Folino Gallo, M. and Falcone, G.: Interpretation of
720 radon anomalies in seismotectonic and tectonic-gravitational settings: the southeastern
721 Crati graben (Northern Calabria, Italy), *Tectonophysics*, 396, 181–193,
722 <https://doi.org/10.1016/j.tecto.2004.11.008>, 2005.

723 Thomas, D.M., Cuff, K.E., Cox, M.E.: The association between ground gas radon
724 variations and geologic activity in Hawaii, *J. Geophys. Res-Sol EA*, 91, 12186-12198,
725 [https://ui.adsabs.harvard.edu/link_gateway/1986JGR....9112186T/doi:10.1029/JB091iB](https://ui.adsabs.harvard.edu/link_gateway/1986JGR....9112186T/doi:10.1029/JB091iB12p12186)
726 [12p12186](https://ui.adsabs.harvard.edu/link_gateway/1986JGR....9112186T/doi:10.1029/JB091iB12p12186). 1986.

- 727 Toutain, J. P., Munoz, M., Poitrasson, F. and Lienard, F.: Springwater chloride ion
728 anomaly prior to a ML = 5.2 Pyrenean earthquake, *Earth Planet Sc. Lett.*, 149, 113–119.
729 [https://doi.org/10.1016/S0012-821X\(97\)00066-6](https://doi.org/10.1016/S0012-821X(97)00066-6), 1997.
- 730 Tsunogai, U. and Wakita, H.: Precursory chemical changes in groundwater: Kobe
731 earthquake, Japan, *Science*, 269, 61-63, <https://doi.org/10.1126/science.269.5220.61>,
732 1995.
- 733 Turcotte, W. L.: Earthquake prediction, *Annu. Rev. Earth Pl. Sc.*, 19, 263 – 281,
734 <https://doi.org/10.1146/annurev.ea.19.050191.001403>, 1991.
- 735 Usta, D., Ateş, Ş., Beyazpırınç, M., Kanar, F., Yıldız, H., Uçar, L., İsmail Akça, İ., Tufan,
736 E. and Örtlek, A. T.: New data on the stratigraphy of the central and northern Amonous
737 mountains (Osmaniye - Gaziantep - K. Maraş). *Bulletin of the Turkish Association of*
738 *Petroleum Geologists*, 27, 57-98, 2015.
- 739 Usta, D., Ateş, Ş., Kanar, F., Beyazpırınç, M., Uçar, L., Yıldız, H., Tufan, E., Akça, İ.,
740 Örtlek, A. T.: Doğu Toroslar'ın jeolojisi ve jeodinamik evrimi projesi (K.Maras, Osmaniye,
741 Gaziantep, Adana, Hatay). General Directorate of Minerals and Exploration of Turkey
742 (MTA) unpublished Report No 13568, 494 pages, Ankara (in Turkish), 2017.
- 743 Uyeda, S., Nagao, T. and Kamogawa, M.: Short-term earthquake prediction: Current
744 status of seismo-electromagnetics, *Tectonophysics*, 470, 205–213.
745 <https://doi.org/10.1016/j.tecto.2008.07.019>, 2008.
- 746 Wakita, H.: Geochemical challenge to earthquake prediction, *P. Natl. Acad. Sci. USA*, 93,
747 9, 3781-3786, DOI: 10.1073/pnas.93.9.3781, 1996.
- 748 Wakita, H., Nakamura, Y. and Sano, Y.: Short-term and intermediate-term geochemical
749 precursors, *Pure and Applied Geophysics*, 125, 267–278,
750 <https://doi.org/10.1007/BF00878999>, 1988.
- 751 **Walia, V., Yang, T.F., Hong, W.L., Lin, S.J., Fu, C.C., Wen, K.L., Chen, C.H.:**
752 **Geochemical variation of soil–gas composition for fault trace and earthquake precursory**
753 **studies along the Hsincheng fault in NW Taiwan, *Applied Radiation and Isotopes* 67,**
754 **1855-1863, <https://doi.org/10.1016/j.apradiso.2009.07.004>, 2009.**
- 755 Wang, C-Y. and Manga, M.: *Water and Earthquakes. Lecture Notes in Earth System*
756 *Sciences*. Springer, 307 pp. <https://doi.org/10.1007/978-3-030-64308-9>. 2021.
- 757 Westaway, R. and Arger, J.: The Gölbaşı basin, southeastern Turkey: a complex
758 discontinuity in a major strike-slip fault zone, *J. Geol. Soc. London*, 153, 729 – 744,
759 <https://doi.org/10.1144/gsjgs.153.5.0729>, 1996.

760 Xiang, Y. and Peng, S.: Hydrochemical and stable isotopes ($\delta^2\text{H}$ and $\delta^{18}\text{O}$) changes
761 of groundwater from a spring induced by local earthquakes, Northwest China. *Frontiers*
762 in *Earth Sciences*, 09 January 2023. *Solid Earth Geophysics*.
763 <https://doi.org/10.3389/feart.2022.1100068>, 2023.

764 Vallianatos, F., Triantis, D., Tzani, A., Anastasiadis, C., Stavrakas, I.: Electric earthquake
765 precursors: from laboratory results to field observations. *Physics and Chemistry of the Earth Parts*
766 *A/B/C* 29, 339-351. <https://doi.org/10.1016/j.pce.2003.12.003>, 2004.

767 Vallianatos, F. and Tzani, A.: Electric current generation associated with the deformation rate of
768 a solid: Preseismic and coseismic signals. *Physics and Chemistry of the Earth Parts A/B/C* 23,
769 933-938. [https://doi.org/10.1016/S0079-1946\(98\)00122-0](https://doi.org/10.1016/S0079-1946(98)00122-0), 1998.

770

771 Virk, H.S., and Singh, B.: Radon anomalies in soil gas and groundwater as earthquake
772 precursor phenomena, *Tectonophysics*, 227, 215-224, [https://doi.org/10.1016/0040-](https://doi.org/10.1016/0040-1951(93)90096-3)
773 [1951\(93\)90096-3](https://doi.org/10.1016/0040-1951(93)90096-3), 1993.

774 Yönlü, Ö., Altunel, E. and Karabacak, V.: Geological and geomorphological evidence for
775 the southwestern extension of the East Anatolian Fault Zone, Turkey. *Earth Planet Sc.*
776 *Lett.*, 469, 1–14, <http://dx.doi.org/10.1016/j.epsl.2017.03.034>, 2017.

777 Yu, H., Liu, L., Ma, Y., Yan, R., Liu, J., Ma, Y., Li, Z., Zhang, X., Zhao, J., Yu, C.: Observed
778 hydrological changes associated with active tectonic blocks before three consecutive
779 earthquakes in Qinghai, China *Scientific Reports* 13, 8988.
780 <https://doi.org/10.1038/s41598-023-36274-2>, 2023.

781 Zeyrek, M., Ertekin, K., Kacmaz, S., Seyis, C., İnan, S.: An ion chromatography method
782 for the determination of major anions in geothermal water samples, *Geostand. Geoanal.*
783 *Res.*, 34, 67-77, <http://dx.doi.org/10.1111/j.1751-908X.2009.00020.x>, 2010.

784