

# A systemic and comprehensive assessment of coastal hazard changes: method and application to France and its overseas territories

Marc Igigabel<sup>1</sup>, Marissa Yates<sup>2,3</sup>, Michalis Vousdoukas<sup>4</sup>, Youssef Diab<sup>5</sup>

<sup>1</sup>Cerema, Technopôle Brest Iroise, 155 rue Pierre Bouguer, BP5, 29280 Plouzané, France

5 <sup>2</sup>LHSV & Cerema, 6 quai Watier, BP 49, 78401 Chatou, France

<sup>3</sup>LHSV, Ecole des Ponts, EDF R&D, 6 quai Watier, BP 49, 78401, Chatou, France

<sup>4</sup>European Commission Joint Research Centre, Via Enrico Fermi 2749, I-21027-Ispra, Italy.

<sup>5</sup>Université Gustave Eiffel, Lab'Urba, 5, Boulevard Descartes, 77430 Champs-sur-Marne, France

*Correspondence to:* Marc Igigabel (marc.igigabel@cerema.fr)

10 **Abstract.** In the context of climate change, height and frequency variations of extreme sea levels (ESL) are studied using  
deterministic and probabilistic approaches. However, this type of approach does not highlight the dynamic effects (waves,  
currents) generated by ~~meteocean~~ metocean events (storms, cyclones, long swells and tsunamis), beyond their effects on sea\_  
levels. In particular, ESL estimates are ~~certainly~~ calculated by considering the main determining physical factors but cannot  
~~report on~~ include all the effects of these factors. Ultimately, this can lead to confusion between ESL and hazard. This article  
15 proposes a systemic assessment method to analyze coastal hazard changes at regional scales, integrating parameters influencing  
sea\_levels, as well as factors describing the geomorphological context (length and shape of the coast, width of the continental  
shelf), ~~meteocean~~ metocean events, and the marine environment (e.g., coral reef state and sea ice extent). French mainland and  
overseas territories were selected to apply the method. The present study highlighted the need to consider not only the sea\_  
level variability, but also the current and future characteristics of ~~meteocean~~ metocean events. The long, concave coasts  
20 bordered by a wide continental shelf appear particularly sensitive to variations in the intensity or trajectory of  
~~meteocean~~ metocean events. Coral reef degradation in the tropics and the decrease in seasonal sea ice extent in the polar regions  
can also significantly change the nearshore hydrodynamics and impacts on the shoreline. These results help to predict the types  
of hazard (shoreline erosion, rapid submersion and/or permanent flooding) that will increase the most in different coastal zones.

## 1 Introduction

25 In 2001, the frequency and intensity of extreme climate events, including marine flooding from tropical and other storms was  
identified by the Intergovernmental Panel on Climate Change (IPCC) as one of the five « reasons for concern » (IPCC, 2001,  
2007). Recent research has improved knowledge of the mechanisms generating ~~sea-level~~ sea level rise (Cazenave and Llovel,  
2010; Cazenave and Le Cozannet, 2013; Hamlington et al., 2020; Fox-Kemper et al., 2021). Frederikse et al. (2020) showed  
that since 1900 the sum of the contributions to sea\_level change is consistent with the trends and multidecadal variability in  
30 observed ~~sea-level~~ sea level on both global and basin scales. However, significant uncertainties remain about the evolution of

hazards at the coastline generated by the marine environment. In this article, these hazards will be referred to as « coastal hazards », with the understanding, however, that the coast may be affected by other sources of hazard (intense precipitation, overflow of streams, etc.).

35 The issue of the evolution of coastal hazards is often addressed through the evolution of extreme sea levels (ESLs) along the coast. For example, Vitousek et al. (2017) warned that the 10 to 20 cm of sea level rise expected no later than 2050, will more than double the frequency of extreme water-level events in the tropics. Vousdoukas et al. (2018) projected a very likely increase of the global average 100-year ESL between 2000 and 2100 of 34–76 cm under a moderate-emission-mitigation-policy scenario and of 58–172 cm under a business as usual scenario. Under these scenarios, a large part of the tropics would  
40 be exposed annually to the present-day 100-year ESL in 2050. These assertions concern the evolution of water levels at the coast, but are not sufficient to describe the evolution of coastal hazards. Yet many authors evaluating changes in coastal risks restrict the representation of the hazard to ESLs (e.g. Hallegatte et al. (2013), Tiggeloven et al. (2020), Rasmussen et al. (2020)). This kind of analysis uses a deterministic approach based on a single factor (the ESL that is expressed by two parameters: maximum water level and frequency) to represent a complex hazard that can only be fully addressed using a  
45 systemic approach that considers multiple factors (Iggabel et al., 2021). In comparison, on land, systemic approaches are used in studying surface runoff and defining strategies in water resource and flood risk management (Shaikh et al., 2022; Verma et al., 2023; Mehta et al., 2023).

To answer this problem in the coastal domain, it is necessary to highlight that during a metocean event, the phenomena of  
50 flooding and coastal erosion are not only determined by the maximum sea level, but also by coastal waves and currents, and overtopping and overflow discharges over flood protection structures (Formentin et al., 2018; Iggabel et al., 2022). Estimating these discharges and hydrodynamic conditions for the duration of an event requires a good understanding of the physical phenomena that generate the hazard. Retrospective analyses of events help to understand correctly the mechanisms that cause the observed flooding or erosion. For example, by simulating Total Water Levels (TWLs) along the Bight extending from  
55 North Carolina to Florida during three historical Tropical Cyclones (TCs) with similar tracks, Hsu et al. (2023) found that the magnitude and duration of the increase in TWLs and wind waves are influenced by TC intensity, translational speed and distance from the shore. In particular, the authors showed that a decrease in TC translation speed led to longer exceedance durations of TWLs, which may result in larger impacts. However, it is not possible to predict deterministically the physical characteristics of future events, nor to assess the corresponding hydrodynamic conditions. To compensate for this, probabilistic  
60 approaches have been developed. For example, using a large number of synthetic hurricanes that consider the natural variability in hurricane frequency, size, intensity and track, Krien et al. (2015) estimate the 100-year and 1000-year surge levels for the archipelago of Guadeloupe. Following the same principle, Krien et al. (2017) estimate 100-year surge levels in Martinique for the present climate or considering a potential sea level rise. These results help to determine the necessary levels of protection structures in the short and medium term. However, a single parameter (the maximum water level) is not enough to characterize

65 the hazard and define all the crisis management measures, particularly when water levels exceed the level of protection or  
when protection structures fail (for example, breaches in levees or dunes). In addition, the accuracy of these estimates decreases  
in the long term due to: (i) high uncertainties in sea levels beyond 2050 (IPCC, 2019); (ii) the increase in the proportion of  
high-intensity cyclones worldwide (Masson-Delmotte et al., 2021); and (iii) environmental instability, notably because of  
geomorphological (e.g., subsidence, coastline retreat) and biological (e.g., degradation of coral reefs and mangroves) changes.  
70 These changes modify hydrodynamic and hydrosedimentary processes on the coast both in the long term and during individual  
events.

With the aim of making progress in the global assessment of coastal hazards in the long term, the guiding principle of this  
paper is to promote the use of the latest advances in research on changes in metocean events and water levels, while also  
75 showing the importance of studying other factors whose evolution is more predictable than storm surge and that may be equally  
important, namely: tidal regimes, geomorphological settings and environmental changes (particularly those modifying  
hydrodynamic conditions at the coast).

Although water levels are not the only parameter to consider, it is necessary to begin by clarifying the definitions of the  
80 different levels to which we will refer regularly.

First, it is necessary to distinguish between global mean sea level (GMSL) and relative sea level (RSL). The GMSL rise can  
be defined as the global change in mean sea level (MSL) relative to the terrestrial reference frame, due to the combined effects  
of change in the volume of the ocean and change in the level of the sea floor (Church et al., 2013). The RSL rise can be defined  
85 as the change in sea level with respect to the land (Lowe et al., 2010).

Second, it is necessary to assess the main factors influencing the evolution of water levels, whether they are GMSL, RSL or  
ESLs.

90 MSL changes, both globally and locally, vary according to seasonal, annual, and longer time scales. These variations may be  
caused by changes in the mass of water in the ocean (e.g., due to melting of glaciers and ice sheets and changes in terrestrial  
water storage), by changes in ocean water density (e.g., volume expansion as the water warms), by changes in the shape of the  
ocean basins (e.g., due to plate tectonics) and changes in the Earth's gravitational and rotational fields, and by local subsidence  
or uplift of the land (Oppenheimer et al., 2019). In the future, changes in the GMSL are expected to be strongly controlled by  
95 the greenhouse gas (GHG) emission scenarios (Oppenheimer et al., 2019; Fox-Kemper et al., 2021) and cryosphere evolution:  
in case of faster melting of the Greenland and Antarctic ice caps, Bamber et al. (2019) estimate the increase in GMSL above  
2 m in the 21st century.

The RSL will depend on both the GMSL and local vertical land movement (VLM). VLM can have natural causes (e.g. isostasy, elastic flexure of the lithosphere, earthquakes and volcanoes, landslides and sedimentation) and, more locally, anthropogenic causes, in particular, soil loading, extraction of hydrocarbons and/or groundwater, drainage, mining activities (Gregory et al., 2019). During the 20th century, the coasts of Tokyo, Shanghai, and Bangkok collapsed by several meters as a result of water extraction and soil loading from the construction of buildings and infrastructure (Nicholls et al., 2008). Subsidence rates of the order of 1 to 10 cm/year are commonly measured in coastal megacities and are exceeding sea level change rates by a factor of ten (Erkens et al., 2015). However, even in the absence of subsidence, the majority of coastlines are expected to experience RSL changes on the order of 30 percent of GMSL rise (Gregory et al., 2019; IPCC, 2019; Dayan et al., 2021).

ESLs are also projected to change due to changes in RSL, tides, wind waves, and storm surges (Vousdoukas et al., 2018). Storm surges are defined by Gregory et al. (2019) as « the elevation or depression of the sea surface with respect to the predicted tide during a storm ». The predominant factor in the evolution of ESLs is the variability in sea levels generated by astronomical tides and storm surges (Buchanan et al., 2016). The combination of tide and storm surge phenomena requires a statistical approach. Thus, the change in ESL events is commonly expressed in terms of an amplification factor and an allowance. The amplification factor denotes the amplification in the average occurrence frequency of a certain extreme event, often referenced to the water level with a 100-year return period estimated from historical data. The allowance denotes the increased height of the water level [m] with a given return period. This allowance equals the regional projection of RSL rise with an additional height related to the uncertainty in the projection (Hunter, 2012). Amplification factors are strongly determined by the local variability in ESL events. Locations where this variability is large due to large storm surges and astronomical tides will experience a moderate amplification of the occurrence frequency of extremes in comparison to locations with small variability. Globally, this contrast between regions with large and small amplification factors becomes clear for projections by mid-century and considerable in the coming centuries (Vitousek et al., 2017). In particular, many coastal areas in the lower latitudes may expect amplification factors of 100 or larger by mid-century, regardless of the GHG emissions scenario. By the end of the century, and in particular if GHG emissions are not reduced, such amplification factors may be widespread along global coastlines (Vousdoukas et al., 2018).

The objective of this paper is to present a comprehensive method to assess the evolution of coastal hazards at regional scales in the context of climate change. The proposed systemic method emphasizes the need to focus on the analysis and interpretation of the modeling results, by putting them into perspective with respect to the biophysical conditions (both current and forecasted).

This method was developed as the result of an empirical process, considering diverse situations. This process has the advantage of demonstrating the wide applicability of the proposed method. The case studies used to develop the method are in France and French territories, but they are located in different latitudes (equator, tropics and temperate zones) where they are exposed

135 to different climates and are characterized by different geomorphological configurations (including continental or island). This study highlights that considering the qualitative factors describing the geomorphological context, metocean events and the marine environment in a systemic approach was necessary to assess the evolution of hazards.

140 To ensure that the method can be applied for operational purposes, the use of freely accessible data has been promoted. However, applications of this method should include a thorough bibliographic analyse or even additional observational or modeling studies at the chosen case study sites. It is important to emphasize here that the application of the method depends strongly on the available data, and therefore it is necessary to gather the best data for each site. Since the quantity and quality of data is not the same everywhere, the uncertainties in the results will also vary and must be addressed.


145 Finally, following the application of the method to the case studies, the results obtained using this systemic approach can contribute to improving predictions of the evolution of different types of coastal hazards (shoreline erosion, rapid submersion and/or permanent flooding). This review of the evolution of sea levels (GMSL, RSL and ELS) highlights the importance of the evolution of coastal hazards. How will these hazards change with evolving water levels? How can other factors such as the geomorphological configuration of the coast, the types of metocean phenomena, and other changes in the marine environment be incorporated into the analyses?

150 The objective of this article is to present a systemic approach to assess the evolution of coastal hazards at regional scales. The method will be applied to France and its overseas territories, which are located in different latitudes (equator, tropics and temperate zones), exposed to different climates, and characterized by different geomorphological configurations (continental or island). Finally, as part of the discussion, the results will be put in perspective with the anticipated effects of climate change on the coast. This analysis allows identifying different types of hazard evolution in coastal areas.

## 155 2 Method for assessing changes in coastal hazards at regional scales

To assess the evolution of coastal hazards, at regional and multidecadal scales, it is necessary to combine a quantitative approach evaluating the parameters characterizing ESLs and a qualitative approach analyzing other factors more difficult to quantify, such as the meteo-oceanic event type, the geomorphological configuration of the coast and other environmental factors (e.g. coral reef degradation in the tropics and decrease in seasonal sea ice extent in the polar regions) that may change the hydraulic actions on the shoreline.

160 Figure 1 shows the three proposed steps. These three stages will be presented in succession. In applying the method, however, it is possible to implement the first two steps in parallel.



**Figure 1: Method for assessing changes in coastal hazards**

165 Before presenting these three steps, it is essential to remember that flood risk, and more generally, all coastal risks, can be defined in at least two alternative ways (FLOODsite, 2009). The first definition considers risk to be the result of the exposure of a vulnerable stake to a hazard, which is reflected in the following formula:

$$\text{risk} = \text{hazard (meteo-oceanic event)} * \text{exposure} * \text{vulnerability (of the society/area/structure)} \quad (1)$$

170 However, in an attempt to quantify risk, and considering that the word « risk » suggests a probability of occurrence, a second definition may be highlighted:

- the two terms « exposure » and « vulnerability » can be substituted by « consequences », with the consequences being generally more quantifiable (for example, in number of fatalities and economic damage) than the previous two terms;
- the hazard can be represented by its probability distribution.

175

This yields the second definition:

risk = probability (of the hazard) \* consequences

(2)

180 This article is mainly devoted to the evaluation of the hazard. Consequences (e.g. erosion and flooding) will only be addressed in the discussion. However, reference to risk definitions is necessary because the expression of hazard depends on the definition of risk.

185 Quantitative assessments of ESL changes (Step 1 of Fig. 1) are probabilistic and therefore fall within the second definition of risk. The challenge of the proposed method is to use these evaluations, in combination with other factors, such as the types of ~~meteocean~~metocean events, the geomorphological configuration of the coast and changes in the marine environment affecting hydraulic conditions. These « additional factors », which are more difficult to quantify, will be studied in a second, parallel phase (step 2 of Fig. 1). The last step is to describe the evolution of the hazard within the framework of a systemic approach. In this step (step 3 of Fig. 1), the first definition of risk is applied to search for a sufficiently synthetic way of presenting the data to produce an overall assessment.

190 Given the multiple effects of climate change on sea levels (GMSL and RSL), atmospheric conditions, and wave conditions, the evolution of the factors and parameters can only be assessed using global development hypotheses. To represent different trajectories, the IPCC identified five scenarios referred to as SSPx-y, where ‘SSPx’ refers to the Shared Socio-economic Pathway (‘SSP’ describing the socio-economic trends underlying the scenario), and ‘y’ refers to the approximate level of radiative forcing (in  $W\ m^{-2}$ ) resulting from the scenario in the year 2100. To determine the most relevant scenario and identify the most appropriate time horizon, an exposure and vulnerability analysis and an assessment of the duration of the project are required. These reflections are an essential prerequisite for the application of the proposed method.~~The application of the proposed method must be preceded by an analysis of exposure and vulnerability, as well as a reflection on the duration of the project. For long term adaptations, a general description of the evolution of the hazard, including the estimated uncertainties, is more appropriate than a forecast (likely imprecise) of the evolution of a parameter at a given date.~~

200 For the assessment of sea levels, Table 1 shows the projections for the GMSL under the SSP5-8.5 scenario at three time scales: 2050, 2100 and 2300.

205 **Table 1: GMSL projections for 2050, 2100 and 2300 for the SSP5-8.5 scenario. Median values and ranges (17th to 83rd percentiles) are shown using the 1986-2005 period as a reference (IPCC, 2019).**

<u>Climate scenario</u>	<u>2050 Median</u>	<u>2050 17-83% range</u>	<u>2100 Median</u>	<u>2100 17-83% range</u>	<u>2300 Median</u>	<u>2300 17-83% range</u>
<u>SSP5-8.5</u>	<u>+ 32 cm</u>	<u>23 cm to 40 cm</u>	<u>+ 84 cm</u>	<u>61 cm to 110 cm</u>	<u>+385 cm</u>	<u>230 cm to 540 cm</u>

210 The GMSL values displayed in Table 1 show high uncertainties in the long term. For example, in 2100, the median value is 0.84 m and the 17th to 83rd percentile range is 0.61 to 1.10 m (the high value is almost double the low value). These GMSL estimates were established for the SSP5-8.5 scenario, but greater differences exist if other scenarios are considered in a complementary manner. Similar (or higher, since other local phenomena must be considered) uncertainties exist for the RSL and the centennial ESL.

215 In conclusion, to make long-term management and adaptation decisions, a general description of the evolution of the hazard, including the estimated uncertainties, is considered more appropriate than a forecast of the evolution of a parameter at a given date, which is likely to be imprecise. Accordingly, conservative assumptions will be used when assessing sea level components in this method.

## 2.1 Step 1: ESL change assessment

220 Tidal simulations show no significant impacts of RSL rise on tidal amplitude during the 21st century at regional scales, although this does not exclude potential local effects (Pickering et al., 2012; Haigh et al., 2020). For example, Idier et al. (2017) showed that notable increases in high tide levels occur in the northern Irish Sea, the southern part of the North Sea and the German Bight, and decreases occur mainly in the western English Channel. Depending on the location, they can account for +/-15% of RSL rise (as long as RSL rise remains smaller than 2 m). Given the strong uncertainties of RSL rise, it is assumed in this study that the tides and tidal range remain constant.

225 ESLs are the result of the combination of the RSL, the tide level, and the storm surges produced by ~~meteo-oceanic~~ events (Calafat et al., 2022). To understand the evolution of ESLs, it is assumed that ~~meteo-oceanic~~ events simultaneously generate atmospheric storm surge (caused by atmospheric depressions, wind) and energetic waves. At the coast, waves also contribute to the total water level by two phenomena:

- 230
- wave set-up: a time-varying (on the order of 30 minutes), wave-driven increase in the mean water level at the coast, resulting from wave dissipation in the nearshore (Bowen et al., 1968);
  - run-up: the time-varying (on the order of the wave period, i.e. a few second, or the period of infragravity-waves, i.e. a few minutes) water level reached by a wave on a beach or a coastal structure, relative to the static water level, measured vertically (run-up may generate overtopping which, unlike overflowing, occurs intermittently).

235 In the proposed method, storm surge and wave set-up will be considered to estimate the ESL, but wave run-up will be neglected because of the challenge associated with modeling it at fine scales (Aucan et al., 2018; Gomes da Silva et al., 2020).

240 To fully account for changes in ESLs, the analysis must include both changes in the average occurrence frequency of a certain extreme event and the increased water level (in absolute terms and as a percentage) with a given return period. For this reason,



the second definition of risk will be used to study the joint probabilities of the parameters impacting the sea level, namely the RSL, the tidal range and the storm surge (including the effects of waves). Following Vousdoukas et al. (2017) changes in the magnitude and frequency of occurrence of the present 100-year ESL (ESL<sub>100</sub>) are evaluated. It is assumed that ESL are driven by the combined effect of MSL, tides ( $\eta_{tide}$ ) and water level fluctuations due to waves and storm surges ( $\eta_{w-ss}$ ). As a result,

245 ESL can be defined as:

$$ESL = MSL + \eta_{tide} + \eta_{w-ss} \quad (3)$$

We must here appreciate the need to use a more detailed method for each particular case, because according to Idier et al. (2019), depending on the type of environments (e.g., morphology, hydrometeorological context), non-linear interactions between various components - tide, surge, wave set-up - can reach several tens of centimeters. For instance, using a numerical modeling approach, focusing on the German Bight area (SE of North Sea) and assuming a sea level rise of 0.54 m, Arns et al. (2015) show that taking into account the interactions between mean sea level, tide and atmospheric surge leads to positive changes in extreme water levels relative to the MSL rise. The largest non-linear increases in the order of +0.15 m occur in the shallow areas of the Wadden Sea. However, introducing a novel approach to statistically assess the non-linear interaction of tide and non-tidal residual, Arns et al. (2020) demonstrate that extreme sea levels are up to 30% (or 70 cm) higher if non-linear interactions are not accounted for. The largest effects of Tide Surge Interaction (TSI) are found for the US East Coast and the Gulf of Mexico, the UK North Sea coastline, and parts of the southern Japanese coast. The highest value, 66 cm, is found in Cromer along the UK North Sea coastline. On the selected coasts, other values are generally between 20 and 50 cm. In conclusion, while statistical methods demonstrate that, globally, non-linear interaction modulates extreme sea levels, numerical modeling shows that non-linear interaction can locally induce increases up to tens of centimeters. Given the high uncertainties regarding the long-term evolution of sea levels and geomorphological changes, we shall consider that equation (3) provides in the general case a conservative estimate of the ESL and, if not, that the other sources of uncertainty are greater than the water level rise related to these non-linear interactions.

250

255

260

The climate extremes contribution  $\eta_{w-ss}$  from waves and storm surge can be estimated as:

$$265 \quad \eta_{w-ss} = SSL + 0.2 H_s \quad (4)$$

where SSL is the storm surge level,  $H_s$  is the significant wave height and  $0.2 H_s$  is a generic approximation of the wave set-up (U.S. Army Corps of Engineers, 2002). This equation is a conservative estimate of the wave set-up, which mostly depends on the local slope, breaking wave height, and wave period (Stockdon et al., 2006), and may be closer to 10% of the breaking wave height. Larger values may only be observed at steep sandy beaches (e.g. Martins et al., 2022) or steep shore platforms (Sheremet et al., 2014; Lavaud et al., 2022). In the current state of the art, numerical simulations of set-up have been carried out only locally (e.g., Lange et al. (2021), van Ormondt et al. (2021)). Such retrospective simulations are computationally expensive (e.g. to simulate accurately set-up, model grid resolution of ~10-50 m may be needed for study sites with large variations in local bathymetry and the wave field). In addition : (i) accurate and high resolution wave setup modeling requires high resolution bathymetric data (Stephens et al., 2011), which are not available at the spatial scales of our case study; (ii)

270

275 ~~significant morphological changes are expected in nearshore areas, especially with rising sea levels, which can have a~~  
~~significant effect on the wave set-up at different timescales (e.g. especially for sandy beaches exposed to waves: Ruggiero et~~  
~~al., 2001 ; Thiébot et al., 2012 ; Brivois et al., 2012). Thus, at large spatial (e.g. global) and long temporal (e.g. 21st century)~~  
~~scales, simplified models such as the one used here (Vousdoukas et al., 2018) may provide a first estimate of the expected~~  
~~wave set-up. This expression is used here since the objective of this work is not to improve existing deterministic methods, but~~  
280 ~~to present a systemic and comprehensive method for assessing the evolution of coastal hazards.~~

~~The tidal range may have a strong influence on the frequency of ESLs, in particular for microtidal coasts (strong amplification~~  
~~factor for low tidal ranges).~~

285 ~~Storm surge and wave setup affect the frequency of ESLs (strong amplification factor for low surges) and can also influence~~  
~~the allowance. For example, through a statistical analysis of tide gauge observations, Calafat et al. (2022) showed that trends~~  
~~in surge extremes and sea level rise both made comparable contributions to the overall change in ESLs in Europe since 1960.~~  
~~However, on the three maritime facades of France, the relative importance of this factor is expected to decline gradually in the~~  
~~21st century due to increases in RSL becoming predominant (Vousdoukas et al., 2017).~~

290 ~~In general, tidal simulations show no significant impacts of RSL rise on tidal amplitude during the 21st century at regional~~  
~~scales, although this does not exclude potential local effects (Haigh et al., 2020). For example, Idier et al. (2017) showed that~~  
~~notable increases in high tide levels occur in the northern Irish Sea, the southern part of the North Sea and the German Bight,~~  
~~and decreases occur mainly in the western English Channel. Depending on the location, they can account for +/-15% of RSL~~  
295 ~~rise (as long as RSL rise remains smaller than 2 m). Given the strong uncertainties on RSL rise, we will assume here that the~~  
~~tides are in a steady state and, consequently, that the tidal range does not change the allowance.~~

## 2.2 Step 2: Evaluation of additional factors

Two types of regional factors can be considered essential for all coastlines: phenomena likely to generate energetic sea states  
300 and the geomorphological configuration. In addition, other environmental factors affecting hydraulic conditions should be  
considered in the tropics (in particular, coral reef state) and at the poles (in particular, sea ice extent).

### 2.2.1 ~~Meteocean~~Metocean event types

Three main coast categories are considered in this method: exposure to no particular phenomenon, to storms, or to cyclones  
(which also implies exposure to storms). On these three categories are superimposed (i) the potential exposure to long swells  
305 generated by a ~~meteo-oceanic~~ event far from the studied coastline (ii) the potential exposure to tsunamis, since the origin of  
tsunamis is not related to the meteorological conditions.

The increase in wave damage could also be assessed, considering local changes in significant wave height (average height of the highest one-third of the waves in a given sea state). ~~However, trends in coastal wave climates are reported, with a low confidence level, in the IPCC (2019) report. Therefore, for example, projections of future extreme significant wave height consistently estimate an increase in the Southern Ocean and a decrease in the north-eastern Atlantic Ocean and Mediterranean Sea.~~ However, these trends will not be explored in detail: the focus will be on the strong differences that already exist between the wave climates of the maritime facades.

### 2.2.2 Geomorphological configurations of the coast

It is necessary to identify characteristics of the coast that influence the different components of the surges.

~~Storm surges are controlled not only by the characteristics of metocean events (e.g. for tropical cyclones (TC), the TC intensity, the distance to the TC eye, the TC heading direction and the TC translation (Hsu et al., 2023)), but also by the width of the continental shelf and the presence of shallow waters relative water depth (Kennedy et al., 2012). First, the analysis of storm surges computed from the national REFMAR database reveals that they are controlled not only by storm tracks but also by the width of the continental shelf and the presence of shallow waters.~~ For example, during the 20-year study period (1998-2018), storm surges hardly reached 1.0 m along the coastlines of the southern Bay of Biscay and the eastern Mediterranean Sea, but exceeded 2.0 m in the English Channel (Dodet et al., 2019). In shallow water, the wind effects can significantly dominate the effect of the atmospheric pressure, and the surge can be strongly amplified by the bathymetry along the coast (Bertin et al., 2012). The width and depth of the continental shelf have an important influence on the magnitude of storm surges.

Secondly, in regions exposed to cyclones, the surges measured on islands are much lower than those measured on continental facades. For example, on the islands of the West Indies, the storm surges rarely exceed 3 m (Krien et al., 2015 and Krien et al., 2017) ~~Cerema, 2019, 2020, 2021~~, whereas in the case of Katrina, which impacted the United States in 2005, surges in eastern Louisiana reached values between 3.05 m and 5.79 m (Graumann, 2006). During the same event, the surges exceeded 8 m at several locations along the Mississippi coast (Dietrich et al., 2010). This difference between the surges observed on continental coasts and island coasts is explained by the fact that, in the case of a hurricane, the impact of the low pressure associated with the storm on surge is minimal in comparison to the water being forced toward the shore by the wind. However, in the case of small islands (e. g. West Indies or La Réunion), the surge is generally reduced: wind-forced water accumulates less on short shorelines than on long shorelines (Durand, 1996). Therefore, the length of the maritime facade appears to be the second determining factor of the surges.

Thirdly, the maximum potential storm surge for a particular location is sensitive to small changes in storm intensity, forward speed, size, angle of approach to the coast, central pressure, and also the shape and characteristics of coastal features such as

340 bays and estuaries ([Flather, 2001](#); [Rego and Li, 2010](#) and [Kennedy et al., 2012](#))<sup>†</sup>: Wind-forced water will accumulate less on convex-shaped shorelines than on concave-shaped shorelines ([Bertin et al., 2012](#); [Krien et al., 2015](#); [Krien et al., 2017](#)). Therefore, the shape of the coast appears to be the third determining factor of the surges.

Consequently, in order to account for the main geomorphological parameters determining the magnitude of the surge, we  
345 propose to use three criteria:

- the length of the coast;
- the width of the continental shelf;
- the convex or concave shape of the coastline.

### 2.2.3 Other environmental factors changing hydraulic conditions

350 Other phenomena induced by climate change can increase the intensity and frequency of extreme events, or increase their effects:

- ocean acidification combines with ocean warming and deoxygenation to impact ecosystems (e. g., coral reefs and oyster beds), which can reduce coastal flood protection. Under conditions expected in the 21st century, global warming and ocean acidification ~~should~~ may compromise carbonate accretion, with corals becoming increasingly  
355 rare on reef systems. The result should be less diverse reef communities and carbonate reef structures that fail to be maintained. Climate change also exacerbates local stresses from declining water quality and overexploitation of key species, driving reefs increasingly toward the tipping point for functional collapse ([Hoegh-Guldberg et al., 2007](#); [Albright et al., 2018](#));
- in polar regions, the decrease in seasonal sea ice extent in the Arctic, together with a longer open water season, provide  
360 less protection from storm impacts ([Forbes, 2011](#)). In addition, a longer fetch allows more energetic waves to form, which can have consequences outside of the polar regions, for example on coasts exposed to southern swells.

These two factors that affect wave formation and propagation, as well as currents, are also considered in the proposed method.

### 2.3 Step 3: global hazard assessment

365 ~~In order to clearly distinguish the current approach from existing analytical approaches (e.g., [Vitousek et al., 2017](#); [Vousdoukas et al., 2018](#)), it is important to note that RSL changes, tidal regimes and storm surge and wave impacts not only influence the evolution of coastal hazards through the sea level alone, but also can have an additional influence by the energy carried by waves and currents. The data used to draw up an overall summary should therefore not be limited to variations in the height and frequency of the ELSs. It is also necessary to incorporate the parameters necessary for their evaluation, and the other~~

---

<sup>†</sup><https://www.nhc.noaa.gov/surge/>

370 ~~factors identified, namely the meteocean event type, the geomorphological configuration, and the other environmental factors~~  
~~changing hydrodynamics at the shoreline.~~

~~Based on this principle~~ Following step 1 and step 2, the overall assessment of the hazard should be produced using the following factors and parameters (evaluated on the basis of the SSP5-8.5 scenario):

- 375
- RSL height change: ( $\Delta$ RSL);
  - tidal range;
  - surge with a 100-year return period;
  - ESL height variation ( $\Delta$ ESL);
  - ESL percentage change ( $\% \Delta$ ESL);
- 380
- return period of the present day 100-year ESL;
  - geomorphological configuration of the coast;
  - ~~meteocean~~ metocean event type; and
  - other environmental factors affecting the hydrodynamics at the shoreline.

385 In this method, each factor should not be considered independently. On the contrary, the joint effects of the different factors should be assessed to understand the dynamics of the system. However, the diversity of the different observed contexts prevents proposing a quantitative calibration and weighting system. This is the main reason why this method is called « systemic », in reference to the study of a system (the coast). However, this method should not be called « systematic ». Even though the analysis framework incorporates multiple factors, the expert in charge of the study of a particular site should consider the qualitative and quantitative factors together. In summary, this method aims to help experts to make a structured evaluation.

### 390 **3 Application of the proposed method to French coasts**

The proposed method is applied to France and its overseas coasts (cf. Fig. 2). In the following, the islands of Saint-Martin, Saint-Barthélemy, Guadeloupe and Martinique may be referred to as the « West Indies ».




Figure 2: A map showing the French territories considered in this study. *Credit : Cerema, after Hoshie.*

In the following, the islands of Saint Martin, Saint Barthélemy, Guadeloupe and Martinique may be referred to as the « West Indies ».

The choice of the scenario and the time scales should ~~take into account~~ consider the existence of the high human and economical issues~~the high stakes~~ at the considered-studied coasts (coastal cities, port and industrial facilities) and the strong uncertainties about the contributions of ice caps to the rise in water levels (Bamber et al., 2019; Dayan et al., 2021). For these two reasons, we use the SSP5-8.5 scenario. ~~For the assessment of sea levels (GMSL, RSL and ESL), the median values are selected at two time scales: 2050 and 2100. Table 1 shows the projections for the GMSL under the SSP5-8.5 scenario.~~

~~Table 1: GMSL projections for 2050, 2100 and 2300 for the SSP5-8.5 scenario. Median values and ranges for the 17th to 83rd percentiles are shown using the 1986-2005 period as a reference (IPCC, 2019).~~

Climate scenario	2050	2050	2100	2100	2300	2300
	Median	17-83% range	Median	17-83% range	Median	17-83% range
SSP5-8.5	+32 cm	23 cm to 40 cm	+84 cm	61 cm to 110 cm	+385 cm	230 cm to 540 cm

~~The GMSL values displayed in Table 1 show high uncertainties in the long term: for the SSP5-8.5 scenario in 2100, the median value is 0.84 m and the 17th to 83rd percentile range is 0.61 to 1.10 m (the high value is almost double the low value). Similar (or higher, since other local phenomena must be considered) uncertainties exist for the RSL and the centennial ESL.~~

### 415 3.1 ~~±~~ESL change assessment

The evolution of ESLs is assessed using the three components (Eq. 3): (1) RSL, (2) tidal range and (3) combined storm surge and wave setup. The framework developed by Vousdoukas et al. (2018) is used to evaluate the contributions of each of these components, of which the baseline values are calculated in global reanalyses of waves and storm surges. Then, CMIP5 models are used to estimate future relative changes to the meteorological water levels. Lastly, changes in sea level, the astronomical tide, and meteorological water levels are combined to produce the ESL values.~~Projections of waves and storm surges were based on hydrodynamic simulations driven by atmospheric forcing from six Coupled Model Intercomparison Project Phase 5 (CMIP5) climate models.~~

#### 3.1.1 RSL: projections

Table 2 shows the median values of the RSL rise projections for the period 1995-2014 at selected points along the French coastline. For the SSP5-8.5 scenario, the median values of the RSL rise on the French coasts are fairly uniform: between 0.16 m and 0.20 m in 2050, and between 0.76 m and 0.92 m in 2100.

**Table 2: Projections of RSL rise at selected points along the French coastline compared to the period 1995–2014 for the SSP5-8.5 scenario**

	Projection of RSL rise (m)	
	2050	2100

Calais	0.19	0.86
Le Havre	0.19	0.87
Saint-Malo	0.19	0.87
Brest	0.19	0.83
La Rochelle	0.16	0.76
Saint-Jean-de-Luz	0.17	0.79
Port Vendres	0.16	0.76
Sète	0.16	0.76
Marseille	0.17	0.78
Saint-Pierre (Saint-Pierre-et-Miquelon)	0.19	0.83
Pointe-à-Pitre (Guadeloupe)	0.20	0.90
Cayenne (French Guiana)	0.20	0.90
Pointe des Galets (La Réunion)	0.19	0.92
Papeete (French Polynesia)	0.20	0.91

430

	<u>Geographic coordinates</u>		<u>Projection of RSL rise (m)</u>	
	<u>Latitude</u>	<u>Longitude</u>	<u>2050</u>	<u>2100</u>
<u>Calais</u>	<u>50.972122801049395</u>	<u>1.8400588679271384</u>	<u>0.19</u>	<u>0.86</u>
<u>Le Havre</u>	<u>49.485774012966544</u>	<u>0.0897840202510471</u>	<u>0.19</u>	<u>0.87</u>
<u>Saint-Malo</u>	<u>48.641170797005124</u>	<u>-2.0313888026241402</u>	<u>0.19</u>	<u>0.87</u>
<u>Brest</u>	<u>48.368792381060274</u>	<u>-4.4887286031866935</u>	<u>0.19</u>	<u>0.83</u>
<u>La Rochelle</u>	<u>46.14879125637811</u>	<u>-1.1691588857635562</u>	<u>0.16</u>	<u>0.76</u>
<u>Saint-Jean-de-Luz</u>	<u>43.39842588942395</u>	<u>-1.676715829943722</u>	<u>0.17</u>	<u>0.79</u>
<u>Port Vendres</u>	<u>42.52411089711506</u>	<u>3.1143835886292806</u>	<u>0.16</u>	<u>0.76</u>
<u>Sète</u>	<u>43.39330679508365</u>	<u>3.699492031443802</u>	<u>0.16</u>	<u>0.76</u>
<u>Marseille</u>	<u>43.29569227471328</u>	<u>5.352448215467127</u>	<u>0.17</u>	<u>0.78</u>
<u>Saint-Pierre (Saint-Pierre-et-Miquelon)</u>	<u>46.786272435631275</u>	<u>-56.16190646722868</u>	<u>0.19</u>	<u>0.83</u>
<u>Pointe-à-Pitre (Guadeloupe)</u>	<u>16.23300045952869</u>	<u>-61.53571250198115</u>	<u>0.20</u>	<u>0.90</u>
<u>Cayenne (French Guiana)</u>	<u>4.93572687841612</u>	<u>-52.340676954198194</u>	<u>0.20</u>	<u>0.90</u>



<u>Pointe des Galets (La Réunion)</u>	<u>-20.936178918145654</u>	<u>55.280686667086655</u>	<u>0.19</u>	<u>0.92</u>
<u>Papeete (French Polynesia)</u>	<u>-17.53479287238126</u>	<u>-149.58674796473545</u>	<u>0.20</u>	<u>0.91</u>

### 3.1.2 Tidal range

Using the standard tidal classification system [\(Masselink and Short, 1993\)](#), the French coastline can be divided into three categories based on the tidal range data provided by SHOM (source: Data.shom.fr):

- microtidal coastline (tidal range < 2 m): the coasts of the Mediterranean with a tidal amplitude between 0.2 and 0.5 m and the coasts of the West Indies, La Réunion and French Polynesia where the tidal amplitudes are less than 1 m; in addition, the coasts of Saint-Pierre-et-Miquelon, where the amplitude reaches 1.7 m;
- mesotidal coastline (tidal range between 2 and 4 m): the coasts of French Guiana (the maximum amplitude reaches 2.9 m);
- macrotidal coastline (tidal range > 4 m): on the coasts of the Atlantic, English Channel and North Sea, with significant differences shown in Fig. 3.




Figure 3: Maximum tidal range (source: Data.shom.fr)

### 445 3.1.3 Waves and storm surges

Information on surges will be presented for France and its overseas territories.

In Metropolitan France, there are significant variations along the coastline, as shown by the estimates of the 100-year return period storm surge presented in Table 3.

**Table 3: Estimate of the 100-year return period storm surge, calculated as the average value of statistical adjustments of a Pareto distribution (GPD) and an exponential law (Cerema, 2018) applied to storm peaks. The values of Calais, Le Havre and La Rochelle have been corrected to ~~take into account~~ consider historical events and the comparative study of frequency analysis approaches for extreme storm surges (Hamdi et al., 2014).**

Tide gauge	100-year return period surge (m)
Calais	1.6
Le Havre	1.6
Saint-Malo	1.1
Brest	1.0
La Rochelle	1.7
Saint-Jean-de-Luz	0.6
Port-Vendres	0.9
Sète	1.1
Marseille	1.3

Although the relative differences are significant, they remain small in absolute terms, ~~since these coasts are only exposed to storms and not to cyclones~~. As mentioned in the presentation of the method, the large width of the continental shelf in the North Channel and the positioning at the center of the gulf or bay (e.g., Le Havre, Saint Malo, La Rochelle, Sète and Marseille) or in a strait (e.g., Calais) explain the higher values observed at these locations.

Overseas, the Cerema (2019, 2020, 2021) provides information on the storm surge measured, observed and modelled in La Réunion, French Guiana, Martinique, Guadeloupe, Saint-Martin and Saint-Barthélemy and Saint-Pierre-et-Miquelon. Surges measured by tide gauges commonly reach values of 0.5 to 1 m. Nevertheless, these measurements are often taken in sheltered areas (e.g., ports). Additional increases in the water level along the open coast may be more or less significant depending on the geomorphological configuration. For example, during the passage of Cyclone Irma on September 6, 2017, an instantaneous surge of 2.0 m was measured at the Saint-Martin tide gauge<sup>2</sup>. The surge, modeled by Météo-France, was more than 3 m on the northern coasts of Saint-Martin (Marigot Bay, Grand Case) and Gustavia (Saint-Barthélemy), but hardly more than 1.2 m on the island's almost straight coastline (De la Torre ~~Y.~~, 2017). Similarly, in September 1989, during the passage of Cyclone Hugo on the Guadeloupe archipelago, there was evidence that the ~~sea level~~ sea level had increased by 2 to 3 m along the coast (Pagney, 1991; Saffache et al., 2003; Krien et al., 2015).

Finally, on the islands of La Reunion and French Polynesia, in addition to cyclones, southern swells represent one of the main hazards (Lecacheux et al., 2012). Continuous water height and wave measurements from the French national observation

<sup>2</sup> [https://data.shom.fr/donnees/refmar/SAINT\\_MARTIN](https://data.shom.fr/donnees/refmar/SAINT_MARTIN)

services ReefTEMPS (<https://www.reeftemps.science/>) and DYNALIT (<https://www.dynalit.fr/>) show that on coastlines directly exposed to swell (e.g. the Hermitage in La Réunion), 10-year return period events generate, at the breaking point, waves with  $H_s \sim 7$  to 8 m, which induce a 1.2 m setup. Infragravity waves with  $H_s \sim 1$  m can be superimposed on this set-up (Bertin et al., 2020). These values are lower in ports (e.g. Saint Gilles).

475 ~~This~~ ~~These~~ ~~information~~ ~~observations~~ confirms the need, as a complement to the measurement and modeling of surges, to study the geomorphological configuration of the coastline in order to understand the very significant differences that may appear.

### 3.1.4 ESL projection (intensity and frequency)

The framework developed by Vousdoukas et al. (2018) provides estimates of the ESL allowance ~~along overseas coasts of for~~ France ~~and its overseas territories~~ (Table 4). The results obtained in 2100 for the SSP5-8.5 scenario shows that the ESL increase in absolute value is relatively homogeneous on the various maritime facades ~~of metropolitan France~~: 0.89 to 1.00 m for the English Channel, 0.74 to 0.77 m for the Bay of Biscay and 0.75 to 0.78 m for the western Mediterranean. However, the percentage variations are very different, ranging between: 16 and 22% for the English Channel, 20 and 24% for the Bay of Biscay and 52 and 63% for the western Mediterranean, in 2100. The increase will therefore be much more noticeable in the Mediterranean.

485 ~~Along overseas coasts, the estimated ESL increase (absolute value) is relatively homogeneous, ranging between 0.16 and 0.24 m in 2050 and 0.85 and 0.95 m in 2100. The percent change in ESL is expected to be in the range of 8% to 18% by 2050, which is likely to be noticeable in terms of hazard intensity, particularly in Guadeloupe and French Polynesia, where the relative increases are estimated to be the largest. By 2100, the increases are projected to be around 50 to 75% (with the exception of Saint-Pierre-et-Miquelon).~~

490

**Table 4: Summary of the projected absolute and relative changes of the 100-year event ESL ( $\Delta$ ESL and % $\Delta$ ESL) ~~for the SSP5-8.5 scenario under SSP5-8.5~~, during the years 2050 and 2100.**

	SSP5-8.5 - 2050		SSP5-8.5 - 2100	
	$\Delta$ ESL (m)	% $\Delta$ ESL	$\Delta$ ESL (m)	% $\Delta$ ESL
Calais	0.21	4.3	0.94	19.2
Le Havre	0.23	4.4	1.00	19.6
Saint-Malo	0.21	3.8	0.89	16.0
Brest	0.22	5.0	0.96	21.5
La Rochelle	0.16	4.3	0.77	19.9
Saint-Jean-de-Luz	0.16	5.3	0.74	24.1
Port Vendres	0.16	13.3	0.75	60.7
Sète	0.18	12.5	0.76	52.5
Marseille	0.18	14.6	0.78	63.2

<u>Saint-Pierre (Saint-Pierre-et-Miquelon)</u>	<u>0.24</u>	<u>8.4</u>	<u>0.92</u>	<u>31.8</u>
<u>Pointe-à-Pitre (Guadeloupe)</u>	<u>0.22</u>	<u>14.1</u>	<u>0.91</u>	<u>58.1</u>
<u>Cayenne (French Guiana)</u>	<u>0.18</u>	<u>10.6</u>	<u>0.85</u>	<u>49.2</u>
<u>Pointe des Galets (La Réunion)</u>	<u>0.16</u>	<u>9.3</u>	<u>0.88</u>	<u>52.5</u>
<u>Papeete (French Polynesia)</u>	<u>0.23</u>	<u>17.8</u>	<u>0.95</u>	<u>74.7</u>

495 The analysis of ESL height changes should be supplemented by a frequency analysis based on Table 5. The forecasts show that sea levels with a 100-year return period could occur on an annual basis by the end of this century in mainland France (except in the center of the Atlantic facade, around La Rochelle). Some regions are projected to experience an even higher increase in the frequency of occurrence of ESL, most notably along the Mediterranean, where the present day 100-year ESL is projected to occur about ten times a year. The higher increase in the Mediterranean is closely related to the low variability of ~~sea level~~ sea levels on these microtidal coastlines (Vitousek et al., 2017).

500 For the overseas coasts, in 2100, the present day 100-year ESL is projected to occur more frequently, approximately ten times a year, except for Saint-Pierre-et-Miquelon where this frequency would be about three times a year. The increase in frequency is likely to be noticeable in 2050, especially in French Guiana, French Polynesia and Saint-Pierre-et-Miquelon, where the return periods of the present day 100-year ESL are estimated to be only 7, 16 and 18 years, respectively.

505 **Table 5: Return period (in years) of the present day 100-year ESL for the SSP5-8.5 scenario under SSP5-8.5 in the years 2050 and 2100.**

	Projected return period of the current 100-year ESL in:	
	2050	2100
Calais	22.81	0.75
Le Havre	26.56	0.87
Saint-Malo	27.69	0.81
Brest	20.61	0.73
La Rochelle	36.16	2.59
Saint-Jean-de-Luz	33.87	0.63
Port Vendres	27.89	0.10
Sète	30.02	0.56
Marseille	26.88	0.10
<u>Saint-Pierre</u> <u>(Saint-Pierre-et-Miquelon)</u>	<u>17.69</u>	<u>0.31</u>
<u>Pointe-à-Pitre (Guadeloupe)</u>	<u>35.14</u>	<u>0.10</u>
<u>Cayenne (French Guiana)</u>	<u>7.54</u>	<u>0.10</u>

<u>Pointe des Galets (La Réunion)</u>	<u>57.53</u>	<u>0.10</u>
<u>Papeete (French Polynesia)</u>	<u>15.86</u>	<u>0.10</u>

The framework developed by Vousdoukas et al. (2018) provides estimates of the ESL allowance along overseas coasts (cf. Table 6). The estimated ESL increase (absolute value) is relatively homogeneous for the overseas coasts, ranging between 0.16 and 0.24 m in 2050 and 0.85 and 0.95 m in 2100. The percent change in ESL is expected to be in the range of 8% to 18% by 2050, which is likely to be noticeable in terms of hazard intensity, particularly in Guadeloupe and French Polynesia, where the relative increases are estimated to be the largest. By 2100, the increases are projected to be around 50 to 75% (with the exception of Saint Pierre et Miquelon).

**Table 6: Summary of the projected absolute and relative changes of the 100-year event ESL ( $\Delta$ ESL and % $\Delta$ ESL) for the SSP5-8.5 scenario, during the years 2050 and 2100.**

	SSP5-8.5-2050		SSP5-8.5-2100	
	$\Delta$ ESL (m)	% $\Delta$ ESL	$\Delta$ ESL (m)	% $\Delta$ ESL
Saint Pierre (Saint Pierre et Miquelon)	0.24	8.4	0.92	31.8
Pointe à Pitre (Guadeloupe)	0.22	14.1	0.91	58.1
Cayenne (French Guiana)	0.18	40.6	0.85	49.2
Pointe des Galets (La Réunion)	0.16	9.3	0.88	52.5
Papeete (French Polynesia)	0.23	17.8	0.95	74.7

For the overseas coasts, the present day 100-year ESL is projected to occur more frequently, approximately ten times a year, except for Saint Pierre et Miquelon where this frequency would be about three times a year (Table 7). The increase in frequency is likely to be noticeable in 2050, especially in French Guiana, French Polynesia and Saint Pierre et Miquelon, where the return periods of the present day 100-year ESL are estimated to be only 7, 16 and 18 years, respectively.

**Table 7: Return period (in years) of the present day 100-year ESL for the SSP5-8.5 scenario in the years 2050 and 2100.**

	2050	2100
Saint Pierre (Saint Pierre et Miquelon)	17.69	0.31
Pointe à Pitre (Guadeloupe)	35.14	0.10
Cayenne (Guyane)	7.54	0.10
Pointe des Galets (La Réunion)	57.53	0.10
Papeete (French Polynesia)	15.86	0.10

## 3.2 Evaluation of additional factors

Two regional factors can be considered essential for all coastlines: the type of meteocean event and the geomorphological configuration. In addition, other environmental factors affecting the hydraulic conditions should also be considered at the tropics and poles.

### 530 3.2.1 ~~Meteocean~~Metocean event types

For metropolitan France, the main type of metocean event affecting the selected coasts is storm. It is important to note, however, that tropical-like storms in the Mediterranean Sea, also called « Mediterranean hurricanes » or « medicanes », are occasionally observed. These storms sometimes attain hurricane intensity and can severely affect the coast (Fita et al., 2007). More frequent and higher intensity extreme values are present in the south of France (Patlakas et al., 2020). Classifying the intensity of storms and cyclones as "weak", "moderate" or "severe", Romero and Emanuel (2016) project a higher number of moderate and severe medicanes (at the expense of weak storms) and an increased occurrence of medicanes in the western Mediterranean in the period (2081–2100) compared to the period (1986–2005).

540 Overseas, the metocean event types are storms and cyclones ~~for~~in the West Indies (Krien et al., 2015; Krien et al., 2017) and Saint-Pierre-et-Miquelon (Catto and Batterson, 2011; Han et al., 2012; Masson, 2014), and ~~;~~ storms, cyclones, long swells and tsunamis (moderate exposure) for La Réunion (Lecacheux et al., 2012; Sahal and Morin, 2012; Quentel et al., 2013; Allgeyer et al., 2017) and French Polynesia (Larrue and Chiron, 2010). In Guiana, no phenomenon generating energetic sea states are identified.

### 3.2.2 Geomorphological configuration of the coast

545 The geomorphological configuration of each coast is evaluated in Table 6 using three criteria:

- the length of the coast,
- the width of the continental shelf, and
- the convex or concave shape of the coastline.

550 In the framework of a systemic approach, all of the available knowledge of the study sites is used to assess the surge potential associated with the geomorphological configuration. This knowledge includes information from bibliographical references and data obtained from bathymetric maps provided by the SHOM (Table 6) (Table 8) allows estimating qualitatively the surge potential associated with each geomorphological configuration.

555 Table 68: Assessment of the surge potential associated with the geomorphological configuration of the study sites (source: Data.shom.fr)



	<u>Bibliographical references</u>	Order of magnitude of coastline length (km)	Width of the continental shelf	Shape of the coastline	Surge potential associated with the geomorphological configuration
English Channel – North Sea	<u>Le Gorgeu and Guitonneau (1954), Hequette (2010), Bardet et al. (2011), Haigh et al. (2011), Weisse et al. (2012), Idier et al. (2012), Maspataud et al. (2013), Hamdi et al. (2014), Vousdoukas et al. (2016), Latapy et al. (2017), Vousdoukas et al. (2017), Hamdi et al. (2018), and DREAL Nord Pas-de-Calais (2024)</u>	1000	Very large (>200 km)	Alternation of concave and convex shapes	Very high
Bay of Biscay	<u>Allgeyer et al. (2013), Bertin et al. (2014), Hamdi et al. (2014), Bertin et al. (2015), Hamdi et al. (2015), Bulteau et al. (2015), Vousdoukas et al. (2016) and Vousdoukas et al. (2017), Garnier et al. (2018), and Khan et al. (2023).</u>	1000	Large (100 to 200 km)	Concave	High
<del>West</del> Mediterranean	<u>Ullmann et al. (2007), Fita et al. (2007), Campins et al. (2011), Conte and Lionello (2013), Cavicchia et al. (2014), Androulidakis et al. (2015), Vousdoukas et al. (2016), Romero and Emanuel (2016), Vousdoukas et al. (2017), Muis et al. (2020), Elkut et al. (2021), Patlakas et al. (2021), and Toomey et al. (2022).</u>	1000	Normal to the West (~ <del>around</del> 50 km), nonexistent to the East	Concave to the West and convex to the East	High to the West Low to the East
Saint-Pierre-et-Miquelon	<u>Catto and Batterson (2011), Han et al. (2012) and Masson (2014)</u>	10	Very large (>200 km)	Alternation of concave and convex shapes	High
Guadeloupe	<u>Pagney (1991), Zahibo et al. (2007), Dorville and Zahibo (2010), Lin and Chavas (2012), and Krien et al. (2015)</u>	5 to 20	Narrow (<20 km)	Alternation of concave and convex shapes	Low
French Guyana	<u>Gratiot et al. (2007), Chevalier et al. (2008), and Thiéblemont et al. (2023)</u>	1000	Large (~ <del>about</del> 100 km)	<del>Linear</del> <u>Mainly linear</u> or slightly convex, <del>except for estuaries</del>	Moderate



La Réunion	<a href="#">Lecacheux et al. (2012)</a> , <a href="#">Sahal and Morin (2012)</a> , <a href="#">Quentel et al. (2013)</a> , and <a href="#">Allgeyer et al. (2017)</a>	20	Nonexistent	Mainly linear or convex ( <del>except 2 bays in the North West</del> )	Very low
French Polynesia	<a href="#">Pirazzoli and Montaggioni (1988)</a> , <a href="#">Aubanel et al. (1999)</a> , <a href="#">Larrue and Chiron (2010)</a> , <a href="#">Webb and Kench (2010)</a> , <a href="#">Becker et al. (2012)</a> , <a href="#">Yates et al. (2013)</a> , <a href="#">Le Cozannet et al. (2013)</a> , <a href="#">Martinez-Asensio et al. (2019)</a> , and <a href="#">Barriot et al. (2023)</a>	1 to 20	Nonexistent	Mainly linear or convex ( <del>with many small-scale exceptions</del> )	Very low

In metropolitan France, the geomorphological configurations (i.e. long coastline, wide continental shelf, linear and/or concave shape) generally favor the development of substantial surges. The main exception is the eastern Mediterranean facade, which has convex-shaped coasts without a continental shelf (extension in the sea of the Alpine massif). Another exception is the southern part of the Bay of Biscay, where the concave shape and the length of the coastline favour the surge, while the small width of its continental shelf is a limiting factor.

For the overseas territories, the analysis reveals very contrasting situations:

- Guiana has a broad continental shelf, but its coastline is linear or convex, which is less conducive to surges;
- Saint-Pierre-et-Miquelon is in a situation similar to that of metropolitan France, with a wide continental shelf and the alternation of convex and concave forms of the coast. Although the length of the coastline is smaller, the close proximity of the large island of Newfoundland, which has a 200 km coastline, can contribute to large surges;
- The other islands are far from continents and have continental shelves that are narrow (e.g., Guadeloupe) or non-existent (e.g., La Réunion and French Polynesia), which is, with the short coastline length, a factor limiting the surge magnitude. For La Réunion and French Polynesia, the surges are even more limited by the linear or convex shape of the coastline.

In summary, considering only the geomorphological configuration, the coasts of metropolitan France (except the east of the Mediterranean facade and the Southern part of the Bay of Biscay), French Guiana and Saint-Pierre-et-Miquelon have all the characteristics favoring large surges. Smaller surges are expected on all criteria in La Réunion and French Polynesia, and based on all criteria except the shape of the coast in the West Indies.

### 3.2.3 Other environmental factors influencing the hydraulic conditions

La Réunion and French Polynesia are exposed to long southern swells and will therefore experience direct changes in the swell climate related to the decrease in seasonal sea ice extent.

580 The islands of the West Indies, La Réunion and French Polynesia are bordered by coral reefs whose degradation by ocean changes resulting from climate change (e.g. increases in water temperature and ocean acidification) may be a major factor in aggravating coastal hazards.

### 3.3 Integrated hazard evolution analysis

585 The evolution of coastal hazards can be assessed by considering both the quantitative sea level change estimates and the qualitative geomorphological configuration, ~~meteocean~~metocean event and environmental factor analysis for France (Table 97) and its overseas territories (Table 108).

#### 3.3.1 Hazard evolution in the European territory of France

590 **Table 79: Summary of the main factors influencing the hazard evolution on the mainland France's coasts. The data relative to ESLs refer to a present day 100-year ESL. The reference period for the RSL projection is 1995-2014. The reference period for the ESL projection is 1980-2014. The ~~slightly detrimental least important~~ factors of the evolution of the hazard are in italics, the ~~important detrimental~~ factors in bold and the ~~highly detrimental most important~~ factors in capital and bold. In the first column, these same fonts describe the estimated global evolution of the hazard for each coast.**

	Surge potential associated with the geomorphological configuration	Reference <del>meteocean</del> metocean event	Tidal range	100-year surge (m)	$\Delta$ RSL (m) under SSP5-8.5	$\Delta$ ESL (m) under SSP5-8.5	% $\Delta$ ESL under SSP5-8.5	Return period of the present day 100-year ESL under SSP5-8.5 (year)
<b>English Channel – North Sea</b>	<b>VERY HIGH</b>	Storm	<i>Macrotidal (7 to 14 m)</i>	1.0 to 1.6	In 2050 : 0.19 In 2100 : 0.83 to 0.87	In 2050 : 0.21 to 0.23 In 2100 : 0.89 to 1.00	In 2050 : 3.8 to 5 In 2100 : 16 to 21.5	<b>In 2050 : 20 to 28</b> <b>In 2100 : 0.73 to 0.87</b>
<b>Bay of Biscay</b>	<b>High to the North</b>	Storm	<i>Macrotidal (5 m to 7 m)</i>	0.6 to 1.7	In 2050 : 0.16 to 0.17 In 2100 : 0.76 to 0.79	In 2050 : 0,16 In 2100 : 0.74 to 0.77	In 2050 : 4.3 to 5.3 In 2100 : 19.9 to 24.1	<b>In 2050 : 33 to 36</b> <b>In 2100 : 0.63 to 2.6</b>
	Moderate to the South							
<b>WEST MEDITERRANEAN</b>	<b>High to the West</b>	<b>Storm and moderate exposure to cyclone</b>	<b>MICROTIDAL (&lt; 2 m)</b>	0.9 to 1.3	In 2050 : 0.16 to 0.17 In 2100 : 0.76 to 0.78	In 2050 : 0,16 to 0.18 In 2100 : 0.75 to 0.78	<b>IN 2050 : 12.5 TO 14.6</b> <b>IN 2100 : 52.5 TO 63.2</b>	<b>IN 2050 : 26 TO 30</b> <b>IN 2100 : 0.10 TO 0.56</b>
<b>East Mediterranean</b>	<i>Low to the East</i>							

595 By 2100, for the SSP5-8.5 scenario, the increases in RSL and ESL on the three maritime facades are rather similar in absolute terms, with median values between 0.74 and 1.0 m. The effect of these increases is nevertheless significantly different for each maritime facade, depending on the type of ~~meteocean~~metocean event, tidal range, and surge magnitude:

- In the English Channel and the North Sea, the macrotidal range and storm occurrence produce high variability of sea levels. The increase of 0.9 to 1.0 m of the 100-year ESL represents only 16 to 22%. The allowance is therefore relatively low. On the other hand, the increase in the RSL is sufficient for a present day 100-year ESL to have a return period of less than one year (amplification factor greater than 100);
- In the Bay of Biscay, the situation is quite similar, but with a smaller tidal range and surge magnitude (especially on the coast of the Atlantic Pyrenees, where the maximum tides are in the order of 5 m and the 100-year return period surge in the order of 0.6 m). Under these conditions, the 0.75 m increase in the centennial ESL corresponds to a relative increase of 20 to 24%, and above all, a current centennial ESL is likely to occur on average once or twice a year (amplification factor between 50 and 100);
- In the Mediterranean, the situation is clearly aggravated by the very low variability of the sea level (microtidal regime and small surges). As a result, the 100-year ESL increase of 0.77 m corresponds to a large relative increase (greater than 50%) and the present day 100-year ESL is likely to occur between 2 and 10 times per year (amplification factor between 500 and 1000).

By 2050, the 100-year ESL will rise by 5% in the North Channel and the Bay of Biscay. In the Mediterranean, this increase will already be of the order of 20%. However, the evolution of the situation will be especially noticeable in the increased frequency of ESLs (by a factor of 3 to 4).

In addition to this quantitative analysis, the qualitative analysis shows that the geomorphological configurations of the three metropolitan maritime facades (except on the eastern half of the Mediterranean coast and the southern part of the Bay of Biscay) favor the formation of storm surges, by the width of the continental shelves, and the length and shape of the coastlines. On these coasts, a change in the characteristics of storms could therefore lead to a sharp increase in surges. In particular, an increased occurrence and intensity of medicanes in the western Mediterranean, as projected by Romero and Emanuel (2016), would aggravate strongly coastal hazards. Analysis of the surge observed in 2010 during the passage of storm Xynthia in the Bay of Biscay demonstrates the vulnerability of coastal areas to unusual ~~meteocean~~metocean phenomena (Bertin et al., 2012). The storm's low pressure center track propagated to the northeast, contrasting with other storm tracks usually propagating to the east or southeast. Despite its low intensity, this storm caused the large marine submersions in France (Cerema, 2016).

Finally, it should be recalled that the amplification factors related to ESLs (which are expected to reach values between 50 and 1000 in 2100 on the coasts of metropolitan France) must be distinguished from the evolution of extreme events such as storms, cyclones and tsunamis, whose frequency and intensity is not projected to change as significantly as ESLs. However, even if the frequency and intensity of ~~meteocean~~metocean events remain constant, an increase in the ESL frequency and intensity would increase the effects of these events on the coasts.

Table 810: Same as Table 79 for the coastlines of French overseas territories.

	Surge potential associated with the geomorphological configuration	Other regional factor of change	Reference <del>meteocean</del> metocean event	Tidal range	100-year surge (m)	ΔRSL (m) under SSP5-8.5	ΔESL (m) under SSP5-8.5	%ΔESL under SSP5-8.5	Return period of the present day 100-year ESL under SSP5-8.5 (year)
Saint-Pierre (Saint-Pierre-et-Miquelon)	High	None	Storm and moderate exposure to cyclone	MICROTIDAL (< 2 M)	1 to 2	In 2050 : 0,19 In 2100 : 0.83	In 2050 : 0.24 In 2100 : 0.92	In 2050 : 8.4 In 2100 : 31.8	IN 2050 : 18 IN 2100 : 0.3
POINTE-A-PITRE (GUADELOUPE)	Low	Coral reef degradation	STORM AND CYCLONE	MICROTIDAL (< 2 M)	1 to 3	In 2050 : 0,20 In 2100 : 0,90	In 2050 : 0.22 In 2100 : 0.91	In 2050 : 14.1 In 2100 : 58.1	IN 2050 : 35 IN 2100 : 0.1
Cayenne (French Guiana)	Moderate	None	Trade winds (without storm)	Mesotidal (2 to 4 m)	0.4	In 2050 : 0,20 In 2100 : 0,90	In 2050 : 0.18 In 2100 : 0.85	In 2050 : 10.6 In 2100 : 49.2	IN 2050 : 7,5 IN 2100 : 0.1
POINTE DES GALETS (LA REUNION)	Very low	Coral reef degradation Decrease in seasonal sea ice extent	STORM CYCLONE +LONG SWELL +MODERATE EXPOSURE TO TSUNAMIS	MICROTIDAL (< 2 M)	1 to 3	In 2050 : 0,19 In 2100 : 0,92	In 2050 : 0.16 In 2100 : 0.88	In 2050 : 9.3 In 2100 : 52.5	IN 2050 : 58 IN 2100 : 0.1
PAPEETE (FRENCH POLYNESIA)	Very low	Coral reef degradation Decrease in seasonal sea ice extent	STORM CYCLONE +LONG SWELL +MODERATE EXPOSURE TO TSUNAMIS	MICROTIDAL (< 2 M)	1 to 3	In 2050 : 0,20 In 2100 : 0,91	In 2050 : 0.23 In 2100 : 0.95	IN 2050 : 17.8 IN 2100 : 74.7	IN 2050 : 16 IN 2100 : 0.1

In 2100, the quantitative analysis shows that for the SSP5-8.5 scenario:

- The coasts of the West Indies, La Réunion, and French Polynesia are microtidal (<2 m), but the passage of the cyclones can generate large surges (up to 3 m). The allowance (50-75% increase) and amplification (increase in the frequencies of the current centennial ESL by about 1000) of these coasts are similar to those of the Mediterranean coast of France. However, the change in the intensity and frequency of the ESLs must be considered more carefully

in these territories because of the greater damage that tropical cyclones can cause compared to storms (and medicanes) that reach the European territory of France;

- 640
- The coast of Saint-Pierre-et-Miquelon is also microtidal (<2 m). While strong winds that regularly blow on these islands can produce large surges, the variability in sea levels remains low because of the microtidal regime. The situation is therefore similar to that of the Mediterranean (including the moderate exposure to cyclones), with a smaller estimated increase in ESLs, as shown by the allowance (about 32%) and amplification (about 300);
  - The coast of French Guiana is mesotidal (maximum range of 2.9 m), and the maximum recorded surge is only 0.4 m.
- 645 The increase in the amplitude and frequency of ESLs should be of the same order of magnitude as that of the Mediterranean coast of France, the West Indies and La Réunion. However, this coast is located close to the equator and is not exposed to ~~meteo-oceanic~~ events that generate strong swells. The increase in coastal hazards therefore is expected to be less intense than in the other overseas territories.

650 By 2050, the evolution of the hazard is likely to be perceptible primarily because of the increase in the frequency of ESLs by a factor of 2 for La Réunion, 3 for the West Indies, 5 for Saint-Pierre-et-Miquelon, 6 for French Polynesia, and 13 for French Guiana. The most significant changes are for the territories with low variability in sea levels that are subject to storms ~~(Saint-Pierre-et-Miquelon whose centennial ESL will increase by 8%)~~ and to cyclones (Saint-Pierre-et-Miquelon, La Réunion, the West Indies and French Polynesia, with centennial ESL increases of 8, 9, 14 and 18%, respectively). In this comparison, French

655 Polynesia shows both the greatest increase in frequency and intensity of the 100-year ESL. French Guiana is also expected to experience negative changes in the ESL by 2050, but since it is not exposed to storm waves, this territory is not expected to be exposed to catastrophic ~~meteocean~~metocean events.

In addition, the qualitative analysis shows that the geomorphological configurations of the overseas territories (except for Saint-Pierre-et-Miquelon and French Guiana) do not favor the formation of storm surges. The situation is aggravated in the

660 West Indies by the degradation of coral reefs and on La Réunion and French Polynesia (islands exposed to cyclones and southern swells) by the degradation of coral reefs and the decrease in seasonal sea ice extent.

Finally, as for the coasts of metropolitan France, it is necessary to distinguish between changes in the frequency and intensity of ESLs and extreme ~~meteocean~~metocean events. Even if extreme ~~meteocean~~metocean events are not expected to increase as significantly as ESLs, an increase in ESLs will lead to an increase in coastal hazards.

## 665 4 Discussion

The application of the proposed method leads to increased understanding of coastal hazards, ~~taking into account~~considering the hydrodynamic forcing and the geomorphological context. These results, ~~considered~~ in conjunction with anticipated coastal responses to the effects of climate change, can shed light on the evolution of the hazard on the terrestrial domain.

#### 4.1 Influence of local factors affecting the coastline

670 Understanding the evolution of coastal hazards requires evaluating slow changes that occur over decadal timescales (e.g. RSL rise, ocean acidification and warming, morphological evolution), in conjunction with extreme events that occur on daily time scales, such as tsunamis, cyclones, storms (Irigibel et al., 2021). However, while some factors can be ~~taken into account~~considered at the regional level (and were therefore ~~considered by~~included in the proposed method), others can only be considered at finer spatial scales because of the complex interactions between ~~sea-level~~sea level rise and the morphological evolution of coastal areas. In order to assess local hazard evolution, one must ~~consider~~study local effects, such as subsidence, especially in deltas (Syvitski, 2008). Second, the coastal zone is a buffer zone where a multitude of processes and feedbacks may also be important:

- 680 • in estuaries, changes in coastal morphology (bathymetry, shoreline topography, and anthropogenic development) can influence (positively or negatively) extreme events, including changes in the spread of tidal waves and storm surges (Talke et al., 2020);
- 685 • along sandy coasts, coastal morphology is also likely to change as a function of the local morphological and sedimentary characteristics, anthropogenic development, climatic changes (temperature, precipitation and wind), wave conditions, and frequency of events, which has an important impact- on the ability of a system to recover between energetic events (Masselink et al., 2016). In addition, wave direction is also an important parameter related to longshore sediment transport, which can change the shape of a coastline (Ruggiero et al., 2010; Casas-Prat and Sierra, 2010);
- 690 • in the polar regions, accelerating permafrost thaw is promoting rapid erosion of ice-rich sediments (Lantuit et al., 2011). Melting of ice and associated thaw subsidence may induce instability of various infrastructure components. Arctic RSL rise and sea surface warming have the potential to substantially contribute to this thawing (Lamoureux et al., 2015).

RSL rise can greatly increase coastal hazards, because an increase in water depth, assuming no change in bathymetry, may cause increases in nearshore wave conditions. The coastal impacts of ESLs are largely due to increased wave impacts, potentially driving morphological changes and erosion, as well as coastal protection failure and overwash/inundation (CIRIA et al., 2013, Vousdoukas et al., 2017). Moreover, as indicated by Krien et al. (2017), coastal ecosystems such as mangroves, coral reefs or seagrass beds may not be able to adapt to human activities and climate change (e.g., Waycott et al., 2009; Wong et al., 2014), which could have large impacts on coastal hazards (e.g., Alongi, 2008; Wong et al., 2014; IPCC, 2019). Ideally, all of these phenomena should be considered in assessing coastal hazard evolution and in defining the adaptation measures for a particular location. While this may be too difficult to achieve for a regional study, it is necessary ~~to take into account~~ at the local level.

## 4.2 Understanding hazard evolution on land

On many coasts, accelerated ~~sea level~~sea level rise is likely to result in permanent submersion of unprotected lowlands. More frequent and intense episodic coastal flooding could also occur with future changes in the wave climate and storm surges. In some locations, this may result in chronic coastline erosion (Ranasinghe, 2016). Thus, depending on the coastline, distinct phenomena may appear, which may be classified, using the proposed method (characterisation of the hazard on a regional scale in the maritime domain) and the insights from existing studies (characterization of phenomena affecting the coastal fringe locally).

By neglecting adaptive actions, hazard evolution may be categorized into one of the following three main scenarios (defined by the sea level variability and the types of ~~meteoecean~~metocean events, knowing that other factors also influence the hazard evolution and must be considered subsequently):

- coastlines with low variability in sea levels that are not exposed to ~~meteoecean~~metocean events (e.g. French Guiana): a RSL rise will likely generate a small increase in the risk of rapid submersion by the sea. Nevertheless, rapid flooding by river floods or by intense precipitation remains possible. In addition, the gradual rise of the RSL is likely to cause increased erosion (which, in general, results in coastline retreat) and/or to the progressive and permanent flooding of lowlands (i.e. below sea level);
- coastlines with high ~~sea level~~sea level variability that are exposed to ~~meteoecean~~metocean events (e.g., the Atlantic coastline of metropolitan France): RSL rise is likely to cause increased erosion and the degradation of protection systems (i.e. anthropogenic structures, such as levees, seawalls, groynes and breakwaters; and natural features such as beaches, dunes, salt marshes, mangroves and coral reefs) over the years, even in the absence of extreme events. During an extreme event (due to the conjunction of a storm, cyclone or tsunami with a high spring tide), the risk may be increased both by the regular weakening of the protection systems and by the increased impacts of waves related to the rise of the RSL;
- coastlines with low ~~sea level~~sea level variability that are exposed to ~~meteoecean~~metocean events (e.g. Saint-Pierre-et-Miquelon, West Indies, La Réunion, French Polynesia): RSL rise is likely to cause, in the short term, the three phenomena described in the previous cases: increased risk of coastal erosion, increased risk of rapid flooding, and permanent flooding of lowlands.

In addition to these three main scenarios, there are coastlines with high variability in sea levels and with relatively low exposure to ~~meteoecean~~metocean events, which will likely show slower increases in coastal hazards.

## 5 Conclusion

Changes in sea levels (~~GMSL~~, RSL and ESL) are subject to high uncertainties, at the global scale (economic development, GHG emission scenarios, ice sheet response) and at regional or local scales (response of coastlines to climatic changes and

changes in wave climate). Quantified estimates of sea level changes are necessary, but should also be put into perspective with factors describing the state and evolution of the coastal environment. The proposed method for evaluating coastal hazard evolution is based on this principle, emphasizing the importance of considering sea level changes, as well as the geomorphological configuration, ~~meteocean~~metocean event types, and, where appropriate, additional marine environment changes affecting hydrodynamic conditions (in particular, in the tropics or at the poles).

The application of this method to selected French coastlines demonstrated the expected differences in future coastal hazard evolution.

The study of the variability of sea levels with a quantitative approach confirmed the results of previous studies, namely the largest increase of ESLs on coasts with a small tidal range and storm surges. This is the case for the French overseas territories and, in metropolitan France, for the Mediterranean facade. The increase in ESLs, in intensity and frequency, should be limited on the coasts with large tidal ranges (French facades of the Atlantic Ocean, the English Channel and the North Sea).

In addition to the results obtained for ESLs by quantitative approaches, the qualitative approach makes it possible to assess more comprehensively the evolution of coastal hazards.

First, the type of ~~meteocean~~metocean event is essential: coasts not exposed to extreme events (storms, cyclones, long swells or tsunamis), even if they are microtidal (e.g. Guiana), should experience a moderate hazard increase compared to exposed coasts. All other characteristics being equal (geomorphological configurations, environmental changes and tidal conditions), the hazard will increase more rapidly on the coasts exposed to cyclones (the West Indies, La Réunion, and French Polynesia, Saint-Pierre-et-Miquelon and the Mediterranean facade) than on the coasts exposed to storms (North Sea, English Channel and Bay of Biscay~~Mediterranean facade~~).

Secondly, certain geomorphological configurations (long maritime facades, concave in shape and bordered by a wide continental shelf) favor the generation of surges during extreme events. These criteria help to identify the coastlines most vulnerable to changes in the characteristics of extreme events (especially their trajectories). This result is important because quantitative approaches are based on climate change scenarios, which therefore represent only part of the possible evolutions. The application of quantitative and qualitative approaches for French territories makes it possible to distinguish several scenarios: some coastlines benefit from a high variability in sea levels, which is a positive feature to limit the effects of the rise in RSL, but, on the other hand, have geomorphological characteristics favoring surges (e.g. Atlantic, Channel and North Sea coasts). Other shorelines are expected to be disadvantaged by both low sea level variability (small tidal range and moderate surges) and a geomorphological configuration that favors surges (e.g., the western portion of the Mediterranean coastline and the concave shorelines of the West Indies). Finally, some coastlines have the disadvantage of small sea level variability, but



their geomorphological configuration does not favor surges (e.g. the eastern part of the Mediterranean coastline). This analysis highlights the need to study not only the water level variability, but also the current and future exposure to ~~meteocean~~metocean events.

770

Third, the coral reef degradation in the tropics and the decrease in seasonal sea ice extent in the polar regions can also significantly change the nearshore hydrodynamics and impacts on the shoreline.

775

These results represent a substantial step forward, not only because they are achieved through a comprehensive and systemic approach that requires a step back, but also because the proposed method uses factors for which the uncertainty in the projections is the lowest: the geomorphological configuration of the ~~coast~~continental shelf at a regional scale will be invariable over the centuries, regional tidal ~~conditions~~regime will be stable, climate change impacts on coral reefs (Albright et al., 2018; Hoegh-Guldberg et al., 2018) and seasonal sea ice extent (Forbes, 2011) are fairly well known. However, local climate changes and, more particularly, changes in storm and cyclone trajectories are an area where uncertainties remain particularly high (Seneviratne et al., 2021).

780

Ultimately, the impacts of sea level changes on human communities are evaluated by considering additionally the local-scale effects on natural features and anthropogenic structures. These analyses aim to predict the types of hazard (shoreline erosion, rapid submersion and/or permanent flooding) that will increase the most in different coastal zones (shoreline erosion, rapid submersion and/or permanent flooding). Shorelines are expected to be marked by a quasi-generalized increase in the risks of erosion of natural features and weakening of anthropogenic structures. However, regional and local-scale differences are expected depending on the submersion type: coastlines with high ~~sea level~~sea level variability will experience mainly rapid submersion effects, whereas coastlines with low ~~sea level~~sea level variability will experience mainly permanent flooding of low-lying land, and, rapid submersion when exposed to extreme ~~meteocean~~metocean events.

785

790

To go beyond the systemic approach proposed here, a follow-up study could be completed using: (i) a consensus evaluation based on bibliographic studies (IPCC approach) of each of the identified determining factors ; (ii) or a machine learning approach to explore the added benefits of artificial intelligence (AI). In addition to large-scale hydrodynamic model outputs and other environmental data, analyses may integrate deep learning method outputs. Machine learning can be a powerful tool in reducing greenhouse gas emissions and helping society adapt to a changing climate (Kaack et al., 2021; Rolnick et al., 2022). AI has already provided interesting results in the field of hydrology (Kumar et al., 2023) and coastal flooding (Tiggeloven et al., 2021; Jones et al., 2023).

795

800

In parallel with research on systemic approaches, the assessment of water levels remains an essential topic. Future efforts could benefit from recent developments on ocean modeling like the new generation Global Tide and Surge Model Version 3.0

(GTSMv3.0), which can be used to simulate dynamically tides, storm surges, and changes in MSL, including interaction effects (Muis et al., 2020); or the new global, fully coupled, unstructured model of Mentaschi et al. (2023), coupling waves, storm surge and tides.

## 6 Code availability

805 The Delft3D-FM code is currently being made available in <http://oss.deltares.nl>. The WW3 model description is available in: <https://polar.ncep.noaa.gov/waves/wavewatch/>. The code applied for the non-stationary extreme value statistics (Mentaschi et al., 2016) is available in: <https://github.com/menta78/tsEva>.

## 7 Data availability

810 The global ESL data that support the findings of this study are available in the LISCoAsT repository of the JRC data collection (<http://data.jrc.ec.europa.eu/collection/LISCOAST>) though this link: <http://data.jrc.ec.europa.eu/dataset/jrc-liscoast-10012>, with the identifiers: <https://doi.org/10.2905/jrc-liscoast-10012>; PID: <http://data.europa.eu/89h/jrc-liscoast-10012>

## 8 Author contribution

MI, MY and YD conceived the paper. MV developed the model code and performed the simulations. MI prepared the manuscript with contributions from all co-authors.

## 815 9 Competing interests

The authors declare that they have no conflict of interest.

## 10 Acknowledgments

820 The authors would like to thank Xavier Bertin, CNRS research director at the Littoral ENvironment et Sociétés (LIENSs) laboratory for his comments about the manuscript and for sharing of water height and wave data from the national observation services ReefTEMPS and DYNALIT.

The authors would also like to thank the Naval Hydrographic and Oceanographic Service (in French: Service hydrographique et océanographique de la Marine or SHOM), as well as Anne-Laure Tiberi-Wadier (Cerema) and Xavier Kergadallan (Cerema) for their contributions to bibliographic research.

825

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors. All authors declare that they have no conflicts of interest.

## 11 References

830 Albright, R., Takeshita, Y., Koweek, D., Ninokawa A., Wolfe K., Rivlin T., et al.: Carbon dioxide addition to coral reef waters suppresses net community calcification, *Nature* 555, 516–519, <https://doi.org/10.1038/nature25968>, 2018.

[Allgeyer, S., Daubord, C., Hébert, H., Loevenbruck, A., Schindelé, F., and Madariaga, R.: Could a 1755-like tsunami reach the French Atlantic coastline? constraints from twentieth century observations and numerical modeling. \*Pure and Applied Geophysics\*, 170\(9–10\), 1415–1431, 2013.](#)

835 [Allgeyer, S., Quentel, É., Hébert, H. et al.: Tsunami Hazard in La Réunion Island \(SW Indian Ocean\): Scenario-Based Numerical Modelling on Vulnerable Coastal Sites. \*Pure Appl. Geophys.\* 174, 3123–3145. <https://doi.org/10.1007/s00024-017-1632-9>, 2017.](#)

840 [Alongi, D. M.: Mangrove forests: resilience, protection from tsunamis, and response to global climate change. \*Estuar. Coast. Shelf S.\*, 76, 1–13, 2008.](#)

Almar, R., Ranasinghe, R., Bergsma, E.W.J., Diaz, H., Melet, A., Papa, F., et al.: A global analysis of extreme coastal water levels with implications for potential coastal overtopping. *Nat Commun*, 12, 3775. <https://doi.org/10.1038/s41467-021-24008-9>, 2021.

845 [Androulidakis, Y. S., Kombiadou, K. D., Makris, C. V., Baltikas, V. N., and Krestenitis, Y. N.: Storm surges in the Mediterranean sea: Variability and trends under future climatic conditions. \*Dynamics of Atmospheres and Oceans\*, 71, 56–82. <https://doi.org/10.1016/j.dynatmoce.2015.06.001>, 2015.](#)

850 [Arns A., Wahl T., Dangendorf S., Jensen J.: The impact of sea level rise on storm surge water levels in the northern part of the German Bight. \*Coast Eng\* 96:118–131. <https://doi.org/10.1016/j.coastaleng.2014.12.002>, 2015](#)

855 [Arns, A., Wahl, T., Wolff, C., Vafeidis A., Haigh I., Woodworth P., Niehüser S. and Jensen J.: Non-linear interaction modulates global extreme sea levels, coastal flood exposure, and impacts. \*Nat Commun\* 11, 1918. <https://doi.org/10.1038/s41467-020-15752-5>, 2020.](#)

Aubanel, A., Marquet, N., Colombani, J.M., and Salvat, B.: Modifications of the shore line in the Society islands (French Polynesia). Ocean and Coastal Management, 42, 419–438, 1999.

860

Aucan J., Hoeke R. K., Storlazzi C. D., Stopa J., Wandres M., and Lowe R.: Waves do not contribute to global sea-level rise. Nature Climate Change, 9(1):2–2, December 2018.

865

Bamber, L. J., Oppenheimer, M., Kopp, R. E. and Cooke, R. M.: Ice sheet contributions to future sea-level rise from structured expert judgment. Proceedings of the National Academy of Sciences, p. 5., 2019.

Bardet, L., Duluc, C.-M., Rebour, V., and L’Her, J.: Regional frequency analysis of extreme storm surges along the French coast, Nat. Hazards Earth Syst. Sci., 11, 1627–1639, <https://doi.org/10.5194/nhess-11-1627-2011>, 2011.

870

Barriot, J-P., Zhang, F., Ducarme, B., Wöppelmann, G., André, G. & Gabillon, A.: A database for sea-level monitoring in French Polynesia. Geoscience Data Journal, 10, 368–384. Available from: <https://doi.org/10.1002/gdj3.172>, 2023.

875

Becker, M., Meyssignac, B., Letetrel, C., Llovel, W., Cazenave, A., and Delcroix, T.: Sea level variations at tropical Pacific islands since 1950. Global and Planetary Change, 80–81, 85–98. doi: 10.1016/j.gloplacha.2011.09.004, 2012.

Bertin, X., Bruneau, N., Breilh, J.-Fr., Fortunato, A. and Karpytchev, M.: Importance of wave age and resonance in storm surges: The case Xynthia, Bay of Biscay, Ocean Modelling, Volume 42, Pages 16-30, ISSN 1463-5003, <https://doi.org/10.1016/j.ocemod.2011.11.001>, 2012.

880

Bertin X., Li K., Roland A., Zhang Y.J., Breilh J.-F. and Chaumillon E.: A modeling-based analysis of the flooding associated with Xynthia, central Bay of Biscay. Coast Eng 94:80–89, 2014.

885

Bertin, X., Li K., Roland A., and Bidlot J.-R.: The contribution of short-waves in storm surges: two case studies in the Bay of Biscay, Continental Shelf Research, 96, 1-15, doi:10.1016/j.csr.2015.01.005, 2015.

Bertin X., Martins K., de Bakker A., Chataigner T., Guérin T., et al.: Energy Transfers and Reflection of Infragravity Waves at a Dissipative Beach Under Storm Waves. Journal of Geophysical Research. Oceans, 125 (5), pp.e2019JC015714, 2020.

890

Bowen, A. J., Inman, D. L. and Simmons, V. P.: Wave ‘set-down’ and set-up, J. Geophys. Res., 73, 2569–2577, doi:10.1029/JB073i008p02569, 1968.

Brivois O., Idier D., Thiébot J., Castelle B., Le Cozannet G., Calvete D.: On the use of linear stability model to characterize the morphological behaviour of a double bar system. Application to Truc Vert beach (France). CR Geosci 344:277–287. <https://doi.org/10.1016/j.crte.2012.02.004>, 2012.

895

Buchanan, M.K., Kopp R.E., Oppenheimer M. and Tebaldi C.: Allowances for evolving coastal flood risk under 41 uncertain local sea-level rise, *Climatic Change*, 137(3–4), doi:10.1007/s10584-016-1664-7, 2016.

Bulteau, T., Idier, D., Lambert, J., and Garcin, M.: How historical information can improve estimation and prediction of extreme coastal water levels: application to the Xynthia event at La Rochelle (France), *Nat. Hazards Earth Syst. Sci.*, 15, 1135–1147, <https://doi.org/10.5194/nhess-15-1135-2015>, 2015.

900

Calafat, F.M., Wahl, T., Tadesse, M.G. and Sparrow, S.N.: Trends in Europe storm surge extremes match the rate of sea-level rise, *Nature* 603, 841–845. <https://doi.org/10.1038/s41586-022-04426-5>, 2022).

905

Campins, J., Genovés, A., Picornell, M. A., and Jansà, A.: Climatology of Mediterranean cyclones using the era-40 dataset. *International Journal of Climatology*, 31(11), 1596–1614. <https://doi.org/10.1002/joc.2183>, 2011.

Casas-Prat, M. and Sierra, J. P.: Trend analysis of the wave storminess: the wave direction, *Adv. Geosci.*, 26, 89–92. <https://doi.org/10.5194/adgeo-26-89-2010>, 2010.

910

Catto N. R., Batterson M. J.: Igor and other hurricane and extratropical transitions in Newfoundland: geomorphologic and landscape impacts. *Geohydro 2011* 2011:1–4, 2011.

Cavicchia, L., Von Storch, H., and Gualdi, S.: A long-term climatology of medicanes. *Climate Dynamics*, 43, 1183, 1195. <https://doi.org/10.1007/s00382-013-1893-7>, 2014.

915

Cazenave, A. and Llovel, W.: Contemporary sea level rise. *Ann. Rev. Marine Sci.* 2, 145–173, 2010.

920

Cazenave, A. and Le Cozannet, G.: Sea level rise and its coastal impacts, *Earth's Future*, 2, 15–34. doi:10.1002/2013EF000188, 2013.

Cerema: Étude des systèmes de protection contre les submersions marines, Collection : Références, ISBN : 978-2-37180-118-9, 442 pages, 2016.

925

- Cerema: Les niveaux marins extrêmes. Ports de métropole, Collection: Données, ISBN : 978-2-37180-308-4, 2018.
- 930 Cerema: Dynamiques et évolution du littoral, Fascicule 11: Synthèse des connaissances de Saint-Pierre-et-Miquelon, Collection: Connaissances, ISBN: 978-2-37180-341-1, 2020.
- Cerema: Dynamiques et évolution du littoral, Fascicule 12: Synthèse des connaissances de Saint-Barthélemy et Saint-Martin, Collection: Connaissances, ISBN: 978-2-37180-342-8, 2020.
- 935 Cerema: Dynamiques et évolution du littoral, Fascicule 13: Synthèse des connaissances de l'archipel de Guadeloupe, Collection: Connaissances, ISBN: 978-2-37180-316-9, 2020.
- Cerema: Dynamiques et évolution du littoral, Fascicule 14: Synthèse des connaissances de la Martinique, Collection: Connaissances, ISBN: 978-2-37180-343-5, 2020.
- 940 Cerema: Dynamiques et évolution du littoral, Fascicule 15: Synthèse des connaissances de la Guyane, Collection: Connaissances, ISBN: 978-2-37180-344-2, 2021.
- Cerema: Dynamiques et évolution du littoral, Fascicule 17: Synthèse des connaissances de La Réunion, Collection: Connaissances, ISBN: 978-2-37180-345-9, 2020.
- 945 [Chevalier C., Froidefond J.-M. and Devenon J.-L.: Numerical analysis of the combined action of littoral current, tide and waves on the suspended mud transport and on turbid plumes around French Guiana mudbanks, Continental Shelf Research, Volume 28, Issues 4–5, Pages 545-560, ISSN 0278-4343, https://doi.org/10.1016/j.csr.2007.09.011, 2008.](https://doi.org/10.1016/j.csr.2007.09.011)
- 950 Church J. A., Clark P. U., Cazenave A., Gregory J. M., Jevrejeva S., Levermann A., Merrifield M. A., Milne G. A., Nerem R. S., Nunn P. D., Payne A. J., Pfeffer W. T., Stammer D., Unnikrishnan A. S.: Sea level change, In: Climate change 2013: the physical science basis, Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change, Cambridge University Press, Cambridge, United Kingdom, pp 1137–1216, 2013.
- 955 CIRIA, MEDDE, USACE: The International Levee Handbook, C731, CIRIA, UK, London, ISBN : 978-0-86017-734-0, 1332 pages, 2013.

- 960 Conte, D., and Lionello, P.: Characteristics of large positive and negative surges in the Mediterranean sea and their attenuation in future climate scenarios. *Global and Planetary Change*, 111, 159–173. <https://doi.org/10.1016/j.gloplacha.2013.09.006>, 2013.
- Dayan H., Le Cozannet G., Speich S. and Thiéblemont R.: High-End Scenarios of Sea-Level Rise for Coastal Risk-Averse Stakeholders. *Front. Mar. Sci.* 8:569992, doi: 10.3389/fmars.2021.569992, 2021.
- 965 De la Torre, Y.: Impacts du cyclone Irma sur le littoral des « îles du Nord » à Saint-Martin et Saint-Barthélemy, Rapport BRGM/RP-67291-FR, 29 p., 20 ill., 2 ann., 2017.
- Dietrich, J. C., Bunya, S., Westerink, J. J., Ebersole, B. A., Smith, J. M., Atkinson, J. H., et al.: A high resolution coupled riverine flow, tide, wind, wind wave and storm surge model for southern Louisiana and Mississippi: part II — synoptic description and analyses of Hurricanes Katrina and Rita. *Monthly Weather Review* 138, 378–404, 970 <https://doi.org/10.1175/2009MWR2907.1>, 2010.
- Dodet, G., Bertin, X., Bouchette, F., Gravelle, M., Testut, L., and Wöppelmann, G.: Characterization of Sea-level Variations Along the Metropolitan Coasts of France : Waves, Tides, Storm Surges and Long-term Changes, *Journal of Coastal Research*, 975 88(sp1), 10-24. <https://doi.org/10.2112/SI88-003.1>, 2019.
- Dorville, J.-F. M. and Zahibo, N.: Hurricane Omar Waves Impact on the West Coast of the Guadeloupe Island, October 2008. *Open Oceanography Journal*, 4, 83–91, 2010.
- 980 DREAL Nord Pas-de-Calais: Détermination de l'aléa de submersion marine intégrant les conséquences du changement climatique en région Nord – Pas de Calais. Phase 1: Compréhension du fonctionnement du littoral, [https://www.hauts-de-france.developpement-durable.gouv.fr/IMG/pdf/50292\\_-\\_sub\\_npc\\_-\\_phase\\_1\\_-\\_version\\_4.pdf](https://www.hauts-de-france.developpement-durable.gouv.fr/IMG/pdf/50292_-_sub_npc_-_phase_1_-_version_4.pdf), last access: 08 February 2024.
- 985 Durand, F.: Hydrodynamique sédimentaire sur le plateau insulaire de la Martinique, Thèse doctorale, Université Bordeaux 1. 214 p., 1996.
- Elkut, A. E., Taha, M. T., Abu Zed, A. B. E., Eid, F. M., and Abdallah, A. M.: Wind-wave hindcast using modified ECMWF era-interim wind field in the Mediterranean sea. *Estuarine, Coastal and Shelf Science*, 252, 107267. 990 <https://doi.org/10.1016/j.ecss.2021.107267>, 2021.

Erkens, G., Bucx, T., Dam, R., De Lange, G. and Lambert, J.: Sinking Coastal Cities. Proceedings of the International Association of Hydrological Sciences, 372, 189-198, <https://doi.org/10.5194/piahs-372-189-2015>, 2015.

995 [Fita, L., Romero, R., Luque, A., Emanuel, K., and Ramis, C.: Analysis of the environments of seven Mediterranean tropical-like storms using an axisymmetric, nonhydrostatic, cloud resolving model. Natural Hazards and Earth System Sciences, 7, 41, 56. <https://doi.org/10.5194/nhess-7-41-2007>, 2007.](#)

1000 [Flather, R.A.: Storm surges. In: Steele, J., Thorpe, S., Turekian, K. \(Eds.\), Encyclopedia of Ocean Sciences. Academic, San Diego, California, pp.2882–2892, 2001.](#)

FLOODsite: Language of risk. Project definitions, Report T34-02-01, Second edition, 55 pp. e-publication on [www.floodsite.net](http://www.floodsite.net). Available online: [www.floodsite.net](http://www.floodsite.net), 2009.

1005 [Forbes, D.L.: State of the Arctic coast 2010: scientific review and outlook. Land-Ocean Interactions in the Coastal Zone, Institute of Coastal Research, <http://arcticcoasts.org/files/sac/state%20of%20the%20arctic%20rept-exec%20summary%20res.pdf>, 2011.](#)

1010 [Formentin, S.M. and Zanuttigh, B.: A new method to estimate the overtopping and overflow discharge at over-washed and breached dikes. Coast. Eng., 140, 240–256. <https://doi.org/10.1016/j.coastaleng.2018.08.002>, 2018](#)

Fox-Kemper B, Hewitt HT, Xiao C et al.: Chapter 9: ocean, cryosphere and sea level change. In: Masson-Delmotte V, Zhai P, Pirani A et al (eds) Climate Change 2021: the physical science basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Intergovernmental Panel on Climate Change (IPCC), 2021.

1015 [Frederikse, T., Landerer, F., Caron, L. et al.: The causes of sea-level rise since 1900, Nature 584, 393–397, <https://doi.org/10.1038/s41586-020-2591-3>, 2020.](#)

1020 [Garnier, E., Ciavola, P., Armaroli, C., Spencer, T., and Ferreira, O.: Historical analysis of storms events: case studies in France, England, Portugal and Italy, Coast. Eng., 134, 10–23, <https://doi.org/10.1016/j.coastaleng.2017.06.014>, 2018.](#)

[Gomes da Silva, P., Coco, G., Garnier, R., Klein, A.H., 2020. On the prediction of runup, setup and swash on beaches. Earth-Sci. Rev. 204, 103148. <http://dx.doi.org/10.1016/j.earscirev.2020>.](#)



- 1025 [Gratiot N., Gardel A. and Anthony E. J.: Trade-wind waves and mud dynamics on the French Guiana coast, South America: Input from ERA-40 wave data and field investigations, \*Marine Geology\*, Volume 236, Issues 1–2, 2007, Pages 15-26, ISSN 0025-3227, <https://doi.org/10.1016/j.margeo.2006.09.013>, 2007.](#)
- Graumann, A., Houston, T., Lawrimore, J. H., Levinson, D. H., Lott, N., McCown, S. et al.: Hurricane Katrina, A  
1030 Climatological Perspective. October 2005, Updated August 2006. NOAA’s National Climatic Data Center.  
<https://www1.ncdc.noaa.gov/pub/data/techrpts/tr200501/tech-report-200501z.pdf>, 2006.
- Gregory, J.M., Griffies, S.M., Hughes, C.W. et al.: Concepts and Terminology for Sea Level: Mean, Variability and Change,  
Both Local and Global. *Surv Geophys* 40, 1251–1289, <https://doi.org/10.1007/s10712-019-09525-z>, 2019.
- 1035 [Haigh I., Nicholls R. and Wells N.: Rising sea levels in the English Channel 1900 to 2100. \*Marit Eng\* 164\(MA2\):81–92, 2011.](#)
- Haigh, I.D., Pickering, M.D., Green, J.A.M., Arbic, B.K., Arns, A., Dangendorf, S., et al.: The Tides They Are a-Changin’: A  
comprehensive review of past and future non-astronomical changes in tides, their driving mechanisms and future implications,  
1040 *Reviews of Geophysics*, 57, e2018RG000636, <https://doi.org/10.1029/2018RG000636>, ~~2020~~2019.
- Hallegatte, S., Green, C., Nicholls, R. J., and Corfee-Morlot, J.: Future flood losses in major coastal cities, *Nature Climate  
Change*, 3, 802–806, <https://doi.org/10.1038/nclimate1979>, 2013.
- 1045 [Hamdi, Y., Bardet, L., Duluc, C.-M., and Rebour, V.: Extreme storm surges: a comparative study of frequency analysis approaches, \*Nat. Hazards Earth Syst. Sci.\*, 14, 2053–2067, <https://doi.org/10.5194/nhess-14-2053-2014>, 2014.](#)
- [Hamdi, Y., Bardet, L., Duluc, C.-M., and Rebour, V.: Use of historical information in extreme-surge frequency estimation: the case of marine flooding on the La Rochelle site in France, \*Nat. Hazards Earth Syst. Sci.\*, 15, 1515–1531, <https://doi.org/10.5194/nhess-15-1515-2015>, 2015.](#)
- 1050 [Hamdi, Y., Garnier, E., Giloy, N., Duluc, C.-M., and Rebour, V.: Analysis of the risk associated with coastal flooding hazards: a new historical extreme storm surges dataset for Dunkirk, France, \*Nat. Hazards Earth Syst. Sci.\*, 18, 3383–3402, <https://doi.org/10.5194/nhess-18-3383-2018>, 2018.](#)
- 1055 [Hamlington, B. D., Gardner, A. S., Ivins, E., Lenaerts, J. T. M., Reager, J. T., Trossman, D. S., et al. \(2020\). Understanding of contemporary regional sea-level change and the implications for the future. \*Reviews of Geophysics\*, 58, e2019RG000672. <https://doi.org/10.1029/2019RG000672>](#)

1060 Han, G., Ma, Z., Chen, D., deYoung B. and Chen N.: Observing storm surges from space: Hurricane Igor off Newfoundland. Sci Rep 2, 1010. <https://doi.org/10.1038/srep01010>, 2012.

Hequette A.: Les risques naturels littoraux dans le Nord-Pas-de-Calais, France, VertigO, hors ser. 8, 2010.

1065 Idier, D., Dumas, F., and Muller, H.: Tide-surge interaction in the English Channel, Nat. Hazards Earth Syst. Sci., 12, 3709–3718, <https://doi.org/10.5194/nhess-12-3709-2012>, 2012.

Hoegh-Guldberg, O., Mumby, P. J., Hooten, A. J., Steneck, R. S., Greenfield P., Gomez E., et al.: Coral Reefs Under Rapid Climate Change and Ocean Acidification. *Science*, 80(318), 2007.

1070 Hsu, C.-E., Serafin, K. A., Yu, X., Hegermiller, C. A., Warner, J. C., and Olabarrieta, M.: Total water levels along the South Atlantic Bight during three along-shelf propagating tropical cyclones: relative contributions of storm surge and wave runup, Nat. Hazards Earth Syst. Sci., 23, 3895–3912, <https://doi.org/10.5194/nhess-23-3895-2023>, 2023.

1075 Hunter, J.: A simple technique for estimating an allowance for uncertain sea level rise, *Clim. Change*, 113(2), 239–252, doi:10.1007/s10584-011-0332-1, 2012.

Idier, D., Paris, F., Le Cozannet, G., Boulahya, F., Dumas, F.: Sea-level rise impacts on the tides of the European Shelf, *Continental Shelf Research*, Volume 137, 56-71, <https://doi.org/10.1016/j.csr.2017.01.007>, 2017.

1080 Idier D., Bertin X., Thompson P., and Pickering M. D.: Interactions between mean sea level, tide, surge, waves and flooding: Mechanisms and contributions to sea level variations at the coast. *Surveys in Geophysics*, 40(6):1603–1630, 2019.

Igigabel, M., Diab, Y. and Yates M.: Exploring Methodological Approaches for Strengthening the Resilience of Coastal Flood Protection System. *Front. Earth Sci.* 9:756936. doi: 10.3389/feart.2021.756936, 2021.

1085 Igigabel, M., Nédélec, Y., Bérenger, N., Flouest, N., Bernard, A., Chassé, P., et al.: Guidelines for Analysing Coastal Flood Protection Systems After a Submersion. *Water* 14, 15. doi:10.3390/w14010015, 2022.

IPCC: Climate Change 2001: Third Assessment Report (TAR), 2001.

1090

IPCC: Climate Change 2007: Fourth Assessment Report (AR4), 2007.

1095 IPCC: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp., 2014.

1100 IPCC: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate [H.- O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, M. Nicolai, A. Okem, J. Petzold, B. Rama, N. Weyer (eds.)], Available online: <https://www.ipcc.ch/srocc/> (accessed on 20 December 2021), 2019.

1105 IPCC: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)], Cambridge University Press, 2021.

[Kaack, L., Donti, P., Strubell, E. & Rolnick, D. Artificial intelligence and climate change: Opportunities, considerations, and policy levers to align AI with climate change goals \(2021\). https://eu.boell.org/en/2020/12/03/artificial-intelligence-and-climate-change](https://eu.boell.org/en/2020/12/03/artificial-intelligence-and-policy-levers-to-align-ai-with-climate-change-goals)

1110 [Kennedy, A.B., Westerink, J.J., Smith, J.M., Hope, M.E., Hartman, M., Taflanidis, A.A., Tanaka, S., Westerink, H., Cheung, K.F., Smith, T., Hamann, M., Minamide, M., Ota, A., Dawson, C.: Tropical cyclone inundation potential on the Hawaiian Islands of Oahu and Kauai. Ocean Model. 52–53, 54–68. http://dx.doi.org/ 10.1016/j.ocemod.2012.04.009, 2012.](http://dx.doi.org/10.1016/j.ocemod.2012.04.009)

1115 [Khan, M. J. U., Beld, I., Wöppelmann, G., Testut, L., Latapy, A. and Pouvreau, N.: Extension of a high temporal resolution sea level time series at Socoa \(Saint-Jean-de-Luz, France\) back to 1875. Earth System Science Data. 15. 5739-5753. 10.5194/essd-15-5739-2023, 2023.](https://doi.org/10.5194/essd-15-5739-2023)

1120 [Krien, Y., Dudon, B., Roger, J., and Zahibo, N.: Probabilistic hurricane-induced storm surge hazard assessment in Guadeloupe, Lesser Antilles, Nat. Hazards Earth Syst. Sci., 15, 1711–1720, https://doi.org/10.5194/nhess-15-1711-2015, 2015.](https://doi.org/10.5194/nhess-15-1711-2015)

[Krien, Y., Dudon, B., Roger, J., Arnaud, G., and Zahibo, N.: Assessing storm surge hazard and impact of sea level rise in the Lesser Antilles case study of Martinique, Nat. Hazards Earth Syst. Sci., 17, 1559–1571, https://doi.org/10.5194/nhess-17-1559-2017, 2017.](https://doi.org/10.5194/nhess-17-1559-2017)

1125 [Kumar, V., Kedam, N., Sharma, K. V., Mehta, D. J., & Caloiero, T.: Advanced Machine Learning Techniques to Improve Hydrological Prediction: A Comparative Analysis of Streamflow Prediction Models. Water, 15\(14\), 2572, 2023](https://doi.org/10.3390/w15142572)

Lamoureux, S., Forbes, D., Bell, T. and Manson, G.: The impact of climate change on infrastructure in the western and central Canadian Arctic. In: From Science to Policy in the Western and Central Canadian Arctic: an Integrated Regional Impact Study (IRIS) of Climate Change and Modernization [Stern, G.A. and A. Gaden (eds.)], ArcticNet, Quebec, pp. 300–341, 2015.

[Lange, A.M.Z., Fiedler, J.W., Becker, J.M., Merrifield, M.A., Guza, R.T.: Estimating runup with limited bathymetry. Coast. Eng., 172, 104055, 2021.](#)

1135 Lantuit, H., Atkinson, D., Overduin P. P., Grigoriev, M., Rachold, V., Grosse, G. and Hubberten, H.: Coastal erosion dynamics on the permafrost-dominated Bykovsky Peninsula, north Siberia, 1951–2006, Polar Res., 30(1), 7341. <https://doi.org/10.3402/polar.v30i0.7341>, 2011.

1140 [Larrue S. and Chiron T.: « Les îles de Polynésie française face à l'aléa cyclonique », Vertigo, Volume 10 Numéro 3 URL : http://journals.openedition.org/vertigo/10558 ; DOI : https://doi.org/10.4000/vertigo.10558, 2010.](#)

[Latapy, A., Arnaud, H., Pouvreau, N., and Weber N.: Reconstruction of sea level changes in Northern France for the past 300 years and their relationship with the evolution of the coastal zone, in: Coast 2017, Bordeaux, https://doi.org/10.13140/RG.2.2.14180.07041, 2017.](#)

1145 Lavaud, L., Bertin, X., Martins, K., Pezerat, M., Coulombier, T. and Dausse, D.: Wave dissipation and mean circulation on a shore platform under storm wave conditions. Journal of Geophysical Research: Earth Surface, 127, e2021JF006466. <https://doi.org/10.1029/2021JF006466>, 2022.

1150 [Lecacheux, S., Pedreros, R., Le Cozannet, G., Thiebot, J., De La Torre, Y. and Bulteau, T.: A method to characterize the different extreme waves for islands exposed to various wave regimes: a case study devoted to Reunion Island, Nat. Hazards and Earth Syst. Sci., 12\(7\), 2425-2437, 2012.](#)

1155 [Le Cozannet, G., Garcin, M., Petitjean, L., Cazenave, A., Becker, M., Meyssignac, B., Walker, P., Devilliers, C., Le Brun, O., Lecacheux, S., Baills, A., Bulteau, T. Yates, M., and Wöppelmann, G.: Exploring the relation between sea level rise and shoreline erosion using sea level reconstructions: an example in French Polynesia. Journal of Coastal Research 65\(sp2\), 2137-2142. https://doi.org/10.2112/SI65-361.1, 2013.](#)

1160 [Le Gorgeu, V. and Guitonneau, R. : Reconstruction de la Digue de l'Est à Dunkerque, Coast. Eng., 5, 555–586, https://icce-ojs-tamu.tdl.org/icce/index.php/icce/article/viewFile/2043/1716, 1954.](#)

Lin, N. and Chavas, D.: On hurricane parametric wind and applications in storm surge modeling. J. Geophys. Res., 117, D09120, doi:10.1029/2011JD017126, 2012.

1165 Lowe J. A., Woodworth P. L., Knutson T., McDonald R. E., McInnes K. L., Woth K., von Storch H., Wolf J., Swail V., Bernier  
N. B., Gulev S., Horsburgh K. J., Unnikrishnan A. S., Hunter J. R. and Weisse R.: Past and future changes in extreme sea  
levels and waves. In: Woodworth PL, Aarup T, Wilson WS, Church JA (eds) Understanding sea-level rise and variability.  
Wiley, New York, 2010.

170 Martinez-Asensio, A., Wöppelmann, G., Ballu, V., Becker, M., Testut, L., Magnan, A.K. et al.: Relative Sea-level rise and the  
influence of vertical land motion at tropical Pacific Islands. Global and Planetary Change, 176, 132–143,  
<https://www.sciencedirect.com/science/article/pii/S0921818118306751>, 2019.

Martins K., Bertin X., Mengual B., Pezerat M., Lavaud L., Guérin T. and Zhang Y.J.: Wave-induced mean currents and setup  
over barred and steep sandy beaches. Ocean Modelling, Volume 179, November 2022, 102110, 2022.

175 Maspataud A., Ruz M.-H. and Vanhée S.: Potential impacts of extreme storm surges on a low-lying densely populated  
coastline: the case of Dunkirk area, Northern France. Nat Hazards 66:1327–1343, 2013.

180 Masson, A.: The extratropical transition of Hurricane Igor and the impacts on Newfoundland. Nat Hazards 72, 617–632.  
<https://doi.org/10.1007/s11069-013-1027-x>, 2014.

185 Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M.  
I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B.:  
Climate Change 2021: The Physical Science Basis, Contribution of Working Group I to the Sixth Assessment Report of the  
[Intergovernmental Panel on Climate Change.](https://www.ipcc.ch/report/ar6/wg1/downloads/report/IPCC_AR6_WGI_TS.pdf)  
[https://www.ipcc.ch/report/ar6/wg1/downloads/report/IPCC\\_AR6\\_WGI\\_TS.pdf](https://www.ipcc.ch/report/ar6/wg1/downloads/report/IPCC_AR6_WGI_TS.pdf), 2021.

190 Masselink, G. and Short, A. D.: "The effect of tidal range on beach morphodynamics and morphology: a conceptual beach  
model". Journal of Coastal Research. 9 (3): 785–800. ISSN 0749-0208, 1993.

Masselink, G., Castelle, B., Scott T., Dodet, G., Suarez, S., Jackson, D. and Floc'h, F.: Extreme wave activity during  
2013/2014 winter and morphological impacts along the Atlantic coast of Europe, Geophys. Res.Lett., 43, 2135–2143,  
doi:10.1002/2015GL067492, 2016.

195 Mehta, D., Hadvani, J., Kanthariya, D., and Sonawala, P.: Effect of land use land cover change on runoff characteristics using curve number: A GIS and remote sensing approach. International Journal of Hydrology Science and Technology, 16(1), 1-16, 2023.

200 Mentaschi, L., Vousdoukas, M., Voukouvalas, E., Sartini, L., Feyen L., Besio, G., Alfieri, L.: Non-stationary extreme value analysis: a simplified approach for Earth science applications, Hydrol. Earth Syst. Sci. Discuss., 1–38, 2016.

Mentaschi L, Vousdoukas MI, García-Sánchez G, Fernández-Montblanc T, Roland A, Voukouvalas E, Federico I, Abdolali A, Zhang YJ and Feyen L: A global unstructured, coupled, high-resolution hindcast of waves and storm surge. Front. Mar. Sci. 10:1233679. doi: 10.3389/fmars.2023.1233679, 2023

205 Moss, R., Babiker, M., Brinkman, S., Calvo, E., Carter, T., Edmonds, J., et al.: Towards new scenarios for analysis of emissions, climate change, impacts and response strategies. IPCC Expert Meeting Report, 25 pp, 19–21 September, 2007, Noordwijkerhout, The Netherlands, <https://www.ipcc.ch/site/assets/uploads/2018/05/expert-meeting-ts-scenarios-1-1.pdf>, 2008.

210 Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., van Vuuren, D. P., et al.: The next generation of scenarios for climate change research and assessment, Nature, 463(7282), 747–756. <https://doi.org/10.1038/nature08823>, 2010.

215 Muis, S., Apecechea, M. I., Dullaart, J., de Lima Rego, J., Madsen, K. S., Su, J., et al.: A high-resolution global dataset of extreme sea levels, tides, and storm surges, including future projections. Frontiers in Marine Science, 7, 263. <https://doi.org/10.3389/fmars.2020.00263>, 2020.

220 Mentaschi, L., Vousdoukas, M., Voukouvalas, E., Sartini, L., Feyen L., Besio, G., Alfieri, L.: Non-stationary extreme value analysis: a simplified approach for Earth science applications, Hydrol. Earth Syst. Sci. Discuss., 1–38, 2016.

225 Moss, R., Babiker, M., Brinkman, S., Calvo, E., Carter, T., Edmonds, J., et al.: Towards new scenarios for analysis of emissions, climate change, impacts and response strategies. IPCC Expert Meeting Report, 25 pp, 19–21 September, 2007, Noordwijkerhout, The Netherlands, <https://www.ipcc.ch/site/assets/uploads/2018/05/expert-meeting-ts-scenarios-1-1.pdf>, 2008.

~~Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., van Vuuren, D. P., et al.: The next generation of scenarios for climate change research and assessment, *Nature*, 463(7282), 747–756. <https://doi.org/10.1038/nature08823>, 2010.~~

1230

Nicholls, R. J., Hanson, S., Herweijer, C., and Patmore, N.: Ranking port cities with high exposure and vulnerability to climate extremes, <https://www.oecd-ilibrary.org/content/workingpaper/011766488208>, 2008.

1235

Oppenheimer M., Glavovic B., Hinkel J. et al.: Chapter 4: sea level rise and implications for low lying islands, coasts and communities. In: Pörtner H.O., Roberts D., Masson-Delmotte V. et al (eds). IPCC Special Report on the Ocean and Cryosphere in a Changing Climate. Intergovernmental Panel on Climate Change (IPCC), 2019.

1240

Pagney, F.: Genèse et dynamique de l'ouragan Hugo sur la Guadeloupe, In *Annales de géographie* (pp. 152- 165), Armand Colin, (1991).

~~Patlakas, P., Stathopoulos, C., Tsalis, C., and Kallos, G.: Wind and wave extremes associated with tropical-like cyclones in the Mediterranean basin. *International Journal of Climatology*, 41(S1), E1623–E1644. <https://doi.org/10.1002/joc.6795>, 2021.~~

1245

~~Pickering M.D., Wells N.C., Horsburgh K.J., Green J.A.M.: The impact of future sea-level rise on the European Shelf tides. *Continental Shelf Research*, Volume 35, 2012, Pages 1-15, ISSN 0278-4343, <https://doi.org/10.1016/j.csr.2011.11.011>, 2012.~~

~~Pirazzoli, P. A. and Montaggioni, L. F.: Holocene sea level changes in French Polynesia, *Paleogeography, paleoclimatology paleoecology*, 68 (2-4), 153-175, 1988.~~

1250

Planton, S., Le Cozannet, G., Cazenave, A., Costa, S., Douez, O., Gaufres, P. et al.: Le climat de la France au XXI<sup>e</sup> siècle. Vol 5: Changement climatique et niveau de la mer: de la planète aux côtes françaises. [Rapport de recherche] Ministère de l'Écologie et du Développement Durable, 2015.

1255

~~Quentel, E., Loevenbruck, A., Hébert, H., and Allgeyer, S.: Tsunami hazard in La Réunion island from numerical modeling of historical events, *Nat. Hazards Earth Syst. Sci. Discuss.*, 1, 1823–1855, <https://doi.org/10.5194/nhessd-1-1823-2013>, 2013.~~

Raftery, A. E., Zimmer, A., Frierson, D. M. W., Startz, R. and Liu, P.: Less than 2 °C warming by 2100 unlikely, *Nat. Clim. Chang.*, 7, 637–641, <https://doi.org/10.1038/nclimate3352>, 2017.

- 1260 Ranasinghe, R.: Assessing climate change impacts on open sandy coasts: A review. *Earth-Science Reviews*, Volume 160, Pages 320-332, ISSN 0012-8252, <https://doi.org/10.1016/j.earscirev.2016.07.011>, 2016
- Rasmussen, D. J., Kulp, S., Kopp, R. E., Oppenheimer M. and Strauss B. H.: Popular extreme sea level metrics can better communicate impacts, *Climatic Change* 170, 30, <https://doi.org/10.1007/s10584-021-03288-6>, 2022.
- 1265 [Rego, J.L., LI, C.: Nonlinear terms in storm surge predictions: effect of tide and shelf geometry with case study from Hurricane Rita. \*J. Geophys. Res.\* 115, C06020, 2010.](#)
- [Rolnick, D. et al. Tackling Climate Change with Machine Learning. \*ACM Comp. Surv.\* 55, 42:1–42:96 \(2022\).](#)
- 1270 [Romero, R., and Emanuel, K.: Climate change and hurricane-like extratropical cyclones: Projections for north Atlantic polar lows and medicanes based on cmip5 models. \*Journal of Climate\*, 30\(1\), 279–299. <https://doi.org/10.1175/JCLI-D-16-0255.1>, 2016.](#)
- 1275 [Ruggiero, P., Komar, P.D., McDougal, W.G., Marra, J.J., Beach, R.A.: Wave runup, extreme water levels and erosion properties backing beaches. \*J. Coast. Res.\*, 17, 407–419, 2001.](#)
- Ruggiero, P., Buijsman, M., Kaminsky, G. ~~and~~ Gelfenbaum, G.: Modeling the effects of wave climate and sediment supply variability on large-scale shoreline change. *Marine Geology* 273, 127-140, 2010.
- 1280 [Saffache, P., Marc, J.-V., and Huyghes-Belrose, V.: Les cyclones en Guadeloupe : quatre siècles cataclysmiques, IBIS Rouge Editions, Martinique, 2003.](#)
- [Sahal, A. and Morin, J.: Effects of the October 25, 2010, Mentawai tsunami in La Réunion Island \(France\): observations and crisis management. \*Nat Hazards\* 62, 1125–1136. <https://doi.org/10.1007/s11069-012-0136-2>, 2012.](#)
- 1285 [Shaikh, M. M., Lodha, P., Lalwani, P. and Mehta, D.: Climatic projections of Western India using global and regional climate models. \*Water Practice & Technology\*, 17\(9\), 1818-1825, 2022.](#)
- 1290 Seneviratne, S.I., Zhang X., Adnan M., Badi W., Dereczynski C., Di Luca A. et al.: Weather and Climate Extreme Events in a Changing Climate. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K.



- 1295 Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1513–1766, doi:10.1017/9781009157896.013, 2021.
- Sheremet, A., Staples, T., Arduhin, F., Suanez, S. & Fichaut, B.: Observations of large infragravity runup at Banneg Island, France, *Geophysical Research Letters*, 41, 976–982, <https://doi.org/10.1002/2013GL058880>, 2014.
- 1300 Slangen, A.B.A., Adloff, F., Jevrejeva, S., Leclercq, P.W., Marzeion, B., Wada, Y., and Winkelmann R.: A review of recent updates of sea-level projections at global and regional scales, *Surv. Geophys.*, 38, no. 1, 385–406, doi:10.1007/s10712-016-9374-2, 2017.
- 1305 [Stephens, S.A., Coco, G., Bryan, K.R.: Numerical simulations of wave setup over barred beach profiles: Implications for predictability. \*J. Waterw. Port Coast. Ocean Eng.\* 137 \(4\), 175–181. \[http://dx.doi.org/10.1061/\\(ASCE\\)WW.1943-5460.0000076\]\(http://dx.doi.org/10.1061/\(ASCE\)WW.1943-5460.0000076\), 2011.](https://doi.org/10.1061/(ASCE)WW.1943-5460.0000076)
- 1310 [Stockdon, H.F., Holman, R.A., Howd, P.A., Sallenger, A.H.: Empirical parameterization of setup, swash, and runup. \*Coast. Eng.\* 53, 573–588. <https://doi.org/10.1016/j.coastaleng.2005.12.005>, 2006.](https://doi.org/10.1016/j.coastaleng.2005.12.005)
- Syvitski, J.P.M.: Deltas at risk, *Sustainability Science* 3, 23–32, <https://doi.org/10.1007/s11625-008-0043-3>, 2008.
- Talke, S. A., Familkhalili, R., Helaire, L. T., Jay, D. A., Orton, P. M. & Ralston, D. K.: The influence of human induced landscape and bathymetry changes on tides, surge and extreme water levels. In *Ocean Sciences Meeting 2020. AGU*, 2020.
- 1315 [Thiéblemont, R., Le Cozannet, G., D’Anna, M. et al.: Chronic flooding events due to sea-level rise in French Guiana. \*Sci Rep\* 13, 21695. <https://doi.org/10.1038/s41598-023-48807-w>, 2023.](https://doi.org/10.1038/s41598-023-48807-w)
- 1320 [Thiébot J., Idier D., Garnier R., Falquès A., Ruessink G.: The influence of wave direction on the morphological response of a double sandbar system. \*Cont Shelf Res\* 32:71–85. <https://doi.org/10.1016/j.csr.2011.10.014>, 2012](https://doi.org/10.1016/j.csr.2011.10.014)
- 1325 Tiggeloven, T., de Moel, H., Winsemius, H. C., Eilander, D., Erkens, G., Gebremedhin, E., Diaz Loaiza, A., Kuzma, S., Luo, T., Iceland, C., Bouwman, A., van Huijstee, J., Ligtoet, W., and Ward, P. J.: Global-scale benefit–cost analysis of coastal flood adaptation to different flood risk drivers using structural measures, *Natural Hazards and Earth System Sciences*, 20, 1025–1044, <https://doi.org/10.5194/nhess-20-1025-2020>, 2020.

Tiggeloven, T., Couasnon, A., van Straaten, C. et al.: Exploring deep learning capabilities for surge predictions in coastal areas. Sci Rep 11, 17224. <https://doi.org/10.1038/s41598-021-96674-0>, 2021.

1330 Toomey, T., Amores, A., Marcos, M., Orfila, A., & Romero, R.: Coastal hazards of tropical-like cyclones over the Mediterranean Sea. Journal of Geophysical Research: Oceans, 127, e2021JC017964. <https://doi.org/10.1029/2021JC017964>, 2022.

1335 Ullmann A., Pirazzoli P. A. and Tomasin A.: Sea surges in Camargue: Trends over the 20th century. Cont Shelf Res 27:922–934, 2007.

U.S. Army Corps of Engineers: Coastal Engineering Manual, U.S. Army Corps of Engineers, Washington, DC., 2002.

1340 van Ormondt, M., Roelvink D., and van Dongeren A.: "A Model-Derived Empirical Formulation for Wave Run-Up on Naturally Sloping Beaches" Journal of Marine Science and Engineering 9, no. 11: 1185. <https://doi.org/10.3390/jmse9111185>, 2021.

van Vuuren, D.P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., et al.: The representative concentration pathways: an overview. Climatic Change 109, 5. <https://doi.org/10.1007/s10584-011-0148-z>, 2011.

1345 Verma, S., Verma, M. K., Prasad, A. D., Mehta, D., Azamathulla, H. M., Muttill, N., and Rathnayake, U.: Simulating the Hydrological Processes under Multiple Land Use/Land Cover and Climate Change Scenarios in the Mahanadi Reservoir Complex, Chhattisgarh, India. Water, 15(17), 3068, 2023.

1350 Vitousek S., Barnard P. L., Fletcher C. H., Frazer N., Erikson L., and Storlazzi C. D.: Doubling of coastal flooding frequency within decades due to sea-level rise. Scientific Reports, 7(1), 2017.

Vousdoukas, M., Voukouvalas, E., Annunziato, A., Giardino, A., and Feyen, L.: Projections of extreme storm surge levels along Europe. Climate Dynamics, 47, 3171, 3190. <https://doi.org/10.1007/s00382-016-3019-5>, 2016.

1355 Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Verlaan, M., and Feyen, L.: Extreme sea levels on the rise along Europe's coasts. Earth's Future, 5, 304–323. doi:10.1002/2016EF000505, 2017.

- 1360 Vousdoukas, M.I., Mentaschi, L., Voukouvalas, E., Verlaan, M., Jevrejeva, S., Jackson L. P. & Feyen, L.: Global probabilistic  
projections of extreme sea levels show intensification of coastal flood hazard. *Nat Commun* 9, 2360.  
<https://doi.org/10.1038/s41467-018-04692-w>, 2018.
- 1365 [Waycott, M., Duarte, C. M., Carruthers, T. J. B., Orth, R. J., Dennison, W. C., Olyarnik, S., Calladine, A., Fourqurean, J. W., Heck, K. L., Hughes, A. R., Kendrick, G. A., Kenworthy, W. J., Short, F. T., and Williams, S. L.: Accelerating loss of seagrasses across the globe threatens coastal ecosystems. \*P. Natl. Acad. Sci. USA\*, 106, 12377–12381, 2009.](#)
- [Webb, A.P. and Kench, P.S.: The dynamic response of reef islands to sea level rise: Evidence from multi-decadal analysis of island change in the Central Pacific. \*Global and Planetary Change\*, 72, 234–246, 2010.](#)
- 1370 [Weisse R., von Storch H., Niemeier H.D. and Knaack H.: Changing North Sea storm surge climate: an increasing hazard? \*Ocean Coast Manag\* 68:58–68, 2012.](#)
- 1375 [Wong, P. P., Losada, I. J., Gattuso, J.-P., Hinkel, J., Khattabi, A., K. L. McInnes, Saito, Y., and Sallenger, A.: Coastal systems and low-lying areas, in: \*Climate Change 2014: Impacts, Adaptation, and Vulnerability, Part A: Global and Sectoral Aspects, Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change\*, edited by: Field, C. B., Barros, V. R., Dokken, D. J., Mach, K. J., Mastrandrea, M. D., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y. O., Genova, R. C., Girma, B., Kissel, E. S., Levy, A. N., MacCracken, S., Mastrandrea, P. R., and White, L. L., Cambridge University Press, Cambridge, UK and New York, NY, USA, 361–409, 2014.](#)
- 1380 [Yates M., Le Cozannet G., Garcin, M., Salaï, E. and Walker, P.: Multi-decadal shoreline change on Manihi and Manuae, French Polynesia. \*Journal of Coastal Research\*, 2013.](#)
- [Zahibo, N., Pelinovsky, E., Talipova, T., Rabinovich, A., Kurkin, A., and Nikolkina, I.: Statistical analysis of cyclone hazard for Guadeloupe, Lesser Antilles, \*Atmos. Res.\*, 84, 13–29, 2007.](#)