



# A systemic and comprehensive assessment of coastal hazard changes: method and application to France and its overseas territories

Marc Igigabel<sup>1</sup>, Marissa Yates<sup>2,3</sup>, Michalis Vousdoukas<sup>4</sup>, Youssef Diab<sup>5</sup>

<sup>1</sup>Cerema, Technopôle Brest Iroise, 155 rue Pierre Bouguer, BP5, 29280 Plouzané, France

5 <sup>2</sup>LHSV & Cerema, 6 quai Watier, BP 49, 78401 Chatou, France

<sup>3</sup>LHSV, Ecole des Ponts, EDF R&D, 6 quai Watier, BP 49, 78401, Chatou, France

<sup>4</sup>European Commission Joint Research Centre, Via Enrico Fermi 2749, I-21027-Ispra, Italy.

<sup>5</sup>Université Gustave Eiffel, Lab'Urba, 5, Boulevard Descartes, 77430 Champs-sur-Marne, France

*Correspondence to:* Marc Igigabel (marc.igigabel@cerema.fr)

10 **Abstract.** In the context of climate change, height and frequency variations of extreme sea levels (ESL) are studied using  
deterministic and probabilistic approaches. However, this type of approach does not highlight the dynamic effects (waves,  
currents) generated by meteocean events, beyond their effects on sea-levels. In particular, ESL estimates are certainly  
calculated by considering the main determining physical factors but cannot report on all the effects of these factors.  
Ultimately, this can lead to confusion between ESL and hazard. This article proposes a systemic assessment method to  
15 analyze coastal hazard changes at regional scales, integrating parameters influencing sea-levels, as well as factors describing  
the geomorphological context (length and shape of the coast, width of the continental shelf), meteocean events (storms,  
cyclones and tsunamis), and the marine environment (e.g., coral reef state and sea ice extent). French mainland and overseas  
territories were selected to apply the method. We have thus highlighted the need to study not only the sea-level variability,  
but also the current and future characteristics of meteocean events. The long, concave coasts bordered by a wide continental  
20 shelf appear particularly sensitive to variations in the intensity or trajectory of meteocean events. Coral reef degradation in  
the tropics and the decrease in seasonal sea ice extent in the polar regions can also significantly change the nearshore  
hydrodynamics and impacts on the shoreline. These results help to predict the types of hazard (shoreline erosion, rapid  
submersion and/or permanent flooding) that will increase the most in different coastal zones.

## 1 Introduction

25 In 2001, the frequency and intensity of extreme climate events, including marine flooding from tropical and other storms was  
identified by the Intergovernmental Panel on Climate Change (IPCC) as one of the five « reasons for concern » (IPCC, 2001,  
2007). Recent research has improved knowledge of the mechanisms generating sea-level rise. Frederikse et al. (2020)  
showed that since 1900 the sum of the contributions to sea-level change is consistent with the trends and multidecadal  
variability in observed sea-level on both global and basin scales. However, significant uncertainties remain about the  
30 evolution of hazards at the coastline generated by the marine environment. In this article, these hazards will be referred to as



« coastal hazards », with the understanding, however, that the coast may be affected by other sources of hazard (intense precipitation, overflow of streams, etc.).

The issue of the evolution of coastal hazards is often addressed through the evolution of extreme sea levels (ESLs) along the coast. For example, Vitousek et al. (2017) warned that the 10 to 20 cm of sea-level rise expected no later than 2050, will more than double the frequency of extreme water-level events in the tropics. Vousdoukas et al. (2018) projected a very likely increase of the global average 100-year ESL between 2000 and 2100 of 34–76 cm under a moderate-emission-mitigation-policy scenario and of 58–172 cm under a business as usual scenario. Under these scenarios, a large part of the tropics would be exposed annually to the present-day 100-year ESL in 2050. These assertions concern the evolution of water levels at the coast, but are not sufficient to describe the evolution of coastal hazards. Yet many authors evaluating changes in coastal risks restrict the representation of the hazard to ESLs (e.g. Hallegatte et al. (2013), Tiggeloven et al. (2020), Rasmussen et al. (2020)). This kind of analysis uses a deterministic approach based on a single factor (the ESL that is expressed by two parameters: maximum water level and frequency) to represent a complex hazard that can only be fully addressed using a systemic approach that considers multiple factors (Igigabel et al., 2021).

First, it is necessary to distinguish between global mean sea level (GMSL) and relative sea level (RSL). The GMSL rise can be defined as the global change in mean sea level (MSL) relative to the terrestrial reference frame, due to the combined effects of change in the volume of the ocean and change in the level of the sea floor (Church et al., 2013). The RSL rise can be defined as the change in sea-level with respect to the land (Lowe et al., 2010).

Second, it is necessary to assess the main factors influencing the evolution of water levels, whether they are GMSL, RSL or ESLs.

MSL changes, both globally and locally, vary according to seasonal, annual, and longer time scales. These variations may be caused by changes in the mass of water in the ocean (e.g., due to melting of glaciers and ice sheets and changes in terrestrial water storage), by changes in ocean water density (e.g., volume expansion as the water warms), by changes in the shape of the ocean basins (e.g., due to plate tectonics) and changes in the Earth's gravitational and rotational fields, and by local subsidence or uplift of the land (Oppenheimer et al., 2019). In the future, changes in the GMSL are expected to be strongly controlled by the greenhouse gas (GHG) emission scenarios (Oppenheimer et al., 2019; Fox-Kemper et al., 2021) and cryosphere evolution: in case of faster melting of the Greenland and Antarctic ice caps, Bamber et al. (2019) estimate the increase in GMSL above 2 m in the 21st century.

The RSL will depend on both the GMSL and local vertical land movement (VLM). VLM can have natural causes (e.g. isostasy, elastic flexure of the lithosphere, earthquakes and volcanoes, landslides and sedimentation) and, more locally,



65 anthropogenic causes, in particular, soil loading, extraction of hydrocarbons and/or groundwater, drainage, mining activities  
(Gregory et al., 2019). During the 20th century, the coasts of Tokyo, Shanghai, and Bangkok collapsed by several meters as  
a result of water extraction and soil loading from the construction of buildings and infrastructure (Nicholls et al., 2008).  
Subsidence rates of the order of 1 to 10 cm/year are commonly measured in coastal megacities and are exceeding sea-level  
change rates by a factor of ten (Erkens et al., 2015). However, in the absence of subsidence, the majority of coastlines are  
70 expected to experience RSL changes on the order of 30 percent of GMSL rise (Gregory et al., 2019; IPCC, 2019; Dayan et  
al., 2021).

ESLs are also projected to change due to changes in RSL, tides, wind waves, and storm surges (Vousdoukas et al., 2018).  
Storm surges are defined by Gregory et al. (2019) as « the elevation or depression of the sea surface with respect to the  
75 predicted tide during a storm ». The predominant factor in the evolution of ESLs is the variability in sea levels generated by  
astronomical tides and storm surges (Buchanan et al., 2016). The combination of tide and storm surge phenomena requires a  
statistical approach. Thus, the change in ESL events is commonly expressed in terms of an amplification factor and an  
allowance. The amplification factor denotes the amplification in the average occurrence frequency of a certain extreme  
event, often referenced to the water level with a 100-year return period estimated from historical data. The allowance denotes  
80 the increased height of the water level [m] with a given return period. This allowance equals the regional projection of RSL  
rise with an additional height related to the uncertainty in the projection (Hunter, 2012). Amplification factors are strongly  
determined by the local variability in ESL events. Locations where this variability is large due to large storm surges and  
astronomical tides will experience a moderate amplification of the occurrence frequency of extremes in comparison to  
locations with small variability. Globally, this contrast between regions with large and small amplification factors becomes  
85 clear for projections by mid-century and considerable in the coming centuries (Vitousek et al., 2017). In particular, many  
coastal areas in the lower latitudes may expect amplification factors of 100 or larger by mid-century, regardless of the GHG  
emissions scenario. By the end of the century, and in particular if GHG emissions are not reduced, such amplification factors  
may be widespread along global coastlines (Vousdoukas et al., 2018).

90 This review of the evolution of sea levels (GMSL, RSL and ELS) highlights the importance of the evolution of coastal  
hazards. How will these hazards change with evolving water levels? How can other factors such as the geomorphological  
configuration of the coast, the types of meteocean phenomena, and other changes in the marine environment be incorporated  
into the analyses?

95 The objective of this article is to present a systemic approach to assess the evolution of coastal hazards at regional scales.  
The method will be applied to France and its overseas territories, which are located in different latitudes (equator, tropics and  
temperate zones), exposed to different climates, and characterized by different geomorphological configurations (continental



or island). Finally, as part of the discussion, the results will be put in perspective with the anticipated effects of climate change on the coast. This analysis allows identifying different types of hazard evolution in coastal areas.

## 100 2 Method for assessing changes in coastal hazards at regional scales

To assess the evolution of coastal hazards, at regional and multidecadal scales, it is necessary to combine a quantitative approach evaluating the parameters characterizing ESLs and a qualitative approach analyzing other factors more difficult to quantify, such as the meteo-oceanic event type, the geomorphological configuration of the coast and other environmental factors (e.g. coral reef degradation in the tropics and decrease in seasonal sea ice extent in the polar regions) that may change the hydraulic actions on the shoreline.

Figure 1 shows the three proposed steps. These three stages will be presented in succession. In applying the method, however, it is possible to implement the first two steps in parallel.




Figure 1: Method for assessing changes in coastal hazards



110 Before presenting these three steps, it is essential to remember that flood risk, and more generally, all coastal risks, can be defined in at least two alternative ways (FLOODsite, 2009). The first definition considers risk to be the result of the exposure of a vulnerable stake to a hazard, which is reflected in the following formula:

$$\text{risk} = \text{hazard (meteo-oceanic event)} * \text{exposure} * \text{vulnerability (of the society/area/structure)} \quad (1)$$

115 However, in an attempt to quantify risk, and considering that the word « risk » suggests a probability of occurrence, a second definition may be highlighted:

- the two terms « exposure » and « vulnerability » can be substituted by « consequences », with the consequences being generally more quantifiable (for example, in number of fatalities and economic damage) than the previous two terms;
- 120 • the hazard can be represented by its probability distribution.

This yields the second definition:

$$\text{risk} = \text{probability (of the hazard)} * \text{consequences} \quad (2)$$

125 This article is mainly devoted to the evaluation of the hazard. Consequences (e.g. erosion and flooding) will only be addressed in the discussion. However, reference to risk definitions is necessary because the expression of hazard depends on the definition of risk.

Quantitative assessments of ESL changes (Step 1 of Fig. 1) are probabilistic and therefore fall within the second definition of risk. The challenge of the proposed method is to use these evaluations, in combination with other factors, such as the types of  
130 meteocean events, the geomorphological configuration of the coast and changes in the marine environment affecting hydraulic conditions. These « other factors », which are more difficult to quantify, will be studied in a second, parallel phase (step 2 of Fig. 1). The last step is to describe the evolution of the hazard within the framework of a systemic approach. In this step (step 3 of Fig. 1), the first definition of risk is applied to search for a sufficiently synthetic way of presenting the data to  
135 produce an overall assessment.

Given the multiple effects of climate change on sea levels (GMSL and RSL), atmospheric conditions, and wave conditions, the evolution of the factors and parameters can only be assessed using global development hypotheses. To represent different trajectories, the IPCC identified five scenarios referred to as SSPx-y, where ‘SSPx’ refers to the Shared Socio-economic  
140 Pathway (‘SSP’ describing the socio-economic trends underlying the scenario), and ‘y’ refers to the approximate level of radiative forcing (in  $\text{W m}^{-2}$ ) resulting from the scenario in the year 2100. The application of the proposed method must be preceded by an analysis of exposure and vulnerability, as well as a reflection on the duration of the project. For long-term

adaptations, a general description of the evolution of the hazard, including the estimated uncertainties, is more appropriate than a forecast (likely imprecise) of the evolution of a parameter at a given date.

## 145 2.1 Step 1: ESL change assessment

ESLs are the result of the combination of the RSL, the tide level, and the storm surges produced by meteo-oceanic events. To understand the evolution of ESLs, it is assumed that meteo-oceanic events simultaneously generate atmospheric storm surge (caused by atmospheric depressions, wind) and energetic waves. At the coast, waves also contribute to the total water level by two phenomena:

- 150 • wave set-up: a time-varying (on the order of 30 minutes), wave-driven increase in the mean water level at the coast, resulting from wave dissipation in the nearshore (Bowen et al., 1968);
- run-up: the time-varying (on the order of the wave period, i.e. a few second, or the period of infragravity-waves, i.e. a few minutes) water level reached by a wave on a beach or a coastal structure, relative to the static water level, measured vertically (run-up may generate overtopping which, unlike overflowing, occurs intermittently).

155

In the proposed method, storm surge and wave set-up will be taken into account to estimate the ESL, but wave run-up will be neglected because of the challenge associated with modeling it at fine scales.

To fully account for changes in ESLs, the analysis must include both changes in the average occurrence frequency of a certain extreme event and the increased water level (in absolute terms and as a percentage) with a given return period. For this reason, the second definition of risk will be used to study the joint probabilities of the parameters impacting the sea level, namely the RSL, the tidal range and the storm surge (including the effects of waves). Following Vousdoukas et al. (2017) changes in the magnitude and frequency of occurrence of the present 100-year ESL ( $ESL_{100}$ ) are evaluated. It is assumed that ESL are driven by the combined effect of MSL, tides ( $\eta_{tide}$ ) and water level fluctuations due to waves and storm surges ( $\eta_{w-ss}$ ). As a result, ESL can be defined as:

165

$$ESL = MSL + \eta_{tide} + \eta_{w-ss} \quad (3)$$

The climate extremes contribution  $\eta_{w-ss}$  from waves and storm surge can be estimated as:

$$\eta_{w-ss} = SSL + 0.2 H_s \quad (4)$$

170

where SSL is the storm surge level,  $H_s$  is the significant wave height and  $0.2 H_s$  is a generic approximation of the wave set-up (U.S. Army Corps of Engineers, 2002). This equation is a conservative estimate of the wave set-up, which mostly depends on the local slope, breaking wave height, and wave period, and may be closer to 10% of the breaking wave height. Larger values may only be observed at steep sandy beaches (e.g. Martins et al., 2022) or steep shore platforms (Sheremet et



175 al., 2014; Lavaud et al., 2022). This expression is used here since the objective of this work is not to improve existing  
deterministic methods, but to present a systemic and comprehensive method for assessing the evolution of coastal hazards.

The tidal range may have a strong influence on the frequency of ESLs, in particular for microtidal coasts (strong  
amplification factor for low tidal ranges).

180

Storm surge and wave setup affect the frequency of ESLs (strong amplification factor for low surges) and can also influence  
the allowance. For example, through a statistical analysis of tide gauge observations, Calafat et al. (2022) showed that trends  
in surge extremes and sea-level rise both made comparable contributions to the overall change in ESLs in Europe since 1960.  
However, on the three maritime facades of France, the relative importance of this factor is expected to decline gradually in  
185 the 21st century due to increases in RSL becoming predominant (Vousdoukas et al., 2017).

In general, tidal simulations show no significant impacts of RSL rise on tidal amplitude during the 21st century at regional  
scales, although this does not exclude potential local effects (Haigh et al., 2020). For example, Idier et al. (2017) showed that  
notable increases in high tide levels occur in the northern Irish Sea, the southern part of the North Sea and the German Bight,  
190 and decreases occur mainly in the western English Channel. Depending on the location, they can account for  $\pm 15\%$  of RSL  
rise (as long as RSL rise remains smaller than 2 m). Given the strong uncertainties on RSL rise, we will assume here that the  
tides are in a steady state and, consequently, that the tidal range does not change the allowance.

## 2.2 Step 2: Evaluation of additional factors

195 Two types of regional factors can be considered essential for all coastlines: phenomena likely to generate energetic sea states  
and the geomorphological configuration. In addition, other environmental factors affecting hydraulic conditions should be  
considered in the tropics (in particular, coral reef state) and at the poles (in particular, sea ice extent).

### 2.2.1 Meteocean event types

Three main coast categories are considered in this method: exposure to no particular phenomenon, to storms, or to cyclones  
200 (which also implies exposure to storms). On these three categories are superimposed (i) the potential exposure to long swells  
generated by a meteo-oceanic event far from the studied coastline (ii) the potential exposure to tsunamis, since the origin of  
tsunamis is not related to the meteorological conditions.

The increase in wave damage could also be assessed, considering local changes in significant wave height (average height of  
205 the highest one-third of the waves in a given sea state). Trends in coastal wave climates are reported, with a low confidence  
level, in the IPCC (2019) report: for example, projections of future extreme significant wave height consistently estimate an



increase in the Southern Ocean and a decrease in the north-eastern Atlantic Ocean and Mediterranean Sea. However, these trends will not be explored in detail: the focus will be on the strong differences that already exist between the wave climates of the maritime facades.

## 210 2.2.2 Geomorphological configurations of the coast

It is necessary to identify characteristics of the coast that influence the different components of the surges.

215 First, the analysis of storm surges computed from the national REFMAR database reveals that they are controlled not only by storm tracks but also by the width of the continental shelf and the presence of shallow waters. For example, during the 20-year study period (1998-2018), storm surges hardly reached 1.0 m along the coastlines of the southern Bay of Biscay and the eastern Mediterranean Sea, but exceeded 2.0 m in the English Channel (Dodet et al., 2019). In shallow water, the wind effects can significantly dominate the effect of the atmospheric pressure, and the surge can be strongly amplified by the bathymetry along the coast (Bertin et al., 2012). The width and depth of the continental shelf have an important influence on the magnitude of storm surges.

220

Secondly, in regions exposed to cyclones, the surges measured on islands are much lower than those measured on continental facades. For example, on the islands of the West Indies, the storm surges rarely exceed 3 m (Cerema, 2019, 2020, 2021), whereas in the case of Katrina, which impacted the United States in 2005, surges in eastern Louisiana reached values between 3.05 m and 5.79 m (Graumann, 2006). During the same event, the surges exceeded 8 m at several locations along the Mississippi coast (Dietrich et al., 2010). This difference between the surges observed on continental coasts and island coasts is explained by the fact that, in the case of a hurricane, the impact of the low pressure associated with the storm on surge is minimal in comparison to the water being forced toward the shore by the wind. However, in the case of small islands (e. g. West Indies or La Réunion), the surge is generally reduced: wind-forced water accumulates less on short shorelines than on long shorelines (Durand, 1996). Therefore, the length of the maritime facade appears to be the second determining factor of the surges.

230

Thirdly, the maximum potential storm surge for a particular location is sensitive to small changes in storm intensity, forward speed, size, angle of approach to the coast, central pressure, and also the shape and characteristics of coastal features such as bays and estuaries<sup>1</sup>. Wind-forced water will accumulate less on convex-shaped shorelines than on concave-shaped shorelines. Therefore, the shape of the coast appears to be the third determining factor of the surges.

235

---

<sup>1</sup><https://www.nhc.noaa.gov/surge/>



Consequently, in order to account for the main geomorphological parameters determining the magnitude of the surge, we propose to use three criteria:

- 240
- the length of the coast;
  - the width of the continental shelf;
  - the convex or concave shape of the coastline.

### 2.2.3 Other environmental factors changing hydraulic conditions

245 Other phenomena induced by climate change can increase the intensity and frequency of extreme events, or increase their effects:

- ocean acidification combines with ocean warming and deoxygenation to impact ecosystems (e. g., coral reefs and oyster beds), which can reduce coastal flood protection. Under conditions expected in the 21st century, global warming and ocean acidification should compromise carbonate accretion, with corals becoming increasingly rare on reef systems. The result should be less diverse reef communities and carbonate reef structures that fail to be maintained. Climate change also exacerbates local stresses from declining water quality and overexploitation of key species, driving reefs increasingly toward the tipping point for functional collapse (Hoegh-Guldberg et al., 2007; Albright et al., 2018);
  - in polar regions, the decrease in seasonal sea ice extent in the Arctic, together with a longer open water season, provide less protection from storm impacts (Forbes, 2011). In addition, a longer fetch allows more energetic waves to form, which can have consequences outside of the polar regions, for example on coasts exposed to southern swells.
- 255

These two factors that affect wave formation and propagation, as well as currents, are also considered in the proposed method.

### 260 2.3 Step 3: global hazard assessment

In order to clearly distinguish the current approach from existing analytical approaches (e.g., Vitousek et al., 2017; Vousdoukas et al., 2018), it is important to note that RSL changes, tidal regimes and storm surge and wave impacts not only influence the evolution of coastal hazards through the sea level alone, but also can have an additional influence by the energy carried by waves and currents. The data used to draw up an overall summary should therefore not be limited to variations in the height and frequency of the ELSs. It is also necessary to incorporate the parameters necessary for their evaluation, and the other factors identified, namely the meteocean event type, the geomorphological configuration, and the other environmental factors changing hydrodynamics at the shoreline.

265



Based on this principle, the overall assessment of the hazard should be produced using the following factors and parameters  
270 (evaluated on the basis of the SSP5-8.5 scenario):

- RSL height change: ( $\Delta$ RSL);
- tidal range;
- surge with a 100-year return period;
- ESL height variation ( $\Delta$ ESL);
- 275 • ESL percentage change ( $\% \Delta$ ESL);
- return period of the present day 100-year ESL;
- geomorphological configuration of the coast;
- meteocean event type; and
- other environmental factors affecting the hydrodynamics at the shoreline.

### 280 3 Application of the proposed method to French coasts

The proposed method is applied to France and its overseas coasts (cf. Fig. 2).



**Figure 2: A map showing the French territories considered in this study. Credit : Cerema, after Hoshie.**

285

In the following, the islands of Saint-Martin, Saint-Barthélemy, Guadeloupe and Martinique may be referred to as the « West Indies ».



290 The choice of the scenario and the time scales should take into account the existence of the high stakes at the considered  
coasts (coastal cities, port and industrial facilities) and the strong uncertainties about the contributions of ice caps to the rise  
in water levels (Bamber et al., 2019; Dayan et al., 2021). For these two reasons, we use the SSP5-8.5 scenario. For the  
assessment of sea levels (GMSL, RSL and ESL), the median values are selected at two time scales: 2050 and 2100. Table 1  
shows the projections for the GMSL under the SSP5-8.5 scenario.

295 **Table 1: GMSL projections for 2050, 2100 and 2300 for the SSP5-8.5 scenario. Median values and ranges for the 17th to 83rd  
percentiles are shown using the 1986-2005 period as a reference (IPCC, 2019).**

Climate scenario	2050 Median	2050 17-83% range	2100 Median	2100 17-83% range	2300 Median	2300 17-83% range
SSP5-8.5	+ 32 cm	23 cm to 40 cm	+ 84 cm	61 cm to 110 cm	+385 cm	230 cm to 540 cm

300 The GMSL values displayed in Table 1 show high uncertainties in the long term: for the SSP5-8.5 scenario in 2100, the  
median value is 0.84 m and the 17th to 83rd percentile range is 0.61 to 1.10 m (the high value is almost double the low  
value). Similar (or higher, since other local phenomena must be considered) uncertainties exist for the RSL and the  
centennial ESL.

### 3.1 : ESL change assessment

305 The evolution of ESLs is assessed using the three components (Eq. 3): (1) RSL, (2) tidal range and (3) combined storm surge  
and wave setup. The framework developed by Vousdoukas et al. (2018) is used to evaluate the contributions of each of these  
components. Projections of waves and storm surges were based on hydrodynamic simulations driven by atmospheric forcing  
from six Coupled Model Intercomparison Project Phase 5 (CMIP5) climate models.

#### 3.1.1 RSL: projections

310 Table 2 shows the median values of the RSL rise projections for the period 1995-2014 at selected points along the French  
coastline. For the SSP5-8.5 scenario, the median values of the RSL rise on the French coasts are fairly uniform: between  
0.16 m and 0.20 m in 2050, and between 0.76 m and 0.92 m in 2100.



**Table 2: Projections of RSL rise at selected points along the French coastline compared to the period 1995–2014 for the SSP5-8.5 scenario**

	Projection of RSL rise (m)	
	2050	2100
Calais	0.19	0.86
Le Havre	0.19	0.87
Saint-Malo	0.19	0.87
Brest	0.19	0.83
La Rochelle	0.16	0.76
Saint-Jean-de-Luz	0.17	0.79
Port Vendres	0.16	0.76
Sète	0.16	0.76
Marseille	0.17	0.78
Saint-Pierre (Saint-Pierre-et-Miquelon)	0.19	0.83
Pointe-à-Pitre (Guadeloupe)	0.20	0.90
Cayenne (French Guiana)	0.20	0.90
Pointe des Galets (La Réunion)	0.19	0.92
Papeete (French Polynesia)	0.20	0.91

320

### 3.1.2 Tidal range

Using the standard tidal classification system, the French coastline can be divided into three categories based on the tidal range data provided by SHOM (source: Data.shom.fr):

- microtidal coastline (tidal range < 2 m): the coasts of the Mediterranean with a tidal amplitude between 0.2 and 0.5 m and the coasts of the West Indies, La Réunion and French Polynesia where the tidal amplitudes are less than 1 m; in addition, the coasts of Saint-Pierre-et-Miquelon, where the amplitude reaches 1.7 m;
- mesotidal coastline (tidal range between 2 and 4 m): the coasts of French Guiana (the maximum amplitude reaches 2.9 m);
- macrotidal coastline (tidal range > 4 m): on the coasts of the Atlantic, English Channel and North Sea, with significant differences shown in Fig. 3.

325

330




Figure 3: Maximum tidal range (source: Data.shom.fr)

### 3.1.3 Waves and storm surges

Information on surges will be presented for France and its overseas territories.



In Metropolitan France, there are significant variations along the coastline, as shown by the estimates of the 100-year return period storm surge presented in Table 3.

340 **Table 3: Estimate of the 100-year return period storm surge, calculated as the average value of statistical adjustments of a Pareto distribution (GPD) and an exponential law (Cerema, 2018) applied to storm peaks. The values of Calais, Le Havre and La Rochelle have been corrected to take into account historical events and the comparative study of frequency analysis approaches for extreme storm surges (Hamdi et al., 2014).**

Tide gauge	100-year return period surge (m)
Calais	1.6
Le Havre	1.6
Saint-Malo	1.1
Brest	1.0
La Rochelle	1.7
Saint-Jean-de-Luz	0.6
Port-Vendres	0.9
Sète	1.1
Marseille	1.3

Although the relative differences are significant, they remain small in absolute terms, since these coasts are only exposed to storms, and not to cyclones. As mentioned in the presentation of the method, the large width of the continental shelf in the North Channel and the positioning at the center of the gulf or bay (e.g., Le Havre, Saint Malo, La Rochelle, Sète and 345 Marseille) or in a strait (e.g., Calais) explain the higher values observed at these locations.

Overseas, the Cerema (2019, 2020, 2021) provides information on the storm surge measured, observed and modelled in La Réunion, French Guiana, Martinique, Guadeloupe, Saint-Martin and Saint-Barthélemy and Saint-Pierre-et-Miquelon. Surges 350 measured by tide gauges commonly reach values of 0.5 to 1 m. Nevertheless, these measurements are often taken in sheltered areas (e.g., ports). Additional increases in the water level along the open coast may be more or less significant depending on the geomorphological configuration. For example, during the passage of Cyclone Irma on September 6, 2017, an instantaneous surge of 2.0 m was measured at the Saint-Martin tide gauge<sup>2</sup>. The surge, modeled by Météo-France, was more than 3 m on the northern coasts of Saint-Martin (Marigot Bay, Grand Case) and Gustavia (Saint-Barthélemy), but 355 hardly more than 1.2 m on the island's almost straight coastline (De la Torre Y., 2017). Similarly, in September 1989, during the passage of Cyclone Hugo on the Guadeloupe archipelago, there was evidence that the sea-level had increased by 2 to 3 m along the coast (Pagney, 1991).

Finally, on the islands of La Reunion and French Polynesia, in addition to cyclones, southern swells represent one of the main hazards. Continuous water height and wave measurements from the French national observation services ReefTEMPS

<sup>2</sup> [https://data.shom.fr/donnees/refmar/SAINT\\_MARTIN](https://data.shom.fr/donnees/refmar/SAINT_MARTIN)



360 and DYNALIT show that on coastlines directly exposed to swell (e.g. the Hermitage in La Réunion), 10-year return period events generate, at the breaking point, waves with  $H_s \sim 7$  to 8 m, which induce a 1.2 m setup. Infragravity waves with  $H_s \sim 1$  m can be superimposed on this set-up. These values are lower in ports (e.g. Saint Gilles).

This information confirms the need, as a complement to the measurement and modelling of surges, to study the geomorphological configuration of the coastline in order to understand the very significant differences that may appear.

### 365 3.1.4 ESL projection (intensity and frequency)

The framework developed by Vousdoukas et al. (2018) provides estimates of the ESL allowance along overseas coasts of France (Table 4). The results obtained in 2100 for the SSP5-8.5 scenario shows that the ESL increase in absolute value is relatively homogeneous on the various maritime facades: 0.89 to 1.00 m for the English Channel, 0.74 to 0.77 m for the Bay of Biscay and 0.75 to 0.78 m for the western Mediterranean. However, the percentage variations are very different, ranging  
370 between: 16 and 22% for the English Channel, 20 and 24% for the Bay of Biscay and 52 and 63% for the western Mediterranean, in 2100. The increase will therefore be much more noticeable in the Mediterranean.

**Table 4: Summary of the projected absolute and relative changes of the 100-year event ESL ( $\Delta$ ESL and % $\Delta$ ESL) under SSP5-8.5, during the years 2050 and 2100.**

	SSP5-8.5 - 2050		SSP5-8.5 - 2100	
	$\Delta$ ESL (m)	% $\Delta$ ESL	$\Delta$ ESL (m)	% $\Delta$ ESL
Calais	0.21	4.3	0.94	19.2
Le Havre	0.23	4.4	1.00	19.6
Saint-Malo	0.21	3.8	0.89	16.0
Brest	0.22	5.0	0.96	21.5
La Rochelle	0.16	4.3	0.77	19.9
Saint-Jean-de-Luz	0.16	5.3	0.74	24.1
Port Vendres	0.16	13.3	0.75	60.7
Sète	0.18	12.5	0.76	52.5
Marseille	0.18	14.6	0.78	63.2

375

The analysis of ESL height changes should be supplemented by a frequency analysis based on Table 5. The forecasts show that sea levels with a 100-year return period could occur on an annual basis by the end of this century in mainland France (except in the center of the Atlantic facade, around La Rochelle). Some regions are projected to experience an even higher increase in the frequency of occurrence of ESL, most notably along the Mediterranean, where the present day 100-year ESL  
380 is projected to occur about ten times a year. The higher increase in the Mediterranean is closely related to the low variability of sea-levels on these microtidal coastlines.



**Table 5: Return period (in years) of the present day 100-year ESL under SSP5-8.5 in the years 2050 and 2100.**

	Projected return period of the current 100-year ESL in:	
	2050	2100
Calais	22.81	0.75
Le Havre	26.56	0.87
Saint-Malo	27.69	0.81
Brest	20.61	0.73
La Rochelle	36.16	2.59
Saint-Jean-de-Luz	33.87	0.63
Port Vendres	27.89	0.10
Sète	30.02	0.56
Marseille	26.88	0.10

385

The framework developed by Vousdoukas et al. (2018) provides estimates of the ESL allowance along overseas coasts (cf. Table 6). The estimated ESL increase (absolute value) is relatively homogeneous for the overseas coasts, ranging between 0.16 and 0.24 m in 2050 and 0.85 and 0.95 m in 2100. The percent change in ESL is expected to be in the range of 8% to 18% by 2050, which is likely to be noticeable in terms of hazard intensity, particularly in Guadeloupe and French Polynesia, where the relative increases are estimated to be the largest. By 2100, the increases are projected to be around 50 to 75% (with the exception of Saint-Pierre-et-Miquelon).

390

**Table 6: Summary of the projected absolute and relative changes of the 100-year event ESL ( $\Delta$ ESL and  $\%$  $\Delta$ ESL) for the SSP5-8.5 scenario, during the years 2050 and 2100.**

	SSP5-8.5 - 2050		SSP5-8.5 - 2100	
	$\Delta$ ESL (m)	$\%$ $\Delta$ ESL	$\Delta$ ESL (m)	$\%$ $\Delta$ ESL
Saint-Pierre (Saint-Pierre-et-Miquelon)	0.24	8.4	0.92	31.8
Pointe-à-Pitre (Guadeloupe)	0.22	14.1	0.91	58.1
Cayenne (French Guiana)	0.18	10.6	0.85	49.2
Pointe des Galets (La Réunion)	0.16	9.3	0.88	52.5
Papeete (French Polynesia)	0.23	17.8	0.95	74.7

395

For the overseas coasts, the present day 100-year ESL is projected to occur more frequently, approximately ten times a year, except for Saint-Pierre-et-Miquelon where this frequency would be about three times a year (Table 7). The increase in frequency is likely to be noticeable in 2050, especially in French Guiana, French Polynesia and Saint-Pierre-et-Miquelon, where the return periods of the present day 100-year ESL are estimated to be only 7, 16 and 18 years, respectively.





400

**Table 7: Return period (in years) of the present day 100-year ESL for the SSP5-8.5 scenario in the years 2050 and 2100.**

	2050	2100
Saint-Pierre (Saint-Pierre-et-Miquelon)	17.69	0.31
Pointe-à-Pitre (Guadeloupe)	35.14	0.10
Cayenne (Guyane)	7.54	0.10
Pointe des Galets (La Réunion)	57.53	0.10
Papeete (French Polynesia)	15.86	0.10

### 3.2 Evaluation of additional factors

Two regional factors can be considered essential for all coastlines: the type of meteocean event and the geomorphological configuration. In addition, other environmental factors affecting the hydraulic conditions should also be considered at the tropics and poles.

#### 3.2.1 Meteocean event types

The types of marine-weather events affecting the selected coasts are storms for metropolitan France and Saint-Pierre-et-Miquelon; cyclones for the West Indies; cyclones, long swells and tsunamis (moderate exposure) for La Réunion and French Polynesia. In Guiana, no phenomenon generating energetic sea states are identified.

#### 3.2.2 Geomorphological configuration of the coast

The geomorphological configuration of each coast is evaluated using three criteria:

- the length of the coast,
- the width of the continental shelf, and
- the convex or concave shape of the coastline.

The corresponding data, obtained from bathymetric maps provided by the SHOM (Table 8) allows estimating qualitatively the surge potential associated with each geomorphological configuration.

420



425 **Table 8: Assessment of the surge potential associated with the geomorphological configuration of the study sites (source: Data.shom.fr)**

	Order of magnitude of coastline length (km)	Width of the continental shelf	Shape of the coastline	Surge potential associated with the geomorphological configuration
English Channel – North Sea	1000	Very large (>200 km)	Alternation of concave and convex shapes	Very high
Bay of Biscay	1000	Large (100 to 200 km)	Concave	High
West Mediterranean	1000	Normal to the West (around 50 km), nonexistent to the East	Concave to the West and convex to the East	High to the West Low to the East
Saint-Pierre-et-Miquelon	10	Very large (>200 km)	Alternation of concave and convex shapes	High
Guadeloupe	5 to 20	Narrow (<20 km)	Alternation of concave and convex shapes	Low
French Guyana	1000	Large (about 100 km)	Linear or slightly convex, except for estuaries	Moderate
La Réunion	20	Nonexistent	Mainly linear or convex (except 2 baies in the North West)	Very low
French Polynesia	1 to 20	Nonexistent	Mainly linear or convex (with many small-scale exceptions)	Very low

430 In metropolitan France, the geomorphological configurations (i.e. long coastline, wide continental shelf, linear and/or concave shape) generally favor the development of substantial surges. The main exception is the eastern Mediterranean facade, which has convex-shaped coasts without a continental shelf (extension in the sea of the Alpine massif). Another exception is the southern part of the Bay of Biscay, where the concave shape and the length of the coastline favour the surge, while the small width of its continental shelf is a limiting factor.



For the overseas territories, the analysis reveals very contrasting situations:

- 435
- Guiana has a broad continental shelf, but its coastline is linear or convex, which is less conducive to surges;
  - Saint-Pierre-et-Miquelon is in a situation similar to that of metropolitan France, with a wide continental shelf and the alternation of convex and concave forms of the coast. Although the length of the coastline is smaller, the close proximity of the large island of Newfoundland, which has a 200 km coastline, can contribute to large surges;
  - The other islands are far from continents and have continental shelves that are narrow (e.g., Guadeloupe) or non-
- 440
- existent (e.g., La Réunion and French Polynesia), which is, with the short coastline length, a factor limiting the surge magnitude. For La Réunion and French Polynesia, the surges are even more limited by the linear or convex shape of the coastline.

In summary, considering only the geomorphological configuration, the coasts of metropolitan France (except the east of the Mediterranean facade and the Southern part of the Bay of Biscay), French Guiana and Saint-Pierre-et-Miquelon have all the

445

characteristics favoring large surges. Smaller surges are expected on all criteria in La Réunion and French Polynesia, and based on all criteria except the shape of the coast in the West Indies.

### 3.2.3 Other environmental factors influencing the hydraulic conditions

La Réunion and French Polynesia are exposed to long southern swells and will therefore experience direct changes in the swell climate related to the decrease in seasonal sea ice extent.

450

The islands of the West Indies, La Réunion and French Polynesia are bordered by coral reefs whose degradation by ocean changes resulting from climate change (e.g. increases in water temperature and ocean acidification) may be a major factor in aggravating coastal hazards.

### 3.3 Integrated hazard evolution analysis

The evolution of coastal hazards can be assessed by considering both the quantitative sea level change estimates and the

455

qualitative geomorphological configuration, meteocean event and environmental factor analysis for France (Table 9) and its overseas territories (Table 10).

460



### 3.3.1 Hazard evolution in the European territory of France

465 **Table 9: Summary of the main factors influencing the hazard evolution on the mainland France’s coasts. The data relative to ESLs refer to a present day 100-year ESL. The reference period for the RSL projection is 1995-2014. The reference period for the ESL projection is 1980-2014. The least important factors of the evolution of the hazard are in italics, the important factors in bold and the most important factors in capital and bold. In the first column, these same fonts describe the estimated global evolution of the hazard for each coast.**

	Surge potential associated with the geomorphological configuration	Reference meteocean event	Tidal range	100-year surge (m)	$\Delta$ RSL (m) under SSP5-8.5	$\Delta$ ESL (m) under SSP5-8.5	% $\Delta$ ESL under SSP5-8.5	Return period of the present day 100-year ESL under SSP5-8.5 (year)
<b>English Channel – North Sea</b>	<b>VERY HIGH</b>	Storm	<i>Macrotidal (7 to 14 m)</i>	1.0 to 1.6	In 2050 : 0.19 In 2100 : 0.83 to 0.87	In 2050 : 0.21 to 0.23 In 2100 : 0.89 to 1.00	In 2050 : 3.8 to 5 In 2100 : 16 to 21.5	<b>In 2050 : 20 to 28</b> <b>In 2100 : 0.73 to 0.87</b>
<b>Bay of Biscay</b>	<b>High to the North</b> Moderate to the South	Storm	<i>Macrotidal (5 m to 7 m)</i>	0.6 to 1.7	In 2050 : 0.16 to 0.17 In 2100 : 0.76 to 0.79	In 2050 : 0,16 In 2100 : 0.74 to 0.77	In 2050 : 4.3 to 5.3 In 2100 : 19.9 to 24.1	<b>In 2050 : 33 to 36</b> <b>In 2100 : 0.63 to 2.6</b>
<b>WEST MEDITERRANEAN</b>	<b>High to the West</b>	Storm	<b>MICROTIDAL (&lt; 2 m)</b>	0.9 to 1.3	In 2050 : 0.16 to 0.17 In 2100 : 0.76 to 0.78	In 2050 : 0,16 to 0.18 In 2100 : 0.75 to 0.78	<b>IN 2050 : 12.5 TO 14.6</b> <b>IN 2100 : 52.5 TO 63.2</b>	<b>IN 2050 : 26 TO 30</b> <b>IN 2100 : 0.10 TO 0.56</b>
<b>East Mediterranean</b>	<i>Low to the East</i>							

470 By 2100, for the SSP5-8.5 scenario, the increases in RSL and ESL on the three maritime facades are rather similar in absolute terms, with median values between 0.74 and 1.0 m. The effect of these increases is nevertheless significantly different for each maritime facade, depending on the type of meteocean event, tidal range, and surge magnitude:

- 475 • In the English Channel and the North Sea, the macrotidal range and storm occurrence produce high variability of sea levels. The increase of 0.9 to 1.0 m of the 100-year ESL represents only 16 to 22%. The allowance is therefore relatively low. On the other hand, the increase in the RSL is sufficient for a present day 100-year ESL to have a return period of less than one year (amplification factor greater than 100);
- 480 • In the Bay of Biscay, the situation is quite similar, but with a smaller tidal range and surge magnitude (especially on the coast of the Atlantic Pyrenees, where the maximum tides are in the order of 5 m and the 100-year return period surge in the order of 0.6 m). Under these conditions, the 0.75 m increase in the centennial ESL corresponds to a relative increase of 20 to 24%, and above all, a current centennial ESL is likely to occur on average once or twice a year (amplification factor between 50 and 100);
- In the Mediterranean, the situation is clearly aggravated by the very low variability of the sea level (microtidal regime and small surges). As a result, the 100-year ESL increase of 0.77 m corresponds to a large relative increase



485 (greater than 50%) and the present day 100-year ESL is likely to occur between 2 and 10 times per year  
(amplification factor between 500 and 1000).

By 2050, the 100-year ESL will rise by 5% in the North Channel and the Bay of Biscay. In the Mediterranean, this increase  
will already be of the order of 20%. However, the evolution of the situation will be especially noticeable in the increased  
490 frequency of ESLs (by a factor of 3 to 4).

In addition to this quantitative analysis, the qualitative analysis shows that the geomorphological configurations of the three  
metropolitan maritime facades (except on the eastern half of the Mediterranean coast and the southern part of the Bay of  
Biscay) favor the formation of storm surges, by the width of the continental shelves, and the length and shape of the  
495 coastlines. On these coasts, a change in the characteristics of storms could therefore lead to a sharp increase in surges.  
Analysis of the surge observed in 2010 during the passage of storm Xynthia in the Bay of Biscay demonstrates the  
vulnerability of coastal areas to unusual meteocean phenomena (Bertin et al., 2012). The storm's low pressure center track  
propagated to the northeast, contrasting with other storm tracks usually propagating to the east or southeast. Despite its low  
intensity, this storm caused the large marine submersions in France (Cerema, 2016).

500 Finally, it should be recalled that the amplification factors related to ESLs (which are expected to reach values between 50  
and 1000 in 2100 on the coasts of metropolitan France) must be distinguished from the evolution of extreme events such as  
storms, cyclones and tsunamis, whose frequency and intensity is not projected to change as significantly. However, even if  
the frequency and intensity of meteocean events remain constant, an increase in the ESL frequency and intensity would  
505 increase the effects of these events on the coasts.

510

515



### 3.3.2 Hazard evolution in the overseas territories of France

**Table 10: Same as table 9 for the coastlines of French overseas territories.**

	Surge potential associated with the geomorphological configuration	Other regional factor of change	Reference meteocean event	Tidal range	100-year surge (m)	$\Delta$ RSL (m) under SSP5-8.5	$\Delta$ ESL (m) under SSP5-8.5	% $\Delta$ ESL under SSP5-8.5	Return period of the present day 100-year ESL under SSP5-8.5 (year)
<b>Saint-Pierre (Saint-Pierre-et-Miquelon)</b>	High	None	Storm	<b>MICROTIDAL (&lt; 2 M)</b>	1 to 2	In 2050 : 0,19 In 2100 : 0,83	In 2050 : 0,24 In 2100 : 0,92	In 2050 : 8,4 In 2100 : 31,8	<b>IN 2050 : 18</b> <b>IN 2100 : 0,3</b>
<b>POINTE-A-PITRE (GUADELOUPE)</b>	Low	Coral reef degradation	<b>STORM AND CYCLONE</b>	<b>MICROTIDAL (&lt; 2 M)</b>	1 to 3	In 2050 : 0,20 In 2100 : 0,90	In 2050 : 0,22 In 2100 : 0,91	<b>In 2050 : 14,1</b> <b>In 2100 : 58,1</b>	<b>IN 2050 : 35</b> <b>IN 2100 : 0,1</b>
Cayenne (French Guiana)	Moderate	None	Trade winds (without storm)	<b>Mesotidal (2 to 4 m)</b>	<b>0,4</b>	In 2050 : 0,20 In 2100 : 0,90	In 2050 : 0,18 In 2100 : 0,85	<b>In 2050 : 10,6</b> <b>In 2100 : 49,2</b>	<b>IN 2050 : 7,5</b> <b>IN 2100 : 0,1</b>
<b>POINTE DES GALETS (LA REUNION)</b>	Very low	Coral reef degradation Decrease in seasonal sea ice extent	<b>STORM CYCLONE +LONG SWELL +MODERATE EXPOSURE TO TSUNAMIS</b>	<b>MICROTIDAL (&lt; 2 M)</b>	1 to 3	In 2050 : 0,19 In 2100 : 0,92	In 2050 : 0,16 In 2100 : 0,88	<b>In 2050 : 9,3</b> <b>In 2100 : 52,5</b>	<b>IN 2050 : 58</b> <b>IN 2100 : 0,1</b>
<b>PAPEETE (FRENCH POLYNESIA)</b>	Very low	Coral reef degradation Decrease in seasonal sea ice extent	<b>STORM CYCLONE +LONG SWELL +MODERATE EXPOSURE TO TSUNAMIS</b>	<b>MICROTIDAL (&lt; 2 M)</b>	1 to 3	In 2050 : 0,20 In 2100 : 0,91	In 2050 : 0,23 In 2100 : 0,95	<b>IN 2050 : 17,8</b> <b>IN 2100 : 74,7</b>	<b>IN 2050 : 16</b> <b>IN 2100 : 0,1</b>

520 In 2100, the quantitative analysis shows that for the SSP5-8.5 scenario:

- The coasts of the West Indies, La Réunion, and French Polynesia are microtidal (<2 m), but the passage of the cyclones can generate large surges (up to 3 m). The allowance (50-75% increase) and amplification (increase in the frequencies of the current centennial ESL by about 1000) of these coasts are similar to those of the Mediterranean coast of France. However, the change in the intensity and frequency of the ESLs must be considered more carefully in these territories because of the greater damage that cyclones can cause compared to storms that reach the European territory of France;

525



- The coast of Saint-Pierre-et-Miquelon is also microtidal (<2 m). While strong winds that regularly blow on these islands can produce large surges, the variability in sea levels remains low because of the microtidal regime. The situation is therefore similar to that of the Mediterranean, with a smaller estimated increase in ESLs, as shown by the allowance (about 32%) and amplification (about 300);
- The coast of French Guiana is mesotidal (maximum range of 2.9 m), and the maximum recorded surge is only 0.4 m. The increase in the amplitude and frequency of ESLs should be of the same order of magnitude as that of the Mediterranean coast of France, the West Indies and La Réunion. However, this coast is located close to the equator and is not exposed to meteo-oceanic events that generate strong swells. The increase in coastal hazards therefore is expected to be less intense than in the other overseas territories.

By 2050, the evolution of the hazard is likely to be perceptible primarily because of the increase in the frequency of ESLs by a factor of 2 for La Réunion, 3 for the West Indies, 5 for Saint-Pierre-et-Miquelon, 6 for French Polynesia, and 13 for French Guiana. The most significant changes are for the territories with low variability in sea levels that are subject to storms (Saint-Pierre-et-Miquelon whose centennial ESL will increase by 8%) and to cyclones (La Réunion, the West Indies and French Polynesia, with ESL increases of 9, 14 and 18%, respectively). In this comparison, French Polynesia shows both the greatest increase in frequency and intensity of the 100-year ESL. French Guiana is also expected to experience negative changes in the ESL by 2050, but since it is not exposed to storm waves, this territory is not expected to be exposed to catastrophic meteocean events.

In addition, the qualitative analysis shows that the geomorphological configurations of the overseas territories (except for Saint-Pierre-et-Miquelon and French Guiana) do not favor the formation of storm surges. The situation is aggravated in the West Indies by the degradation of coral reefs and on La Réunion and French Polynesia (islands exposed to cyclones and southern swells) by the degradation of coral reefs and the decrease in seasonal sea ice extent.

Finally, as for the coasts of metropolitan France, it is necessary to distinguish between changes in the frequency and intensity of ESLs and extreme meteocean events. Even if extreme meteocean events are not expected to increase, an increase in ESLs will lead to an increase in coastal hazards.

#### 4 Discussion

The application of the proposed method leads to increased understanding of coastal hazards, taking into account the hydrodynamic forcing and the geomorphological context. These results, considered in conjunction with anticipated coastal responses to the effects of climate change, can shed light on the evolution of the hazard on the terrestrial domain.



#### 4.1 Influence of local factors affecting the coastline

Understanding the evolution of coastal hazards requires evaluating slow changes that occur over decadal timescales (e.g. RSL rise, ocean acidification and warming, morphological evolution), in conjunction with extreme events that occur on daily time scales, such as tsunamis, cyclones, storms (Igigabel et al., 2021). However, while some factors can be taken into account at the regional level (and were therefore considered by the proposed method), others can only be considered at finer spatial scales because of the complex interactions between sea-level rise and the morphological evolution of coastal areas. In order to assess local hazard evolution, one must consider local effects, such as subsidence, especially in deltas (Syvitski, 2008). Second, the coastal zone is a buffer zone where a multitude of processes and feedbacks may also be important:

- in estuaries, changes in coastal morphology (bathymetry, shoreline topography, and anthropogenic development) can influence (positively or negatively) extreme events, including changes in the spread of tidal waves and storm surges (Talke et al., 2020);
- along sandy coasts, coastal morphology is also likely to change as a function of the local morphological and sedimentary characteristics, climatic changes (temperature, precipitation and wind), wave conditions, and frequency of events, which has an important impact on the ability of a system to recover between energetic events (Masselink et al., 2016). In addition, wave direction is also an important parameter related to longshore sediment transport, which can change the shape of a coastline (Ruggiero et al., 2010; Casas-Prat and Sierra, 2010);
- in the polar regions, accelerating permafrost thaw is promoting rapid erosion of ice-rich sediments (Lantuit et al., 2011). Melting of ice and associated thaw subsidence may induce instability of various infrastructure components. Arctic RSL rise and sea surface warming have the potential to substantially contribute to this thawing (Lamoureaux et al., 2015).
- RSL rise can greatly increase coastal hazards, because an increase in water depth, assuming no change in bathymetry, may cause increases in nearshore wave conditions. The coastal impacts of ESLs are largely due to increased wave impacts, potentially driving morphological changes and erosion, as well as coastal protection failure and overwash/inundation (CIRIA et al., 2013, Vousdoukas et al., 2017).

Ideally, all of these phenomena should be considered in assessing coastal hazard evolution and in defining the adaptation measures for a particular location. While this may be too difficult to achieve for a regional study, it is necessary to take into account at the local level.

#### 4.2 Understanding hazard evolution on land

On many coasts, accelerated sea-level rise is likely to result in permanent submersion of unprotected lowlands. More frequent and intense episodic coastal flooding could also occur with future changes in the wave climate and storm surges. In





590 some locations, this may result in chronic coastline erosion (Ranasinghe, 2016). Thus, depending on the coastline, distinct phenomena may appear, which may be classified, using the proposed method (characterisation of the hazard on a regional scale in the maritime domain) and the insights from existing studies (characterization of phenomena affecting the coastal fringe locally).

595 By neglecting adaptive actions, hazard evolution may be categorized into one of the following three main scenarios (defined by the sea level variability and the types of meteocean events, knowing that other factors also influence the hazard evolution and must be considered subsequently):

- 600 • coastlines with low variability in sea levels that are not exposed to meteocean events (e.g. French Guiana): a RSL rise will likely generate a small increase in the risk of rapid submersion by the sea. Nevertheless, rapid flooding by river floods or by intense precipitation remains possible. In addition, the gradual rise of the RSL is likely to cause increased erosion (which, in general, results in coastline retreat) and/or to the progressive and permanent flooding of lowlands (i.e. below sea level);
- 605 • coastlines with high sea-level variability that are exposed to meteocean events (e.g., the Atlantic coastline of metropolitan France): RSL rise is likely to cause increased erosion and the degradation of protection systems (i.e. anthropogenic structures, such as levees, seawalls, groynes and breakwaters; and natural features such as beaches, dunes, salt marshes, mangroves and coral reefs) over the years, even in the absence of extreme events. During an extreme event (due to the conjunction of a storm, cyclone or tsunami with a high spring tide), the risk may be increased both by the regular weakening of the protection systems and by the increased impacts of waves related to the rise of the RSL;
- 610 • coastlines with low sea-level variability that are exposed to meteocean events (e.g. West Indies, La Réunion, French Polynesia): RSL rise is likely to cause, in the short term, the three phenomena described in the previous cases: increased risk of coastal erosion, increased risk of rapid flooding, and permanent flooding of lowlands.
- In addition to these three main scenarios, there are coastlines with high variability in sea levels and with relatively low exposure to meteocean events, which will likely show slower increases in coastal hazards.

## 5 Conclusion

615 Changes in sea levels (GMSL, RSL and ESL) are subject to high uncertainties, at the global scale (economic development, GHG emission scenarios, ice sheet response) and at regional or local scales (response of coastlines to climatic changes and changes in wave climate). Quantified estimates of sea level changes are necessary, but should also be put into perspective with factors describing the state and evolution of the coastal environment. The proposed method for evaluating coastal hazard evolution is based on this principle, emphasizing the importance of taking into account sea level changes, as well as



the geomorphological configuration, meteocean event types, and, where appropriate, additional marine environment changes  
620 affecting hydrodynamic conditions (in particular, in the tropics or at the poles).

The application of this method to selected French coastlines demonstrated the expected differences in future coastal hazard  
evolution.

625 The study of the variability of sea levels with a quantitative approach confirmed the results of previous studies, namely the  
largest increase of ESLs on coasts with a small tidal range and storm surges. This is the case for the French overseas  
territories and, in metropolitan France, for the Mediterranean facade. The increase in ESLs, in intensity and frequency,  
should be limited on the coasts with large tidal ranges (French facades of the Atlantic Ocean, the English Channel and the  
North Sea).

630

In addition to the results obtained for ESLs by quantitative approaches, the qualitative approach makes it possible to assess  
more comprehensively the evolution of coastal hazards.

635 First, the type of meteocean event is essential: coasts not exposed to extreme events (storms, cyclones, long swells or  
tsunamis), even if they are microtidal (e.g. Guiana), should experience a moderate hazard increase compared to exposed  
coasts. All other characteristics being equal (geomorphological configurations, environmental changes and tidal conditions),  
the hazard will increase more rapidly on the coasts exposed to cyclones (the West Indies, Réunion, and French Polynesia)  
than on the coasts exposed to storms (Mediterranean facade).

640 Secondly, certain geomorphological configurations (long maritime facades, concave in shape and bordered by a wide  
continental shelf) favor the generation of surges during extreme events. These criteria help to identify the coastlines most  
vulnerable to changes in the characteristics of extreme events (especially their trajectories). This result is important because  
quantitative approaches are based on climate change scenarios, which therefore represent only part of the possible  
evolutions. The application of quantitative and qualitative approaches for French territories makes it possible to distinguish  
645 several scenarios: some coastlines benefit from a high variability in sea levels, which is a positive feature to limit the effects  
of the rise in RSL, but, on the other hand, have geomorphological characteristics favoring surges (e.g. Atlantic, Channel and  
North Sea coasts). Other shorelines are expected to be disadvantaged by both low sea level variability (small tidal range and  
moderate surges) and a geomorphological configuration that favors surges (e.g., the western portion of the Mediterranean  
coastline and the concave shorelines of the West Indies). Finally, some coastlines have the disadvantage of small sea level  
650 variability, but their geomorphological configuration does not favor surges (e.g. the eastern part of the Mediterranean  
coastline). This analysis highlights the need to study not only the water level variability, but also the current and future  
exposure to meteocean events.



655 Third, the coral reef degradation in the tropics and the decrease in seasonal sea ice extent in the polar regions can also significantly change the nearshore hydrodynamics and impacts on the shoreline.

660 These results represent a substantial step forward, not only because they are achieved through a comprehensive and systemic approach that requires a step back, but also because the proposed method uses factors for which the uncertainty in the projections is the lowest: the geomorphological configuration of the coast will be invariable over the centuries, regional tidal conditions will be stable (Haigh et al., 2020), climate change impacts on coral reefs (Albright et al., 2018; Hoegh-Guldberg et al., 2018) and seasonal sea ice extent (Forbes, 2011) are fairly well known. However, local climate changes and, more particularly, changes in storm and cyclone trajectories are an area where uncertainties remain particularly high (Seneviratne et al., 2021).

665 Ultimately, the impacts of sea level changes on human communities are evaluated by considering additionally the local-scale effects on natural features and anthropogenic structures. These analyses aim to predict the types of hazard (shoreline erosion, rapid submersion and/or permanent flooding) that will increase the most in different coastal zones (shoreline erosion, rapid submersion and/or permanent flooding). Shorelines are expected to be marked by a quasi-generalized increase in the risks of erosion of natural features and weakening of anthropogenic structures. However, regional and local-scale differences are  
670 expected depending on the submersion type: coastlines with high sea-level variability will experience mainly rapid submersion effects, whereas coastlines with low sea-level variability will experience mainly permanent flooding of low-lying land, and, rapid submersion when exposed to extreme meteocean events.

## 6 Code availability

The Delft3D-FM code is currently being made available in <http://oss.deltares.nl>. The WW3 model description is available in:  
675 <https://polar.ncep.noaa.gov/waves/wavewatch/>. The code applied for the non-stationary extreme value statistics (Mentaschi et al., 2016) is available in: <https://github.com/menta78/tsEva>.

## 7 Data availability

The global ESL data that support the findings of this study are available in the LISCoAsT repository of the JRC data collection (<http://data.jrc.ec.europa.eu/collection/LISCOAST>) though this link: <http://data.jrc.ec.europa.eu/dataset/jrc-liscoast-10012>, with the identifiers: <https://doi.org/10.2905/jrc-liscoast-10012>; PID: <http://data.europa.eu/89h/jrc-liscoast-10012>  
680 10012



## 8 Author contribution

MI, MY and YD conceived the paper. MV developed the model code and performed the simulations. MI prepared the manuscript with contributions from all co-authors.

## 685 9 Competing interests

The authors declare that they have no conflict of interest.

## 10 Acknowledgments

The authors would like to thank Xavier Bertin, CNRS research director at the Littoral ENvironment et Sociétés (LIENSs) laboratory for his comments about the manuscript and for sharing of water height and wave data from the national  
690 observation services ReefTEMPS and DYNALIT.

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors. All authors declare that they have no conflicts of interest.

## 11 References

695 Albright, R., Takeshita, Y., Koweek, D., Ninokawa A., Wolfe K., Rivlin T., et al.: Carbon dioxide addition to coral reef waters suppresses net community calcification, *Nature* 555, 516–519, <https://doi.org/10.1038/nature25968>, 2018.

Almar, R., Ranasinghe, R., Bergsma, E.W.J., Diaz, H., Melet, A., Papa, F., et al.: A global analysis of extreme coastal water levels with implications for potential coastal overtopping. *Nat Commun*, 12, 3775. <https://doi.org/10.1038/s41467-021-24008-9>, 2021.  
700

Bamber, L. J., Oppenheimer, M., Kopp, R. E. and Cooke, R. M.: Ice sheet contributions to future sea-level rise from structured expert judgment. *Proceedings of the National Academy of Sciences*, p. 5., 2019.

705 Bertin, X., Bruneau, N., Breilh, J.-Fr., Fortunato, A. and Karpytchev, M.: Importance of wave age and resonance in storm surges: The case Xynthia, Bay of Biscay, *Ocean Modelling*, Volume 42, Pages 16-30, ISSN 1463-5003, <https://doi.org/10.1016/j.ocemod.2011.11.001>, 2012.



710 Bowen, A. J., Inman, D. L. and Simmons, V. P.: Wave ‘set-down’ and set-up, *J. Geophys. Res.*, **73**, 2569–2577,  
doi:10.1029/JB073i008p02569, 1968.

Buchanan, M.K., Kopp R.E., Oppenheimer M. and Tebaldi C.: Allowances for evolving coastal flood risk under 41 uncertain  
local sea-level rise, *Climatic Change*, **137**(3–4), doi:10.1007/s10584-016-1664-7, 2016.

715 Calafat, F.M., Wahl, T., Tadesse, M.G. and Sparrow, S.N.: Trends in Europe storm surge extremes match the rate of sea-  
level rise, *Nature* **603**, 841–845. <https://doi.org/10.1038/s41586-022-04426-5>, 2022).

Casas-Prat, M. and Sierra, J. P.: Trend analysis of the wave storminess: the wave direction, *Adv. Geosci.*, **26**, 89–92.  
720 <https://doi.org/10.5194/adgeo-26-89-2010>, 2010.

Cazenave, A. and Le Cozannet, G.: Sea level rise and its coastal impacts, *Earth’s Future*, **2**, 15–34.  
doi:10.1002/2013EF000188, 2013.

725 Cerema: Étude des systèmes de protection contre les submersions marines, Collection : Références, ISBN : 978-2-37180-  
118-9, 442 pages, 2016.

Cerema: Les niveaux marins extrêmes. Ports de métropole, Collection: Données, ISBN : 978-2-37180-308-4, 2018.

730 Cerema: Dynamiques et évolution du littoral, Fascicule 11: Synthèse des connaissances de Saint-Pierre-et-Miquelon,  
Collection: Connaissances, ISBN: 978-2-37180-341-1, 2020.

Cerema: Dynamiques et évolution du littoral, Fascicule 12: Synthèse des connaissances de Saint-Barthélemy et Saint-Martin,  
Collection: Connaissances, ISBN: 978-2-37180-342-8, 2020.

735 Cerema: Dynamiques et évolution du littoral, Fascicule 13: Synthèse des connaissances de l’archipel de Guadeloupe,  
Collection: Connaissances, ISBN: 978-2-37180-316-9, 2020.

740 Cerema: Dynamiques et évolution du littoral, Fascicule 14: Synthèse des connaissances de la Martinique, Collection:  
Connaissances, ISBN: 978-2-37180-343-5, 2020.

Cerema: Dynamiques et évolution du littoral, Fascicule 15: Synthèse des connaissances de la Guyane, Collection:  
Connaissances, ISBN: 978-2-37180-344-2, 2021.



745 Cerema: Dynamiques et évolution du littoral, Fascicule 17: Synthèse des connaissances de La Réunion, Collection: Connaissances, ISBN: 978-2-37180-345-9, 2020.

Church J. A., Clark P. U., Cazenave A., Gregory J. M., Jevrejeva S., Levermann A., Merrifield M. A., Milne G. A., Nerem R. S., Nunn P. D., Payne A. J., Pfeffer W. T., Stammer D., Unnikrishnan A. S.: Sea level change, In: Climate change 2013: the physical science basis, Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change, Cambridge University Press, Cambridge, United Kingdom, pp 1137–1216, 2013.

CIRIA, MEDDE, USACE: The International Levee Handbook, C731, CIRIA, UK, London, ISBN : 978-0-86017-734-0, 1332 pages, 2013.

755 Dayan H., Le Cozannet G., Speich S. and Thiéblemont R.: High-End Scenarios of Sea-Level Rise for Coastal Risk-Averse Stakeholders. *Front. Mar. Sci.* 8:569992, doi: 10.3389/fmars.2021.569992, 2021.

De la Torre, Y.: Impacts du cyclone Irma sur le littoral des « îles du Nord » à Saint-Martin et Saint-Barthélemy, Rapport BRGM/RP-67291-FR, 29 p., 20 ill., 2 ann., 2017.

Dietrich, J. C., Bunya, S., Westerink, J. J., Ebersole, B. A., Smith, J. M., Atkinson, J. H., et al.: A high resolution coupled riverine flow, tide, wind, wind wave and storm surge model for southern Louisiana and Mississippi: part II — synoptic description and analyses of Hurricanes Katrina and Rita. *Monthly Weather Review* 138, 378–404, 765 <https://doi.org/10.1175/2009MWR2907.1>, 2010.

Dodet, G., Bertin, X., Bouchette, F., Gravelle, M., Testut, L., and Wöppelmann, G.: Characterization of Sea-level Variations Along the Metropolitan Coasts of France: Waves, Tides, Storm Surges and Long-term Changes, *Journal of Coastal Research*, 88(sp1), 10-24. <https://doi.org/10.2112/SI88-003.1>, 2019.

770 Durand, F.: Hydrodynamique sédimentaire sur le plateau insulaire de la Martinique, Thèse doctorale, Université Bordeaux 1. 214 p., 1996.

Erkens, G., Bucx, T., Dam, R., De Lange, G. and Lambert, J.: Sinking Coastal Cities. Proceedings of the International Association of Hydrological Sciences, 372, 189-198, <https://doi.org/10.5194/piahs-372-189-2015>, 2015.



FLOODsite: Language of risk. Project definitions, Report T34-02-01, Second edition, 55 pp. e-publication on [www.floodsite.net](http://www.floodsite.net). Available online: [www.floodsite.net](http://www.floodsite.net), 2009.

780 Forbes, D.L.: State of the Arctic coast 2010: scientific review and outlook. Land-Ocean Interactions in the Coastal Zone, Institute of Coastal Research, <http://arcticcoasts.org/files/sac/state%20of%20the%20arctic%20rept-exec%20summary%20res.pdf>, 2011.

785 Fox-Kemper B, Hewitt HT, Xiao C et al.: Chapter 9: ocean, cryosphere and sea level change. In: Masson-Delmotte V, Zhai P, Pirani A et al (eds) Climate Change 2021: the physical science basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Intergovernmental Panel on Climate Change (IPCC), 2021.

790 Frederikse, T., Landerer, F., Caron, L. et al.: The causes of sea-level rise since 1900, *Nature* 584, 393–397, <https://doi.org/10.1038/s41586-020-2591-3>, 2020.

Graumann, A., Houston, T., Lawrimore, J. H., Levinson, D. H., Lott, N., McCown, S. et al.: Hurricane Katrina, A Climatological Perspective. October 2005, Updated August 2006. NOAA's National Climatic Data Center. <https://www1.ncdc.noaa.gov/pub/data/techrpts/tr200501/tech-report-200501z.pdf>, 2006.

795 Gregory, J.M., Griffies, S.M., Hughes, C.W. et al.: Concepts and Terminology for Sea Level: Mean, Variability and Change, Both Local and Global. *Surv Geophys* 40, 1251–1289, <https://doi.org/10.1007/s10712-019-09525-z>, 2019.

800 Haigh, I.D., Pickering, M.D., Green, J.A.M., Arbic, B.K., Arns, A., Dangendorf, S., et al.: The Tides They Are a-Changin': A comprehensive review of past and future non-astronomical changes in tides, their driving mechanisms and future implications, *Reviews of Geophysics*, 57, e2018RG000636, <https://doi.org/10.1029/2018RG000636>, 2020.

805 Hallegatte, S., Green, C., Nicholls, R. J., and Corfee-Morlot, J.: Future flood losses in major coastal cities, *Nature Climate Change*, 3, 802–806, <https://doi.org/10.1038/nclimate1979>, 2013.

Hoegh-Guldberg, O., Mumby, P. J., Hooten, A. J., Steneck, R. S., Greenfield P., Gomez E., et al.: Coral Reefs Under Rapid Climate Change and Ocean Acidification. *Science*, 80(318), 2007.

810 Hunter, J.: A simple technique for estimating an allowance for uncertain sea level rise, *Clim. Change*, 113(2), 239–252, doi:10.1007/s10584-011-0332-1, 2012.



- Idier, D., Paris, F., Le Cozannet, G., Boulahya, F., Dumas, F.: Sea-level rise impacts on the tides of the European Shelf, *Continental Shelf Research*, Volume 137, 56-71, <https://doi.org/10.1016/j.csr.2017.01.007>, 2017.
- 815 Igigabel, M., Diab, Y. and Yates M.: Exploring Methodological Approaches for Strengthening the Resilience of Coastal Flood Protection System. *Front. Earth Sci.* 9:756936. doi: 10.3389/feart.2021.756936, 2021.
- IPCC: Climate Change 2001: Third Assessment Report (TAR), 2001.
- 820 IPCC: Climate Change 2007: Fourth Assessment Report (AR4), 2007.
- IPCC: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp., 2014.
- 825 IPCC: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate [H.- O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, M. Nicolai, A. Okem, J. Petzold, B. Rama, N. Weyer (eds.)], Available online: <https://www.ipcc.ch/srocc/> (accessed on 20 December 2021), 2019.
- 830 IPCC: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)], Cambridge University Press, 2021.
- 835 Lamoureux, S., Forbes, D., Bell, T. and Manson, G.: The impact of climate change on infrastructure in the western and central Canadian Arctic. In: *From Science to Policy in the Western and Central Canadian Arctic: an Integrated Regional Impact Study (IRIS) of Climate Change and Modernization* [Stern, G.A. and A. Gaden (eds.)], ArcticNet, Quebec, pp. 300–341, 2015.
- 840 Lantuit, H., Atkinson, D., Overduin P. P., Grigoriev, M., Rachold, V., Grosse, G. and Hubberten, H.: Coastal erosion dynamics on the permafrost-dominated Bykovsky Peninsula, north Siberia, 1951–2006, *Polar Res.*, 30(1), 7341. <https://doi.org/10.3402/polar.v30i0.7341>, 2011.





845 Lavaud, L., Bertin, X., Martins, K., Pezerat, M., Coulombier, T. and Dausse, D.: Wave dissipation and mean circulation on a  
shore platform under storm wave conditions. *Journal of Geophysical Research: Earth Surface*, 127, e2021JF006466.  
<https://doi.org/10.1029/2021JF006466>, 2022.

850 Lowe J. A., Woodworth P. L., Knutson T., McDonald R. E., McInnes K. L., Woth K., von Storch H., Wolf J., Swail V.,  
Bernier N. B., Gulev S., Horsburgh K. J., Unnikrishnan A. S., Hunter J. R. and Weisse R.: Past and future changes in  
extreme sea levels and waves. In: Woodworth PL, Aarup T, Wilson WS, Church JA (eds) *Understanding sea-level rise and  
variability*. Wiley, New York, 2010.

855 Martins K., Bertin X., Mengual B., Pezerat M., Lavaud L., Guérin T. and Zhang Y.J.: Wave-induced mean currents and  
setup over barred and steep sandy beaches. *Ocean Modelling*, Volume 179, November 2022, 102110, 2022.

860 Masselink, G., Castelle, B., Scott T., Dodet, G., Suanez, S., Jackson, D. and Floc'h, F.: Extreme wave activity during  
2013/2014 winter and morphological impacts along the Atlantic coast of Europe, *Geophys. Res.Lett.*, 43, 2135–2143,  
doi:10.1002/2015GL067492, 2016.

Mentaschi, L., Vousdoukas, M., Voukouvalas, E., Sartini, L., Feyen L., Besio, G., Alfieri, L.: Non-stationary extreme value  
analysis: a simplified approach for Earth science applications, *Hydrol. Earth Syst. Sci. Discuss.*, 1–38, 2016.

865 Moss, R., Babiker, M., Brinkman, S., Calvo, E., Carter, T., Edmonds, J., et al.: Towards new scenarios for analysis of  
emissions, climate change, impacts and response strategies. IPCC Expert Meeting Report, 25 pp, 19–21 September, 2007,  
Noordwijkerhout, The Netherlands, <https://www.ipcc.ch/site/assets/uploads/2018/05/expert-meeting-ts-scenarios-1-1.pdf>,  
2008.

870 Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., van Vuuren, D. P., et al.: The next generation of  
scenarios for climate change research and assessment, *Nature*, 463(7282), 747–756. <https://doi.org/10.1038/nature08823>,  
2010.

875 Nicholls, R. J., Hanson, S., Herweijer, C., and Patmore, N.: Ranking port cities with high exposure and vulnerability to  
climate extremes, <https://www.oecd-ilibrary.org/content/workingpaper/011766488208>, 2008.



- Oppenheimer M., Glavovic B., Hinkel J. et al.: Chapter 4: sea level rise and implications for low lying islands, coasts and communities. In: Pörtner H.O., Roberts D., Masson-Delmotte V. et al (eds). IPCC Special Report on the Ocean and Cryosphere in a Changing Climate. Intergovernmental Panel on Climate Change (IPCC), 2019.
- 880 Pagney, F.: Genèse et dynamique de l'ouragan Hugo sur la Guadeloupe, In Annales de géographie (pp. 152- 165), Armand Colin. (1991).
- Planton, S., Le Cozannet, G., Cazenave, A., Costa, S., Douez, O., Gaufrès, P. et al.: Le climat de la France au XXI<sup>e</sup> siècle. Vol 5: Changement climatique et niveau de la mer: de la planète aux côtes françaises. [Rapport de recherche] Ministère de  
885 l'Ecologie et du Développement Durable, 2015.
- Raftery, A. E., Zimmer, A., Frierson, D. M. W., Startz, R. and Liu, P.: Less than 2 °C warming by 2100 unlikely, Nat. Clim. Chang., 7, 637–641, <https://doi.org/10.1038/nclimate3352>, 2017.
- 890 Ranasinghe, R.: Assessing climate change impacts on open sandy coasts: A review. Earth-Science Reviews, Volume 160, Pages 320-332, ISSN 0012-8252, <https://doi.org/10.1016/j.earscirev.2016.07.011>, 2016
- Rasmussen, D. J., Kulp, S., Kopp, R. E., Oppenheimer M. and Strauss B. H.: Popular extreme sea level metrics can better communicate impacts, Climatic Change 170, 30, <https://doi.org/10.1007/s10584-021-03288-6>, 2022.
- 895 Ruggiero, P., Buijsman, M., Kaminsky, G. & Gelfenbaum, G.: Modeling the effects of wave climate and sediment supply variability on large-scale shoreline change. Marine Geology 273, 127-140, 2010.
- Seneviratne, S.I., Zhang X., Adnan M., Badi W., Dereczynski C., Di Luca A. et al.: Weather and Climate Extreme Events in  
900 a Changing Climate. In Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1513–1766, doi:10.1017/9781009157896.013, 2021.
- 905 Sheremet, A., Staples, T., Arduin, F., Suanez, S. & Fichaut, B.: Observations of large infragravity runup at Banneg Island, France, Geophysical Research Letters, 41, 976–982, <https://doi.org/10.1002/2013GL058880>, 2014.



- 910 Slangen, A.B.A., Adloff, F., Jevrejeva, S., Leclercq, P.W., Marzeion, B., Wada, Y., and Winkelmann R.: A review of recent updates of sea-level projections at global and regional scales, *Surv. Geophys.*, 38, no. 1, 385-406, doi:10.1007/s10712-016-9374-2, 2017.
- Syvitski, J.P.M.: Deltas at risk, *Sustainability Science* 3, 23–32, <https://doi.org/10.1007/s11625-008-0043-3>, 2008.
- 915 Talke, S. A., Familkhalili, R., Helaire, L. T., Jay, D. A., Orton, P. M. & Ralston, D. K.: The influence of human induced landscape and bathymetry changes on tides, surge and extreme water levels. In *Ocean Sciences Meeting 2020*. AGU, 2020.
- Tiggeloven, T., de Moel, H., Winsemius, H. C., Eilander, D., Erkens, G., Gebremedhin, E., Diaz Loaiza, A., Kuzma, S., Luo, T., Iceland, C., Bouwman, A., van Huijstee, J., Ligtoet, W., and Ward, P. J.: Global-scale benefit–cost analysis of coastal flood adaptation to different flood risk drivers using structural measures, *Natural Hazards and Earth System Sciences*, 20, 920 1025–1044, <https://doi.org/10.5194/nhess-20-1025-2020>, 2020.
- U.S. Army Corps of Engineers: *Coastal Engineering Manual*, U.S. Army Corps of Engineers, Washington, DC., 2002.
- 925 van Vuuren, D.P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., et al.: The representative concentration pathways: an overview. *Climatic Change* 109, 5. <https://doi.org/10.1007/s10584-011-0148-z>, 2011.
- Vitousek, S., Barnard, P. L., Fletcher, C. H., Frazer, N., Erikson, L. and Storlazzi, C. D.: Doubling of coastal flooding frequency within decades due to sea level rise. *Sci. Rep.*, 7(1), 1399. <https://www.nature.com/articles/s41598-017-01362-7>, 930 2017.
- Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Verlaan, M., and Feyen, L.: Extreme sea levels on the rise along Europe’s coasts. *Earth’s Future*, 5, 304–323. doi:10.1002/2016EF000505, 2017.
- 935 Vousdoukas, M.I., Mentaschi, L., Voukouvalas, E., Verlaan, M., Jevrejeva, S., Jackson L. P. & Feyen, L.: Global probabilistic projections of extreme sea levels show intensification of coastal flood hazard. *Nat Commun* 9, 2360. <https://doi.org/10.1038/s41467-018-04692-w>, 2018.