



Influence of data source and copula statistics on estimates of compound extreme water levels in a river mouth environment

Kévin Dubois^{1,2}, Morten Andreas Dahl Larsen^{3,4}, Martin Drews³, Erik Nilsson^{1,2}, Anna Rutgersson^{1,2}

¹Department of Earth Sciences, Uppsala University, Uppsala, 752 36, Sweden

5 ²Centre of Natural Hazards and Disaster Science (CNDS), Uppsala University, Uppsala, 752 36, Sweden

³Department of Technology, Management and Economics, Technical University of Denmark, Lyngby, 2800, Denmark

⁴Danish Meteorological Office, Copenhagen, 2100, Denmark

Correspondence to: Kévin Dubois (kevin.dubois@geo.uu.se)

Abstract.

10 Coastal and riverine floods are major concerns worldwide as they can impact highly populated areas and result in significant economic losses. In a river mouth environment, interacting hydrological and oceanographical processes can enhance the severity of floods. The compound flood risks from high sea levels and high river runoff levels are often estimated using statistical copulas. Here, we systematically investigate the influence of different data sources and the choice of statistical copula on extreme water level estimates. While we focus on the river mouth at Halmstad city (Sweden), the approach presented is
15 easily transferable to other sites. Our results show that the compound occurrence of high sea levels and river runoff may lead to heightened flood risks as opposed to considering them as independent processes and that in the current study, this is dominated by the hydrological driver. We also show that the choice of data sources and copula can strongly influence the outcome of such analyses. Our findings contribute to framing existing studies, which typically only consider selected copulas and data sets, by demonstrating the importance of considering uncertainties.

20 1 Introduction

Floods can cause severe damage and disrupt activities and infrastructures in harbours and coastal communities. Flooding can result from meteorological, hydrological and oceanographic sources such as storm surges, extreme river runoff or precipitation. Storm surges correspond to seawater being pushed by the wind stress and the barometric pressure effect under deep low-pressure weather systems. Heavy precipitation can form under different conditions such as intense cyclonic activity
25 or convective weather. River runoff can have different origins, such as snow melting upstream or intense precipitation. River runoff and precipitation may take some time to drain into the sea, and the flow from land to the sea can therefore be slowed down or even momentarily become blocked when storm surges happen (Wahl et al., 2015).

Settlements and infrastructure located in river mouth environments are inherently susceptible to all of the above. The combination of multiple factors, extreme or not, happening at the same place simultaneously, successively, or consecutively,
30 can potentially lead to larger, compounded floods, and more severe impacts on the environment and society. Compound



flooding can also result when preceding conditions amplify the impact of the event (Andrée et al., 2023; Zscheischler et al., 2020; AghaKouchak et al., 2020). Even if trends over the last forty to sixty years are estimated with high uncertainties, it is likely that extremes including compound events are becoming more severe in Northern Europe in a changing climate (Rutgersson et al., 2021).

35 Couason et al. (2020) highlight the importance of considering interactions between river discharge and storm surge
extremes in river mouth environments. They demonstrate that dependencies between these factors are not random and may
result from relations between weather systems at the synoptic scale with local conditions such as the topography. In
Northwestern Europe, it has been shown that the fluvial flood hazard increases with high sea levels and stronger storms, and
that this may be critical in populated and low-elevation coastal areas (Ganguli et al., 2019). For example, Eilander et al. (2020)
40 find the Baltic Sea and the Kattegat basin to be particularly susceptible to compound flood hazards based on the dependency
between skew surge levels and river discharge. Without considering the occurrence of storm surges, Eilander et al. (2020)
further show that flood depths are underestimated and subsequently so is the estimated number of people exposed to river
floods in this area. Meanwhile, Moftakhari et al. (2017) demonstrate that sea level rise (SLR) is likely to increase the impacts
from compound flooding by 2030 and 2050 under the Representative Concentration Pathways (RCPs) 4.5 and 8.5 for eight
45 major cities around the US coastline.

Compound flooding in coastal areas can also be caused by a combination of heavy precipitation and high sea levels.
Hence, the probability of compound flooding is expected to significantly increase in the Baltic and North Sea areas, where an
event with a current return period (RP) of around sixty years is projected to occur every ten years in 2100 due to the
combination of SLR and increased extreme precipitation (Bevacqua et al., 2019).

50 Not taking compound flooding effects into account may result in an underestimation of the flood hazards in the coastal
zone, including river mouths. Thus, analyzing and understanding these events is of high relevance to coastal communities. In
this study, we evaluate the potential impact of extreme hydrological and oceanographic coastal events on the coastal city of
Halmstad (Sweden), which is a port, industrial, and recreational city at the mouth of the Nissan river. Halmstad is located on
the west coast of Sweden (fig. 1) and is naturally prone to coastal, fluvial, and pluvial flooding. The area is subject to
55 extratropical cyclones (Hoskins and Hodges, 2002; Dacre et al., 2012), resulting in rather high sea levels by storm surges for
the area (Wolski et al., 2014). According to the Swedish Meteorological and Hydrological Institute (SMHI), Halmstad recorded
the highest ever sea level measured in Sweden of 235 cm on the 29th of November 2015 during the wind storm “Gorm”.
Halmstad has also been severely impacted by river floods. While Nissan is the main river crossing the city of Halmstad, smaller
rivers are also present and can create floods, such as Fylleån’s river. Finally, the West Coast of Sweden is found to be one of
60 the areas in Sweden expecting the most significant impacts due to SLR during this century (Hieronymus et al., 2020). Thus,
nearby studies have stressed the necessity to update coastal protection measures along the Swedish (Hieronymus et al., 2022)
and the German Baltic Sea shorelines (Kiesel et al., 2023).



The main goal of the current study is to investigate the impact of different data sources, methodology, and representation of compound extremes on estimates of extreme water levels. Using Halmstad as an example, we explore the potential influence on flood risk assessments related to compound effects from river flooding within the coastal area.





Figure 1: Regional map of southern and central Sweden with a zoom around our study area and the city of Halmstad (Pawlowicz, 2020; Sentinel 2 cloudless, 2021)

70 2 Data and Methods

In the following, compound effects are defined in terms of the interaction between coastal sea level and river discharge when at least one of the two is subject to an “extreme” value. An event is considered in the extreme range when a studied variable reaches its annual maximum value. The correlation between co-occurring events has been studied as it provides insight into the relationship between each set of two variables. The probability of getting an extreme river discharge associated with a high sea level and the opposite permits the assessment of the potential compound effects between those two processes.

Figure 2 presents the main steps of the workflow describing the methodology; this is described in the following sub-sections. Firstly, we analyzed different time-series records of sea level and Nissan River discharge data from models and observations at Halmstad using extreme value theory and a Generalized Extreme Value (GEV) distribution (Coles, 2001) – hereby referred to as the “univariate approach” - to estimate Return Levels (RLs) on every factor independently. Secondly, we defined sets of coupled events based on single variables. Thirdly, we analyzed the correlation between sea level and river discharge events. If this analysis indicated a potential compound event, we studied the co-dependency between the two variables by fitting a statistical copula distribution function (Sadegh et al., 2018). We finally performed a statistical analysis of the compound events to study the differences between each data source and its associated uncertainties.



85 **Figure 2: Workflow describing the methodology used in this paper, starting from the oceanographic and hydrological data (in orange) to the univariate (in blue) and bivariate (in green) approaches used for flood risk analysis.**

2.1 Data

90 An analysis of time-series records of sea level (fig. 3-a) and river discharge (fig. 4-a) at Halmstad was carried out. As mentioned above, a univariate distribution was initially fitted based on the GEV distribution and extreme value theory (Coles, 2001) for each time series collected. The temporal differences in the lengths of each dataset induce substantial differences and associated uncertainties, which dominate in case of extreme RP. Consequently, a moderately extreme 30-year RP event was chosen as the maximum value considered. For comparison, we also consider more frequent events with a 5-year RP.

2.1.1 Sea level data

95 - Observations

Hourly sea level observations data were obtained from SMHI at Halmstad's tide gauge denoted as the station "HALMSTAD SJÖV" with station number 35115 in the open database provided by SMHI (fig. 1). However, to carry out this analysis, the period with sea level observations (titled "obs sea level") was insufficient as only 13 years (from 2009 to 2021) are available; based on the p-value from correlation analysis the associated results are statistically insignificant. To extend this sea level
 100 record, a set of reanalysis data and a machine learning approach have been investigated and used.



- ***Reanalysis***

Hourly sea surface variations (in m) covering the period from 1993 to 2020 with a spatial resolution of approximately two nautical miles have been provided by the Copernicus Marine Environment Monitoring Service's (CMEMS) Baltic Monitoring and Forecasting Centre (BAL MFC) (CMEMS, 2022). This reanalysis uses the ice-ocean model NEMO-NORDIC (Pemberton et al., 2017), and the data are assimilated with the Localized Singular Evolutive Interpolated Kalman (LSEIK) method (Nerger et al., 2005). Data is extracted from the closest grid point to Halmstad's tide gauge. This sea level dataset is named: "reanalysis".

110 - ***Machine learning model***

A probabilistic machine learning method, Quantile Regression Forests (QRF), is used (Meinshausen, 2006). Sea level records from the neighbouring station of Viken (station named "VIKEN", number 2228 in the SMHI database) is used to train the QRF model over an eight year-period, where it is correlated with Halmstad's observed sea level. The resulting sea level estimates at Halmstad include both mean predictions and percentiles to assess uncertainties and variability following the methodology introduced by Dubois et al. (2023). The last three years of available data at Halmstad are used to validate the QRF model, emphasizing extreme events predictions. The QRF model is used to produce a first dataset called: "predicted Halmstad" ("pred Halmstad") and a second one named: "reconstructed Halmstad" ("rec Halmstad").

The predicted Halmstad dataset provides daily sea level (in cm) for the full period of available sea level observations from the station at Viken, here from 1977 to 2021.

120 The reconstructed Halmstad dataset provides daily sea level (in cm). It joins both sets, i.e., combines observations from Halmstad from 2009 to 2021 and the predicted Halmstad data from the QRF model from 1977 to 2009. Thus, it also covers the period from 1977 to 2021. Further attempts to enrich the QRF model by including reanalysis data (i.e., as part of the training) did not improve the predicted sea levels in the reconstructed data sets significantly, which emphasized the need for local observations. These were fortunately available, even if not for the entire more extended period. Similar findings, that is, significant improvements when using local observations as means to train a machine learning of sea level were previously found in this region (e.g. Hieronymus et al., 2019).




Figure 3: Halmstad sea level time series and annual maxima from different sources: observations ("obs sea level"), reconstructed ("rec Halmstad") and predicted ("pred Halmstad") derived from a machine learning model trained on data from the Viken station, and reanalysis ("reanalysis") datasets (panel a). RIs estimated from corresponding GEV fits of each dataset and the associated 95th percentile confidence intervals (background colours). The dots depict empirical data (panel b).

130



135 It is not only the length of the observation period that is short. Also, the reanalysis dataset exhibits a bias and does not predict the observed extreme sea levels. Accordingly, the uncertainty estimated from both univariate GEV analyses are large (fig. 3-b). The predicted and reconstructed data sets yields result with a smaller uncertainty range. Hence, the reconstructed data set, which is based on observations when available, was chosen as the best source of sea level information for the bivariate analysis.

2.1.2 River runoff data

- *Observations*


140 River discharge data were obtained from SMHI at station 2471: “Nissaström” (fig. 1), covering a basin of 2437 km². Observations of daily river discharge in m³/s (obs Nissan) have been provided from 1997 to 2021.

- *E-Hype model*

145 Modelled river discharge data are taken from the Hydrological Predictions for the Environment (HYPE) model, which simulates water flows and quality at different spatial scales (Lindström et al., 2010). Daily temperature and precipitation values are used as dynamic forcing in this model. The European HYPE model: “E-Hype2016_version_16_g” (E-Hype) provides daily river discharge (in m³/s) from 1989 to 2021 (<https://vattenwebb.smhi.se/om-vattenwebb>).

- *S-Hype model*

150 SMHI has set up the HYPE model for Sweden, now used operationally to forecast hydrological conditions over Sweden, such as floods and droughts. It covers all of Sweden (450000 km²), where the country has been divided into sub-basins of 28 km² on average (Strömquist et al., 2012). S-Hype3 model’s data (S-Hype) of daily river discharge (in m³/s) has been provided from 2004 to 2020 (Donnelly et al., 2016).



155

Figure 4: Nissan's river time series and annual maxima from different sources: observations ("obs Nissan"), European HYPE model ("E-Hype") and Swedish HYPE model ("S-Hype") from SMHI (panel a). RIs derived from GEV fits to each dataset are shown with 95th percentile confidence intervals (background colours). The dots are the empirical data (panel b).

160 The available time series associated with the S-Hype model is rather limited, leading to a wide uncertainty band when carrying out the univariate analysis (fig. 4). Conversely, the Nissan observations and E-Hype data sets lead to RIs that are associated

with more bounded uncertainty estimates. In this light, we choose the E-Hype dataset for the bivariate analyses as the data are available over a more extended period. The largest RLs are seen for the E-Hype dataset for RPs above five years.

165 2.1.3 Sets of coupled events

To study compound events in a river mouth environment from both a river discharge and sea level perspective, we defined two different sets of events based on the data discussed above. The first one used sea-level annual maxima (S_n) and associated daily maximum river discharge (q_n) within a defined time period centered on the date of S_n (+/- Δ days). The second one uses river discharge annual maxima (Q_n) and associated hourly maximum sea levels (s_n) within a defined three days window (Couasnon et al., 2020; Moftakhari et al., 2017; Sadegh et al., 2018) centered on the date of Q_n (+/- one day). Each of the four sea level time-series observed and modelled records were then correlated with each of the three river discharge ones, which makes up a total of twelve different datasets (table A2).

2.2 Statistical analysis

175 - *Univariate analysis*

To estimate the extreme values of Nissan's river runoff and Halmstad sea levels, and their associated RPs, a GEV distribution was fitted to the annual extremes separately for each time series record (Coles, 2001; Ahsanullah, 2016). This was done using the MATLAB-based GEV-fitting algorithm, which provides parameter estimates and 95% confidence bands. Here, we do not make any assumption concerning the dependence between the two variables of interest, sea level and river discharge; each variable is modelled independently based on its own marginal distribution.

- *Bivariate analysis*

Initially, the Pearson, Spearman and Kendall's correlation coefficients and the associated p-values were calculated for each of the twelve collated data sets to assess whether there was a relationship between river runoff and sea level. The usual threshold value of 5% was defined as evidence for rejecting the H_0 null hypothesis, that is, the two variables are independent. When p-values were found to be lower than the threshold, the null hypothesis could be rejected, and the two variables similarly found to show significant dependency. However, when p-values are above 5%, H_0 cannot be rejected, so the two variables can be independent.

To represent the compound extremes, we apply copula modelling, which has been found to be useful for representing a joint probability (Hao et al., 2016). The analyses were carried out using the Multihazard Scenario Analysis Toolbox (MhAST), Version 2.0 from Sadegh et al., (2018). The statistical copula method models the dependence structure of the two random variables (Joe, 2014; Sadegh et al., 2017). It links or joins individual univariate distributions into a joint multivariate distribution that has a specified correlation structure (Tootoonchi et al., 2022). The MhAST toolbox fits 25 different copulas



to an input dataset. It first calculates the best possible fitting marginal distribution for each univariate dataset. It then proposes
195 the best copula fit based on the Maximum Likelihood, Akaike Information Criterion (AIC), and the Bayesian Information
Criterion (BIC). The root mean square error (RMSE) and Nash-Sutcliffe efficiency (NSE) values are calculated for each
copula. Here, we evaluated the difference between each of the copula fits. An uncertainty analysis was also carried out using
the MhAST toolbox with a “Weighted Average” and a “Maximum Density” approach (Sadegh et al., 2018). This first approach
reproduces a distribution of potentially compound hazards. Based on the determined copula probability function, random
200 samples are weighted accordingly and then calculated from the critical joint RP. The second approach is based on the “most
likely scenario” and provides an uncertainty range around it by adjusting the parameters of the copula distribution fit.
Two types of Hazard Scenarios (HSs) have commonly been proposed to study the risk of compound floods related to sea level
and river discharge (Salvadori et al., 2016; Mofatkhari et al., 2019; Serinaldi, 2015). The “AND scenario” corresponds to a
scenario where both the river discharge and the sea level are large enough to make a bivariate occurrence hazardous. The “OR
205 scenario” corresponds to a scenario where either the river discharge or the sea level or both are large enough to make a bivariate
occurrence troublesome.

2.3 Methodology

Firstly, a correlation analysis was carried out for each set, as proposed in section 2.2. This analysis investigated the
significance of independence between the sea level and river discharge during extreme occurrences. Then, each set was used
210 as input to the MhAST toolbox, which performed the compound analysis and returned 25 copula fits ranked depending on
different criteria (section 2.2). Among the 25 copulas fitted, only the ones presenting a closed-form joint probability density
function (Sadegh et al., 2017) were further investigated since, in these cases, “most likely scenarios” and their associated
uncertainties can be defined. Chosen RLs were calculated for each copula, and their uncertainties were assessed. Adopting the
“AND scenario” (see above) permitted us to investigate the dependency between sea level and river discharge during extreme
215 events. Conversely, the “OR scenario” was finally preferred when looking at RLs as this looks into the “total” risk of flooding,
whether originating from hydrological, coastal sources or both in combination.

To compare and evaluate the role of copulas and the role played by sea level and river discharge, respectively, a
notion of normalized difference value (NDV) was introduced. We defined it as the normalized difference between the RL
values of interest from the bivariate analyses. Here, we normalize relative to the corresponding E-Hype univariate RL, which
220 yields a representative dimensionless quantity. It should be pointed out that this quantity does not represent the “amplification”
with respect to the univariate case since, as shown by Serinaldi (2015), one cannot compare RLs of different dimensionality.
Suppose we investigate the resulting spread from using different copulas (based on the most likely scenarios within one set
under the 5-year RP); the NDV is calculated as the difference between the maximum and minimum value of the most likely
scenarios for any copula within a specific set divided by univariate 5-years RL derived from the E-Hype data set (section 3.2).
225 When we look into the sensitivity of river discharge datasets, the sea level dataset is fixed, and the NDV is measured as the
normalized difference between the maximum and minimum values of the most likely scenarios of best fits among the three




sets of associated river discharge divided by the corresponding E-Hype univariate RL; and vice-versa, when looking into the sensitivity of sea level datasets. The NDV term indicates the magnitude of change or difference in RL results we can expect when choosing a certain copula or input dataset. Very small NDVs suggest that the corresponding choice of a variable of interest does not strongly influence the results. In contrast, large NDVs indicate that a particular choice results in significant differences.

3 Results and discussion

From the rank correlation analysis, the datasets based on sea level annual maxima (S_n , q_n) did not reveal any significant dependency (i.e., “compoundness”) between sea level and river discharge, and therefore, no copula analysis was done (table A.1). In this case, the univariate analysis seemed to fit best under the proposed conditions of this study. Conversely, the datasets based on river discharge annual maxima (Q_n , s_n) yielded significant dependencies, suggesting a possible compound impact on river discharge. In the following, we mainly focus on the “OR scenario” (see above) as we are interested in the compound flood risk driven regardless of the situation (oceanographic or hydrological).

3.1 Dependency / Interdependency of the variables

Figure 5 and Table A1 show the dependency between the river discharge annual maxima and associated sea level local maxima (Q_n , s_n) event sets as expressed in section 2.1. Figure 5 displays the best copula distribution fit: BB1 from the rec Halmstad / E-Hype set under the “AND scenario” hypothesis for the 2, 5, 15 and 30-year RP. The full lines depict the RLs considering sea level and river discharge as dependent variables (derived from the best copula distribution fit). In contrast, the dashed lines show analogous results when assuming the two variables to be independent. The figure shows that the lines are not superposed, highlighting a dependency. Also, for all RLs presented, each RL from the independent hypothesis (dashed line) is placed below each corresponding RL from the dependent hypothesis (full line), supporting the hypothesis that compound events lead to higher flood risks when considering compound extremes. Therefore, for example, a 30-year RP, when looking at the independent variables, would become a 13-year RP when considering the variables’ dependency. This frequency increase comes from the compound effects and can be highlighted for each RP and copulas tested. Also, the dependency between extreme hydrological conditions and high oceanographic ones stresses the presence of compound effects, which lead to higher levels of river discharge and sea level during such events at the estuary. A probability density function is assigned to each copula fit curve, which permits obtaining a probability of co-occurrences for a possible event along each curve, which is later used to carry out the uncertainty analysis (Sadegh et al., 2018). For example, along the 5-year RP curve, the probability of getting a 5-year RL of 180 m³/s river discharge and 93 cm sea level is higher than getting one of 201 m³/s and 20 cm or one of 101 m³/s and a 112 cm one (fig. A1).



260 **Figure 5: RLs for base case set rec Halmstad / E-Hype. Full lines correspond to the return period isolines of river discharge (y-axis) annual maxima and associated sea level (x-axis) maxima (Q_n, s_n). The dashed lines represent the distribution fit, assuming the independence between both variables. Blue dots show observed data. BB1 copula is used to model the dependence of river discharge annual maxima and associated sea level maxima calculated for each RPs visible in red text (2, 5, 15 and 30 years).**

3.2 Compound hazard potential on river floods

265 The focus is on river discharge RLs as a proxy for fluvial flooding indicators. Figure 6 represents the 5- and 30-year river discharge RLs from the set rec Halmstad / E-Hype under the “OR scenario” hypothesis for each copula tested and its associated uncertainties values from two approaches. The best copula fit based on the different criteria in this case (section 2.2) is BB1 (red diamond). The stars and diamonds represent the maximum density of the calculated RL for each copula, which can be interpreted as the most likely scenario under the bivariate analysis.

270 - ***RP = 5 years***

The 5-year RL from the E-Hype model is 201 m³/s with a 95th percentile confidence interval of 167-246 m³/s under the univariate GEV distribution fit. The BB1 copula fit has a 5-year RL of 220 m³/s. The copulas’ 5-year RLs of “most likely scenarios” differ around 26 m³/s, all between 208 and 234 m³/s. The RL copulas’ uncertainties are displayed with the boxplots from two methods: the “Weighted Average” approach showed with the outlined error bars, and the “Maximum Density” approach showed with the filled error bars. The “Weighted Average” approach gives more considerate uncertainty ranges than

275



the “Maximum Density” (section 2.2). Indeed, for the best copula fit, the “Maximum Density” approach looking at the uncertainty of the “most likely scenarios” results in a narrow band of a maximum of 19 m³/s per copula against a more extensive range of 159 m³/s going from 202 to 361 m³/s with the “Weighted Average” approach. All copulas present a similar pattern. Moreover, the RL uncertainties for the “Maximum Density” approach are all located within the 95th confidence interval of the univariate RL. However, the “Weighted Average” approach gives a 75th percentile of around 255 to 269 m³/s and a nonoutlier maximum of around 324 to 361 m³/s above the 246 m³/s corresponding to the 95th percentile of the univariate GEV fit, indicating the importance of considering bivariate analysis method. The BB1 copula chosen as the best fit here by the different evaluation criteria mentioned in section 2.2 presents neither the smallest nor largest uncertainty band.

285 - ***RP = 30 years***

The 30-year RL from the E-Hype model is 263 m³/s with a 95th percentile confidence interval going from 189 to 431 m³/s under the univariate GEV distribution fit. The BB1 copula fit has a 30-year RL of 278 m³/s. The copulas’ 30-year RL of “most likely scenarios” differ by around 25 m³/s, with all of them between 267 and 292 m³/s, except for the Gaussian copula. For all copulas except the Gaussian one, the “Weighted Average” approach gives a more extensive uncertainty range than the “Maximum Density” one. Indeed, for the best copula fit, the “Maximum Density” approach results in a relatively narrow band of 24 m³/s, going from 272 to 296 m³/s, against a more extensive range of 92 m³/s going from 261 to 353 m³/s with the “Weighted Average” one. All copulas except the Gaussian and the Tawn ones present a similar pattern. Moreover, all RL uncertainties for both uncertainty analysis approaches are within the 95th confidence interval of the univariate RL for the 30-year RP.

295


- ***Sensitivity to the choice of copula***

For both 5- and 30-year RPs, the copulas and their associated uncertainties present a similar pattern with higher RLs when considering the compound effects. Depending on the choice of copulas, the most likely scenarios differ up to 26 m³/s for the 5-year RP and up to 25 m³/s for the 30-year RP, with the Tawn copula giving the minimum value and the Fischer-Kock and FGM copulas giving the maximum value. When only looking at the most likely scenarios values for each copula, they differ in a range approximately equal to 13% and 9.5% for the 5- and 30-year RPs, respectively (table A3). For each copula, the uncertainties’ relative errors based on the “Maximum Density” approach differ from 1% (Gaussian) to 9.9% (Fischer-Kock) and from 2.3% (Joe) to 52% (Gaussian) for the 5- and 30-year RPs respectively; from 63% (Joe) to 75% (Fischer-Kock) and from 16% (Joe) to 35% (Fischer-Kock) for the “Weighted Average” approach. For comparison, the relative errors for the univariate GEV fit are around 41% and 92% for the 5- and 30-year RPs, respectively. When considering the “Maximum Density” uncertainties, all RLs of all copula are in the range of 208-234 m³/s for the 5-year RL and 262-398 m³/s for the 30-year RL (Table A3; fig. 6).

305
These differences in resulting RLs emphasize the importance of the role played by the choice of copulas and the consideration of quantifying uncertainties.



310



315

Figure 6: 5- (a) and 30-year (b) RLs of river discharge from copula fits and univariate GEV fits for rec Halmstad / E-Hype. Each column represents a copula distribution fit. Stars represent the most likely scenario return values from each copula for each set, and the red diamond is the best copula fit. Two uncertainty approaches are displayed as boxplots, giving a statistical summary. Median, first, and third quartiles are represented in each box, whiskers represent minimum/maximum values, and dots represent outliers. Outlined boxplots correspond to the “Weighted Average” approach, and filled ones to the “Maximum Density” approach.

3.3 Sensitivity analysis on compound flood hazard potential -OR scenario-

This section focuses on the impact of data sources on resulting RP statistics, aiming to compare copula analyses considering compound events. As seen in section 2.1, we have twelve possible data sets to analyze for Halmstad city extracted from models and observations. As mentioned in section 2.1, the univariate analysis presents different results, including RL values and confidence intervals for each river runoff time series.

5-year and 30-year univariate RLs of river runoff, respectively, differ by around 9 m³/s and 21 m³/s with values of 202 m³/s and 241 m³/s based on observation gauge (red); 193 m³/s and 252 m³/s based on S-Hype model (blue); 201 m³/s and 263 m³/s based on E-Hype model (green) as displayed in Figure 4-b. However, uncertainties associated with the 95th percentile confidence interval differ vastly from respectively around 86 m³/s and 185 m³/s (observation); 121 m³/s and 811 m³/s (S-Hype); 79 m³/s and 242 m³/s (E-Hype) as displayed with the background colours on the figure.

3.3.1 Dependency / Independency of the variables

Figure 7 presents resulting RLs for combined ranges of each variable set for the 5- and 30-year RPs as in Figure 5, but with results from six different data sources to study the resulting impacts. The dependency is evident for each set, with

330




each full line moved away from its corresponding dashed line, highlighting the dependency and compound effects for any sets tested. The differences between solid and dashed lines in Figure 7 are typically contained within about 20 cm sea level or 25 m³/s river discharge, constituting about 10-15% of the extreme 5- and 30-year RLs for the site with a gap increasing with higher RPs. At first glance, these differences may be perceived as a reasonably small compound effect, but every little increase

335 in extreme situations can have a consequence for society. It should be noted that switching data sources may have a more significant effect on estimated RLs; hence, both method and choice of data are essential.

Some sets behave similarly as their corresponding dashed and full lines are almost superposed as for the sets obs Nissan / pred Halmstad & rec Halmstad or E-Hype / pred Halmstad & rec Halmstad in both 5- and 30-year RPs. This similarity stresses the idea that river discharge predominates over sea-level inputs.

340



345

Figure 7: 5- (a) and 30-year (b) RPs isolines of river discharge (y-axis) annual maxima and associated sea level (x-axis) maxima (Q_n , s_n) for Halmstad. Full lines implement the compound effect, and dashed lines represent fit, assuming independence between both variables. Dots show observed data. The best copula fit is used to model the dependence of (Q_n , s_n) calculated for each set visible in coloured text.

3.3.2 Compound hazard potential on river floods

The most likely scenarios of 5- and 30-year RPs and their associated uncertainties on the different sets are calculated as described in section 2.1. This study focuses on extreme hydrological events associated with oceanographic conditions and, therefore, concentrates on the RLs of river discharge. Figure A2 displays those results for each set in the same way as Figure

350 6. Figure A2-a returns the results of the 5-year RP and Figure A2-b, the 30-year RP analysis.



Under the 5-year RP, lower RLs are found for all obs sea level sets (fig. A2-a), corresponding to the data available and short duration of overlapping periods, with a maximum of 13 years and limited by the extent of sea level observations. Under the 30-year RP, lower RLs are found for the set E-Hype / obs sea level and all S-Hype sets except for S-Hype / obs sea level, which presents the most extreme values.

For the 5- and 30-year RPs, the three sets associated with the E-Hype model, which show statistical significance, lead to similar and higher values, respectively, than all the other sets. The last set showing statistical significance is associated with the S-Hype model and leads to somewhat different results between the 5-year RP, with slightly lower RLs, and the 30-year RP, with generally higher RLs (fig. A2).

All most likely scenario values calculated from the copula analysis under both the 5- and 30-year RPs are within the range of the 95th percentile confidence interval of the univariate GEV distribution fit (fig. A2).

Uncertainties associated with the copula analysis and following the “Maximum Density” approach do not extend too much from the median values and stay within the confidence interval of the univariate GEV distribution (fig. A2) for most of the copulas tested. Under this “Maximum Density” approach and based on the best copula fits, they differ by about 3-8 m³/s for the 5-year RP and 2-9 m³/s for the 30-year RP. Under those same conditions, the uncertainties from the “Weighted Average” approach vary between 65 and 149 m³/s for the 5-year RP and between 37 and 68 m³/s for the 30-year RP. Therefore, uncertainties related to the “Maximum Density” approach associated with the most likely scenarios are relatively small, providing reasonable confidence in such scenarios. Conversely, the “Weighted Average” approach uncertainties provide a confidence interval on possibly more extreme scenarios, which is relevant when communicating RLs.

370

- *Input datasets selection*

Depending on the choice of river time series as initial input, the results of the copula analysis under the 5- and 30-year RPs differ substantially around a maximum of 20 m³/s and 40 m³/s, respectively, with an NDV range of 6-10% and 8.4-15% (fig. 8). This contrasts with the choice of sea level time series as initial input with a maximum difference of around 6 and 12 m³/s equivalent to 1.5-3% and 2.3-4.6% NDV bounds for the 5- and 30-year RPs, respectively, without considering the three sets associated with obs sea level. Those results are based on the most likely scenarios from each best copula fit and did not consider the obs sea level associated sets. It emphasizes that the choice of sea level records has a lower influence than the one of river discharge within this study on compound hydrological extreme events on our example study site (Halmstad). Similar results could be expected for the surrounding area (West coast of Sweden).

380

- *Copula selection*

To evaluate the role played by choice of the copula, we calculated the NDV for each set between the maximum and the minimum values returned by the 18 copulas tested without considering the sets with obs sea level data input as it was too short for bivariate analysis. Among all the different sets, the BB1, the Gaussian and the Clayton copulas are the best ones based on the different statistical criteria (section 2.2). Moreover, when only looking at the sets associated with the same river runoff


385



input, the best copula fit is the same: Clayton for obs Nissan, BB1 for E-Hype except for S-Hype, which has Gaussian as the best fit for S-Hype/rec Halmstad and Clayton for the two other sets.

For most of the sets, the “Fischer-Kock” and the “FGM” copulas give the highest RLs and the “Tawn” and “Joe” copulas give the smallest ones (fig. A2). It results in NDVs between 5.5% and 13% for the 5-year RP and between 3.8% and 9.5% for the 30-year RP. The base case E-Hype / rec Halmstad presents the highest NDVs when comparing with other sets NDVs, which emphasizes that the choice of copula is relatively more important than in other sets (fig. 8). Based on our assumption that this is possibly the best set, in terms of data sources, it stresses the idea that the choice of copula becomes more and more critical when input datasets are long enough and statistically significant.

Therefore, the choice of copula has a similar influence as the choice of river discharge records for each of the nine sets tested here, as the obs sea level has not been considered. For both the 5- and 30-year RPs, the choice of sea level is the least impactful. Under the 5-year RP, the choice of copula is overall the most important before the choice of river discharge, but under the 30-year RP, the choice of river discharge predominates. However, this differs when looking at specific sets’ copula NDVs (fig. 8).



400 **Figure 8: Normalized difference values (%) for evaluating the importance of copula fit and forcing data for both 5- and 30-year RPs as mentioned in section 3.3.2.**



4 Limitations

Observed time-series datasets have a relatively short length, leading to rather high uncertainties once applying the
405 GEV analysis. Similarly, model time-series datasets have inherent uncertainties, which can be challenging to quantify. Various
data sources were assessed for their applicability in bivariate analysis, and direct sea level observations available for only 13
years were a limiting factor. We focus on longer reconstructed time series and other data sources for the principal analysis.
Moreover, assumed stationarity within the datasets can be a limitation while performing the statistical analysis (Kudryavtseva
et al., 2021).

410 A compound analysis is seen as a relatively new approach within this field of study, which also involves some
limitations, such as the quantification of uncertainties within a multivariate analysis that differ widely depending on the choice
of a copula.

5 Summary and conclusions

This study assesses the hydrological and oceanographic processes that may lead to compound flood effects in
415 Halmstad. The method is easily transferable to other regions or sites. In the paper, we stress the importance of the choice of
data sources and copulas for multivariate analysis. Based on our analysis, we conclude that:

- A dependency is found between the annual maxima of river discharge and the corresponding sea level. The opposite
dependency for annual sea level maxima and associated river discharge was not considered significant at this site.
- 420 • All values of the “most likely” scenarios and their uncertainties resulting from the copula analysis are within the range
of the 95th percentile confidence interval of the univariate GEV distribution fit.
- The choice of river time series as initial input influences the results of the copula analysis to a higher degree than the
choice of sea level time series as initial input.
- The choice of copula has a similar magnitude of its influence on return period statistics as the choice of river discharge
425 input for most of the twelve sets tried.
- According to statistical criteria, the Clayton, BB1 and Gaussian (once) copulas performed the best in this study.

Uncertainties in compound flood hazard quantification are essential to consider. They can come from different sources, such
as methodology and data sources. This study stresses the importance of the choice of data sources and copula.



430 5 Appendices

Table A1: Rank correlation (rho) and p-values of the twelve different sets based on (Q_n/S_n) in columns: “ / river” and (q_n/S_n) in columns “ / sea” ; the best set of study is displayed with bold and underlined; p_values above 5% are highlighted in italic.

river	sea level	rank correlation	rho / river	p / river	rho / sea	p / sea	
E-Hype	reanalysis	Pearson	0,4532	0,0154	<i>0,0443</i>	<i>0,8227</i>	
		Kendall	0,3280	0,0141	<i>0,0370</i>	<i>0,7992</i>	
		Spearman	0,4532	0,0163	<i>0,0443</i>	<i>0,8226</i>	
obs Nissan		Pearson	0,2430	0,2478	0,6684	0,0922	
		Kendall	0,2896	0,1594	0,6410	0,0725	
		Spearman	0,2419	0,2478	0,6674	0,0922	
S-Hype		Pearson	0,3799	0,1325	<i>-0,0760</i>	0,7719	
		Kendall	0,3088	0,0914	<i>-0,0588</i>	0,7765	
		Spearman	0,3799	0,1333	<i>-0,0760</i>	0,7729	
E-Hype		observed	Pearson	0,4725	0,1030	0,4056	0,1908
			Kendall	0,3590	0,1000	0,2424	0,3108
			Spearman	0,4725	0,1057	0,4056	0,1926
obs Nissan	Pearson		0,4396	0,1329	<i>-0,1049</i>	0,7456	
	Kendall		0,3333	0,1289	<i>-0,0606</i>	0,8406	
	Spearman		0,4396	0,1350	<i>-0,1049</i>	0,7495	
S-Hype	Pearson		0,6273	0,0388	<i>-0,0490</i>	0,8799	
	Kendall		0,4909	0,0405	<i>-0,0303</i>	0,9466	
	Spearman		0,6273	0,0440	<i>-0,0490</i>	0,8863	
E-Hype	pred Viken		Pearson	0,4439	0,0109	0,2438	0,1788
			Kendall	0,3145	0,0111	0,1452	0,2518
			Spearman	0,4439	0,0116	0,2438	0,1782
obs Nissan		Pearson	0,3400	0,1040	0,2435	0,2516	
		Kendall	0,2319	0,1189	0,1522	0,3128	
		Spearman	0,3400	0,1044	0,2435	0,2505	
S-Hype		Pearson	0,3676	0,1466	0,2843	0,2687	
		Kendall	0,2794	0,1288	0,1912	0,3081	
		Spearman	0,3676	0,1471	0,2843	0,2678	
<i>E-Hype</i>		<i>rec Viken</i>	Pearson	0,4836	0,0044	0,2753	0,1273
			Kendall	0,3523	0,0036	0,1815	0,1500
			Spearman	0,4836	0,0048	0,2753	0,1272
obs Nissan	Pearson		0,3446	0,0916	0,1487	0,4880	
	Kendall		0,2333	0,1076	0,1014	0,5071	



S-Hype	Spearman	<i>0,3446</i>	<i>0,0921</i>	<i>0,1487</i>	<i>0,4863</i>
	Pearson	<i>0,4882</i>	<i>0,0550</i>	<i>0,1373</i>	<i>0,5994</i>
	Kendall	0,3833	0,0413	<i>0,1029</i>	<i>0,5976</i>
	Spearman	<i>0,4882</i>	<i>0,0572</i>	<i>0,1373</i>	<i>0,5986</i>

435 Table A2: Summary report of runs from MhAST copula analysis for the twelve different sets; the best study set is highlighted in bold, italic and underlined.

river	sea level	Copula best fit	number of co-occurring years
obs Nissan	reanalysis	Clayton	24
	obs sea level	Galambos	13
	pred Halmstad	Clayton	24
	rec Halmstad	Clayton	25
<i>E-Hype</i>	reanalysis	BB1	28
	obs sea level	Gaussian	13
	pred Halmstad	BB1	32
	<i>rec Halmstad</i>	<i>BB1</i>	<i>33</i>
S-Hype	reanalysis	Clayton	17
	obs sea level	BB1	11
	pred Halmstad	Clayton	17
	rec Halmstad	Gaussian	16

Table A3: Summary report from the river discharge's results and associated uncertainties from MhAST copula analysis for the E-Hype / rec Halmstad set; the results from the univariate method are highlighted in bold, italic and underlined. The Gaussian copula has not been considered for the analysis of the “most likely scenarios” row.

<i>E-Hype / rec Halmstad</i>	<i>copula distribution fit</i>		<i>5-years RL</i>	<i>30-years RL</i>
<i>univariate</i>		<i>median</i>	<i>201</i>	<i>263</i>
		<i>max 95%</i>	<i>250</i>	<i>431</i>
		<i>min 5%</i>	<i>167</i>	<i>189</i>
		<i>relative error</i>	<i>41,29%</i>	<i>92,02%</i>
<i>"most likely scenarios"</i>	Fischer-Kock / FGM	max copula	234	292
	Tawn	min copula	208	267
	<i>NDV</i>		<i>12,94%</i>	<i>9,51%</i>



uncertainties "Maximum Density" approach	BB1	max without outliers	224	284
	BB1	min without outliers	218	276
	BB1	NDV	2,99%	3,04%
	Joe	max without outliers	217	274
	Joe	min without outliers	210	268
	Joe	NDV	3,48%	2,28%
	Fischer-Kosck	max without outliers	235	296
	Fischer-Kosck	min without outliers	215	272
	Fischer-Kosck	NDV	9,95%	9,13%
	Gaussian	max without outliers	220	398
	Gaussian	min without outliers	218	262
	Gaussian	NDV	1,00%	51,71%
uncertainties "Weighted Average" approach	BB1	max without outliers	351	329
	BB1	min without outliers	202	261
	BB1	NDV	74,13%	25,86%
	Joe	max without outliers	328	303
	Joe	min without outliers	202	261
	Joe	NDV	62,69%	15,97%
	Fischer-Kosck	max without outliers	353	353
	Fischer-Kosck	min without outliers	202	261
	Fischer-Kosck	NDV	75,12%	34,98%
	Gaussian	max without outliers	349	338
	Gaussian	min without outliers	202	261
	Gaussian	NDV	73,13%	29,28%




Figure A1: Set E-Hype / rec Halmstad, best copula fit: BB1. [2, 5, 10, 25, 50, 100] RPs and associated densities. The left and lower panels correspond to marginal RPs curves of each univariate parameter individually, river discharge and sea level (extracted from MhAST software).



445 Fig. A2-a)







Fig. A2-b)





450 **Figure A2: 5- (a) and 30-year (b) RPs values of river discharge from copula fits and univariate GEV fits. Each subplot corresponds to a set of events from an association of river discharge and sea level inputs displayed as a matrix, and in each column, a copula distribution fit where two uncertainty approaches are displayed as error bars. Stars represent the most likely scenarios return values from each copula for each set and each red diamond, the best-fit copula. The two uncertainty approaches are displayed as boxplots that give a statistical summary. Median, First and third quartiles are represented in each box; Whiskers represent minimum and maximum values, and dots represent outliers. Outlined boxplots correspond to the “Weighted Average” approach, and filled ones to the “Maximum Density” approach. The set E-Hype / Rec Halmstad, used as a base case, is highlighted by the red rectangles.**

455 *Code availability.* The code used to generate the figures can be acquired by contacting the first author (kevin.dubois@geo.uu.se).

Data availability. Observations data are available from SMHI at <https://www.smhi.se/data/oceanografi/ladda-ner-oceanografiska-observationer#param=sealevelrh2000,stations=core> for the sea level (accessed on: 14th of October 2021) and <https://vattenweb.smhi.se/station/> for the river discharge (accessed on: 28th of April 2021).

460 *Author contributions.* KD conducted the analysis. KD prepared the manuscript with contributions from all co-authors.

Competing interests. The authors declare that they have no conflict of interests.

Acknowledgements. The work forms part of the project: Extreme events in the coastal zone – a multidisciplinary approach for better preparedness. We further thank Johanna Mård, Christoffer Hallgren and Faranak Tootoonchi for our useful discussions.

465 *Financial support.* This research was funded by the Swedish Research Council FORMAS (Grant No. 2018-01784) and the Centre of Natural Hazards and Disaster Science (CNDS).

References

- Aghakouchak, A., Chiang, F., Huning, L. S., Love, C. A., Mallakpour, I., Mazdizyani, O., Moftakhari, H., Papalexiou, S. M., Ragno, E., and Sadegh, M.: Climate Extremes and Compound Hazards in a Warming World, *Annu. Rev. Earth Planet. Sci.*, 48, 519–548, <https://doi.org/10.1146/annurev-earth-071719-055228>, 2020.
- 470 Ahsanullah, M.: Extreme value distributions, edited by: Chris P. Tsokos, Tampa, U., https://doi.org/10.2991/978-94-6239-222-9_137, 2016.
- Andrée, E., Su, J., Andreas Dahl Larsen, M., Drews, M., Stendel, M., and Madsen, K. S.: The role of preconditioning for extreme storm surges in the western Baltic Sea, *Nat. Hazards Earth Syst. Sci.*, 23, 1817–1834, <https://doi.org/https://doi.org/10.5194/nhess-23-1817-2023>, 2023.



- 475 Bevacqua, E., Maraun, D., Vousdoukas, M. I., Voukouvalas, E., Vrac, M., Mentaschi, L., and Widmann, M.: Higher probability of compound flooding from precipitation and storm surge in Europe under anthropogenic climate change, *Sci. Adv.*, 5, 1–8, <https://doi.org/10.1126/sciadv.aaw5531>, 2019.
- CMEMS: For Baltic Sea Physical Reanalysis Products :, 1–27, <https://doi.org/https://doi.org/10.48670/moi-00013>, 2022.
- Coles, S.: An Introduction to Statistical Modeling of Extreme Values, *Technometrics*, 44, 397–397,
480 <https://doi.org/10.1198/tech.2002.s73>, 2001.
- Couasnon, A., Eilander, D., Muis, S., Veldkamp, T. I. E., Haigh, I. D., Wahl, T., Winsemius, H. C., and Ward, P. J.: Measuring compound flood potential from river discharge and storm surge extremes at the global scale, *Nat. Hazards Earth Syst. Sci.*, 20, 489–504, <https://doi.org/10.5194/nhess-20-489-2020>, 2020.
- Crameri, F., Shephard, G. E., and Heron, P. J.: The misuse of colour in science communication, *Nat. Commun.*, 11, 1–10,
485 <https://doi.org/10.1038/s41467-020-19160-7>, 2020.
- Dacre, H. F., Hawcroft, M. K., Stringer, M. A., and Hodges, K. I.: An extratropical cyclone atlas a tool for illustrating cyclone structure and evolution characteristics, *Bull. Am. Meteorol. Soc.*, 93, 1497–1502, <https://doi.org/10.1175/BAMS-D-11-00164.1>, 2012.
- Donnelly, C., Andersson, J. C. M., and Arheimer, B.: Using flow signatures and catchment similarities to evaluate the E-HYPE
490 multi-basin model across Europe, *Hydrol. Sci. J.*, 61, 255–273, <https://doi.org/10.1080/02626667.2015.1027710>, 2016.
- Dubois, K., Andreas Dahl Larsen, M., Drews, M., Nilsson, E., and Rutgersson, A.: Technical note : Extending sea level time series for extremes analysis with machine learning and neighbouring station data, *Egusph. [preprint]*, 1–14, <https://doi.org/https://doi.org/10.5194/egusphere-2023-1159>, 2023, 2023.
- Eilander, D., Couasnon, A., Ikeuchi, H., Muis, S., Yamazaki, D., Winsemius, H. C., and Ward, P. J.: The effect of surge on
495 riverine flood hazard and impact in deltas globally, *Environ. Res. Lett.*, 15, <https://doi.org/10.1088/1748-9326/ab8ca6>, 2020.
- Ganguli, P. and Merz, B.: Extreme Coastal Water Levels Exacerbate Fluvial Flood Hazards in Northwestern Europe, *Sci. Rep.*, 9, 1–14, <https://doi.org/10.1038/s41598-019-49822-6>, 2019.
- Hao, Z. and Singh, V. P.: Review of dependence modeling in hydrology and water resources, *Prog. Phys. Geogr.*, 40, 549–578, <https://doi.org/10.1177/0309133316632460>, 2016.
- 500 Hieronymus, M. and Kalén, O.: Sea-level rise projections for Sweden based on the new IPCC special report: The ocean and cryosphere in a changing climate, *Ambio*, 49, 1587–1600, <https://doi.org/10.1007/s13280-019-01313-8>, 2020.
- Hieronymus, M. and Kalén, O.: Should Swedish sea level planners worry more about mean sea level rise or sea level extremes?, *Ambio*, 51, 2325–2332, <https://doi.org/10.1007/s13280-022-01748-6>, 2022.
- Hieronymus, M., Hieronymus, J., and Hieronymus, F.: On the application of machine learning techniques to regression
505 problems in sea level studies, *J. Atmos. Ocean. Technol.*, 36, 1889–1902, <https://doi.org/10.1175/JTECH-D-19-0033.1>, 2019.
- Hoskins, B. J. and Hodges, K. I.: New perspectives on the Northern Hemisphere winter storm tracks, *J. Atmos. Sci.*, 59, 1041–1061, [https://doi.org/10.1175/1520-0469\(2002\)059<1041:NPOTNH>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<1041:NPOTNH>2.0.CO;2), 2002.
- Joe, H.: Dependence modeling with copulas, *Depend. Model. with Copulas*, 1–457, <https://doi.org/10.1201/b17116>, 2014.



- Kiesel, J., Lorenz, M., König, M., Gräwe, U., and Vafeidis, A. T.: Regional assessment of extreme sea levels and associated coastal flooding along the German Baltic Sea coast, *Nat. Hazards Earth Syst. Sci.*, 23, 2961–2985, <https://doi.org/10.5194/nhess-23-2961-2023>, 2023.
- Kudryavtseva, N., Soomere, T., and Männikus, R.: Non-stationary analysis of water level extremes in Latvian waters, Baltic Sea, during 1961–2018, *Nat. Hazards Earth Syst. Sci.*, 21, 1279–1296, <https://doi.org/10.5194/nhess-21-1279-2021>, 2021.
- Lindström, G., Pers, C., Rosberg, J., Strömqvist, J., and Arheimer, B.: Development and testing of the HYPE (Hydrological Predictions for the Environment) water quality model for different spatial scales, *Hydrol. Res.*, 41, 295–319, <https://doi.org/10.2166/nh.2010.007>, 2010.
- Meinshausen, N.: Quantile Regression for Left-Truncated Semicompeting Risks Data, *J. Mach. Learn. Res.*, 67, 701–710, <https://doi.org/10.1111/j.1541-0420.2010.01521.x>, 2011.
- Moftakhari, H. R., Salvadori, G., AghaKouchak, A., Sanders, B. F., and Matthew, R. A.: Compounding effects of sea level rise and fluvial flooding, *Proc. Natl. Acad. Sci. U. S. A.*, 114, 9785–9790, <https://doi.org/10.1073/pnas.1620325114>, 2017.
- Moftakhari, H., Schubert, J. E., Aghakouchak, A., Matthew, R. A., and Sanders, B. F.: Linking statistical and hydrodynamic modeling for compound flood hazard assessment in tidal channels and estuaries, *Adv. Water Resour.*, 128, 28–38, <https://doi.org/10.1016/j.advwatres.2019.04.009>, 2019.
- Nerger, L., Hiller, W., and Schröter, J.: A comparison of error subspace Kalman filters, *Tellus A Dyn. Meteorol. Oceanogr.*, 57, 715, <https://doi.org/10.3402/tellusa.v57i5.14732>, 2005.
- Pawlowicz, R.: M_Map: A mapping package for MATLAB, version 1.4m, [Computer software], available online at www.eoas.ubc.ca/~rich/map.html, 2020.
- Pemberton, P., Löptien, U., Hordoir, R., Höglund, A., Schimanke, S., Axell, L., and Haapala, J.: Sea-ice evaluation of NEMO-Nordic 1.0: A NEMO-LIM3.6-based ocean-sea-ice model setup for the North Sea and Baltic Sea, *Geosci. Model Dev.*, 10, 3105–3123, <https://doi.org/10.5194/gmd-10-3105-2017>, 2017.
- Rutgersson, A., Kjellström, E., Haapala, J., Stendel, M., Danilovich, I., Drews, M., Jylhä, K., Kujala, P., Guo Larsén, X., Halsnæs, K., Lehtonen, I., Luomaranta, A., Nilsson, E., Olsson, T., Särkkä, J., Tuomi, L., and Wasmund, N.: Natural Hazards and Extreme Events in the Baltic Sea region, *Earth Syst. Dyn. Discuss.*, 1–80, <https://doi.org/10.5194/esd-2021-13>, 2021.
- Sadegh, M., Moftakhari, H., Gupta, H. V., Ragno, E., Mazdidasni, O., Sanders, B., Matthew, R., and AghaKouchak, A.: Multihazard Scenarios for Analysis of Compound Extreme Events, *Geophys. Res. Lett.*, 45, 5470–5480, <https://doi.org/10.1029/2018GL077317>, 2018.
- Sadegh, M., Ragno, E., and Aghakouchak, A.: Multivariate Copula Analysis Toolbox (MvCAT): Describing dependence and underlying uncertainty using a Bayesian framework, *Eos, Trans. Am. Geophys. Union*, 64, 929, <https://doi.org/10.1029/eo064i046p00929-04>, 2017.
- Salvadori, G., Durante, F., Michele, C. De, Bernardi, M., and Petrella, L.: A multivariate copula-based framework for dealing with hazard scenarios and failure probabilities, *J. Am. Water Resour. Res.*, 52, 2–2, <https://doi.org/10.1111/j.1752-1688.1969.tb04897.x>, 2016.



- Sentinel-2 cloudless by EOX IT Services GmbH is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. The required attribution including the given links is "Sentinel-2 cloudless -
545 <https://s2maps.eu> by EOX IT Services GmbH (Contains modified Copernicus Sentinel data 2021)". 2021.
- Serinaldi, F.: Dismissing return periods!, *Stoch. Environ. Res. Risk Assess.*, 29, 1179–1189, <https://doi.org/10.1007/s00477-014-0916-1>, 2015.
- Strömqvist, J., Arheimer, B., Dahné, J., Donnelly, C., and Lindström, G.: Water and nutrient predictions in ungauged basins: set-up and evaluation of a model at the national scale, *Hydrol. Sci. J.*, 57, 229–247,
550 <https://doi.org/10.1080/02626667.2011.637497>, 2012.
- Tootoonchi, F., Sadegh, M., Haerter, J. O., Rätty, O., Grabs, T., and Teutschbein, C.: Copulas for hydroclimatic analysis: A practice-oriented overview, *Wiley Interdiscip. Rev. Water*, 9, 1–28, <https://doi.org/10.1002/wat2.1579>, 2022.
- Wahl, T., Jain, S., Bender, J., Meyers, S. D., and Luther, M. E.: Increasing risk of compound flooding from storm surge and rainfall for major US cities, *Nat. Clim. Chang.*, 5, 1093–1097, <https://doi.org/10.1038/nclimate2736>, 2015.
- 555 Wolski, T., Wiśniewski, B., Giza, A., Kowalewska-Kalkowska, H., Boman, H., Grabbi-Kaiv, S., Hammarklint, T., Holfort, J., and Lydeikaite, Ž.: Extreme sea levels at selected stations on the Baltic Sea coast, *Oceanologia*, 56, 259–290, <https://doi.org/10.5697/oc.56-2.259>, 2014.
- Zscheischler, J., Martius, O., Westra, S., Bevacqua, E., Raymond, C., Horton, R. M., van den Hurk, B., AghaKouchak, A., Jézéquel, A., Mahecha, M. D., Maraun, D., Ramos, A. M., Ridder, N. N., Thiery, W., and Vignotto, E.: A typology of
560 compound weather and climate events, *Nat. Rev. Earth Environ.*, 1, 333–347, <https://doi.org/10.1038/s43017-020-0060-z>, 2020.