

Assessing future impacts of tropical cyclones on global banana production

Sophie Kaashoek¹, Žiga Malek^{2,3,1}, Nadia Bloemendaal^{1,4}, Marleen C. de Ruiter¹,

¹Institute for Environmental Studies, Vrije Universiteit Amsterdam, De Boelelaan 1111, 1081HV, Amsterdam, the Netherlands

² Biotechnical faculty, University of Ljubljana, Jamnikarjeva 101, 1000 Ljubljana, Slovenia

³ Novel Data Ecosystems for Sustainability (NODES), International Institute for Applied Systems Analysis (IIASA), Schlossplatz 1, 2361 Laxenburg, Austria

⁴ Royal Netherlands Meteorological Institute (KNMI), Utrechtseweg 297, 3731 GA De Bilt, The Netherlands

Correspondence to: Marleen de Ruiter (m.c.de.ruiter@vu.nl)

Abstract. Tropical cyclones (TCs) are projected to increase in intensity globally, impacting human lives, infrastructure, and important agricultural activities, such as banana production. Banana production is already impacted by TCs in several parts of the world, leading to price volatility and impacted livelihoods of banana producers. While many potential impacts on banana production have already been quantified on a local scale, it remains unclear how bananas could be impacted by TCs across the globe under future climate conditions. To address this, we first looked at the documented impacts of TCs on banana production from different places all around the world. Using spatially explicit data on banana producing regions and future TC occurrence and magnitude, we then identified the spatial distribution and extent of areas where TCs could impact banana production. Our results suggest that considerable portions of global banana production are at risk to be impacted by TCs under future climate conditions, and we show this for different return periods.

Globally, 24.3% of all banana-producing areas are projected to suffer major or complete (>84%) damage under current climate conditions, increasing to 26.5% under future climate scenarios at the 100-year wind speed return period. Additionally, we estimate that 30.1% of global production under current conditions, and 31.1% under future conditions will be majorly or completely damaged at the 100-year return period. The regions predominantly affected in the future are Southeast Asia, China, India and the Caribbean, potentially experiencing substantial disruption in banana production. Our results therefore indicate that considerable efforts in climate adaptation are essential to ensure the stability of global banana supply chains.

1. Introduction

Currently, average global surface temperatures have already increased between 1.1 and 1.2 degrees Celsius above pre-industrial levels (IPCC, 2023). These increasing temperatures can trigger fast changing weather patterns potentially causing more frequent and intense extremes affecting water and food security with high confidence (IPCC, 2021). Particularly tropical cyclones (TCs) are projected to increase in intensity (Knutson et al., 2020). This is predominantly driven by increasing sea-surface temperatures (SSTs), which serve as fuel for an intensifying TC. This is an alarming projection, since TCs can

substantially impact coastal communities, causing potentially large agricultural and financial losses and human casualties. One of the agricultural activities with a high socio-economic significance that is under threat of TCs, is banana production. Globally, bananas are the most consumed fruit (FAO, 2022). In countries all over the world bananas provide a large part of nutritional diversity in the diet of the average consumer. The wide-spread benefits of bananas include its nutritious values; they are rich in calories, are perfectly suitable for dietary diversity and they contain a variety of vitamins & fibers (Petsakos, 2019). A sustainable production of the fruit poses an important source for nutrition to maintain food security and sustain income (Petsakos et al., 2019, Varma & Bebbber, 2019). In Latin American, the Caribbean and Asian countries, bananas remain a significant contributor to the economy and provide a livelihood for millions (FAO, 2019).

To many rural households, bananas generate an important part of their income (Mohan, 2017). In addition, bananas, together with plantain, are an important source of nutrition for millions of people. A reduced availability of bananas leads to higher prices for consumers and can cause a sudden and drastic decrease in income and economic prosperity of tens of thousands of producers in exporting countries (FAO, 2022). Due to our globalized economic system and its strong interdependencies, the potential negative impact on the production system will not only be harmful on a local scale, but can escalate to the global scale (Lesk et al., 2016). Consequently, the partial or full destruction of banana plantations across larger areas would have a large influence on the currently stable and reliable global value chain of bananas (Varma, et al., 2020). In addition, a disruption in banana production will negatively impact the United Nation's Sustainable Development Goals, including the first (no poverty), second (zero hunger) and third (good health and well-being) goals (United Nations, 2015). Banana production is already expected to be more impacted by numerous extreme weather events in the coming decades such as drought, heavy precipitation, and heatwaves (Malek et al. 2022). Yet so far, studies on the impact of extreme weather on agricultural areas have mainly focused on the influence of extreme heat or precipitation deficit on production, thereby lacking a perspective on high- impact low-probability events like TCs (Calberto et al., 2015, Lesk et al., 2016, Varma & Bebbber, 2019, Malek et al, 2022). This is remarkable, since the direct impact of TCs on the agricultural sector can be substantial (FAO, 2015, Kunze, 2021). Research has shown that there is a clear negative direct impact of TCs on banana producing areas (Huigen & Jens, 2006, Robinson & Saúco, 2010, Mohan, 2017). TCs can completely disrupt banana cultivation and it can take up to a year to recover depending on farmers' access to facilities and finance (Mamuye, 2016). For example, the 2021 Hurricanes Eta and Iota caused a 51% decrease of banana yields in Honduras (FAO, 2022. P.2). Due to production disruption, there is less supply increasing the average price of the fruit (Beer et al., 2014). However, due to a lack of available data on TCs, the spatial impact of this hazard for bananas can not be identified on a global scale (Malek et al., 2022). Several indirect risk assessments have been performed after the occurrence of a TC event on macro- or meso-scale (Huigen et al., 2006, Beer et al., 2014, Mohan, 2017). It is expected that in the coming decades banana producers in Southeast Asia, Oceania and the Caribbean will likely be impacted even more by TCs, but there is no evidence of where impacts are expected to occur on a supra-national scale (, Varma & Bebbber, 2019, Malek et al., 2022). This is problematic, as banana cultivators have no reliable projections of the probability to be affected by a TC. When banana producers are provided with better estimates of the likelihood of TC impacts, they will be able to implement adaptation strategies, hereby increasing their resilience (Chavez et al., 2015).

In this paper, we therefore assess how future TCs could impact banana production on a global scale. We first synthesize the documented impacts of TCs on banana production, by reviewing recent studies. Using data on the global distribution of current and future TCs, we identify where banana producing areas will be exposed to TCs. Finally, we quantify and map the extent of banana production that will experience more frequent and damaging TCs. This way, we contribute to the ongoing efforts to better understand the potential impacts of TCs on agriculture.

2. Methods

2.1 Methods overview

We perform a risk assessment by mapping the global TC risk to banana producing areas under baseline (1980-2017) and future (2015–2050 under SSP585 forcing scenario) climate conditions. We follow the risk framework of the UNDRR (2017) that is based on the understanding that risk is the product of hazard (Section 2.2), exposure (Section 2.3), and vulnerability (Section 2.4). We refer to hazard as the probability of occurrence of a certain TC wind speed, exposure as the banana producing area at risk, and vulnerability as the fraction of banana plants that will be damaged by a natural hazard (UNDRR, 2017).

2.2 Tropical cyclone hazard mapping

To represent the hazard component of our risk framework, we use the Synthetic Tropical cyclOne geneRation Model (STORM) baseline and near-future climate TC wind speed return period (RP) maps (Bloemendaal et al., 2020b, Bloemendaal et al., 2022). STORM is the first model that captures future global-scale TC wind speed probability trends derived at high (10 km) spatial resolution. STORM is created using a synthetic, fully statistical modelling approach that simulates new TCs by using statistical information on TC characteristics (frequency, track, and intensity) derived from historical observations. STORM then simulates 10,000 years of synthetic TCs based on baseline climate conditions (representing 1980-2017), thereby providing the necessary data for the calculation of wind speed RPs (in m/s). To analyze changes in TC risk under climate change, we use the STORM future climate dataset (Bloemendaal et al., 2022). This dataset extracts future-climate TC statistics using four high resolution GCM climate models based on baseline and Shared Socioeconomic Pathway 585(SSP) 2015-2050 climate conditions and the STORM algorithm. SSP585 is a high-emissions scenario and assumes that there will be no mitigation policies on greenhouse gas emissions, leading to high radiative forcing (O'Neill et al., 2016). As there is little deviation in terms of TC activity between lower-forcing scenarios and the high emission SSP585 scenario (IPCC, 2021, Bloemendaal et al., 2022). The STORM datasets are generated on a global scale, with the datasets split up over six basins (Eastern Pacific, North Atlantic, North Indian, South Indian, South Pacific, Western Pacific). The STORM present and future wind speed RP datasets provide 10-meter 10-minute maximum wind speeds at various RP levels and at 10 km x 10 km grid cell resolution. From these datasets, we extract the 10-, 100-, and 200-yr wind speed RP maps, to account for both events with higher

95 probabilities but lower intensity, and events with lower probabilities but higher intensities. Note that a high RP corresponds to a low probability of occurrence and thereby a higher risk probability (Ward et al., 2011, Bevacqua et al., 2019).

2.3 Exposure

2.3.1 Banana production and impacts of tropical cyclones

100 In 2023, global banana export volume was 19.2 million tonnes, of which Latin America & Caribbean exported 15 million tonnes of bananas (FAO, 2023). The five largest exporting countries are Ecuador, the Philippines, Costa Rica, Guatemala, Colombia, and the Dominican Republic (FAO, 2023). The most consumed and traded bananas can be divided in two large groups; the ‘dessert’ banana types which are sweet and can be consumed raw or ‘cooking’ banana types who need to be treated before use (FAO, 2019). Cooking bananas are mainly found in Asia and East Africa. The dessert banana is concentrated in Latin America and Asia, including the Cavendish type also in the Caribbean. The harvest of the plantain is found primarily in
105 Africa, Latin America and Asia. The estimated size of the global total banana plantation is 5.6 million ha (Machovina & Feeley, 2013, Coltro & Karaski, 2019, FAO, 2019). In this study, we assume all banana types exhibit the same vulnerability to TC wind speeds. A distinction will not be drawn between the types, as the scope of this study is primarily centered on highly destructive events such a TCs, wherein the resultant damage is anticipated to be equivalent across both types. Furthermore, the absence of systemic data necessitates an equal consideration of cultivator types in evaluating the impacts of such destructive
110 events.

TCs can cause significant damage to banana producing areas in different ways (Table 1). Despite the diverse impacts on banana production, TCs particularly lead to loss through wind speeds, and less so by amount of rainfall (Lin et al., 2020). This is mostly because the fruit easily blows off the plant by strong winds and directly creates crop losses. Humid winds can have a
115 positive influence by helping to pollinate banana flowers, disperse pests and beneficial insects. On the negative side, strong winds can cause physical damage to the fruit itself, either destroying them, or reducing the quality of production, potentially limiting their direct human consumption (Robinson & Saúco, 2010).

Table 1: Types of damage from wind to banana

Type of damage	Description	Source
Recovery period	The harvest period of bananas takes approximately a year, so when banana plantations are damaged the recovery period can take up to 9-12 months	Mamuye, 2016

Height	Due to height of the plant more susceptible to TC compared to other crops	Huigen et al., (2006), Robinson & Saúco, (2010)
Leaves changes production capacity	The producing capacity of the banana is reduced when leaves are damaged.	Mohan (2017)
Productivity	The wind can also cause the fruit to fall off the plant prematurely or affect the productivity of the herb by influencing the layer between the undisturbed air adjacent to the surface and temperature of the leave	Robinson& Saúco (2010)
Low wind speed	Even wind at lower speeds (10 m/s) can cause damage to banana plants, particularly if the plants are already weakened or stressed for other reasons such as pests and diseases. In some cases, this weaker wind may be strong enough to uproot the plants or cause significant damage to the leaves and stems	Mohan (2017)
Dry wind	Dry winds can lead to water shortage impacting growth.	Robinson& Saúco (2010)
Seasonal winds	Strong seasonal winds can cause leave tearing, and slower winds can increase leaf and dust abrasion.	Robinson& Saúco (2010)
High winds	In addition to the physical damage to the plants and fruit, high winds can also cause other issues that can impact banana production. For example, the winds can blow debris and sand around, which can make it more difficult for the plants to grow properly. The winds can also create dust storms, which can reduce visibility and make it harder for farmers to work in the fields	El-Kady (2012)

120

2.3.2 Mapping banana producing areas

To map the banana producing areas susceptible to a TC, we used MAPSPAM data on global spatial distribution for major crops, which includes bananas (IFPRI 2019). These data present recent systematically developed data on banana distribution, by combining a set of statistical and remote sensing data. The spatial resolution of the data is roughly 10x10 km at the equator.

125

To exclude outlier areas in countries, where bananas are generally not produced or are produced in protected environments such as greenhouses (such as the European Union), we applied a threshold of 5 ha within the 10x10km grid cell. Grid cells are

considered relevant only when containing banana-producing areas that surpass the specified 5-hectare threshold (and where banana cultivation covers more than 0.05% of the total grid). We then looked at banana production in tons in these areas, using the same data, as MAPSPAM not only provides data on the spatial distribution of harvested areas, but also production.

2.4 Vulnerability

2.4.1 Vulnerability curve

Vulnerability ascribes the characteristics of the exposed asset that make it susceptible to the effects of a certain hazard (here a TC) (UNDRR, 2017). The vulnerability curve assesses the damage state against the hazard intensity showing the wind speed (m/s) interval and the mean damage ratio (MDR) (Mo, et al., 2016, Yum et al., 2021). Vulnerability curves can be based on empirical, analytical, and expert opinion or judgement methods (Merz et al., 2010). Yet so far, most academically developed vulnerability curves for TCs quantify wind speed damage focusing on the built environment (Sun et al., 2021, Yum et al., 2021). To the best of our knowledge, there is currently no evidence on vulnerability curves of wind-induced damage for bananas. To quantify the level of damage to banana plantations caused by a certain wind speed, we therefore have to develop vulnerability curves. This is done by executing a systemic review of reported past TC events damaging banana plantations. To construct the curves, we use an approach similar to the research by Yum et al. (2021) where the authors use existing white and grey literature reporting on historic TC events, their maximum wind speed and damages. While we acknowledge that TC impacts are often caused by an interplay of wind, precipitation and storm surge, we use wind speed as a proxy for all three hazards as information on the latter two hazards is often lacking or incomplete (see for instance Eberenz et al., (2019) and Yum et al., (2021) for a similar approach for building vulnerability curves). Such an approach based on empirical data allows for a higher accuracy of the damage curve (Merz et al., 2010). To retrieve reports of past events of damaged banana plantation by wind speed, peer-reviewed articles, and news reports about destroyed banana plantations by TC wind were explored by Google Search and the Google Scholar and WorldCat academic databases (published anytime using the following search terms: bananas + storms, hurricanes, (tropical) cyclones, typhoons; and we excluded studies that did not include wind damages. The events were categorized into the time, location, windspeed and % of destroyed plantation. We identified 22 cases where impacts of TC on banana production were quantified from all around the globe (see S1). The wind speeds near the location that were mentioned in the events were taken. When an event had no mention of maximum wind speeds, we used the historical TC database of NOAA (2024) and retrieved the wind speeds and trajectories for the affected region. The wind speed closest to the affected region was taken. This way we were able to identify the wind speed and trajectory of the TC at all locations. The 22 cases were categorized by time, location, windspeed and % of destroyed plantation.

2.4.2 Mean Damage Ratio

A damage function is a common tool used to present the vulnerability of assets, in this case banana producing areas, to hazard, in this case a TC (De Ruiter et al., 2017). All 22 events found were used to calculate the damage ratio of the TC-induced

damaged banana plantation against the corresponding wind speed intensity, similar to the method discussed by Yum et al. (2021). The damage ratio indicates the likelihood of a banana plant surpassing a certain damage level threshold (v_{min}) as the windspeed increases up to a certain critical point. Above the threshold, the exceedance probability increases before it reaches a 100% destroyed plantation (v_{max}), showing the plantation is fully destroyed (Yum et al., 2021). Yum et al. (2021) quantify the damage ratio based on insurance data (using monetary values). In our study, we only assess the physical vulnerability due to the TC wind speed. The equation used is described in Eq. (1):

165

$Damage\ ratio = \frac{X_{destroyed}}{X_{Total}}$	(1)
---	-----

Where $X_{destroyed}$ represents the destroyed plantation area (in ha), and X_{total} represents the total plantation area (in ha). From Equation (1), it follows that higher damage ratios relate to higher vulnerability. The formula is used for the events lacking a damage ratio. To quantify the damage ratio, the spatial destruction (ha destroyed) and the entire scope (ha plantation) needs to be identified. Some events referred to banana plantation losses based on total banana production area in a country. Given that countries have a significant difference in size and spatial distribution of production areas, it is necessary to assess the exact location where an individual TC struck the plantation. Hence, to obtain a robust and reliable database, events are removed when the location could not be determined. 22 damage ratios ($d_1, d_2, \dots, d_{22}$) were identified with corresponding $MDR = \left(\frac{dx + dx + \dots, dx}{Total\ dx} \right) \cdot 100$ wind speeds ($v_1, v_2, \dots, v_{22}$) (see S1). The datasets were categorized from lowest to highest wind speed and grouped into three wind-speed intervals, namely: 0-25, 26-43, and above 44 (see Table 2). The wind speed intervals were established based on expert judgement (see Table 1) and the Saffir-Simpson (1974) scale (see Table 2). According to Robinson (2010) and Mohan (2017), evidence of damage to banana plantations was already observed at windspeeds of 10 m/s, hence why we set the lower limit of the curve (V_{min}) at 10 m/s. The resulting vulnerability curve is shown in Figure 1. Similar to other studies (e.g. Yum et al 2021 and Shinozuka et al., 2000), the curve has been smoothed.

180

Besides deriving the vulnerability curve, we can also use our database to define three different types of damage. Based on our database we define three damage categories: moderate damage to a banana plantation occurs up to 25 m/s, between 26 and 43 m/s they experience major damage, and above 44 m/s they are completely damaged. Next, we calculate the mean damage ratio (MDR) for each of these intervals. As such, MDR1 corresponds to the MDR over the moderate (first) damage interval, MDR2 corresponds to the MDR for the major (second) damage interval, and MDR3 corresponds to the complete (third) damage interval (see Table 2). The MDR is calculated using damage ratio (d) and are defined in Eq. (2)


185

$MDR = \left(\frac{(dx + dx + \dots, dx)}{Total\ dx} \right) \cdot 100$	(2)
--	-----

Table 2: Mean Damage Ratio per wind speed interval with corresponding damage level

Wind speed interval (10- min maximum average sustained windspeed in m/s)	Average damage	Damage level
0-10	0%	No Damage
10-25	MDR1: 48%	Moderate damage
26-43	MDR2: 84%	Major damage
44>	MDR3: 96%	Complete destruction

195 **Figure 1: Constructed vulnerability curve of tropical cyclone wind damages to bananas based on a meta-analysis of 22 studies, similar to Yum et al., (2021).**



200


3. Results

3.1 Banana producing areas

We observe an increase in the share of areas at risk and a corresponding increase in damage levels for the three distinct return
205 periods: 10 (RP10), 100 (RP100) and 200-year RP period (RP200). Globally, at RP10, 16.5 % of all bananas producing areas
will potentially be moderately or majorly damaged under baseline climate conditions. At the RP100, 34.5 % will be moderately,
majorly, or completely damaged and 39.1% at RP200. Furthermore, it is evident that there will be a rise in the extent of areas
susceptible to damage from future TCs under the SSP585 scenario (Table 4). At RP10, 11% of banana areas globally will be
subject to increase moderately, and 9.5% to major destruction in the future, presenting a 8% and 51% increase compared to
210 the baseline climate respectively (Tables 3 and 4, Figure 2). In the SSP585 at the RP100 21.7% of areas will be subject to
major damages, and 4.8% to complete destruction, presenting a 3% and 50% increase compared to baseline climate conditions
(Tables 3 and 4, Figure 2).

215

Figure 2: Global distribution of impacted banana producing areas across different damage levels under various return periods for both baseline(1980-2017) and future climates (2015-2050). High resolution images for individual return periods are provided in S4.



235 **Table 3: Global distribution of impacted banana producing areas across different damage levels under various return periods for both baseline (1980-2017) and future climates (2015-2050) (in %).**

	Baseline			Future		
	RP10	RP100	RP200	RP10	RP100	RP200
Moderate	10.2	10.2	10.5	11.0	12.2	13.0
Major	6.3	21.1	22.3	9.5	21.7	23.1
Complete	0.0	3.2	6.3	0.0	4.8	7.5

240 **Table 4: Increase of damaged banana producing area on a worldwide basis comparing baseline (1980-2017) and future climates (2015-2050) across different damage levels under various return periods (in %).**

	RP10	RP100	RP200
Moderate	7.7	19.8	23.4
Major	50.6	2.8	3.6
Complete		50.1	18.6

There are large disparities across the globe when it comes to banana producing areas being affected by TCs (Table 5). At RP100, in the Caribbean, 56.9% is completely damaged, while in South Asia, this value amounts to 0.5% under baseline climate conditions. The strongest changes in TC impacts under future warming are found in Southeast Asia (+22% (major) and 36% (complete) relative increase compared to the baseline climate), China (-7.6% decrease in major, and 289% increase in complete), India (4% decrease in major and no complete damage in the future) and the Caribbean (9.4% increase in major and a 6.5% decrease in complete damage). South America, West and South Africa, will experience minimal impact as TCs are rarely observed in these regions (Table 5).

250 **Table 5: Impacted banana producing area in different regions across major and complete destructed levels under the RP100 for both baseline (1980-2017) and future climates (2015-2050) (in %).** This table focuses on selected banana regions. However, not all regions are affected by tropical cyclones. The presented numbers in the table represent shares or proportions rather than absolute values. To provide a more comprehensive understanding, absolute numbers in hectares can be found in S2 as well as the % damaged banana producing area per countries.

Region	Major		Complete	
	Baseline	Future	Baseline	Future
Caribbean	40.5	44.3	56.9	53.2

Central America	38.0	24.4	0.0	2.9
South America	1.8	1.9	0.0	0.0
West Africa	0.3	0.3	0.0	0.0
East Africa	29.5	30.4	0.0	3.4
South Africa	0.0	0.9	0.0	0.0
Middle East and North Africa	12.3	16.7	0.0	0.0
East Asia	66.8	61.7	2.7	10.7
South Asia	39.6	38.1	0.5	0.0
Southeast Asia	24.5	29.9	8.0	10.9
Oceania	21.0	21.2	0.0	5.2

3.2 Banana production

There is an increase in impacted banana production under the SSP585 scenario for the time period 2015-2050 . We observe an increased share of damaged areas and an increase in damage level for the three different RPs, namely RP10, RP100 and RP200. At RP200 the global share of major damages to banana production increases by 6.1% of total global production, but the total amount of banana production completely destroyed decreases by 1.9% to 5.1% of total global production.

Table 6: Global distribution of impacted banana production across different damage levels under various return periods for both baseline (1980-2017) and future climates (2015-2050) (in %).

	Present			Future		
	RP10	RP100	RP200	RP10	RP100	RP200
Moderate	8.5	14.9	12.5	10.2	16.8	11.7
Major	7.8	27.1	32.9	10.1	27.2	34.9
Complete	0.0	3.0	5.2	0.0	4.0	5.1

Our results show that the regions predominantly affected by TCs are Asia and the Caribbean, experiencing substantial disruption in banana production. Banana production will be subject to complete damage in many parts of the Caribbean, totally amounting to 58.4% of production under the future climate. While this number represents a decrease compared to the baseline climate (68.1% complete damage), our results indicate that the Caribbean will still be most impacted across all regions. East and Southeast Asia, hosting considerable shares of global banana production, are projected to experience 13.3% and 8.8%

complete damage towards the future, respectively. Despite other areas experiencing minor shares of production being impacted by complete damage, many areas are projected to experience major damages from more than 20% of banana production under the future climate: East, South and Southeast Asia, Caribbean and Oceania (Table 7). Such large shares indicate that the stability of the current banana supply chains is under threat, without considerable efforts in climate change adaptation.

Table 7: Impacted banana production in different regions across major and complete destructed levels under the 100RP for both baseline (1980-2017) and future climates (2015-2050) (in %). This table focuses on selected banana-producing regions and acknowledges that not all regions are affected by tropical cyclones. The presented numbers in the table represent shares or proportions rather than absolute values. To provide a more comprehensive understanding, absolute numbers in tons can be found in S3 as well as the % damaged banana producing area per countries.

Region	Major		Complete	
	Baseline	Future	Baseline	Future
Caribbean	19.5	29.2	68.1	58.4
Central America	12.4	8.9	0.0	1.3
South America	1.0	1.0	0.0	0.0
West Africa	0.2	0.2	0.0	0.0
East Africa	3.2	2.9	0.0	0.8
South Africa	0.0	0.2	0.0	0.0
Middle East and North Africa	10.0	15.5	0.0	0.0
East Asia	75.5	66.1	3.0	13.3
South Asia	42.0	42.3	0.7	0.0
Southeast Asia	38.6	43.4	7.9	8.8
Oceania	19.1	20.5	0.0	0.9

Notably, under RP100 and the RP200, a substantial increase is observed in class 3, while significant increments are evident in class 1 and class 2 for RP10 (Table 6). Conversely, a decline in production is seen for RP200. These changes can be attributed to the intensification of TCs in a warming climate, leading to alterations in their trajectories and a higher likelihood of targeting different areas for impact.

Table 6: Increase of globally damaged banana production between baseline (1980-2017) and future (2015-2050) climate conditions (in %) across different damage levels and return periods

	RP10	RP100	RP200
Moderate	20.2	12.7	-6.4

Major	30.2	0.4	6.1
Complete	0.0	30.1	-3.5

Interestingly, we observe a slight decrease in damage under RP100 and R200 for the moderate and complete damage classes in some regions (Table 6). This is predominantly due to a decrease in damage in south Asia (India), which, in turn, is driven by a decrease in TC wind speeds at the RP200. This decrease is caused by a projected landward shift in the TC genesis region in the Bay of Bengal. Consequently, TCs are projected to form closer to the coast, meaning there is less time for a TC to intensify to higher wind speeds before it makes landfall and weakens again (see also Figure 4 in Bloemendaal et al 2022). This results in a decrease in wind speed at higher RPs, leading to lower banana plantation damages in this region.

4. Discussion

4.1 Hazard modelling and impacts on banana production

This study presents the first attempt to map the spatial extent of potential current and future tropical cyclone impacts on banana areas and production globally. However, it potentially underestimates impacts on banana production, by not considering indirect impacts of TCs on banana production. Waterlogging could lead to root rot, and excessive water could lead to favorable conditions for pests and diseases which are a major limitation to banana production (Aguilar et al., 2003). However, whether TCs would lead to such indirect impacts is very context specific, meaning that even within the same grid cell of our study we can expect different indirect impacts, depending on the slope, soil, terrain orientation, previously drained wetlands, to name a few. To account for such indirect impacts, both more detailed TC data, as well as data on soil drainage are necessary. In addition, TCs can impact electricity, refrigeration and road infrastructure, furthermore leading to disruptions in the supply chains and potential inability to harvest and store the crops (Koks et al., 2019).

The STORM future climate datasets were generated based on the high-end SSP585 scenario. While one can discuss the likelihood of this emission scenario, as current developments are steering away from this scenario, the average climate conditions over the 2015 – 2050 time period do not differ substantially between the different forcing scenarios. We therefore believe that the SSP585 input dataset as was used for Bloemendaal et al., (2022) can be seen as a good proxy for changes in TC characteristics over the aforementioned time period. While this study does not consider TC impacts beyond 2050, we alert readers that the average climate conditions past 2050 do start to deviate and that the approach as used in Bloemendaal et al., (2022) does not hold then.

There are also limitations in the STORM model itself that could affect the outcomes of this study. First of all, TC intensity in STORM is solely modelled as having a direct relationship with sea-surface temperatures. In reality TC intensity is also largely influenced by vertical wind shear; an effect absent in STORM. While in general STORM validates well against observations

325 (Bloemendaal et al., 2020c), there are regions where vertical wind shear in reality plays a critical role in governing TC intensity. One of these regions is the Bay of Bengal; for this region, TC intensity and associated RPs tend to be over- and underestimated, respectively. As a consequence, our impact assessment for this region can be overestimated. Secondly, STORM uses the parametric wind field model from Holland (1980) to translate point data to a 2-dimensional wind field. This model assumes asymmetry in the wind field to arise from background flow; in reality, these asymmetries can also be induced by enhanced
330 wind shear or interaction with land. This may result in slightly altered wind speed RPs. Lastly, TC decay after landfall is modeled through an empirical inland decay function (Kaplan and DeMaria, 1995). This decay function was derived based on USA landfalls, and hence may perform less well elsewhere.

Lastly, the authors acknowledge that the referred STORM wind module does not take elevation effects into account. This means that TCs windspeeds can be over- or underestimated over land, as wind speeds can increase or decrease depending on
335 the orientation of for instance a mountain ridge (Bloemendaal et al., 2020a). Second, STORM uses a lower bound at 18m/s wind speed (Bloemendaal et al., 2022). However, it was found that banana plants can already be damaged by speeds above 10 m/s, meaning we cannot assess damages up to 18 m/s wind speed. Our results therefore potentially underestimate damages to banana production. Third, STORM models TCs on a basin scale, meaning that there is no transition of TCs across basins. This can be problematic in regions that can be affected by TCs originating from multiple basins, such as Central America. Lastly,
340 TCs can also cause damage to banana plantations through excessive rainfall, runoff and landslides. These factors are, however, not modelled in STORM. Inclusion of such factors can potentially alter the risk estimates that were presented in this study.

4.3 Vulnerability of banana production systems

We used an empirical approach to construct the vulnerability curve using a meta-analysis of 22 studies, to increase accuracy of the curve. However, limited data availability can impact the results, and in turn the results can be difficult to transfer in
345 space and time.

Damage assessments rely on numerous assumptions and require more thorough validation, According to Merz, et al., (2010) hazard evaluation often overshadows damage assessment. (). Consequently, many damage levels can be considered inconsistent and subject to bias. The vulnerability curve that we developed is based on 22 TC events damaging banana plantations and expert knowledge. When no wind speed was named in the article, we used the NOAA database. In two events,
350 the trajectory did not exactly cross the affected location. In this case, we took the closest measurement. We hereby acknowledge that this is an overestimate for those affected hurricanes. Furthermore, the vulnerability curve is based on the understanding that all banana producing areas have the same features irrespective of the location or country's political or economic situation. We did not include assumptions on the type of banana, or plantation characteristics such as soil. slope and elevation. We acknowledge that this influences the damage level, and suggest to use this components for future studies (??). Furthermore,
355 the type of the plantation identified in the grid cell can vary (Coltro & Karaski, 2019). Bananas that are produced in a mixed crop system have a larger resilience to TC damage, than those produced on a monoculture farm and should therefore be evaluated separately (Huigen et al., 2006, Robinson & Saúco, 2010, Mohan, 2013). Next, it should be noted that because the

constructed vulnerability curve was based on a meta-analysis of reported damages in academic and grey literature, there can be a bias towards the more damaging events as these could have been covered more in literature.

360 We only included physical damage parameters in the development of the vulnerability curve, meaning that future research should also focus on socio-economic and cultural aspects of banana producers. To achieve this, future research should focus on local scales and thereby increasing the accuracy of the vulnerability curves by including more input parameters, among other local terrain and soil characteristics, production system, and implemented adaptation. Such case-studies can consider the climatic conditions and reliance on banana production at the specific location, as well as socio-economic situation of the countries involved. For instance, t. . Lastly, future studies could also assess low probability events as they often show higher damage levels. Even though we already indicate high damage levels under high probability RPs, the effects of the low probabilities will be even larger and should be identified (Ward et al., 2011).

5. Conclusion

370 Identifying the locations and spatial extent of potential impacts of tropical cyclones on banana production is necessary for adaptation , and more resilient banana supply chains. We find that the banana producing areas are at greater risk to be affected by future TCs, which is especially valid for some of the major banana producing regions (Caribbean, South, East and Southeast Asia). We identified the most impacted banana production per TC-prone region for the baseline and for future climate at different return period levels. Our findings indicate that significant portions of global banana production are at risk from tropical cyclones (TCs) under future climate conditions, and we demonstrate this for various return periods. Globally, 24.3% of all banana-producing areas are projected to suffer majorly or completely (>84%) damage under current climate conditions, increasing to 26.5% under future climate scenarios at the 100-year wind speed return period. Additionally, we project that 30.1% of global production under current conditions, and 31.1% under future conditions will be majorly or completely damaged at the 100-year return period. The regions that are most likely to be affected in the future include Southeast Asia, China, India, and the Caribbean, with substantial disruptions in banana production expected. Consequently, our results underscore the urgent need for significant climate adaptation efforts to maintain the stability of global banana supply chains.

Our results can support the identification of vulnerable areas and can therefore be used as a starting point to target and implement adaptation measures, such as awareness raising efforts and early warning systems. This will hopefully contribute to a reduction in the impacts of future TCs to banana plantations.

385

Competing interests

The contact author has declared that none of the authors has any competing interests.

Acknowledgements

390 MCdR received support from the MYRIAD-EU project, which received funding from the European Union's Horizon
2020 research and innovation programme under grant agreement No [101003276](#), and by the Netherlands Organisation for
Scientific Research (NWO) (VENI; grant no. VI.Veni.222.169).

395

References

- 400 Aguilar, E.A., Turner, D.W., Gibbs, D.J., Armstrong, W., Sivasithamparam, K., 2003. Oxygen distribution and movement,
respiration and nutrient loading in banana roots (*Musa* spp. L.) subjected to aerated and oxygen-depleted environments. *Plant*
Soil 253, 9
- Beer, T., Abbs, D., & Alves, O. (2014). Concatenated hazards: tsunamis, climate change, tropical cyclones and floods. In
Tsunami Events and Lessons Learned (pp. 255-270). Springer, Dordrecht.
- 405 Bevacqua, E., Maraun, D., Voudoukas, M. I., Voukouvalas, E., Vrac, M., Mentaschi, L., & Widmann, M. (2019). Higher
probability of compound flooding from precipitation and storm surge in Europe under anthropogenic climate change. *Science*
advances, 5(9), eaaw5531.
- 410 Bloemendaal, N., De Moel, H., Muis, S., Haigh, I. D., & Aerts, J. C. (2020a). Estimation of global tropical cyclone wind speed
probabilities using the STORM dataset. *Scientific data*, 7(1), 1-11.
- Bloemendaal, N. De Moel, H. Muis, S. Haigh, I. D. Aerts, J. C. J. H (2020b). STORM tropical cyclone wind speed return
periods. ResearchData. [Data set] Retrieved from; <https://doi.org/10.4121/12705164.v3>
- 415 Bloemendaal, N., Haigh, I. D., de Moel, H., Muis, S., Haarsma, R. J., & Aerts, J. C. (2020c). Generation of a global synthetic
tropical cyclone hazard dataset using STORM. *Scientific data*, 7(1), 1-12.
- Bloemendaal, N., de Moel, H., Martinez, A.B., Muis, S., Haigh, I.D., van der Wiel, K., Haarsma, R.J., Ward, P.J., Roberts,
420 M.J., Dullaart, J.C. and Aerts, J.C., 2022. A globally consistent local-scale assessment of future tropical cyclone risk. *Science*
advances, 8(17), p.eabm8438.
- Botzen, W. J., Bouwer, L. M., Scussolini, P., Kuik, O., Haasnoot, M., Lawrence, J., & Aerts, J. C. (2019). Integrated disaster
risk management and adaptation. In *Loss and damage from climate change* (pp. 287-315). Springer, Cham.
- 425 Calberto, G., Staver, C., & Siles, P. (2015). An assessment of global banana production and suitability under climate change
scenarios. *Climate change and food systems: global assessments and implications for food security and trade*. Rome: Food
Agriculture Organization of the United Nations (FAO).

- 430 Chavez, E., Conway, G., Ghil, M., & Sadler, M. (2015). An end-to-end assessment of extreme weather impacts on food security. *Nature Climate Change*, 5(11), 997-1001.
- Coltro, L., & Karaski, T. U. (2019). Environmental indicators of banana production in Brazil: Cavendish and Prata varieties. *Journal of cleaner production*, 207, 363-378.
- 435 Eberenz, S., Luthi, S., Bresch., David N. (2020). Regional tropical cyclone impact functions for globally consistent risk assessments. *Natural Hazards and Earth System Sciences*, 21(1), 393-415.
- El-Kady, A. E. (2012). The impact of dust storms on crop production in arid regions: A review. *Journal of Arid Land Studies*,
440 22(2), 29-37.
- FAO. (2017). FAOSTAT Database. Banana <<http://www.fao.org/faostat/en/banana>>
- FAO. (2019). Banana facts and figures. Retrieved January 30, 2023, from [https://www.fao.org/economic/est/est-](https://www.fao.org/economic/est/est-commodities/oilcrops/bananas/bananafacts/en/#.Y7K8BXbMJD8)
445 [commodities/oilcrops/bananas/bananafacts/en/#.Y7K8BXbMJD8](https://www.fao.org/economic/est/est-commodities/oilcrops/bananas/bananafacts/en/#.Y7K8BXbMJD8)
- FAO (2020). Voora, V., Larrea, C., & Bermudez, S. Global market report: bananas. International Institute for Sustainable Development.
- 450 FAO. (2022) Banana Market Review 2021. Rome. Retrieved from <https://www.fao.org/3/cc1610en/cc1610en.pdf>
- FAO. (2023). Banana Market Review 2022. Rome. Retrieved January 30, 2023, from
<https://www.fao.org/3/cc6952en/cc6952en.pdf>
- 455 FAO (2023). BananaMarket Review Preliminary results 2023. Rome. Retrieved July 7, 2023, from
<https://openknowledge.fao.org/server/api/core/bitstreams/db25a6ba-e947-4755-b5f0-cb1ee643f266/content>
- Huigen, M. G., & Jens, I. C. (2006). Socio-economic impact of super typhoon Harurot in San Mariano, Isabela, the Philippines. *World Development*, 34(12), 2116-2136.
- 460 Holland, G. J. An Analytic Model of the Wind and Pressure Profiles in Hurricanes. *Mon Weather Rev* **108**, 1212–1218
[https://doi.org/10.1175/1520-0493\(1980\)108<1212:aamotw>2.0.co;2](https://doi.org/10.1175/1520-0493(1980)108<1212:aamotw>2.0.co;2) (1980).

- 465 IFPRI. 2019. International Food Policy Research Institute, 2019, "Global Spatially-Disaggregated Crop Production Statistics Data for 2010 Version 2.0", <https://doi.org/10.7910/DVN/PRFF8V>, Harvard Dataverse, V4
- IPCC, Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change (2021)
- 470 IPCC, Climate Change 2021: Seneviratne, S.I., X. Zhang, M. Adnan, W. Badi, C. Dereczynski, A. Di Luca, S. Ghosh, I. Iskandar, J. Kossin, S. Lewis, F. Otto, I. Pinto, M. Satoh, S.M. Vicente-Serrano, M. Wehner, and B. Zhou, 2021: Weather and Climate Extreme Events in a Changing Climate. In Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1513–1766, doi:10.1017/9781009157896.013.
- 480 IPCC, Climate Change 2021: Fox-Kemper, B., H.T. Hewitt, C. Xiao, G. Aðalgeirsdóttir, S.S. Drijfhout, T.L. Edwards, N.R. Golledge, M. Hemer, R.E. Kopp, G. Krinner, A. Mix, D. Notz, S. Nowicki, I.S. Nurhati, L. Ruiz, J.-B. Sallée, A.B.A. Slangen, and Y. Yu, 2021: Ocean, Cryosphere and Sea Level Change. In Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1211–1362, doi:10.1017/9781009157896.011.
- 485 Islam, S., Cenacchi, N., Sulser, T. B., Gbegbelegbe, S., Hareau, G., Kleinwechter, U., ... & Wiebe, K. (2016). Structural approaches to modeling the impact of climate change and adaptation technologies on crop yields and food security. *Global Food Security*, 10, 63-70
- 490 Kaplan, J. & DeMaria, M. A Simple Empirical Model for Predicting the Decay of Tropical Cyclone Winds after Landfall. *J. Appl. Meteorol.* 34, 2499-2512, [https://doi.org/10.1175/1520-0450\(1995\)034<2499:ASEMFP>2.0.CO;2](https://doi.org/10.1175/1520-0450(1995)034<2499:ASEMFP>2.0.CO;2) (1995).
- Khanduri, A. C., & Morrow, G. C. (2003). Vulnerability of buildings to windstorms and insurance loss estimation. *Journal of wind engineering and industrial aerodynamics*, 91(4), 455-467.
- 495 Kim, J. M., Kim, T., & Son, K. (2017). Revealing building vulnerability to windstorms through an insurance claim payout prediction model: a case study in South Korea. *Geomatics, Natural Hazards and Risk*, 8(2), 1333-1341.

500 Knutson, T., Camargo, S. J., Chan, J. C., Emanuel, K., Ho, C. H., Kossin, J., ... & Wu, L. (2020). Tropical cyclones and climate
change assessment: Part II: Projected response to anthropogenic warming. *Bulletin of the American Meteorological Society*,
101(3), E303-E322.

505 Koks, E.E., Rozenberg, J., Zorn, C., Tariverdi, M., Vousdoukas, M., Fraser, S.A., Hall, J.W., Hallegatte, S., 2019. A global
multi-hazard risk analysis of road and railway infrastructure assets. *Nat. Commun.* 10, 2677. <https://doi.org/10.1038/s41467-019-10442-3>

510 Lesk, C., Rowhani, P., & Ramankutty, N. (2016). Influence of extreme weather disasters on global crop production. *Nature*,
529(7584), 84-87

515 Lin, Y. C., Wang, W. H., Lai, C. Y., & Lin, Y. Q. (2020). Typhoon type index: A New index for understanding the rain or
wind characteristics of typhoons and its application to agricultural losses and crop vulnerability. *Journal of Applied
Meteorology and Climatology*, 59(5), 973-989

520 Machovina, B., & Feeley, K. J. (2013). Climate change driven shifts in the extent and location of areas suitable for export
banana production. *Ecological economics*, 95, 83-95

525 Malek, Ž., Loeffen, M., Feurer, M., & Verburg, P. H. (2022). Regional disparities in impacts of climate extremes require
targeted adaptation of Fairtrade supply chains. *One Earth*, 5(8), 917-931.

530 Mamuye, N. (2016). Statistical Analysis of Factor Affecting Banana Production in GamoGofa District, Southern Ethiopia.
Engineering and Applied Sciences, 1(1), 5-12.

535 Merz, B., Kreibich, H., Schwarze, R., & Thielen, A. (2010). Review article" Assessment of economic flood damage". *Natural
Hazards and Earth System Sciences*, 10(8), 1697-1724.

540 Mendelsohn, R., Emanuel, K., Chonabayashi, S., & Bakkensen, L. (2012). The impact of climate change on global tropical
cyclone damage. *Nature climate change*, 2(3), 205-209.

- De Moel, H., Jongman, B., Kreibich, H., Merz, B., Penning-Rowsell, E., & Ward, P. J. (2015). Flood risk assessments at different spatial scales. *Mitigation and Adaptation Strategies for Global Change*, 20(6), 865-890.
- 535 Mo, W., Fang, W., Wu, P., & Tong, X. (2016). Development of vulnerability curves to typhoon hazards based on insurance policy and claim dataset. *EGU General Assembly Conference Abstracts*, pp. EPSC2016-3360).
- Mohan, P. (2017). The economic impact of hurricanes on bananas: a case study of Dominica using synthetic control methods. *Food policy*, 68, 21-30
- 540 NOAA National Hurricane Center. (2024). Historical Hurricane Tracks [Dataset]. NOAA. <https://coast.noaa.gov/hurricanes/#map=4/32/-80>
- O'Neill, B. C., Tebaldi, C., Van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., ... & Sanderson, B. M. (2016). The scenario model intercomparison project (ScenarioMIP) for CMIP6. *Geoscientific Model Development*, 9(9), 3461-3482.
- 545 Petsakos, A., Prager, S. D., Gonzalez, C. E., Gama, A. C., Sulser, T. B., Gbegbelegbe, S., ... & Hareau, G. (2019). Understanding the consequences of changes in the production frontiers for roots, tubers and bananas. *Global food security*, 20, 180-188.
- 550 Robinson, J. C., & Saúco, V. G. (2010). *Bananas and plantains* (Vol. 19). Cabi.
- Simpson, R. H. Saffir, The hurricane disaster—Potential scale. *Weatherwise* 27, 169–186 (1974)
- 555 Shinozuka, M., Feng, M. Q., Lee, J., & Naganuma, T. (2000). Statistical analysis of fragility curves. *Journal of engineering mechanics*, 126(12), 1224-1231.
- Sun, Y., Wu, T., & Cao, Z. (2021). Wind vulnerability analysis of standing seam roof system with consideration of multistage performance levels. *Thin-Walled Structures*, 165, 107942.
- 560 UN world conference on disaster risk reduction, 2015 March 14–18, Sendai, Japan. Geneva: United Nations Office for Disaster Risk Reduction; 2015. Retrieved January 30, 2023, from http://www.wcdrr.org/uploads/Sendai_Framework_for_Disaster_Risk_Reduction_2015-2030.pdf.

565 UNDRR (2015) Sendai framework for disaster risk reduction 2015–2030. In: UN world conference on disaster risk reduction,
2015 March 14–18, Sendai, Japan. Geneva: United Nations Office for Disaster Risk Reduction; 2015. Retrieved January 30,
2023, from http://www.wcdrr.org/uploads/Sendai_Framework_for_Disaster_Risk_Reduction_2015-2030.pdf

UNICEF. (2022). IN BRIEF TO THE STATE OF FOOD SECURITY AND NUTRITION IN THE WORLD 2022. FAO,
570 IFAD, UNICEF, WFP and WHO.

Varma, V., & Bebbler, D. P. (2019). Climate change impacts on banana yields around the world. *Nature climate change*, 9(10),
752-757.

575 Varma, V., Thompson, W., Duarte, S. B., Krütli, P., Six, J., & Bebbler, D. P. (2020). Mapping the impacts of hurricanes Maria
and Irma on banana production area in the Dominican Republic. *bioRxiv*.

Ward, P. J., De Moel, H., & Aerts, J. C. J. H. (2011). How are flood risk estimates affected by the choice of return-periods?.
Natural Hazards and Earth System Sciences, 11(12), 3181-3195.

580 Wheeler, T., & Von Braun, J. (2013). Climate change impacts on global food security. *Science*, 341(6145), 508-513.

Yum, S. G., Kim, J. M., & Wei, H. H. (2021). Development of vulnerability curves of buildings to windstorms using insurance
data: An empirical study in South Korea. *Journal of Building Engineering*, 34, 101932. C.G.

585