

Probabilistic Seismic Hazard Assessment of Sweden

Niranjan Joshi^{1,2}, Björn Lund^{1,2}, and Roland Roberts^{1,2}

¹Department of Earth Sciences, Uppsala University, Sweden

²Centre of Natural Hazards and Disaster Science, Uppsala University, Sweden

Correspondence: Niranjan Joshi (niranjan.joshi@geo.uu.se)

Abstract. Assessing seismic hazard in stable continental regions (~~SCR~~) such as Sweden poses unique challenges compared to active seismic regions. With diffuse seismicity, low seismicity rate, few large magnitude earthquakes and little strong motion data, estimating recurrence parameters and determining appropriate attenuation relationships is challenging. This study presents a probabilistic seismic hazard assessment of Sweden based on a recent earthquake catalogue, which includes a ~~large-high~~ number of events with magnitudes ranging from ~~5.9 to~~ $M_W-1.4$ to M_W 5.9, enabling recurrence parameters to be calculated for ~~more source areas~~ ~~smaller source zones~~ than in previous studies, and with less uncertainty. Recent ground motion models developed specifically for stable continental regions, including Fennoscandia, are used in ~~logic trees~~ ~~the logic tree~~ accounting for their uncertainty and the hazard is calculated using the OpenQuake engine. The results are presented in the form of mean peak ground acceleration (PGA) maps at 475 and 2500 year return periods and hazard curves for four seismically active areas in Sweden. We find the highest hazard in Sweden in the northernmost part of the country, in the post-glacial fault province. This is in contrast to previous studies, which have not considered the relatively high seismic activity on the post-glacial faults. We also find ~~relatively high hazard~~ hazard to be relatively high along the northeast coast and in southwestern Sweden, whereas the southeast of Sweden and the mountain region to the northwest have low hazard. For a 475 year return period we estimate the highest ~~PGAs~~ PGA values to be 0.04-~~0.05~~-0.06g, in the far north, and for a 2500 year return period it is 0.1-0.15g in the same area. Significant uncertainties remain to be addressed in regards to the ~~SCR~~-intraplate seismicity in Sweden, including the homogenization of small local magnitudes with large moment magnitudes, the occurrence of large events in areas with little prior seismicity and the uncertainties surrounding the potential for large earthquakes on the post-glacial faults in northern Fennoscandia.

1 Introduction

Earthquakes are known to cause some of the most fatal natural disasters. As short term prediction of where and when large earthquakes occur is currently not routinely possible, and as ~~the disaster development in the event is very rapid~~earthquakes can rapidly evolve into disasters, earthquake risk reduction strategies must focus on the mitigation and preparation phases of the disaster cycle (Elliott, 2020). One key component of this is producing an accurate seismic hazard assessment in terms of probable ground motions, which can help identify areas of high hazard and allow for designing appropriate risk reduction strategies.

At plate boundaries, earthquake activity is high and large events can often be associated with identified fault zones. Conversely, stable continental regions, such as Sweden, tend to show a more diffuse geographically distributed pattern of seismicity (Schulte and Mooney, 2005; England and Jackson, 2011). ~~Occurrence~~, occurrence rates are low (e.g. Lund et al., 2021) and areas of high hazard can be difficult to define (e.g. Calais et al., 2016). However, as England and Jackson (2011) show, the risk posed by earthquakes in such intraplate areas is far from negligible. They showed that of the 130 earthquakes with a thousand or more casualties occurring over a 120 year period prior to 2012, about 100 took place in continental interiors. These intraplate earthquakes also caused approximately 75% more total deaths than the plate boundary events.

~~Historical~~ Based on their analysis of historical and instrumental data ~~until~~ before 2005, Bödvarsson et al. (2006) suggest that Sweden can expect on the order of one magnitude 5 earthquake every century and a magnitude 6 earthquake every millennium (Bödvarsson et al., 2006). As the rate of occurrence of these higher magnitude earthquakes is low, the national authority in charge of building codes (Swedish National Board of Housing, Building and Planning - Boverket) has no current plans to implement Eurocode 8 in Sweden (Boverket, personal communication 2020) and there is no official national seismic hazard map. The general increase in risk awareness in society, ~~particularly after large~~ especially after particularly disastrous earthquakes (Tan and Maharjan, 2018), has, however, led to an increased number of requests for seismic hazard information from building companies and corporate site planning projects. For sensitive infrastructure sites such as nuclear power plants, ~~nuclear waste repositories~~ repositories of spent nuclear fuel, dams and mines, seismic hazard estimates are required and there is therefore a need to better define the hazard spatially, estimate potential ground motions and investigate associated uncertainties.

In this paper, we review earlier estimates of seismic hazard in Sweden, with a special emphasis on the recent release of the European Seismic Hazard Model 2020 (Danciu et al., 2021). We then discuss the development of a new probabilistic seismic hazard model for Sweden, which will take into account the large amount of smaller earthquakes recorded by the Fennoscandian seismic networks in the last two decades (e.g. Lund et al., 2021; Veikkolainen et al., 2021; Ottemöller et al., 2018) as well as recent developments in ground motion models (Fülöp et al., 2020; Weatherill and Cotton, 2020). ~~The hazard is calculated using~~ We use the OpenQuake engine (Pagani et al., 2014) ~~and we produce hazard maps~~, to develop hazard maps with mean estimates for peak ground acceleration (PGA) corresponding to return periods of 475 and 2500 years, and hazard curves for Sweden's most seismically active areas. Finally, we discuss the uncertainties inherent in the models and the potential need for additional approaches to seismic hazard assessments in Sweden.

2 Earthquake activity in Sweden

Sweden is located in a stable continental region (SCR) with very few damaging earthquakes (Bödvarsson et al., 2006; Lund et al., 2021). Most of the country is part of the Fennoscandian shield of the East European Craton, see Figure 1, with Archean and Early Paleoproterozoic rocks in the north that ~~are~~ were, to a large extent, reworked during the Paleoproterozoic Svecofennian/Svecofennian orogeny (Stephens and Bergman-Weihed, 2020). The eastern part of the country is dominated by Svecofennian/Svecofennian rocks whereas the southwest contains Meso- and Neoproterozoic rock of the Sveconorwegian orogeny (Stephens and Bergman-Weihed, 2020). The Gulf of Bothnia in the east, and the Baltic Sea further southeast, com-

prise Mesoproterozoic to Ediacaran and Lower Paleozoic sedimentary platformal cover rocks of varying thickness. Onshore
60 Finland comprises Archean, Svecokarelian/Svecofennian and Mesoproterozoic rocks whereas further south, the Baltic coun-
tries are covered by sedimentary rocks deposited on the East European Craton (Bogdanova et al., 2008). In western Sweden,
older basement and Paleozoic cover rocks are affected by the Caledonian orogeny, which created the present day Swedish
mountain chain (Stephens and Bergman-Weiheid, 2020). Further west, Norway is composed mostly of rocks from the Cale-
donian and Sveconorwegian orogens onshore and passive margin deposits offshore (e.g. Corfu et al., 2014; Bingen et al.,
65 2021). To the south, the craton is separated from Phanerozoic Europe by deformation zones through southernmost Sweden (the
Sorgenfrei-Tornquist zone) and Denmark (the Trans European Suture Zone), over which major changes in lithospheric, crustal
and sediment cover thicknesses occur (e.g. Sandersen et al., 2021). The opening of the Atlantic some 60 Ma ago is the latest
large scale tectonic event to significantly affect Fennoscandia (Stephens and Bergman-Weiheid, 2020).

Fennoscandia has been subject to repeated glaciations during the Quaternary. The latest, the Weichselian ice sheet, reached
70 its maximum approximately 20,000 years before present (BP) (Svendsen et al., 2004), before retreating and disappearing
around 10 ka BP. Large fault scarps in northern Fennoscandia, see Figure 1, have been dated to the final deglaciation phase
(Lagerbäck and Sundh, 2008; Smith et al., 2021) and are inferred to have resulted from earthquakes with moment magnitudes
up to M_W 8 (e.g. Lindblom et al., 2015). Glacial isostatic adjustment is still ongoing in Fennoscandia and GNSS measurements
show that all of Sweden is rebounding, with a maximum uplift velocity of 10.3 mm/year on the northeast coast, decreasing to
75 about 1 mm/year in southernmost Sweden (Vestøl et al., 2019).

The seismicity in Fennoscandia has been the subject of many studies, Gregersen et al. (2021) provide a recent review of
the whole region whereas Lund et al. (2021) focus on recent observations of earthquakes in Sweden, which we briefly review
here. The expansion of the Swedish National Seismic Network (SNSN) from 6 to 65 stations during the first decade of the 21st
century has provided a wealth of new earthquake observations, increasing the total number of recorded events from about 1,600
80 prior to the year 2000 to about 12,500 in 2021-2021, see Figures 1 and 2. Most of the events are small, with the magnitude of
completeness estimated at local magnitude M_L 0.5 within the network. Areas of Earthquakes in Sweden are mostly strike-slip,
and focal depths vary between near-surface and 35 km depth. Areas of relatively high seismic activity include the southwestern
part of Sweden within Sweden include the southwest across Lake Vänern, along the northeast the northeast along the coast
and in the far north along the post-glacial faults, see Figure 1. The southwest and the northeast coast seismicity agrees well
85 with what was known already from the pre-instrumental period (e.g. Kjellén, 1910), the new data have, however, revealed that
activity locally cluster along specific zones and is less diffuse than previously thought. The new observations have significantly
changed how seismic activity in the north is viewed. We now know that seismicity is strongly correlated with the mapped
post-glacial fault (PGF) scarps, indicating that the faults are still very active although event magnitudes are generally low
(Lindblom et al., 2015) (M_L -1 to 3.5, Lindblom et al. (2015)). One reason for the lack of early observations along the PGFs is
90 the low population density in the inland. On the Pärvie fault (Figure 1) for example, there is only one event in the catalogue prior
to 1967, a M_L 2.6 event in 1927. The largest recorded Pärvie event had M_L 3.7 in 1967 (Ahjos and Uski, 1992). Conversely,
the Burträsk PGF (Figure 1), on the northeast coast, has earthquake reports going back to the 18th century, with the largest

events having magnitudes around M_L 4-4.5 (Ahjos and Uski, 1992) and the recent data shows that it is the most seismically active region in Sweden.

95 The largest historically recorded earthquake on land in Fennoscandia was the 1819 Lurøy M_W 5.9 event in Nordland, Norway, located about 80 km from the Swedish border (Mäntyniemi et al., 2020). In 1759, ~~a M_S 5.6 event~~ an event with surface-wave magnitude 5.6 occurred in the waters between Sweden and Denmark (Muir Wood, 1989), and in 1904 a M_W 5.4 event took place just off the Swedish west coast close to the Norwegian border (Bungum et al., 2016). Additional M_W 5+ events have occurred in the regions around Sweden, such as the 2004 Kaliningrad earthquake doublet (Gregersen et al., 2007),
100 the 1892 M_W 5.2 western and the 1834 M_W 5.0 southern Norway events (Ahjos and Uski, 1992). Recently, Olesen et al. (2021) trenched the post-glacial Stuoragurra fault in northern Norway and found evidence of an event with magnitude about 7 occurring as recently as 700 years ago. This indicates that the PGFs may still be capable of generating very large earthquakes.

The driving forces of Fennoscandian seismicity have been a debated subject, see Gregersen et al. (2021) for a recent review. As glacial rebound dominates the deformation rates of the Fennoscandian crust, both vertically and horizontally, GNSS mea-
105 surements cannot resolve any tectonic-driven deformation (Vestøl et al., 2019). However, evidence such as focal mechanisms and in-situ stress measurements points to ridge-push from the opening of the Atlantic as the main driving force for seismicity, potentially moderated by glacially induced stress, sediment load or other regional to local stresses (Gregersen et al., 2021).

3 Previous seismic hazard assessments for Sweden

Here we review earlier seismic hazard assessments for Sweden. We will refer to the intercept and slope of the earthquake
110 frequency-magnitude distribution as the a- and b-values (Gutenberg and Richter, 1944), M_c as the magnitude of completeness of a frequency-magnitude distribution and M_{max} as the maximum possible magnitude for an area.

The studies by Båth (1979) and Slunga (1979) were the first to estimate seismic hazard in Sweden. They only had access to some 25 years of instrumental data and thus relied to a large extent on macroseismic data, which is limited in parts of Sweden due to the low population density. Båth (1979) estimated recurrence rates for earthquakes of varying magnitudes in a 2°-by-2°
115 grid across Fennoscandia and calculated earthquake risk according to the formula put forward by Lomnitz (1974), describing earthquake risk as “the probability $R(D/T)$ that a shock of mean return period T occurs during a design period D ”. He found the highest hazard on the Norwegian west coast ~~For Sweden, he and, for Sweden,~~ estimated a 5% probability in 10 years ~~of an (a return period of 195 years) of an event of $M_L \geq 5$ event and above~~ on both the southwest and northeast coasts. Slunga (1979), investigating the seismic hazard at sites of the then four Swedish nuclear power plants, estimated accelerations of 0.05 – 0.20 g
120 for an annual probability of 10^{-5} . The development of the FENCAT catalogue (Ahjos and Uski, 1992), a continuously updated joint regional catalogue of earthquakes in Northern Europe since 1375, enabled later studies to use a larger, more homogeneous data set for Fennoscandia.

FENCAT data until 1987 were used for the first Swedish probabilistic seismic hazard assessment (PSHA)~~work~~, directed at site-specific assessments for Sweden’s four nuclear power plants (SKI, 1992). The study included Fennoscandian earthquakes
125 south of latitude 61° ~~were used for seismicity rate information in various combinations of seismic source areas~~ to estimate the

seismicity rate. The lack of strong motion data from Fennoscandia prompted SKI (1992) to turn to existing Japanese “Standard Response Spectra for rock sites” for ground motion models (SKI, 1992). The Japanese spectra were modified to fit Swedish earthquake source properties and crustal attenuation properties. The envelope ground response spectra resulting from the SKI (1992) study are still in use in the Swedish nuclear industry today, and ~~for a probability of exceedance of the estimated PGA is~~
130 ~~0.11g for a hard rock site at an annual probability of~~ 10^{-5} ~~per year~~ and a damping of 5% ~~the estimated PGA is 0.11g for a hard rock site~~ (SKI, 1992; Larsson and Larsson, 2018). In a review, Lund et al. (2017) found that SKI (1992) may underestimate the rate of events larger than approximately magnitude M_W 5.2 and recommended an update with modern data and methodologies.

Kijko et al. (1993) used FENCAT and a model that accounts for varying magnitudes of completeness through time in order to quantify probabilistic seismic hazard in Sweden by estimating the recurrence rates of earthquakes. They found that Sweden
135 can expect 5.5 earthquakes per year with magnitudes equal to or greater than 2.4, with the maximum expected earthquake magnitude in southern Sweden being 4.9 for a 615-year time span, and a maximum magnitude of 4.3 in northern Sweden over 331 years. Mäntyniemi et al. (1993) expanded the Kijko et al. (1993) study to all of Fennoscandia and also included probabilities for non-exceedance of various magnitudes in 1 and 50 years in defined subregions. The results for Sweden were similar to that of Kijko et al. (1993).

140 The ~~Wahlström and Grünthal (2000) and follow-up Wahlström and Grünthal (2001)~~ Wahlström and Grünthal (2000, 2001) studies perform full probabilistic seismic hazard assessments for Fennoscandia using the FENCAT catalogue. Wahlström and Grünthal (2001) apply two different magnitude homogenization schemes. They used two large regions for magnitude completeness estimates, and six different regional and non-regionalized seismic source ~~area models~~ zones. Combining the recurrence parameters with two different ground motion models in a logic tree, Wahlström and Grünthal (2001) produce a
145 hazard map for Fennoscandia, with 90% probability of non-exceedance of median horizontal PGA in 50 years, and hazard curves and hazard deaggregations for Sweden, Norway, Denmark and Finland. Their results indicate that the Norwegian west and northwest coasts have the highest hazard in Fennoscandia. The highest hazard in Sweden was found in the southwest, close to the Norwegian border, with an estimated PGA of 0.03g for 475 years return period and 0.08g for 4745 years return period.

Mäntyniemi et al. (2001) produced a seismic hazard map of Fennoscandia by using the site-specific technique of Kijko and
150 Graham (1998, 1999) and applying it to a grid of Fennoscandian locations, similar to a smoothed seismicity approach (e.g. Danciu et al., 2021). They estimate ~~the recurrence parameters (a and b values) and the maximum magnitude, and~~ M_{max} values from the data in a selected specific area around a grid point and use those with a ground motion model to calculate the hazard in terms of ground motion probabilities. Mäntyniemi et al. (2001) used a 0.5° by 0.5° grid with a constant b-value of 0.84 ± 0.01 and an M_{max} of 5.8 over the entire region, while the a-value varied with the seismicity included at each grid point.
155 Ground motion models by Uski and Tuppurainen (1996) and EMSC (1999) were used and the resulting hazard map shows PGA for a 475 year return period, where the highest hazard was obtained at the northwestern coast of Norway with 0.038g (note that western Norway around Bergen was not included in the analysis). The highest PGA in Sweden reached 0.022g both in southwestern Sweden and on the northeast coast.

Bodare and Kulhánek (2006) used the results in Kijko et al. (1993) and Mäntyniemi et al. (1993) together with a relationship
160 between M_L and the acceleration at the epicentre (Båth, 1980) to infer that seismic hazard at hydropower dams in Northern

Sweden is unlikely to exceed 0.04g in the next 50 – 100 years. In two large PSHA projects for the nuclear industry in Finland, analysis of seismicity and definition of seismic source ~~areas-zones~~ was undertaken for Sweden and Finland. The first, the Fennovoima project, assembled seismologists and geologists from Finland and Sweden to perform a full site-specific PSHA for a proposed new nuclear power plant in northern Finland (Korja and Kosonen, 2015; Saari et al., 2015). Using a similar methodology, a PSHA revision was later performed for the sites of the existing nuclear power plants in Finland (Korja and Kihlman, 2016). The hazard results of these studies have not been made public. On a larger scale, the Global Seismic Hazard Assessment Program (GSHAP, Grünthal and Group, 1999) and the European Seismic Hazard Models 2013 and 2020 (ESHM13 and ESHM20; Woessner et al., 2015; Danciu et al., 2021, respectively) have estimated seismic hazard in Sweden. ESHM20 is described in some detail in Section 3.1 below.

170 3.1 Sweden in the European Seismic Hazard Model 2020

The most recent evaluation of seismic hazard in Sweden comes from the 2020 European Seismic Hazard Model (ESHM20) (Danciu et al., 2021), which assesses earthquake hazard in the Euro-Mediterranean region and builds on the 2013 version of the model (ESHM13; Woessner et al., 2015). ESHM20 applies a fully probabilistic framework homogeneously across the entire pan-European region, while accounting for cross country-border issues. The earthquake data come from the European PreInstrumental earthquake CAlogue (EPICA, Rovida and Antonucci, 2021), which spans the years 1000 to 1899, and from an extension of the European-Mediterranean Earthquake Catalogue (EMEC, Grünthal et al., 2013) such that it spans the time 1900 to the end of 2014. Magnitudes are harmonized to the moment magnitude M_W , the catalogue is declustered and only events with $M_W \geq 3.5$ are included in the magnitude-frequency analysis. For Sweden and the surrounding region within 300 km of Sweden's borders and economic zone, the unified ESHM20 earthquake catalogue contains 212 events, from 1497 to 2014, with magnitudes $3.5 \leq M_W \leq 5.8$, [see Figure 2](#).

In order to estimate the recurrence parameters of the magnitude-frequency distribution at the most appropriate spatial distribution, ESHM20 uses a hierarchy of seismic source ~~areas-zones~~. Fennoscandia is assigned to a single maximum magnitude zone, within which M_{max} is uniform, ~~and~~. [It is divided into](#) two “completeness zones”, ~~CSZ, where reporting in each zone where reporting~~ is assumed to be ~~spatially~~-homogeneous through time such that the temporal variation in ~~the magnitude of completeness~~, M_c , is the same [all through the zone](#). At the next levels, Fennoscandia is divided into four “tectonic” source ~~areas-zones~~, TSZ, and about 20 smaller “area” source zones, ASZ, based on geology, seismicity and an assumption of homogeneous seismicity rate. In these zones, ~~recurrence parameters (a- and b-values)~~ are calculated using a doubly truncated Gutenberg-Richter distribution, using an automatic maximum likelihood method based on the earthquakes within the zone (Danciu et al., 2017), for zones with more than 30 earthquakes. For ASZs with less than 30 events, the b-value from the TSZ is re-used and the a-value is rescaled from the TSZ with the ratio of the number of complete events in the ASZ and TSZ. If the ASZ does not contain any events above the magnitude of completeness, an a-value is assigned by rescaling the TSZ value with the area ratio of ASZ to TSZ. Uncertainties are estimated using random sampling and discrete approximation of probability distribution methods, giving 16th, 50th and 84th percentile estimates of the median a- and b-values. A tapered Pareto distribution is also defined to describe the recurrence parameters, primarily to provide a faster decaying alternative closer to

195 ~~the maximum magnitude~~ M_{\max} . For Sweden and a 300 km border region, only two ASZs have enough events for separate estimates of the recurrence parameters, all other use the TSZ b-values and rescaled a-values. All seismogenic source models are implemented through a logic-tree, with the median, 5th and 95th percentiles describing the branches of individual area source Gutenberg-Richter parameters, with 0.6, 0.2 and 0.2 as the respective branch weights. M_{\max} in Fennoscandia has three branches, where M_{\max} is 6.3, 6.6 and 6.9, with weights 0.5, 0.4 and 0.1, respectively.

200 ESHM20 uses a scaled backbone approach for implementing ground motion models (GMMs). This is done by selecting a relevant GMM from the literature, and modifying it based on the observations and knowledge of the region's crustal properties to account for the uncertainty in the expected ground motions. The scaled backbone GMM for the cratonic region is obtained by generating median ground motions on hard rock for various magnitude ($4.0 \leq M_w \leq 8.0$) and distance ranges ($1.0 \leq R_{rup}(km) \leq 500km$), starting with equally weighing 21 Central and Eastern
205 United States (CEUS) GMMs and fitting to a parametric GMM of the same form as Kotha et al. (2020) (defined for active shallow crustal regions in ESHM20), mapped into a scaled backbone logic tree (Weatherill and Cotton, 2020). These ground motions are defined for very hard rock conditions ($V_{S,30} = 3,000$ m/s) ~~conditions~~ and re-scaled to reference $V_{S,30} = 800$ m/s rock ~~condition~~ conditions using the CEUS site amplification models of Stewart et al. (2020) and Hashash et al. (2020). ~~They~~ Weatherill and Cotton (2020) calibrate the model's epistemic uncertainties and propose a logic tree for the application of the
210 model to cratonic regions where one branch is the new parametric Craton model and one branch an adaption of the Shallow Crustal seismicity model used for Central Europe (Kotha et al., 2020), only retaining the mid and upper (i.e. higher velocity) twigs of that branch. ~~For~~ This GMM logic tree combination in used for most of Sweden, Finland and the Baltic, ~~this logic tree combination for cratonic regions is used,~~ while the Shallow Crustal logic tree is used for most of Norway and Denmark.

The final hazard map shows that for Fennoscandia, the seismic hazard is highest along the Norwegian west coast. The
215 Swedish northeast coast and the coast of the Baltic countries also have relatively high hazard whereas most of Denmark, the interior of north and central Sweden and south-central Finland have the lowest hazard. The highest estimated mean PGA for Sweden ~~was~~ is 0.025g for a return period of 475 years, and 0.11g for a return period of 2500 years.

4 Methods

Probabilistic seismic hazard assessment (PSHA) is a widely used technique to estimate seismic hazard which we utilize in
220 this study. It quantifies the rate at which ground-motions are expected for all possible earthquake scenarios (Cornell, 1968); ~~The method can be broadly divided into three steps,~~ based on earthquake recurrence data and crustal attenuation properties. Here we detail how our study implements the steps involved: (1) Preparing earthquake data ~~and associating it with seismic source areas and fault zones,~~ with a calculation, which involves merging data from different catalogues, removing dependent events and homogenising the earthquake magnitude; (2) Defining seismic source zones and associating earthquakes to them;
225 (3) Calculation of recurrence parameters ~~;~~ (2) for each seismic source zone; (4) Defining the seismic source model logic tree; (5) Selecting appropriate ground motion models that represent the attenuation of seismic waves ~~;~~ (3) and defining the ground

[motion model logic tree](#); (6) Estimating hazard probabilistically by accounting for the epistemic uncertainties through a logic tree. ~~Here we detail how our study implements these steps.~~

4.1 Data preparation

230 A number of seismic networks in Fennoscandia collect seismic data and report to FENCAT and other international compilations: The Swedish National Seismic Network (SNSN, Lund et al., 2021), The Norwegian National Seismic Network (NNSN, Ottemöller et al., 2018) and NORSAR (Norsar, 1971), The Finnish National Seismic Network (FNSN, Veikkolainen et al., 2021; Ahjos and Uski, 1992) and Oulu University (Sodankylä Geophysical Observatory / University of Oulu, 1980), the Geological Survey of Denmark and Greenland (GEUS, 2023) and the Geological Survey of Estonia (Soosalu et al., 2022).

235 ~~We compile a homogenized high-quality earthquake catalogue of events in Sweden and a surrounding 300 km zone, using FENCAT, the full SNSN catalogue and the ESHM20 unified earthquake catalogue. The~~ [Location and magnitude uncertainties for the recorded earthquakes have decreased through time, with increased population density in the pre-instrumental period and with improved network configurations in later times. There are no uncertainties associated with location and magnitude estimates in the FENCAT, but since around 2010 the SNSN has average uncertainties of about 2-3 km in the epicenters and 0.1](#)

240 [-0.2 in the magnitudes. Since the rate of recorded non-tectonic seismic events such as mine, quarry, construction and military blasts and mining induced events exceeds the earthquake rate by up to a factor ten or more in much of Fennoscandia, the Fennoscandian seismic network analysts have a high focus on classification, nowadays aided by machine learning algorithms \(Kortström et al., 2016; Eggertsson et al., 2024\). During FENCAT data merging, additional checks for non-tectonic events are carried out, such that the final catalogue is as free as possible from such events. In the older data there are, however, probably](#)

245 [still both man-made and frost related events present that are difficult to identify \(Uski et al., 2015\). The earthquake](#) data is processed with the methodologies developed in Uski et al. (2015), comprising (i) data merging and cleaning, (ii) declustering, and (iii) magnitude homogenization. We briefly describe these steps below.

4.1.1 Data merging and cleaning

The version of the FENCAT catalogue at our disposal spans the year 1375 until the end of 2014 (Korja and Kihlman, 2016).

250 We added the semi-reviewed FENCAT between 2015-01-01 and 2020-09-30 (Finnish National Seismic Network, 2023). Both these catalogues contain earthquakes from all over Fennoscandia and [some other areas in](#) Northern Europe. The SNSN does not report all small microearthquakes to FENCAT, so we also added the entire 2000-08-15 to 2022-02-28 SNSN catalogue to the data set. ~~In addition, we extracted a few events in the region of interest~~ [As we need data from a zone surrounding Sweden out to 300 km distance, we also extracted eight events](#) from the unified ESHM20 catalogue (Danciu et al., 2021) ~~-that were~~

255 [not present in our version of FENCAT.](#) Duplicates were removed from the resulting list of events and remaining non-tectonic events ~~such as quarry, industrial or military blasts, rock bursts, mine collapses etc~~ [that could be identified](#) were removed from the data. As FENCAT contains events far from Sweden, such as along the Mid-Atlantic ridge, we removed these ~~events~~ together with events in Britain, central Poland, Russia east of 40° longitude and Svalbard prior to further processing. The ~~final~~ [joint](#)

earthquake catalogue comprises 24,215 events within the tectonic source ~~areas-zones~~ (Section 4.2) of the hazard assessment, see ~~Figure 4~~ with earthquake magnitudes between M_W -1.4 and M_W 6.1.

4.1.2 Declustering

Probabilistic seismic hazard assessment generally assumes that the earthquakes included in the assessment occur independently of each other, following a Poisson process (Baker et al., 2021). In reality, many events occur as a consequence of other events, such as foreshocks, aftershocks and swarm events. Separating the catalogue into dependent and independent events is referred to as declustering, and numerous techniques exist for this purpose; see review in van Stiphout et al. (e.g. 2012). As discussed in detail in Uski et al. (2015), most declustering methods are designed for plate boundary seismicity, with large events and significant Omori-type aftershock activity. In intraplate Fennoscandia, on the contrary, events are generally small and even moderately large events can have few and small aftershocks, inconsistent with both the Omori and Båth aftershock relationships. Following Uski et al. (2015), we implemented a declustering scheme using a Gardner and Knopoff (1974) windowing procedure based on conservative estimates of location error and manual inspection of clusters. We set the spatial (radius) and temporal distances between an event and its potential aftershock/swarm member to 10 km and 30 days for events with magnitude larger than 1.5, and to 5 km and 15 days for smaller events. We did not consider the magnitude of completeness or the relative size of events in the clustering. The largest event in each cluster was considered as the independent event and forwarded to the declustered catalogue.

The results were checked manually and also by comparing the inter-event times of subsets of the declustered catalogue ~~with-to those from~~ synthetic Poissonian catalogues ~~;~~ (produced using the observed average rates and ~~the-an~~ average of 1,000 simulated inter-event time sequences). The subsets were defined in time intervals where the average yearly rate is approximate similar and visual inspection of the subset inter-event time distributions (Figure A1) shows that the declustering process to a large degree removes the non-Poissonian short inter-event time events in the original catalogue.

In agreement with Mäntyniemi (1996) and Uski et al. (2015), we find that the fraction of dependent events in Fennoscandia is low. Our catalogue retains 19,943 independent events ~~;~~ (Figure 2) implying approximately 18% dependent events. The results show that clustering in Sweden is highest along the very active post-glacial faults in the north, especially the Pärvie and Burträsk faults, and is also significant along the northeast coast and in some areas south of Lake Vänern (Figure A2). The largest cluster contains 199 events and occurred in the Kouvola area in southern Finland, known for its shallow swarm activity. In their study of a 500 km radius area around Pyhäjoki on the northeast coast of Finland, Uski et al. (2015) found a smaller fraction of dependent events ~~of-only~~ 11%, ~~a difference to our result which is likely due to the fact that~~ than we find here. Although their area of interest ~~did not include~~ included most of the seismicity of Finland and Sweden north of Uppsala (Fig. 1), the difference is probably due to it not including the highly active west coast and Nordland regions in Norway, where swarm activity is frequently observed (e.g. Shiddiqi et al., 2022). ~~We-Comparing to ESHM20, we~~ note that the ESHM20 earthquake catalogue only contains a total of 360 events, all with $M_W \geq 3.5$, for the area covered by our declustered catalogue; ~~all-with ≥ 3.5 .~~

4.1.3 Magnitude homogenization

Seismic hazard assessments should be based on homogeneous earthquake size estimates, as commonly represented by the moment magnitude, M_W , scale (Hanks and Kanamori, 1979), which does not saturate at large magnitudes. Most earthquakes in the FENCAT catalogue, however, have magnitude estimates either from macroseismic observations or from local magnitude scales, M_L , from the various network operators, with scales varying slightly over time and institute. Uski et al. (2015) constructed a consistent moment based magnitude scale, $M_W(HEL)$, based on the local Helsinki magnitude scale, $M_L(HEL)$, earthquakes in SNSN after 2000-08-15 which all have estimates of the scalar seismic moment (Rögnvaldsson and Slunga, 1993) and other moment estimates from individual studies. They provided relationships between the various magnitude scales and $M_L(HEL)$, $M_L(HEL)$ and $M_W(HEL)$, and related $M_L(HEL)$ - $M_W(HEL)$ to the scalar seismic moment M_0 as

$$M_L(HEL)M_W(HEL) = 0.83 \log_{10}(M_0) - 7.98 \quad \text{for } \log_{10}(M_0) \leq 13.5$$

$$M_L(HEL)M_W(HEL) = 0.59 \log_{10}(M_0) - 4.73 \quad \text{for } \log_{10}(M_0) > 13.5$$

We will use the $M_W(HEL)$ magnitude homogenisation scheme for all events in this study and for brevity denote $M_W(HEL)$ as M_W . Uski et al. (2015) points out that the relation is only defined by observations to $\log_{10}(M_0) \approx 17.5$ ($M_W \approx 5.6$), but that it agrees well with the original Hanks and Kanamori (1979) to approximately $\log_{10}(M_0) = 19$, or $M_W = 6.6$.

The final declustered and magnitude homogenized catalogue has 19,943 events, with magnitudes ranging between $-1.4 \leq M_W \leq 5.9$. There are 341 events with $M_W \geq 3.5$ and 182 events with $M_W \geq 4.0$ in the tectonic source zones, indicating that our magnitude homogenization scheme produces slightly lower magnitudes than the one used by ESHM20 for Fennoscandia, which has 360 events with $M_W \geq 3.5$ in the same area. We will only use events recorded from the year 1875 onwards, which is approximately when the Swedish Geological Society started more systematic investigations of earthquake reports (Kjellén, 1910). Figure 3 shows earthquake magnitude-density plotted as a function of time for Sweden and the Swedish economic zone and highlights the significant improvement in completeness after the year 2000. We also note the lack of reported $M_W \geq 3.2$ events between 1936 and 1962, the reason for which we do not know, and the very few recorded events with $M_W \geq 4$. Prior to 1950 the data stems almost entirely from macroseismic observations, Figure 3 shows that this has produced a lower magnitude threshold at approximately $M_W 2-2$, which was the lowest magnitude given to macroseismically observed events.

4.2 Seismic source areas/zones

Seismic source areas (SSAs/zones (SSZs)) are defined based on geological and tectonic features, the clustering of seismicity and homogeneity of the seismicity rate. In addition, the observation of earthquake activity in the zone should be homogeneous over time such that variations of the magnitude of completeness, i.e. variations in M_c are the same within each zone are the same. These source areas. The source zones form the basis upon which recurrence parameters, maximum magnitude and ground motion models are calculated and assigned. During the Finnish nuclear industry projects (Korja and Kosonen, 2015; Korja and Kihlman, 2016), Fennoscandian seismologists and geologists came together to define appropriate cross-border SSAs-SSZs for

these site-specific PSHAs. Work on harmonized Fennoscandian ~~SSAs~~ SSZs was continued as part of the review for ESHM20
325 and further developed in workshops in 2022 and 2023 for this study and ongoing studies in Finland and Norway.

As this study focuses on Sweden, we use earthquakes in the area of Sweden and a 300 km wide area surrounding the national
boundaries and, in the sea, the Swedish Economic Zone. Following the concept of super zones in ESHM20, we define both
large scale Tectonic Source Zones (TSZ) and Area Source Zones (ASZ), see Figure 4. The TSZs capture large-scale aspects
of the crust, such as the regional orogenies and deformation structures, while attempting to ensure homogeneous seismicity
330 patterns. Given the relatively large number of earthquakes in the TSZs, they provide large scale recurrence parameters which
we use for ASZs with insufficient number of events. We define four TSZs: Tectonic source zone T1 comprises the Archean
and Svecokarelian rocks of eastern and northeastern Sweden and northern Finland, including the seismicity along the Swedish
northeast coast, most of the Fennoscandian post-glacial faults, the Kuusamo seismicity in Finland and the scattered seismicity
in southeastern Sweden. Zone T2 spans the Sveconorwegian rocks in southwestern Sweden and southern Norway plus the Oslo
335 graben, which are all areas of relatively high seismicity. Western and northern Norway with the Caledonides and the adjacent
ocean areas make up zone T3, including the intense seismicity along the west coast and in the Nordland region. T4 is the least
seismically active area, with the Svecokarelian and Archean rocks of central and southern Finland, and the Baltic countries on
the sediment covered East European Platform.

The smaller zones, ASZs, are delineated based on active deformational features, such as post-glacial faults, smaller scale
340 geological and tectonic features and distinct seismicity patterns, see Figure 4. For zones that extend outside the 300 km limit,
we include all events in the zones in the calculations in order to increase the significance of the results. Uncertainty in the
location of earthquakes are on the order of a few kilometers for the post-2000 data, which will have a minimal effect on the
allocation of events to ASZs. For the pre-instrumental data the location uncertainties can be much larger, especially for event
in the sea areas, but as most zones are large the problem is likely small and we do not attempt to consider these uncertainties
345 in the allocation to ASZs.

4.3 Maximum magnitude

An estimate of the maximum possible earthquake magnitude, M_{\max} , in each seismic zone is used both when calculating the
recurrence parameters, using a doubly truncated Gutenberg-Richter distribution (e.g. Weichert, 1980), and when performing
the actual hazard calculation. In stable continental regions like Fennoscandia, estimation of M_{\max} is complicated by low
350 seismicity rates and short observation times, lack of well-defined active fault zones and general uncertainty about the length,
or existence, of an earthquake cycle (e.g. Wheeler, 2016; Calais et al., 2016). A number of different approaches have therefore
been taken in assessing M_{\max} for PSHA purposes. Two frequently used methods are extreme value type statistics (EVTS)
based on observational data (e.g. Kijko and Graham, 1998; Kijko and Singh, 2011; Beirlant et al., 2019; Zöller, 2022) and
the so called EPRI method of combining global tectonically analogous datasets with local observations (Johnston et al., 1994;
355 Wheeler, 2016).

The largest observed earthquake in Fennoscandia is the Lurøy, Norway, event of 1819, with a magnitude ~~that was revised
by Mäntyniemi et al. (2020) from 5.8 to~~ M_W 5.9 (Mäntyniemi et al., 2020). Using two varieties of EVTS, Mäntyniemi et al.

(2001) found Fennoscandian M_{\max} of 5.84 ± 0.50 and 5.94 ± 0.52 , close to the observed maximum magnitude (excluding the post-glacial faults). As noted by Kijko and Singh (2011), the EVTS methods often underestimate M_{\max} . Wahlström and Grünthal (2001) used the EPRI approach and found a five point probability ~~distributions~~ distribution of M_{\max} for ~~onland~~ onshore Fennoscandia, from M_{\max} 5.85 to 7.70, with equal probabilities of 0.2. ESHM13 and ESHM20 both use the same EPRI based approach to M_{\max} for Fennoscandia, with a resulting distribution of M_{\max} of 6.3, 6.6 and 6.9, with probabilities 0.5, 0.4 and 0.1, respectively.

The post-glacial faults of northern Fennoscandia are still seismically active (e.g. Lindblom et al., 2015), although the main ruptures are inferred to have occurred at the end of the latest deglaciation some 10,000 years ago (e.g. Smith et al., 2021). They have surface lengths of 40 - 150 km (Smith et al., 2021), indicating potential magnitudes of up to M_W 8 (Lindblom et al., 2015). Such large events have been inferred to be triggered by the glacial isostatic stress contribution from the late deglacial phase (e.g. Lund, 2015), but as the recent trenching of the Stuuragurra fault in northern Norway indicate ruptures of magnitude 7 as late as 700 years ago (Olesen et al., 2021), the potential for very large earthquakes in the current stress field needs to be considered.

Here we combine the EPRI approach of ESHM20 with the post-glacial fault information and use an M_{\max} distribution of 6.3, 6.6, 7.0 and 7.5, with probabilities 0.4, 0.4, 0.15 and 0.05, respectively. The large magnitudes mostly affect the hazard on long time scales, as discussed further in Section 6.

4.4 Calculating recurrence parameters

Recurrence parameters for each source zone are calculated assuming that the mainshocks in the catalogue follow a Poisson distribution that can be represented by the Gutenberg-Richter (Gutenberg and Richter, 1944) relationship

$$\log_{10} N(M) = a - bM \quad (1)$$

where $N(M)$ is the cumulative number of earthquakes per year with (moment) magnitude greater than, or equal to, M , a is the activity rate and b represents the ratio of low-magnitude events to high magnitude events. The relationship breaks down for events with magnitudes smaller than some completeness magnitude M_c , where not all events are recorded. Estimating recurrence parameters is challenging in a low seismicity area like Fennoscandia, where larger earthquakes are rare and population density low, ~~such that there are very few recorded events.~~ Very few events have been recorded prior to the installation of more sensitive seismic networks in the 1960s and 1970s. ~~Although the first seismograph in Fennoscandia was installed in Sweden in 1904, the~~ The completeness magnitude of the national earthquake catalogue has varied during most the 20th century ~~from about M4 between about M3~~ to about M2. As Figure 3 shows, due to the rapid expansion of the seismic network in the early 2000, the completeness magnitude for Sweden with within the seismic network has dropped to below M1. However, once the catalogue is subdivided into tectonic or area source zones, the number of earthquakes and M_c vary significantly with time. ~~Therefore,~~ a and we estimate M_c separately for each zone. A time-varying completeness algorithm (Weichert, 1980), as implemented by the OpenQuake (OQ) engine (Pagani et al., 2014), was used to estimate the recurrence parameters. For quality control, we also used Aki (1965)'s maximum likelihood method, as modified by Tinti and Mulargia (1987) ~~in certain time intervals.~~ We do not

take into account uncertainties in individual magnitudes in the recurrence calculations but consider these to be accounted for by the uncertainties in the recurrence parameters themselves. Recurrence parameters, with associated uncertainties, are estimated

for the four tectonic source zones and for area source zones with sufficient data.

As is well established, calculation of a- and b-values with small sample sizes or a small magnitude range may lead to significant bias in the estimates, outside the formal uncertainty limits (e.g. Geffers et al., 2022). We therefore require that a zone contains more than 200 events for a recurrence parameter calculation. For area source zones sparse in data we follow the ESHM20 methodology and assign them the b-value from the corresponding tectonic source zone, while an a-value is calculated by re-scaling the tectonic zone a-value by the ratio of the number of events with magnitude above M_c in the two zones, N_{ASZ}/N_{TSZ} . If there are no events in the zone, the tectonic a-value is scaled by the areal ratio of the two zones.

In most zones we use data from the late 1800 or early 1900 onward. In the far north, however, where there is very little macroseismic data, our records start with the increase in station density in the 1960s. The number of complete events in each tectonic source zone range from 262 in T4 to 7347 in T1. The a-values range from 1.857 in T4 to 3.267 in T3, and the b-values from 0.842 in T2 to 0.912 in T1, see Table 1. In the ASZs, recurrence parameters could only be calculated for 12 out of the 31 zones (zones 2, 3, 4, 9, 13, 14, 15, 18, 23, 24, 30 and 31) due to the sparsity of data in most zones. For zones within 300 km of Sweden, the a-values for the ASZs range between 0.041 for zone 20 in T4 to 2.504 for zone 2 in T1. The b-values range between 0.8 for zone 18 in T1, to 1.051 in ASZ 2, see Table 2, where we list the ASZs in Sweden. Uncertainties are calculated using the OQ implementation of the Weichert algorithm. Zone 23 has the largest b-value uncertainty of about 5%, followed by zone 4 at around 4%. Due to the large number of events with magnitude above M_c in tectonic zone T1, the b-value uncertainty is only ± 0.007 , which we find unreasonably small compared to how the b-value changes when M_c is varied. We therefore set the minimum b-value uncertainty to ± 0.01 . In zones with too few events for a separate calculation of recurrence parameters, we assign the same uncertainty as what we find for the tectonic zone calculations to the a- and b-values. As the a-value is scaled by the number of complete events, this implies that the a-value uncertainty becomes relatively **very**-large in some ASZs.

4.5 Source model logic tree

The source model logic tree is based on the earthquake catalogue, the zonation, the derived recurrence parameters for each zone and their uncertainties, and the maximum magnitude information, see Figure 5. The first branching level represents each individual area source zone, followed by three branches representing the maximum likelihood estimate of the doubly truncated GR distribution and two standard deviations, representing the 5th and 95th percentiles. Four further branching levels are applied to each model for the M_{max} values of 6.3, 6.6, 7.0 and 7.5, each with 0.4, 0.4, 0.15 and 0.05 as their respective weights. The overall structure of the source model logic tree is as implemented in the ESHM20 calculations for the region (Danciu et al., 2021). With three branches each for the a-, b- values and four for M_{max} , a total of twelve branches are defined for each ASZ, Figure 5.

4.6 Ground Motion Models

The attenuation characteristics of the crust are of utmost importance for a seismic hazard assessment as they determine how the ground motions from an earthquake decay with distance. In stable continental regions like Fennoscandia, such ground motion models are difficult to construct as the low seismicity rate of the region, the region's size and the relatively sparse seismic networks imply that there is usually very little strong motion data available. Notwithstanding, Fülöp et al. (2020) used weak motion data from events in Finland and Sweden between 2006 and 2018, with magnitudes $1.5 \leq M_L \leq 4.2$, in the development of a Fennoscandian Ground Motion Model (GMM), Fenno-G16. In order to make up for the lack of large events, they added data from an eastern Canadian subset of the NGA-East data (Goulet et al., 2014), with largest magnitude M_W 6.76, and implemented Fenno-G16 based on an adjustment of the backbone curves of the G16 equation developed by Graizer (2016) for Central and Eastern North America (CENA). The released version of Fenno-G16 contains only one branch in the logic tree and uses a $V_{S,30}$ for very hard rock of 2800 m/s.

The recently developed scaled backbone GMM for cratonic regions in ESHM20 (Weatherill and Cotton, 2020), described in Section 3.1, also partly incorporated NGA-East data and models (Goulet et al., 2018) (Goulet et al., 2021) and used the Fülöp et al. (2020) data set to aid in deciding the weights for the two main branches of cratonic (0.8) and shallow crustal (0.2) in the logic tree.

In this study, we use the two recently developed cratonic GMMs Fenno-G16 (Fülöp et al., 2020) and ESHM20 cratonic (Weatherill and Cotton, 2020) together with the shallow crustal seismicity model used for much of Europe in ESHM20 (Kotha et al., 2020). In order to study the effects of the different GMMs on the hazard estimates, we use our earthquake data set and seismic source zones and calculate hazard at a specific site for each of the following logic tree combinations:

1. The native ESHM20 logic trees for shallow-default and cratonic regions, for a $V_{S,30} = 800$ m/s, which is the ESHM20 default.
2. The native ESHM20 logic tree for shallow-default regions and the ESHM20 cratonic logic tree with $V_{S,30}$ set to 3 km/s.
3. The ESHM20 logic tree for the shallow-default regions and the single-branch Fenno-G16 GMM for the cratonic regions. $V_{S,30} = 3$ km/s for basement sites and $V_{S,30} = 800$ m/s for the sedimentary rocks in Skåne, Öland and Gotland.
4. The ESHM20 logic tree for the shallow-default regions and for the cratonic regions two branches, one with a modified ESHM20 cratonic logic tree, weight 0.6, where the 20% branch with the shallow crustal GMM has been removed, ~~weight 0.6~~, and one with the single-branch Fenno-G16, weight 0.4, see Figure 6. $V_{S,30} = 3$ km/s for the cratonic regions. Figure 6 shows the ESHM20 five branch implementation for epistemic uncertainty of the GMM, σ_μ (Weatherill and Cotton, 2020), where the ϵ_{ii} and weights assigned to each branch are determined by a discrete approximation of the Gaussian distribution depending on the number of standard deviations to include (Danciu et al., 2021; Miller and Rice, 1983). The epistemic uncertainty of the site amplification factor, $\sigma_{\mu,S}$, is similarly included with three branches, however, as we do not adjust the site amplification to $V_{S,30} = 800$ m/s for cratonic areas these branches cancel.

Figure 7 shows how the hazard curves at the location of Uppsala (Fig. 1) vary with the different GMM combinations. The mean hazard estimates are rather similar, we see that the Fenno-G16 model consistently produces slightly higher PGAs at the same probability of exceedance (PoE) than the ESHM20 models and that using a $V_{S,30}$ of 3 km/s consistently lower the hazard compared to a $V_{S,30}$ of 800 m/s, as expected. The uncertainty distribution for Fenno-G16 is generally narrower than those for the ESHM20 models, for which the lower 0.16 fractile limit extends to significantly lower PGA for the same PoE. Model 54, with the combination of Fenno-G16 and ESHM20, has a mean hazard similar to Fenno-G16 at higher PoE but moves toward the midway point between the two at higher PoE. The 0.16 and 0.84 fractile distributions behave similarly.

We choose model 54 as our preferred ground motion model for cratonic areas in this study, as it combines the two most recent cratonic GMMs, incorporates data and models from eastern North America and takes into account seismic observations in Fennoscandia. The weights between Fenno-G16 and the ESHM20 GMMs were determined such that we obtain a larger epistemic uncertainty distribution than Fenno-G16 while somewhat reducing the ESHM20 distribution to lower hazard.

Referring to the zone numbers in Figure 4, we use our cratonic model 54 ground motion logic tree for tectonic zones T1 and T4, encompassing much of Sweden, Finland and the Baltic countries. In tectonic zones T2 and T3 we use the ESHM20 shallow-default logic tree, except for in area source zones 24, 28 and 30, which all use the cratonic combination. As described in Section 2, Swedish surface rock mostly consists of Proterozoic bedrock which in places is covered by a few meters to tens of meters of glacial sediments. There have been no large scale studies of $V_{S,30}$ in Sweden but generally speaking, many single houses and larger constructions are built on bedrock or using piling to bedrock. We therefore use a $V_{S,30}$ of 3 km/s for all cratonic zones, similar to the 2800 m/s used by Fülöp et al. (2020) and corroborated by e.g. Sadeghisorkhani et al. (2020). The two Baltic islands of Öland and Gotland are covered in sand- and limestone (Rosberg and Erlström, 2019), for these areas we use the ESHM20 default of $V_{S,30} = 800$ m/s. In all zones with the shallow-default GMM we also use the default ESHM20 $V_{S,30} = 800$ m/s.

4.7 Definition of OpenQuake parameters

We use the OpenQuake(OQ) engine (v3.12) (Pagani et al., 2014) to estimate the yearly-annual probability of exceedance of peak ground acceleration at the 0.0021 and 0.0004 levels, corresponding to return periods of 475 and 2500 years. The OQ-engine uses the classical PSHA approach based on the methodology proposed by Cornell (1968) and McGuire (1976). The engine allows for flexibility in modelling seismic sources, using predefined GMMs or implementing new GMMs, and characterizes epistemic uncertainty through a logic tree. The engine incorporates the OpenSHA (Field et al., 2003) calculation structures and workflows, which we use on an equally spaced 5 km calculation grid.

Area sources were defined as per the OQ NRML schema for each ASZ (Schorlemmer et al., 2011). We follow ESHM20 and define the minimum magnitude

The minimum magnitude, M_{\min} , used for the hazard computation, as 4.5 for all sources, as that is taken as the magnitude where damages may start to appear. ESHM20 uses a M_{\min} of 4.5 but we follow Mosca et al. (2022) and use M_{\min} 4.0 in order to account for the impulsive nature and high-frequency content of smaller events in a slowly attenuating crust, such as the

Swedish. The damage potential of a M_W 4.0 earthquake was demonstrated in the Folkestone 2007 M_W 4.0 event, which caused significant non-structural but also some structural damage (Sargeant et al., 2008).

We define a depth distribution where ruptures take place, using 5, 10, 20 and 30 km depth weighted as 0.15, 0.35, 0.35 and
490 0.15. This accounts for the fact that although most events occur down to about 20 km depth, some areas have significantly
deeper earthquakes (e.g. Veikkolainen et al., 2017). Point ruptures are used as we ~~include no~~ do not include specific faults in
the model ~~and for the added benefit of reduced calculation times~~. Truncated Gutenberg-Richter distributions are defined by the
lower and upper magnitude bounds M_{\min} and M_{\max} , and the previously calculated a- and b- values with their uncertainties are
used in different branches of the logic tree. Each seismic source area requires the definition of orientations and faulting styles
495 of ruptures, quantified by the strike, dip and rake of nodal planes. We use the strike and dip values adopted by ESHM20 in their
calculations and choose a strike of 0 degrees, a dip of 90 degrees. We define two rake values of 0 or 180 degrees, each with a
weight of 0.5 in the probability distribution

The calculation parameters for the OQ engine are summarized as follows: twelve end-branches for area source models of
the shallow crust and cratonic regions; 18,072 equally spaced grid points (5 km spacing) defined on land within Swedish
500 land borders; a $V_{S,30} = 3$ km/s; point ruptures; a weighted depth distribution; predefined nodal plane orientations, vertical
~~plane~~ planes striking north with a rake of 0 or 180; 25 intensity measures that cover PGA ground motion discretization levels
between 0.005 and 3 g; and calculations for the complete enumerated logic tree.

5 Results

We present ground motion exceedance levels for mainshock earthquakes in terms of peak ground acceleration (PGA), using
505 our source ~~model~~ and ground motion model logic tree implementations as discussed in Sections 4.5 and 4.6. PSHA results are
commonly presented as hazard maps showing PGA with a 10% probability of exceedance (PoE) in 50 years, as that level is
used for building codes, e.g. in Eurocode 8. A 10% exceedance level in 50 years is equivalent to a statistical (Poissonian) return
period of 475 years. ESHM20 uses annual PoEs, where $2.1 \cdot 10^{-3}$ is equivalent to 10% in 50 years and $4 \cdot 10^{-4}$ equivalent to the
previously used 2% in 50 years, slightly altering the return period from 2475 to 2500 years. We follow ESHM20 and present
510 our results as hazard maps for return periods of 475 and 2500 years, and as hazard curves for four seismically active regions in
Sweden. Ground motion exceedance levels for spectral acceleration periods of 0.2s and 1s for a return period of 475-years are
presented in the supplementary.

5.1 Seismic hazard ~~map~~ maps for Sweden

Seismic hazard maps for Sweden with 475 and 2500 year return times are shown in Figure 8. As usual in probabilistic seismic
515 hazard assessment, hazard closely follows the observed earthquake distribution. The maps show the mean horizontal PGA, and
we see that for a 475 year return period the highest hazard is estimated in the northernmost part of Sweden, in the area of the
post-glacial faults and at the northern shores of the Bay of Bothnia, where hazard reaches ~~0.05~~ 0.06g. Much of the northeastern
coast of Sweden shows a mean PGA of 0.03 - 0.04g and the southwest of Sweden shows a mean PGA 0.02 - 0.03g. The

southeast and northwest of Sweden have an estimated mean hazard of less than 0.02g. For the 2500 year hazard map, hazard is similarly distributed, but the highest mean PGA is now 0.15g in the northeast of Sweden, followed by the northeast coast at about 0.1 - 0.125g and the southwest of Sweden between 0.075 - 0.1g. As before, the southeastern and the northwestern regions show the lowest hazard of less than 0.05g. The spectral acceleration maps, in supplementary Figure A3, show similar results as the PGA map for the spatial distribution of hazard. We find, however, that the mean hazard at SA(0.2s) is generally higher than the mean PGA by about 0.01g in all of the country. At SA(1.0s) the mean hazard is everywhere below 0.01g.

We note that the earthquake activity in the Oslo graben in Norway (ASZ 25), and even the activity on the Norwegian west coast, affect the hazard levels in southwestern Sweden to some degree. Interestingly, we do not see the same influence from the Nordland region in Norway ~~where seismic activity~~ (ASZ 4) into Sweden, although seismic activity there is high and it is the region where the largest known earthquake in Fennoscandia occurred in 1819. This is likely due to the relatively high b-value we estimate for ASZ 4 (b = 0.933), and the fact that many of the events occur in swarms which are removed in the declustering.

We complement our maps with a disaggregation analysis for the location in the northeast with the highest hazard in our study and the location in the southwest where Wahlström and Grünthal (2001) found the highest hazard in Sweden. Figure 9 shows that the main contribution to the hazard is primarily from small earthquakes at 10 - 50 km distance, similar for both locations. We note that our study predicts a higher hazard contribution from larger earthquakes than does the Wahlström and Grünthal (2001) analysis.

5.2 Hazard curves for seismogenic areas

The regions in Sweden with the highest seismic activity can broadly be defined as: the southwest around Lake Vänern, the northeast coast and the post-glacial fault province in the north. In Figure 10 we illustrate the seismic hazard at four locations in these regions, using hazard curves with the annual probability of exceedance versus PGA, ~~at four locations in these regions~~: the post-glacial Pärvie and Burträsk faults, the Hälsingland area around Hudiksvall and Västergötland south of Lake Vänern, see stars in Figure 1. We see that the hazard curves are similar for all four regions, the Pärvie and Burträsk faults having slightly higher short term PGA estimates and Västergötland the lowest hazard overall, although all four are within the uncertainty limits of each other. We restrict our curves to a lower limit of an annual PoE of 10^{-5} , with the 0.16 and 0.84 fractile estimates ranging between 0.3g and 0.9g at this limit. Comparing to the Uppsala curves in Figure 7, we note that the short term hazard is similar in Uppsala and Västergötland, but that the longer term hazard is significantly lower in Uppsala.

6 Discussion

The result of our seismic hazard assessment, as illustrated in the maps in Figure 8, differ from many of the previous assessments for Sweden in that the estimated hazard is largest in ~~northernmost Sweden~~ the far north. This is due to the new emerging understanding of the seismic activity on the post-glacial faults (PGFs) in northernmost Fennoscandia (Lindblom et al., 2015). The expansion of the Swedish national seismic network in the early 2000s, from 2 to 22 stations north of 63.5° latitude (Lund et al., 2021), has enabled observations of previously unknown seismic activity rates on these faults (Lindblom et al., 2015).

Almost 3700 earthquakes have been observed in the region in the last two decades (Figure 2), all but two have magnitudes below 3 which is consistent with the few observations prior to the network expansion. The very low population density and limited reporting possibilities in earlier times probably explains the almost complete lack of observations in e.g. the Pärvie area prior to 1967. Unfortunately, there are no conclusive data on current slip rates on the PGFs. For example, the InSAR study by Mantovani and Scherneck (2013) inferred vertical displacements at a few segments along the Pärvie fault, but did not conclude any slip rates. This complicates the inclusion of the post-glacial faults explicitly in a PSHA.

The inference that the last 20 years of earthquake data in northern Sweden indicates that the area has [a relatively high seismic hazard](#) is interesting in the light of recent work on the PGFs in the region. Smith et al. (2021) found repeated ruptures on the Lainio PGF, with the latest recurring after deglaciation, and ~~(Olesen et al., 2021)~~ [Olesen et al. \(2021\)](#) identified ruptures of magnitude 7 as recently as 700 years ago on the Stuoragurra fault in northern Norway. Taken together, all observations suggest that seismic hazard should be taken into careful consideration in the region, especially since the region is home to major mining operations, large hydroelectric power dams and an increasing number of wind energy installations.

6.1 Estimation of recurrence parameters

With Fennoscandia being a low-seismicity region, estimating the recurrence parameters is a major challenge in the seismic hazard assessment process. The problem is multifaceted and involves the low seismicity rate, the few large events and the magnitude homogenization process. As the rate of earthquakes large enough to be felt is low (Lund et al., 2021), there is little data prior to the installation of an improved seismic network in the 1960s. That network consisted of only six analogue stations, so the more significant increase in earthquake observations came with the modern SNSN, from the year 2000 onward, when the completeness magnitude fell below M1 (Figure 3) for a large part of the country (Lund et al., 2021). Many of the observed larger events occurred in earlier times and therefore have macroseismic magnitudes with significant uncertainty. Later events have a variety of local magnitudes, which complicates the magnitude homogenization procedure (Uski et al., 2015). In addition, the magnitude estimation for the many small events in our dataset comes with appreciable uncertainty, e.g. due to low station density, which further complicates the magnitude homogenization procedure.

As discussed in Section 4.4, we use the Weichert (1980) method, as implemented by OpenQuake, to calculate the recurrence parameters, using data reaching back to 1875. We also use Aki's method (Aki, 1965), as revised by Tinti and Mulargia (1987), with the post-2000 data to add confidence in the results. We found that varying M_{\max} during parameter estimation has no significant effect on the results. However, the results are sensitive to variations in the magnitude of completeness, a change in M_c by 0.2 can lead to a significant change in a- or b-value, beyond the formal uncertainties of the original results. The magnitude of completeness is non-trivial to estimate, manually or using some curvature method, especially for the very sparse historical data. It is thus likely that uncertainties in a- and b-values are sometimes larger than the formal estimates, which led us to set a minimum b-value uncertainty to 0.01. This was needed for tectonic zone T1, which contains 7347 events above M_c and therefore produces a very small formal uncertainty.

We note that in their investigation of the sensitivity of site-specific PSHA in Finland to variations in the input parameters, Fülöp et al. (2023) found that M_{\max} variations do not affect the PGA results significantly but is more important at low ground

585 motion frequencies at an annual frequency of exceedance of 10^{-5} . ~~As expected, they~~ They also point out that a- and b-value variations have a large effect on the results, in agreement with our conclusions.

6.2 Comparisons with previous seismic hazard studies

As discussed above, the most significant difference between our seismic hazard model and previous studies is the spatial distribution of hazard in northern Sweden. The Wahlström and Grünthal (2001) study found the largest hazard in Sweden
590 focused in the area surrounding the 1904 M_W 5.4 earthquake on the west coast, with a median PGA of 0.03 - 0.035g for a 475 year return period. This is slightly higher than our estimate of about 0.025g in that area. Similar to us, they found that southeastern and northwestern Sweden, east of the mountain range, has the lowest hazard, below 0.015g. Along the northeast coast they estimate a PGA of 0.015 - 0.02g, lower than our approximately 0.03g, and in the northernmost region of Sweden their 475 year PGA estimates vary between 0.01 - 0.02g, significantly lower than our 0.03 - ~~0.05~~0.06g. We note that Wahlström
595 and Grünthal (2001) use the median hazard for their estimates. As we use mean estimates, our hazard values are likely to be generally somewhat higher than theirs.

The Mäntyniemi et al. (2001) hazard map is similar to the one from Wahlström and Grünthal (2001). They find high hazard in the southwest and along the northeast coast, with the 475 year PGA reaching 0.02 - 0.025g in both regions, similar to our results. In the north, their hazard estimate is lower than ours, at 0.01 - 0.015g. We note that both (Wahlström and Grünthal,
600 2001) and (Mäntyniemi et al., 2001) estimate a higher hazard in the northern Swedish mountain region than us, indicating that their area source zone definition is such that the high seismicity in Norwegian Nordland produces significant hazard in Sweden in their model, or that they estimate lower b-values than we do in that region.

The 2013 European seismic hazard model (ESHM13, Woessner et al., 2015) found the highest hazard in Sweden to be focused in a narrow region across the southernmost part of the country and up along the west coast, with a maximum PGA for
605 475 year return period at approximately 0.025g in the south. This is significantly more than our estimate of less than 0.01g in that area. Along the southwest coast of Sweden the hazard in ESHM13 is about 0.02g, similar to ours, while the rest of the country has a hazard between 0.001 - 0.02g, again significantly less hazard in the north than in our model.

The most recent iteration of the European seismic hazard model (ESHM20, Danciu et al., 2021) changed the distribution of hazard in Sweden significantly as compared to ESHM13 (see Lund et al. (2024) for an in-depth comparison). Hazard estimates
610 were lower in the south and southwest and higher in south-central parts of the country and up the northeast coast, following the coastline around the northern Bay of Bothnia. This agrees with the distribution of hazard in our model, except that in our model the hazard is lower in the southeast. We illustrate the differences between our model and ESHM20 for 475 and 2500 years return periods in Figure 11, which shows that our model produces higher hazard than ESHM20 for both return periods in most of the country. Again, the most notable difference is in the north, where ESHM20 uses the recurrence parameters
615 from the large tectonic zones but where we have very much more data (Figure 2), such that we can better constrain the a- and b-values. Comparing recurrence parameters we note that the ESHM20 area source zone NOAS376 in northern Fennoscandia, encompassing many of the PGFs, is approximately 30% larger in size than our ASZs 13, 14 and 15 combined. The ESHM20 b-value is 0.91 ± 0.032 in that area, which is within the one-sigma uncertainties of the b-values in our ASZ 13 and 14, and just

above the uncertainty limits of ASZ 15, see Table 2. The ESHM20 a-value 1.584 ± 0.0059 is re-scaled from the tectonic zone
620 and significantly lower, considering the much larger area, than our a-values around 1.9 - 2.0 (Table 2) for the individual zones
13, 14 and 15. We see in Figure 11 that our PGA is more than 0.04g higher for a 475 year return period than the ESHM20
estimate in some areas in the north. For a 2500 year return period the difference is 0.125g. Further south along the northeast
coast our hazard is still higher by some 0.02g, a difference which decreases towards the southeast and along the southeast coast
the hazard in our model is smaller than in ESHM20. In the southwest, our model estimates higher hazard than ESHM20 by
625 about 0.015g. We note that ESHM20 and our model agree well along the northwestern border, where none of the models see
much influence of the Norwegian coastal seismicity.

6.3 Seismic hazard and risk in Sweden

Comparing to surrounding regions, the seismic hazard we find in northern Sweden is only surpassed by the hazard in western
Norway around Bergen and in the [Rana-Nordland](#) region in northwestern Norway. Our new results for northern Sweden will
630 be compared to results obtained in neighbouring countries in order to calibrate the hazard along the borders. Probabilistic
seismic hazard projects are ongoing in both Norway and Finland, and we aim to further the work on a joint homogenized
Fennoscandian seismic hazard model in the near future.

Our results show that the hazard correlates well with the observed seismicity. We note, however, that most of the seismic
activity is of low magnitude and that there is not a clear correlation between the observed larger events, above e.g. M_W 4,
635 and more intense microearthquake activity. The Tornquist zone in southern Sweden, for example, has experienced five events
with $M_W \geq 3.7$ in the last century but has a low seismicity rate whereas the Burträsk fault, the most seismically active area in
Sweden, has ~~seen no~~ [only seen two](#) such events during that time. There was a M_W 3.9 event in west-central Sweden in 2014
in an area with only a few prior, small events (Lund et al., 2014), and the large magnitude 5 events in Kaliningrad in 2004
(Gregersen et al., 2007) occurred in an area with very few previously observed events. These surprise events, and the sometimes
640 poor correlation between the rate of small events to large, points to the difficulty of PSHA in stable continental regions. Adding
to the problem is the largely unknown level of temporal and spatial stationarity of earthquakes in these regions.

The large post-glacial faults of northern Fennoscandia pose an interesting problem to seismic hazard assessment. Previously
inferred to have ruptured only once, at the time of final deglaciation of the region (e.g. Lagerbäck and Sundh, 2008), the main
question ~~concerned~~ [has been](#) whether or not the faults could rupture again without an intervening major glaciation. Glacial
645 isostatic adjustment modelling (see review in e.g. Lund, 2015) shows that glacially induced stresses combine with the tectonic
stress state and bring the PGFs towards instability at deglaciation. Due to the slow strain accumulation in the Fennoscandian
Shield, repeated large ruptures on the same faults segments have been considered unlikely with repeat times less than 100,000
years (Korja and Kosonen, 2015). However, as alluded to above, more recent investigations have found that the PGF systems
have seen multiple ruptures along the faults and also multiple ruptures at the same location, see reviews for Fennoscandia in
650 Steffen et al. (2021). Including the PGFs in a PSHA is currently only possible as an increase in M_{max} for area source zones
including PGFs ~~with a weight that~~ [since it has not been possible to detect any ongoing surface slip on the faults. The weight
on such a high](#) M_{max} [branch in the logic tree](#) is difficult to assess due to the current lack of data. We have seen that ~~this a high~~

M_{\max} branch does not have a large effect on the estimated PGA but is more likely to affect the longer periods of ground motion (Fülöp et al., 2023).

655 A more appropriate mechanism, at this time, to take the PGFs into account is through deterministic seismic hazard assessment (DSHA). The faults are mapped in high resolution (e.g. Smith et al., 2021) and the depth extent of current seismicity is known for some PGFs (e.g. Lindblom et al., 2015), making it possible to use scaling relations to estimated potential rupture magnitudes. While a deterministic scenario may have a very low probability of occurrence, a DSHA will provide additional insights from a risk reduction perspective and potentially highlight events that may dominate the seismic risk in a certain area
660 or for a specific site, and therefore must be considered (McGuire, 2001). Performing site-specific seismic hazard, and risk, assessment in the post-glacial fault province in northern Fennoscandia for critical infrastructure, such as dams where low annual probabilities must be considered, a combined PSHA/DSHA approach may be the most appropriate.

7 Summary

This study provides a probabilistic seismic hazard model of Sweden based on updated earthquake catalogues from the Swedish
665 national seismic network (Lund et al., 2021) and neighbouring networks, and recent results and methods from the European seismic hazard model ESHM20 (Danciu et al., 2021). The hazard is calculated using the OpenQuake engine (Pagani et al., 2014), using earthquakes since 1875 represented through 31 area sources zones, and recently developed ground motion models (Fülöp et al., 2020; Weatherill and Cotton, 2020; Kotha et al., 2020). A weighted logic tree approach is used to represent the uncertainties in the recurrence parameter estimates, M_{\max} and the ground motion models. The resulting hazard estimates are
670 calculated as peak ground-acceleration (PGA) values and mapped in Figure 8 for 475 and 2500 years return periods.

The updated earthquake data allow us to use a large number of smaller events that have not previously been available, or used, in seismic hazard assessment. This makes it possible to better constrain the recurrence parameters (a- and b-values) in much of the country, especially for the post-glacial fault province in northern Sweden.

We find a seismic hazard distribution in Sweden which is significantly different to that of earlier studies. The new data
675 from the north indicates that hazard in northernmost Sweden is higher than elsewhere in country, both at 475 and 2500 years return period. For a return period of 475 years, mean PGA along the post-glacial faults and the northern coastal tip of the Bay of Bothnia is estimated to be about 0.04 to ~~0.05~~0.06g. For the rest of the north-eastern coast, mean PGA is estimated to be 0.02-0.03g. The lowest PGA estimates in the country, 0.02g and lower, are found in the ~~south-eastern and north-western parts of the country~~southeast and northwest. Similar trends are seen for the 2500 year PGA estimates, with the highest mean PGA
680 of 0.15g seen in the ~~north-east~~northeast, 0.1-0.125g for the post-glacial faults and the north-eastern coast, 0.075-0.1g for the ~~south-west~~southwest and less than 0.05g for the ~~south-east and the north-west~~southeast and the northwest. Hazard curves for four locations in some the most active areas of the country, the Pärvie and Burträsk faults, the Hälsingland area and south of Lake Vänern, show similar hazard in these areas, with slightly higher hazard along the post-glacial faults.

Compared to ESHM20 (Danciu et al., 2021), our model shows somewhat higher hazard in the more seismically active areas
685 of Sweden but slightly lower hazard in the less active areas. The main difference is in the north, where we find significantly

higher hazard. We note that our declustered, homogenized earthquake catalogue contains 19,943 events, with magnitudes down to just below zero, compared to 360 events in the same area for ESHM20, all with $M_W \geq 3.5$.

Challenges remain in seismic hazard estimation for Fennoscandia. There are significant uncertainties in magnitude homogenization, for the smaller magnitude events compared to the larger, in the estimate of recurrence parameters, as larger events occur infrequently, and in the estimation of ground motions models for the Fennoscandian Shield, as sparse to non-existing data for larger events and short distances implies that both epistemic and aleatory uncertainties are difficult to estimate. On a larger scale, the occurrence of large events in areas with little prior seismicity and the uncertainties surrounding the potential for large earthquakes on the post-glacial faults in northern Fennoscandia pose problems for seismic hazard and risk assessment, and require continued studies.

695 *Data availability.* TEXT

The earthquake catalogue was prepared based on data obtained from the Swedish ~~national seismic network~~ [National Seismic Network](#) (SNSN, 1904), the Norwegian ~~national seismic network~~ [National Seismic Network](#) (University of Bergen, 1982), the Finnish ~~national seismic network~~ [National Seismic Network](#) (FNSN, 1980), The Estonian seismic network (Geological Survey of Estonia (GSE), 1998) and the Geological Survey of Denmark and Greenland (GEUS, 2023).

700 *Code and data availability.* TEXT

The QGIS (<https://www.qgis.org/en/site/>) software was used to delineate and visualize the zones. The OpenQuake engine (<https://github.com/gem/oq-engine>) and its included suite of data preparation programs were used to calculate recurrence parameters, create input files, and run the hazard calculations themselves. The hazard maps were plotted using the plotting function build into OpenQuake and pygmt, and the hazard curves were plotted using matplotlib.

705 *Author contributions.* TEXT

Niranjan Joshi: Data preparation, processing, investigation and visualization, writing. Björn Lund : Conceptualization, methodology, supervision, data preparation and analysis, review, editing and writing. Roland Roberts: Conceptualization, methodology, supervision, review, editing.

Competing interests. TEXT

710 The authors declare no competing interests.

Acknowledgements. The authors gratefully acknowledge discussions with Fennoscandian colleagues and members of the ESHM20 team. We also thank the FENCAT team at the Institute of Seismology, Helsinki University, and the Norwegian National Seismic Network for earthquake data and advice on magnitudes.

References

- 715 Ahjos, T. and Uski, M.: Earthquakes in northern Europe in 1375–1989, *Tectonophysics*, 207, 1–23, [https://doi.org/10.1016/0040-1951\(92\)90469-M](https://doi.org/10.1016/0040-1951(92)90469-M), 1992.
- Aki, K.: Maximum likelihood estimate of b in the formula $\log N = a - bM$ and its confidence limits, *Bull. Earthquake Res. Inst., Tokyo Univ.*, 43, 237–239, <https://cir.nii.ac.jp/crid/1573387450038851840>, 1965.
- Baker, J., Bradley, B., and Stafford, P.: *Seismic hazard and risk analysis*, Cambridge University Press, Cambridge, England, 2021.
- 720 Beirlant, J., Kijko, A., Reynkens, T., and Einmahl, J.: Estimating the maximum possible earthquake magnitude using extreme value methodology, *Nat. Hazards*, 98, 1091–1113, <https://doi.org/10.1007/s11069-017-3162-2>, 2019.
- Bingen, B., Viola, G., Möller, C., Vander Auwera, J., Laurent, A., and Yi, K.: The Sveconorwegian orogeny, *Gondwana Reserach*, 90, 273–313, <https://doi.org/https://doi.org/10.1016/j.gr.2020.10.014>, 2021.
- Bodare, A. and Kulhánek, O.: Dam safety. Earthquake hazard for dams in Sweden, *Tech. Rep. 06:72, Elforsk*, <https://energiforsk.se/>, 2006.
- 725 Bogdanova, S., Bingen, B., Gorbatshev, R., Kheraskova, T., Kozlov, V., Puchkov, V., and Volozh, Y.: The East European Craton (Baltica) before and during the assembly of Rodinia, *Precambrian Research*, 160, 23–45, <https://doi.org/10.1016/j.precamres.2007.04.024>, testing the Rodinia Hypothesis: Records in its Building Blocks, 2008.
- Bungum, H., Pettenati, F., Schweitzer, J., Sirovich, L., and Faleide, J. I.: The 23 October 1904 MS 5.4 Oslofjord Earthquake: Reanalysis Based on Macroseismic and Instrumental Data, *Bull. Seismol. Soc. Am.*, 99, 2836–2854, <https://doi.org/10.1785/0120080357>, 2016.
- 730 Båth, M.: Seismic risk in Fennoscandia, *Tectonophysics*, 57, 285–295, [https://doi.org/10.1016/0040-1951\(79\)90152-5](https://doi.org/10.1016/0040-1951(79)90152-5), 1979.
- Båth, M.: Intensity relations for Swedish earthquakes, *Tectonophysics*, 67, 163–173, [https://doi.org/10.1016/0040-1951\(80\)90171-7](https://doi.org/10.1016/0040-1951(80)90171-7), 1980.
- Bödvarsson, R., Lund, B., Roberts, R., and Slunga, R.: Earthquake activity in Sweden. Study in connection with a proposed nuclear waste repository in Forsmark or Oskarshamn, *Tech. rep., Swedish Nuclear Fuel and Waste Management Co.*, 2006.
- Calais, E., Camelbeeck, T., Stein, S., Liu, M., and Craig, T.: A new paradigm for large earthquakes in stable continental plate interiors, *Geophysical Research Letters*, 43, 10 621–10 637, <https://doi.org/10.1002/2016GL070815>, 2016.
- 735 Corfu, F., Gasser, D., and Chew, D. M.: New perspectives on the Caledonides of Scandinavia and related areas: introduction, *Geological Society, London, Special Publications*, 390, 1–8, <https://doi.org/10.1144/sp390.28>, 2014.
- Cornell, C. A.: Engineering seismic risk analysis, *Bull. Seismol. Soc. Am.*, 58, 1583–1606, <https://doi.org/10.1785/BSSA0580051583>, 1968.
- Danciu, L., Şeşetyan, K., Demircioglu, M., Gülen, L., Zare, M., Basili, R., Elias, A., Adamia, S., Tsereteli, N., Yalçın, H., Utkucu, M., Khan, M. A., Sayab, M., Hessami, K., Rovida, A. N., Stucchi, M., Burg, J.-P., Karakhanian, A., Babayan, H., Avanesyan, M., Mammadli, T., Al-Qaryouti, M., Kalafat, D., Varazanashvili, O., Erdik, M., and Giardini, D.: The 2014 Earthquake Model of the Middle East: seismogenic sources, *Bulletin of Earthquake Engineering*, 16, 3465–3496, <https://doi.org/10.1007/s10518-017-0096-8>, 2017.
- 740 Danciu, L., Nandan, S., Reyes, C. G., Basili, R., Weatherill, G., Beauval, C., Rovida, A., Vilanova, S., Sesetyan, K., Bard, P.-Y., Cotton, F., Wiemer, S., and Giardini, D.: The 2020 update of the European Seismic Hazard Model: Model Overview, *EFEHR Technical Report 001, v1.0.0*, <https://doi.org/10.12686/a15>, 2021.
- Eggertsson, G., Lund, B., Roth, M., and Schmidt, P.: Earthquake or blast? Classification of local-distance seismic events in Sweden using fully connected neural networks, *Geophy. J. Int.*, 236, 1728–1742, <https://doi.org/10.1093/gji/ggae018>, 2024.
- Elliott, J. R.: Earth Observation for the Assessment of Earthquake Hazard, Risk and Disaster Management, *Surveys in Geophysics*, 41, 1323–1354, <https://doi.org/10.1007/s10712-020-09606-4>, 2020.
- 750 EMSC: Euro-Mediterranean Seismological Centre Newsletter, 15, 3pp, 1999.

- England, P. and Jackson, J.: Uncharted seismic risk, *Nature Geoscience*, 4, 348–349, <https://doi.org/10.1038/ngeo1168>, 2011.
- Field, E. H., Jordan, T. H., and Cornell, C. A.: OpenSHA: A Developing Community-modeling Environment for Seismic Hazard Analysis, *Seismol. Res. Lett.*, 74, 406–419, <https://doi.org/10.1785/gssrl.74.4.406>, 2003.
- Finnish National Seismic Network: Interactive map tool for browsing earthquake data, <https://www.seismo.helsinki.fi/EQ-search/query.php>, 2023.
- FNSN: Institute of Seismology, University of Helsinki, HE: The Finnish National Seismic Network, GFZ Data Services, <https://doi.org/10.14470/UR044600>, 1980.
- Fülöp, L., Jussila, V., Aapasuo, R., Vuorinen, T., and Mäntyniemi, P.: A Ground-Motion Prediction Equation for Fennoscandian Nuclear Installations, *Bull. Seismol. Soc. Am.*, 110, 1211–1230, <https://doi.org/10.1785/0120190230>, 2020.
- 760 Fülöp, L., Mäntyniemi, P., Malm, M., Toro, G., Crespo, M., Schmitt, T., Burck, S., and Välikangas, P.: Probabilistic seismic hazard analysis in low-seismicity regions: an investigation of sensitivity with a focus on Finland, *Natural Hazards*, 116, 111–132, <https://doi.org/https://doi.org/10.1007/s11069-022-05666-4>, 2023.
- Gardner, J. and Knopoff, L.: Is the sequence of earthquakes in Southern California, with aftershocks removed, Poissonian?, *Bull. Seismol. Soc. Am.*, 64, 1363–1367, 1974.
- 765 Geffers, G.-M., Main, I., and Naylor, M.: Biases in estimating b-values from small earthquake catalogues: how high are high b-values?, *Geophys. J. Int.*, 229, 1840–1855, <https://doi.org/10.1093/gji/ggac028>, 2022.
- Geological Survey of Estonia (GSE): Estonian Seismic Network, <https://www.egt.ee/en/fields-activity-and-objectives/geology-and-environment/seismic-research-and-monitoring>, 1998.
- GEUS: Registrerede jordskælv i Danmark, <https://www.geus.dk/natur-og-klima/jordskaelv-og-seismologi/registrerede-jordskaelv-i-danmark>, 2023.
- 770 Goulet, C., Kishida, T., Ancheta, T., Cramer, C., Darragh, R., Silva, W., Hashash, Y., Harmon, J., Stewart, J., Wooddell, K., and Young, R. R.: PEER NGA-East database, Tech. Rep. 2014/17, Pacific Earthquake Engineering Research Center, University of California, Berkeley, USA, 2014.
- Goulet, C., Bozorgnia, Y., Abrahamson, N., Kuehn, N., Atik, L. A., Youngs, R., Graves, R., and Atkinson, G.: Central and Eastern North America ground-motion characterisation: NGA-east final report, Tech. Rep. 2018/08, Pacific Earthquake Engineering Research Center, University of California, Berkeley, USA, 2018.
- 775 Goulet, C. A., Bozorgnia, Y., Kuehn, N., Al Atik, L., Youngs, R. R., Graves, R. W., and Atkinson, G. M.: NGA-East Ground-Motion Characterization model part I: Summary of products and model development, *Earthquake Spectra*, 37, 1231–1282, <https://doi.org/10.1177/87552930211018723>, 2021.
- 780 Graizer, V.: Ground-motion prediction equations for Central and Eastern North America, *Bull. Seismol. Soc. Am.*, 106, 1600–1612, <https://doi.org/10.1785/0120150374>, 2016.
- Gregersen, S., Wiejacz, P., Dębski, W., Domanski, B., Assinovskaya, B., Guterch, B., Mäntyniemi, P., Nikulin, V., Pacesa, A., Puura, V., Aronov, A., Aronova, T., Grünthal, G., Husebye, E., and Sliupa, S.: The exceptional earthquakes in Kaliningrad district, Russia on September 21, 2004, *Phys. Earth Planet. Int.*, 164, 63–74, <https://doi.org/10.1016/j.pepi.2007.06.005>, 2007.
- 785 Gregersen, S., Lindholm, C., Korja, A., Lund, B., Uski, M., Oinonen, K., Voss, P. H., and Keiding, M.: Seismicity and Sources of Stress in Fennoscandia, p. 177–197, Cambridge University Press, <https://doi.org/10.1017/9781108779906.014>, 2021.
- Grünthal, G. and Group, T. G. R. W.: Seismic hazard assessment for Central, North and Northwest Europe: GSHAP Region 3, *Annals of Geophysics*, 42, <https://doi.org/10.4401/ag-3783>, 1999.

- Grünthal, G., Wahlström, R., and Stromeyer, D.: The SHARE European Earthquake Catalogue (SHEEC) for the time period 1900–2006 and its comparison to the European-Mediterranean Earthquake Catalogue (EMEC), *J. Seismol.*, 17, 1339–1344, <https://doi.org/10.1007/s10950-013-9379-y>, 2013.
- Gutenberg, B. and Richter, C. F.: Frequency of earthquakes in California, *Bull. Seismol. Soc. Am.*, 34, 185–188, <https://doi.org/10.1785/BSSA0340040185>, 1944.
- Hanks, T. and Kanamori, H.: A moment magnitude scale, *J. Geophys. Res.*, 84, 2348–2350, <https://doi.org/10.1029/JB084iB05p02348>, 1979.
- Hashash, Y. M. A., Ilhan, O., Harmon, J. A., Parker, G. A., Stewart, J. P., Rathje, E. M., Campbell, K. W., and Silva, W. J.: Nonlinear site amplification model for ergodic seismic hazard analysis in Central and Eastern North America, *Earthquake Spectra*, 36, 69–86, <https://doi.org/10.1177/8755293019878193>, 2020.
- Johnston, A., Coppersmith, K., Kanter, L., and Cornell, C.: The Earthquakes of Stable Continental Regions - Assessment of large earthquake potential, Tech. Rep. TR-102261-V1, Electric Power Research Institute, Palo Alto, California, <http://www.epri.com/search/Pages/results.aspx?k=stable%20continental%20regions>, (last accessed August 2023), 1994.
- Kijko, A. and Graham, G.: Parametric-historic procedure for probabilistic seismic hazard analysis. Part I: Estimation of maximum regional magnitude M_{max} , *Pure Appl. Geophys.*, 152, 413–442, 1998.
- Kijko, A. and Graham, G.: Parametric-historic procedure for probabilistic seismic hazard analysis. Part II: Assessment of seismic hazard at specified site, *Pure Appl. Geophys.*, 154, 1–22, 1999.
- Kijko, A. and Singh, M.: Statistical Tools for Maximum Possible Earthquake Magnitude Estimation, *Acta Geophysica*, 59, 674–700, <https://doi.org/10.2478/s11600-011-0012-6>, 2011.
- Kijko, A., Skordas, E., Wahlström, R., and Mäntyniemi, P.: Maximum Likelihood Estimation of Seismic Hazard for Sweden, *Natural Hazards*, 7, 41–57, <https://doi.org/10.1007/BF00595678>, 1993.
- Kjellén, R.: Sveriges jordskalf: Försök till en seismisk landsgeografi, Zachrissons Boktryckeri AB, Göteborg, Sweden, in Swedish, 1910.
- Korja, A. and Kihlman, S.: Seismic source areas in central Fennoscandia, Tech. Rep. S-64, Institute of Seismology, University of Helsinki, Helsinki, Finland, 2016.
- Korja, A. and Kosonen, E.: Seismotectonic framework and seismic source area models in Fennoscandia, northern Europe, Tech. Rep. S-63, Institute of Seismology, University of Helsinki, Helsinki, Finland, 2015.
- Kortström, J., Uski, M., and Tiira, T.: Automatic classification of seismic events within a regional seismograph network, *Computers & Geosciences*, 87, 22–30, <https://doi.org/https://doi.org/10.1016/j.cageo.2015.11.006>, 2016.
- Kotha, S. R., Weatherill, G., Bindi, D., and Cotton, F.: A regionally-adaptable ground-motion model for shallow crustal earthquakes in Europe, *Bulletin of Earthquake Engineering*, 18, 4091–4125, <https://doi.org/10.1007/s10518-020-00869-1>, 2020.
- Lagerbäck, R. and Sundh, M.: Early Holocene faulting and paleoseismicity in northern Sweden, Tech. Rep. C 836, Geological Survey of Sweden, Uppsala, Sweden, 2008.
- Larsson, J. and Larsson, E.: ESHM13 – En ny PSHA-modell för Europa betraktad ur ett svenskt perspektiv, Tech. Rep. 2018:27, Swedish Radiation Safety Authority (SSM), 2018.
- Lindblom, E., Lund, B., Tryggvason, A., Uski, M., Bödvarsson, R., Juhlin, C., and Roberts, R.: Microearthquakes illuminate the deep structure of the endglacial Pärvie fault, northern Sweden, *Geophysical Journal International*, 201, 1704–1716, <https://doi.org/10.1093/gji/ggv112>, 2015.
- Lomnitz, C.: Global tectonics and earthquake risk, *Developments in geotectonics*, Elsevier Scientific Pub. Co., <https://cir.nii.ac.jp/crid/1130282272046071680>, 1974.

- Lund, B.: Palaeoseismology of glaciated terrain, in: *Encyclopedia of Earthquake Engineering*, edited by Beer, M., Kougiumtzoglou, I., Patelli, E., and Au, S.-K., pp. 1765–1779, Springer, Berlin Heidelberg, https://doi.org/10.1007/978-3-642-36197-5_25-1, 2015.
- Lund, B., Shomali, H., Buhcheva, D., Högdahl, K., Tryggvason, A., Bödvarsson, R., Schmidt, P., and Jørgensen, P.: The 2014 magnitude 4.1 Sveg earthquake, *Nordic Seminar on Seismology*, Visby, Sweden, 8-10 October, 2014.
- 830 Lund, B., Roberts, R., and Smith, C.: Review of paleo-, historical and current seismicity in Sweden and surrounding areas with implications for the seismic analysis underlying SKI report 92:3, *Tech. Rep. 2017:35*, Swedish Radiation Safety Authority (SSM), 2017.
- Lund, B., Schmidt, P., Shomali, Z. H., and Roth, M.: The Modern Swedish National Seismic Network: Two Decades of Intraplate Microseismic Observation, *Seismol. Res. Lett.*, 92, 1747–1758, <https://doi.org/10.1785/0220200435>, 2021.
- 835 Lund, B., Mäntyniemi, P., Sadegi-Bagherabadi, A., Korja, A., and Lundwall, J.: Comparing European Seismic Hazard Models: ESHM20 versus ESHM13 at nuclear power plant sites in Sweden and Finland, *Tech. Rep. 2024:1013*, Energiforsk, 2024.
- Mantovani, M. and Scherneck, H.-G.: DInSAR investigation in the Pärvie end-glacial fault region, Lapland, Sweden, *Int. J. Remote Sens.*, 34, 8491–8502, <https://doi.org/10.1080/01431161.2013.843871>, 2013.
- McGuire, R. K.: FORTRAN computer program for seismic risk analysis, <https://doi.org/10.3133/ofr7667>, 1976.
- 840 McGuire, R. K.: Deterministic vs. probabilistic earthquake hazards and risks, *Soil Dynamics and Earthquake Engineering*, 21, 377–384, [https://doi.org/10.1016/s0267-7261\(01\)00019-7](https://doi.org/10.1016/s0267-7261(01)00019-7), 2001.
- Miller, A. C. and Rice, T. R.: Discrete Approximations of Probability Distributions, *Management Science*, 29, 352–362, <http://www.jstor.org/stable/2631060>, 1983.
- Mosca, I., Sargeant, S., Baptie, B., Musson, R., and Pharaoh, T.: The 2020 national seismic hazard model for the United Kingdom, *Bull. Earthquake Eng.*, 20, 633–675, <https://doi.org/10.1007/s10518-021-01281-z>, 2022.
- 845 Muir Wood, R.: The Scandinavian earthquakes of 22 December 1759 and 31 August 1819, *Disasters*, 12, 223–236, <https://doi.org/10.1111/j.1467-7717.1988.tb00672.x>, 1989.
- Munier, R., Adams, J., Brandes, C., et al.: International Database of Glacially-Induced Faults, PANGAEA, <https://doi.org/10.1594/PANGAEA.922705>, 2020.
- 850 Mäntyniemi, P.: Some features of Finnish earthquake data, *Geophysica*, 32, 273–289, 1996.
- Mäntyniemi, P., Wahlström, R., Lindholm, C., and Kijko, A.: Seismic hazard in Fennoscandia: A regionalized study, *Tectonophysics*, 227, 205–213, [https://doi.org/10.1016/0040-1951\(93\)90095-2](https://doi.org/10.1016/0040-1951(93)90095-2), 1993.
- Mäntyniemi, P., Kijko, A., and Retief, P.: Parametric-historic procedure for seismic hazard assessment and its application to northern Europe, *Boll. Geofisica*, 42, 41–55, 2001.
- 855 Mäntyniemi, P., Sørensen, M., Tatevossian, T., Tatevossian, R., and Lund, B.: A Reappraisal of the Lurøy, Norway, Earthquake of 31 August 1819, *Seismol. Res. Lett.*, 91, 2462–2472, <https://doi.org/10.1785/0220190363>, 2020.
- Norsar: NORSAR Station Network, <https://doi.org/10.21348/D.NO.0001>, 1971.
- Olesen, O., Olsen, L., Gibbons, S. J., Ruud, B. O., Høgaas, F., Johansen, T. A., and Kværna, T.: Postglacial Faulting in Norway: Large Magnitude Earthquakes of the Late Holocene Age, in: *Glacially-Triggered Faulting*, edited by Steffen, H., Olesen, O., and Sutinen, R., p. 198–217, Cambridge University Press, <https://doi.org/10.1017/9781108779906.015>, 2021.
- 860 Ottemöller, L., Strømme, M. L., and Storheim, B. M.: Seismic monitoring and data processing at the Norwegian National Seismic Network, *Summary of the Bulletin of the International Seismological Centre*, 52, 27–40, 2018.

- Pagani, M., Monelli, D., Weatherill, G., Danciu, L., Crowley, H., Silva, V., Henshaw, P., Butler, L., Nastasi, M., Panzeri, L., Simionato, M., and Vigano, D.: OpenQuake Engine: An Open Hazard (and Risk) Software for the Global Earthquake Model, *Seismol. Res. Lett.*, 85, 692–702, <https://doi.org/10.1785/0220130087>, 2014.
- Rögnvaldsson, S. T. and Slunga, R.: Routine fault plane solutions for local networks: A test with synthetic data, *Bull. Seismol. Soc. Am.*, 83, 1232–1247, 1993.
- Rosberg, J.-E. and Erlström, M.: Evaluation of the Lund deep geothermal exploration project in the Romeleåsen Fault Zone, South Sweden: a case study, *Geothermal Energy*, 7, <https://doi.org/10.1186/s40517-019-0126-7>, 2019.
- Rovida, A. and Antonucci, A.: EPICA - European PreInstrumental Earthquake Catalogue, version 1.1, <https://doi.org/10.13127/EPICA.1.1>, 2021.
- Saari, J., Lund, B., Malm, M., Mäntyniemi, P., Oinonen, K., Tiira, T., Uski, M., and Vuorinen, T.: Evaluating Seismic Hazard for the Hanhikivi Nuclear Power Plant Site, Seismological Characteristics of the Source Areas, Attenuation of Seismic Signal, and Probabilistic Analysis of Seismic Hazard., Tech. Rep. NE-4459, ÅF-Consult Ltd, Finland, 2015.
- Sadeghisorkhani, H., Gudmundsson, O., Li, K. L., Tryggvason, A., Lund, B., and Högdahl, K.: Shear wave structure of southern Sweden from precise phase-velocity measurements of ambient-noise data, *Geophysical Journal International*, 225, 494–511, <https://doi.org/10.1093/gji/ggaa598>, 2020.
- Sandersen, P. B. E., Gregersen, S., and Voss, P. H.: Postglacial Faulting in Norway: Large Magnitude Earthquakes of the Late Holocene Age, in: *Glacially-Triggered Faulting*, edited by Steffen, H., Olesen, O., and Sutinen, R., pp. 263–282, Cambridge University Press, <https://doi.org/10.1017/9781108779906.015>, 2021.
- Sargeant, S., Stafford, P., Lawley, R., Weatherill, G., A-J.S., W., Bommer, J., Burton, P., Free, M., Musson, R., Kuuyor, T., and Rossetto, T.: Observations from the Folkestone, UK, earthquake of 28 April 2007, *Seismol. Res. Lett.*, 79, 672–687, <https://doi.org/10.1785/gssrl.79.5.672>, 2008.
- Schorlemmer, D., Euchner, F., Kästli, P., and Saul, J.: QuakeML: status of the XML-based seismological data exchange format, *Annals of Geophysics*, 54, <https://doi.org/10.4401/ag-4874>, 2011.
- Schulte, S. M. and Mooney, W. D.: An updated global earthquake catalogue for stable continental regions: reassessing the correlation with ancient rifts, *Geophysical Journal International*, 161, 707–721, <https://doi.org/10.1111/j.1365-246x.2005.02554.x>, 2005.
- Shiddiqi, H. A., Ottemöller, L., Rondenay, S., Halpaap, F., Gradmann, S., and Michálek, J.: Crustal structure and intraplate seismicity in Nordland, Northern Norway: insight from seismic tomography, *Geophys. J. Int.*, 230, 813–830, <https://doi.org/10.1093/gji/ggac086>, 2022.
- SKI: Project Seismic Safety. Characterization of seismic ground motions for probabalistic safety analyses of nuclear facilities in Sweden, Tech. Rep. 92:3, Swedish Nuclear Power Inspectorate (SKI), 1992.
- Slunga, R.: Swedish earthquakes and acceleration probabilities, Tech. rep., Swedish Defence Research Agency, Sweden, http://inis.iaea.org/search/search.aspx?orig_q=RN:10480150, fOA-C–20295, 1979.
- Smith, C. A., Mikko, H., and Grigull, S.: Glacially Induced Faults in Sweden: The Rise and Reassessment of the Single-Rupture Hypothesis, in: *Glacially-Triggered Faulting*, edited by Steffen, H., Olesen, O., and Sutinen, R., p. 218–230, Cambridge University Press, Cambridge, <https://doi.org/10.1017/9781108779906>, 2021.
- SNSN: Swedish National Seismic Network, Uppsala University, Uppsala, Sweden. Other/Seismic network, <https://doi.org/10.18159/SNSN>, 1904.

- 900 Sodankylä Geophysical Observatory / University of Oulu : Northern Finland Seismological Network, <https://doi.org/10.14470/SA879454>, 1980.
- Soosalu, H., Uski, M., Komminaho, K., and Veski, A.: Recent Intraplate Seismicity in Estonia, East European Platform, *Seismol. Res. Lett.*, 93, 1800–1811, <https://doi.org/10.1785/0220210277>, 2022.
- Steffen, H., Olesen, O., and Sutinen, R.: *Glacially-Triggered Faulting*, Cambridge University Press, Cambridge, 905 <https://doi.org/10.1017/9781108779906>, 2021.
- Stephens, M. B. and Bergman-Weiheid, J.: Sweden: Lithotectonic Framework, Tectonic Evolution and Mineral Resources, vol. 50, Geological Society, Memoirs, London, <https://doi.org/10.1144/M50-2019-21>, 2020.
- Stewart, J. P., Parker, G. A., Atkinson, G. M., Boore, D. M., Hashash, Y. M. A., and Silva, W. J.: Ergodic site amplification model for central and eastern North America, *Earthquake Spectra*, 36, 42–68, <https://doi.org/10.1177/8755293019878185>, 2020.
- 910 Svendsen, J. I., Alexanderson, H., Astakhov, V. I., Demidov, I., Dowdeswell, J. A., Funder, S., Gataullin, V., Henriksen, M., Hjort, C., Houmark-Nielsen, M., Hubberten, H. W., Ólafur Ingólfsson, Jakobsson, M., Kjær, K. H., Larsen, E., Lokrantz, H., Lunkka, J. P., Lyså, A., Mangerud, J., Matiouchkov, A., Murray, A., Möller, P., Niessen, F., Nikolskaya, O., Polyak, L., Saarnisto, M., Siegert, C., Siegert, M. J., Spielhagen, R. F., and Stein, R.: Late Quaternary ice sheet history of northern Eurasia, *Quaternary Science Reviews*, 23, 1229–1271, <https://doi.org/https://doi.org/10.1016/j.quascirev.2003.12.008>, quaternary Environments of the Eurasian North (QUEEN), 2004.
- 915 Tan, Y. J. and Maharjan, R.: What Googling Trends Tell Us about Public Interest in Earthquakes, *Seismol. Res. Lett.*, 89, 653–657, <https://doi.org/10.1785/0220170116>, 2018.
- Tinti, S. and Mulargia, F.: Confidence intervals of b-values for grouped magnitudes, *Bull. Seismol. Soc. Am.*, 77, 2125–2134, <https://doi.org/10.1785/BSSA0770062125>, 1987.
- University of Bergen: University of Bergen Seismic Network, <https://doi.org/10.7914/SN/NS>, 1982.
- 920 Uski, M. and Tuppurainen, A.: A new local magnitude scale for the Finnish seismic network, *Tectonophysics*, 261, 23–37, 1996.
- Uski, M., Lund, B., and Oinonen, K.: Compilation of a homogenized earthquake catalogue, in: *Evaluating Seismic Hazard for the Hanhikivi Nuclear Power Plant Site, Seismological Characteristics of the Source Areas, Attenuation of Seismic Signal, and Probabilistic Analysis of Seismic Hazard.*, edited by Saari, J., Lund, B., Malm, M., Mäntyniemi, P., Oinonen, K., Tiira, T., Uski, M., and Vuorinen, T., Report NE-4459, p. 123, ÅF-Consult Ltd, Finland, 2015.
- 925 van Stiphout, T., Zhuang, J., and Marsan, D.: Seismicity declustering, Community Online Resource for Statistical Seismicity Analysis, <https://doi.org/10.5078/corssa-52382934>, 2012.
- Veikkolainen, T., Kukkonen, I. T., and Tiira, T.: Heat flow, seismic cut-off depth and thermal modeling of the Fennoscandian Shield, *Geophys. J. Int.*, 211, 1414–1427, <https://doi.org/10.1093/gji/ggx373>, 2017.
- Veikkolainen, T., Kortström, J., Vuorinen, T., Salmenperä, I., Luhta, T., Mäntyniemi, P., Hillers, G., and Tiira, T.: The Finnish National Seismic Network: Toward Fully Automated Analysis of Low-Magnitude Seismic Events, *Seismol. Res. Lett.*, 92, 1581–1591, 930 <https://doi.org/10.1785/0220200352>, 2021.
- Vestøl, O., Ågren, J., Steffen, H., Kierulf, H., and Tarasov, L.: NKG2016LU: a new land uplift model for Fennoscandia and the Baltic Region, *Journal of Geodesy*, 93, 1759–1779, <https://doi.org/10.1007/s00190-019-01280-8>, 2019.
- Wahlström, R. and Grünthal, G.: Probabilistic seismic hazard assessment (horizontal PGA) for Sweden, Finland and Denmark using 935 different logic tree approaches, *Soil Dynamics and Earthquake Engineering*, 20, 45–58, [https://doi.org/https://doi.org/10.1016/S0267-7261\(00\)00037-3](https://doi.org/https://doi.org/10.1016/S0267-7261(00)00037-3), 2000.

- Wahlström, R. and Grünthal, G.: Probabilistic seismic hazard assessment (horizontal PGA) for Fennoscandia using the logic tree approach for regionalization and nonregionalization models, *Seismol. Res. Lett.*, 72, 33–45, <https://doi.org/10.1785/gssrl.72.1.33>, 2001.
- 940 Weatherill, G. and Cotton, F.: A ground motion logic tree for seismic hazard analysis in the stable cratonic region of Europe: regionalisation, model selection and development of a scaled backbone approach, *Bulletin of Earthquake Engineering*, 18, 6119–6148, <https://doi.org/10.1007/s10518-020-00940-x>, 2020.
- Weichert, D. H.: Estimation of the earthquake recurrence parameters for unequal observation periods for different magnitudes, *Bull. Seismol. Soc. Am.*, 70, 1337–1346, <https://doi.org/10.1785/BSSA0700041337>, 1980.
- 945 Wheeler, R. L.: Maximum Magnitude (Mmax) in the Central and Eastern United States for the 2014 U.S. Geological Survey Hazard Model, *Bull. Seismol. Soc. Am.*, 106, 2154–2167, <https://doi.org/10.1785/0120160048>, 2016.
- Woessner, J., Laurentiu, D., Giardini, D., Crowley, H., Cotton, F., Grünthal, G., Valensise, G., Arvidsson, R., Basili, R., Demircioglu, M. B., Hiemer, S., Meletti, C., Musson, R. W., Rovida, A. N., Sesetyan, K., Stucchi, M., and Consortium, T. S.: The 2013 European Seismic Hazard Model: key components and results, *Bulletin of Earthquake Engineering*, 13, 3553–3596, <https://doi.org/10.1007/s10518-015-9795-1>, 2015.
- 950 Zöller, G.: A Note on the Estimation of the Maximum Possible Earthquake Magnitude Based on Extreme Value Theory for the Groningen Gas Field, *Bull. Seismol. Soc. Am.*, 112, 1825–1831, <https://doi.org/10.1785/0120210307>, 2022.




Figure 1. Seismicity, geology, and tectonic map of Fennoscandia. Black dots indicate earthquakes in the declustered catalogue in this study. Yellow stars indicate locations referred to in the text, from north: "P" the Pärvie fault, "B" the Burträsk fault, Hudiksvall "H" the Hälsingland region, "U" Uppsala and "VG" Västergötland, south of Lake Vänern. The Sorgenfrei-Tornquist and Trans European Suture Zones are marked with dashed black lines. Post-glacial faults (red) from Munier et al. (2020). Map modified after Gregersen et al. (2021)




Figure 2. ~~Tectonic source zonation scheme~~ Earthquakes in the Fennoscandian region from our catalogue between 1875 and 2022 (A blue dots) and the ~~area source zonation scheme~~ ESHM20 catalogue 1497 - 2014 where $M_w \geq 3.5$ (B black stars) used in this study. The red line indicates a zone encompassing Sweden that is 300 km. Red lines: post-glacial faults from the Swedish border or economic zone boundaries Munier et al. (2020). Numbers detail the zone numbers.




Figure 3. Magnitude-time density plot of the earthquakes since 1875 in the zone encompassing Sweden that is 300 km from the Swedish border or economic zone boundaries (blue dots within the red zone in Figure 4A) from the declustered and magnitude homogenized catalogue.




Figure 4. Earthquakes in the ESHM20 catalogue (black stars) and our Tectonic source zonation scheme with the fully declustered earthquake catalogue used in this study (blue dots), north of 63 degrees latitude and between 17 and 32.5 degrees east longitude. Red lines: post-glacial faults (B) Area source zonation scheme. The red line indicates a zone encompassing Sweden that is 300 km from the Swedish border or economic zone boundaries. Numbers detail the zone numbers.




Figure 5. Source model logic tree. The numbers indicate the weights of the different branches.




Figure 6. Ground motion model logic tree. The numbers indicate the weights of the different branches. The logic tree is further detailed in Section 4.6.




Figure 7. Mean hazard curves for Uppsala, Sweden, for different GMM implementations, see Section 4.6 for details. Yearly probability of exceedance versus PGA in g. Solid lines-symbols show the mean, while the upper and lower dashed lines show the 84% and 16% fractiles.




Figure 8. Seismic hazard maps for return periods of 475 years (left) and 2500 years (right). Contour lines represent 0.01g for the 475 year map and 0.025, 0.05g for the 2500 year map. Post-glacial faults indicated by black lines.




Figure 9. Disaggregation analysis showing the contribution of earthquakes with different magnitudes at different distances to the seismic hazard. The analysis is carried out for the site with the highest hazard estimate in our study (65.7N, 22.97E), left, and the site in Sweden used by Wahlström and Grünthal (2001) in their study (59.5N, 12.0E), right, both for a return period of 475 years.




Figure 10. Seismic hazard curves for areas of increased seismicity in Sweden (yellow stars in Figure 1). (A) Pärvie, (B) Burträsk, (C) Hälsingland, (D) Västergötland. Annual probability of exceedance versus peak-ground acceleration in g. Mean (red lines), 50% (blue) and 16% and 84% fractiles (dashed).




Figure 11. Seismic hazard maps showing the difference between our and ESHM20 results for return periods of 475 (left) and 2500 (right) years. Contour lines represent 0.01g for the 475 year map and 0.025g for the 2500 year map. Post-glacial faults indicated by black lines.

Table 1. Recurrence Parameters for Tectonic Source Zonation Schemes.

Zone	a-value $\pm \Delta a \pm 1\sigma$	b-value $\pm \Delta b \pm 1\sigma$	Complete events
1	2.926 \pm 0.005	0.912 \pm 0.01	<u>7347</u>
2	2.451 \pm 0.022	0.842 \pm 0.034	<u>366</u>
3	3.267 \pm 0.013	0.895 \pm 0.023	<u>1012</u>
4	1.857 \pm 0.026	0.887 \pm 0.034	<u>262</u>

Table 2. Recurrence Parameters for Area Source Zones in Sweden.

Zone	a-value $\pm \Delta a \pm 1\sigma$	b-value $\pm \Delta b \pm 1\sigma$	Complete events
2	2.504 \pm 0.012	1.051 \pm 0.023	<u>1384</u>
3	1.864 \pm 0.019	0.878 \pm 0.03	<u>503</u>
9	2.267 \pm 0.012	0.957 \pm 0.022	<u>1200</u>
13	2.041 \pm 0.017	0.922 \pm 0.03	<u>621</u>
14	2.011 \pm 0.018	0.88 \pm 0.031	<u>576</u>
15	1.917 \pm 0.019	0.844 \pm 0.03	<u>507</u>
18	1.767 \pm 0.02	0.8 \pm 0.028	<u>465</u>
23	1.676 \pm 0.028	0.891 \pm 0.047	<u>225</u>
24	2.132 \pm 0.028	0.895 \pm 0.047	<u>227</u>
25	1.586 \pm 0.022	0.842 \pm 0.034	<u>50</u>
26	1.068 \pm 0.005	0.912 \pm 0.01	<u>102</u>
27	1.974 \pm 0.022	0.842 \pm 0.034	<u>122</u>
29	1.78 \pm 0.013	0.895 \pm 0.023	<u>33</u>

Appendix A: [Figures](#)




Figure A1. Inter-event time distributions for the Fennoscandia earthquake data set from 2001-01-01 until 2022-02-28 and a magnitude of completeness of 1.0. (A) Full catalogue, (B) Declustered catalogue. Data, blue dots. Average Poisson inter-event times, red line. Lower 2.5% and upper 97.5% confidence limits, dashed red lines.




Figure A2. Map of Fennoscandia with events belonging to clusters resulting from the declustering algorithm. Cluster sizes indicated by the colours of the dots: 2 events: gray, 3-5 events: blue, 6-10: green, 11 - 20: red, above 20: orange. Larger clusters plotted on top of smaller clusters.




Figure A3. ESHM20 seismic hazard maps showing the mean spectral acceleration (0.2s - left and 1.0s - right) for a return period of 475 years. Contour lines at 0.01g (left) and 2500-years 0.0025g (right). Post-glacial faults indicated by black lines.

Magnitude-time density plots of the earthquakes in the (A) ESHM20 and (B) our catalogue north of 63 degrees latitude and between 15 and 25 degrees east longitude.