

# Spatial identification of regions ~~at-risk~~exposed to multi-hazards at pan European level

Tiberiu-Eugen Antofie<sup>1</sup>, Stefano Luoni<sup>2</sup>, Alois Tilloy,<sup>2</sup> Andrea Sibilgia<sup>3</sup>, Sandro Salari<sup>3</sup>, Gustav Eklund<sup>2</sup>, Davide Rodomonti<sup>3</sup>, Christos Bountzouklis<sup>2</sup> and Christina Corbane<sup>2</sup>

- 5 1. formally at European Commission, Joint Research Centre, Ispra, 21027, Italy (currently: independent researcher)  
2. European Commission, Joint Research Centre, Ispra, 21027, Italy  
3. External Consultant for the European Commission (Unisystems)

10 *Correspondence to:* Tiberiu Antofie Eugen (tiberiuantofie@yahoo.com)

**Abstract.** The Disaster Risk Management Knowledge Centre (DRMKC) is developing and hosting a web platform – the Risk Data Hub - intended to improve the access and sharing of curated EU-wide risk data, tools and methodologies for fostering Disaster Risk Management (DRM) related actions. Within the DRMKC’s Risk Data Hub (RDH) development, we integrate-develop a methodology for the identification of regions with multi-hazard ~~potential-impact~~ exposure at pan European level. With this study we present the methodological approach and we stage it as one fundamental development in support of DRM decision-making at national and subnational level.

15  
20 .-We adopt a meta-analysis approach, and by combining the hotspots of exposure to single hazards’ exposure, we address the challenge of statistically ‘insignificant results’ and we provide an objective “statistical proof” of the multi-hazard potential of a region. We support these results through a validation process which considers empirical data on fatalities and disaster events -as explanatory variables.

Presenting an implemented methodology, scalable down to local subnational level, that reveals types of assets ~~at-risk~~exposed to multiple hazards and their location, we take one further step towards the identification of the disaster risk management pathways in multi-hazard analysisassessment.

25 The outcome of this study provides valuable input and will assist national authorities on the integration of multi-hazard analysis in their National Risk Assessments and Disaster Risk Management plans.

## 1. Introduction

35 The results of a ‘‘Needs and Gaps’’ analysis performed as part of the preparation of the European Commission Staff Working Document – ‘‘Overview of Natural and Man-made Disaster Risks the European Union may face’’ (2014<sup>1</sup>, 2017<sup>2</sup>, 2020<sup>3</sup>), concluded that a gap in knowledge and data availability exists for multi-hazard assessments. A number of international frameworks such as Hyogo Framework for Action (UN-ISDR, 2005) or Sendai Framework for Disaster Risk Reduction 2015–2030, have endorsed the multi-hazard approach for Disaster Risk Reduction.

40 It is now well recognized in the research community that for an adequate understanding of disaster risk potential within a region it is essential to move from single hazard to multi-hazard approach (Marzocchi et al., 2009, Kappes et al., 2012, Gill and Malamud, 2014, Tilloy et al., 2019, Ward, P. J., et al., 2022). The hazard interrelations can lead to a combined impact that is different from the sum of each hazards’ impacts separately. In order to assess the potential hazards and the risk to which a region is exposed, some studies combined independent analysis of single hazards (Granger et al., 1999; van Westen et al., 2002; Greiving et al., 2006; Grünthal et al., 2006; Marzocchi et al., 2012, Forzieri et al., 2016) and superposed natural hazards over a region (multi-layer hazards). Others studies considered hazard interactions (Tarvainen et al., 2006; Han et al., 2007; De Pippo et al., 2008; Kappes et al., 2010; van Westen et al., 2014, Liu et al., 2016; Sadegh et al., 2018; Gill et al., 2020; Claassen et al., 2023; Lee et al., 2024). Often, these assessments are based on case studies within limited spatial extension, addressing a limited number of perils/hazards and addressing specific- a limited number of sectors (Ciurean et al., 50 2018, Tilloy et al., 2019).

Our study aligns with the first definition of multi-layer hazards, as we examine the combined exposures of single hazards over a region, recognizing that hazard interrelations can result in an impact distinct from the sum of individual hazards exposures.

This is exemplified by events such as the Portugal wildfires and flash floods in October 2017, where these hazards occurred in relatively close succession, both affecting the same buildings and infrastructure or the floods and the dam failure in summer of 2002 in Czech Republic when floods had caused significant damage to buildings, infrastructure, and agricultural land and subsequent dam failure from August added to the devastation, impacting structures that were already dealing with the effects of the floods.

One development that addresses these challenges, is the ~~DRMKC~~ Risk Data Hub platform (~~DRMKC~~-RDH) of the Disaster Risk Management Knowledge Centre (DRMKC). The platform improves-facilitates the access and sharing of curated European-wide risk data and methodologies being a fundamental tool in support of the DRM<sup>4</sup> and CCA<sup>5</sup> actions at national

---

<sup>1</sup> [EUR-Lex - 52014SC0134 - EN - EUR-Lex \(europa.eu\)](https://eur-lex.europa.eu/lexuris/ui/entry.do?uri=CELEX:52014SC0134:EN)

<sup>2</sup> [https://ec.europa.eu/echo/sites/echo-site/files/swd\\_2017\\_176\\_overview\\_of\\_risks\\_2.pdf](https://ec.europa.eu/echo/sites/echo-site/files/swd_2017_176_overview_of_risks_2.pdf)

<sup>3</sup> [https://ec.europa.eu/echo/sites/echo-site/files/overview\\_of\\_natural\\_and\\_man-made\\_disaster\\_risks\\_the\\_european\\_union\\_may\\_face.pdf](https://ec.europa.eu/echo/sites/echo-site/files/overview_of_natural_and_man-made_disaster_risks_the_european_union_may_face.pdf)

<sup>4</sup> <https://civil-protection-knowledge-network.europa.eu/knowledge-network-science/data-tools>

and subnational level. Within the DRMKC–~~Risk Data Hub~~RDH development, we propose a methodology which is accessible, scalable and replicable even at subnational and local level for the identification of regions ~~exposed at risk~~ to multi-hazards.

65 The multi-hazard methodological approach is the main goal of this study, focused on addressing four major challenges:

1. identification of regions with significant multi-hazard potential;
2. establishment of the ~~impact exposure~~ relation assets – multiple hazards,
3. quantification of multi-hazard ~~potential impact exposure~~ and
4. transferability of the method.

70 These challenges are further constrained by the wide scale of our analysis (European coverage), the alignment to a common hazard definition and their practical implementation on the online web platform, the DRMKC RDH.

Challenge 1 is addressed in this study with a novel methodology that identifies, at pan-European scale, the regions (Local Administrative Units - LAUs) ~~exposed at risk~~ to multi-hazards with high level of statistical significance ( $p$  value < 0.10).

75 ~~Our approach involves a meta-analysis technique which functions as a powerful significance test (Hak et al., 2016). We adopt a meta-analysis approach, and by combining the hotspots of single hazards' exposure, we address the problem of statistically 'insignificant results' and we provide an objective "statistical proof" of the multi-hazard potential of a region. We combine the hotspots of single hazards' exposure and we generate a unified result, effectively addressing the challenge presented by divergent and even contradicting independent results.~~ This is the first study that uses spatial patterns (clusters/hotspots) and meta-analysis for this purpose.

80 Furthermore for challenge 2, we show that the proposed methodology allows for the detection of the regions ~~at risk~~exposed to multi-hazards, differently, as function of the typology of the assets. This is important as it directly reveals relationships between assets types and -threats, valuable for the identification of the disaster risk management pathways in multi-hazard assessment (Ward, P. J., et al., 2022). This is the central aspect of the multi-hazard analysis presented in this study, which considers the relation of single asset (population and the residential built-up respectively) to the multiple hazards: landslide, 85 coastal flood, river flood, earthquake, wildfires and subsidence.

Challenge 3, the quantification of multi-hazard ~~potential impact exposure~~ is addressed, by ~~totaling~~totalling the assets found only for the regions ~~at risk~~exposed to multi-hazards with high level of significance. This areal dimension approach (Hewitt and Burton, 1971), omits a detailed level of study that could more accurately examine the spatial coincidence, trigger relations or cascading effects when quantifying the impacts from multi-hazards. Nevertheless, we argue that our 90 methodology succeeds in describing the “hazardousness” level of a region which offers a generalized spatial understanding of where the specific assets are exposed to multiple hazards and what hazard becomes accountable for a potential impact.

Challenge 4 is addressed by showing that in contrast to other studies, the transferability of the developed methodology is not limited due to the reliance on case-study-specific data and methods. The methodological approach described in this study is

---

<sup>5</sup> [eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:52021DC0082](http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:52021DC0082) (pg. 6)


already implemented on the DRMKC Risk Data Hub platform and uses the existing pan-European data hosted and shared through the platform.

Research on multi-hazard analysis has underscored several critical gaps that request attention for more effective multi-hazard assessments. These gaps could be listed as : data quality, incomplete, outdated, that impacts on accurate multi-hazard assessments (Cutter et al., 2014, Gentile et al., 2022); understanding the complex interactions between hazards (Gill et al., 2016; Lee et al., 2024); temporal dynamics (Fuchs & Thaler, 2018, De Angelis et al., 2022); addressing varying vulnerabilities across hazards (Saaty, 1987; UNISDR, 2004). Additionally, the limited attention given to uncertainty and sensitivity analyses in multi-hazard assessments (Haasnoot et al., 2013, Camus et al., 2021); the inadequate incorporation of climate change considerations into multi-hazard assessments (IPCC, 2014; Gallina et al., 2016, Ghanbari et al., 2021) and the communication challenges conveying multi-hazard risks to stakeholders (Dallo et al., 2020; De Fino et al).

We structure the study as follows, after the *Introduction* we describe the *Data and the methodology* used, in *Results* and based on these identified regions we provide a statistical analysis looking at different socioeconomic features. Furthermore, a *Validation* exercise is performed followed by *Discussions* and *Conclusions*.

## 2. Data and methodologies

The methodological approach is presented in 3 steps: (1) we describe the underlying exposure data and methodology that creates the bases for our single and multiple—hazard analysis, (2) we present the methodological approach used to find the hotspots for single hazards' exposure, (3) we present the metadata analysis methodology used to combine the hotspots of single hazards' exposure and to identify regions with significant multi-hazard potential. A representation of the entire methodological chain is provided in fig. 1.



120

Figure 1 Different steps of the methodological approach developed in this study

## 2.1 The exposure data and methodology

### 2.1.1 The areal dimension


For this study the multi-hazard spatial coincidence is assessed at the level of areal dimension, represented by the Local Administrative Units (LAUs).

125

The LAUs are the finest hierarchical classification of subdividing the European economic territory into regions in which statistics can be provided at a local level. This dataset comes from the statistical office of the European Union (Eurostat) and represents the administrative units of municipalities and communes of Europe, version 2013. In the present study, the LAUs cover the ~~European Union 28~~ EU27+UK and the European Free Trade Association (EFTA) countries.

130

These administrative entities are used as statistical areas for multi-hazard exposure and hotspot analysis as an approach meant to support disaster risk management activities. Administrative directives, organisations and operational services are coordinated at the level of administrative entities and they become of high relevance when linked down to local level, challenging the gap in the scale of policy and scale of practice (Gaillard et al., 2013).



**Figure 2. Local Administrative Units area (km<sup>2</sup>), spatial distribution (left) and mean LAUs area per country (right)**

There are 122 034 LAUs considered as geographical statistical units on which the aggregations and statistical analysis are performed in this study. Their average area is 39.6 km<sup>2</sup>, the maximum area is 20 688 km<sup>2</sup> (Kiruna, SWE) and the minimum is 0.2 km<sup>2</sup> (Thorpe Hamlet, UK). LAUs present heterogeneities across Europe in terms of area covered especially in northern part of Europe (e.g. Scandinavia), even if they are rather homogeneously distributed within the national boundaries (fig. 2).

Despite being a well-established geographic concept, the process of aggregating higher resolution data to larger administrative units comes with a potential source of error known as modifiable areal unit problem (MAUP). The two related issues to the MAUP, largely presented in the literature (Fotheringham and Wong, 1991; Jelinski and Wu, 1996; Openshaw, 1984) are the scaling and the zonation effect (Charlton, 2009). These are generally altering the variance structure of the data when aggregated due to disconnection across scales and to different ways of subdividing the geographical space at the same scale (Stillwell et al., 2014). In order to minimize the MAUP effect, recommended practices (Su, 2011; Kwan, 2012) which are consistent with our approach focus on using smaller areal unit (e.g., LAUs rather than provinces or countries) for data aggregation. It reduces the potential errors of spatial pattern distortion without completely removing it.

### 2.1.2. Input hazard and Exposure data

The exposure data is built on the relationship hazard (*i*) - assets (*ii*):  $\text{exposure/assets at risk} = f(\text{Assets, Hazard})$ . We overlay spatial information about residential built-up and population with data describing hazard areas in order to define the assets exposure to single hazards. We then aggregate the exposure at the level of LAUs. We search for the significant hotspots of assets at risk exposed from single hazards using two types of exposure aggregation:

- 155 - based on absolute values - the sum of the exposed asset  
 - based on relative values - as ratios or share of the exposure from the total amount of asset in a LAUs.

For the exposure to earthquake, due to the continuous spatial extent of the hazard area, we depict the relative aggregation schema using the density (or share of the exposure compared to the total area of the LAUs). The relative aggregation schema intends to address risk management strategies based on the cost efficient-measure while the absolute schema supports the risk management strategies that prioritize the most affected areas and people.

(i) *Hazard Layers*

The hazard layers considered in this study represent areal extension rather than intensity. We do not use a probabilistic assessment but rather a deterministic approach selecting hazards with average temporal (frequency of occurrence) and spatial probability (susceptibility). A review of the hazard datasets and their characteristics is presented in Table 1. The motivations for their selection along with their usage in disaster risk assessments are presented in the sections dedicated to individual hazards in Supplementary material (Section 1 - Hazard data).

**Table 1. Description of the Hazards scenarios and datasets considered and their characteristics**

Component	Scenario	Description	Spatial resolution	Data source
River flood	1 event in 200yr RP	Areal extent of the river flood prone areas	100m	EFAS (European Flood Awareness System), KULTURisk project
Landslide	High and very high susceptibility classes	Physical characteristics of various terrain factors that provides high predisposition to landslide occurrence (ELSUS 100 layer)	200m	<a href="#">ESDAC (European Soil Data Centre)</a>
Coastal inundation	1 event in 200yr RP	Areal extent of coastal inundation as extreme total water level (TWL) result of the contributions from the mean sea level (MSL), the tide and the combined effect of waves and storm surge.	100m	HELIX project, JRC CoastalRiskandGAP-PESETAII projects
Earthquake	PGA $\geq$ 0.18 (g) for a probability of exceedance of 10% in 50 years (475yr RP)	Areal extent of PGA $\geq$ 0.18 (g) , equivalent of 'Moderate', 'Moderate to heavy' 'Heavy', 'Very heavy' potential damage level of USG Intensity Scale	1000m	SHARE project
Subsidence (from)	Soils with clay content	Areal Extent of fine and very fine soil	1000m	ESDAC, IPL project

drought)	greater than 35%.	texture (particle < 2 mm size) and with clay content greater than 35%.		
<del>WildfireForest</del> fire	Wildland–Urban Interface area (WUI)	WUI areas within 10 km limit range from the historical burned areas (2000-2016)	100m	EFFIS based

170

(ii) *Assets layers*

As assets layer, we use the residential built-up from the European Settlement Map (ESM) (Florczyk, et. al., 2015) and residential population from the Global Human Settlement Layer (GHSL) (Freire, et al., 2015). These are two main groups of assets that are present currently across all types of analysis within the DRMKC Risk Data Hub. The residential built-up is represented as built-up area (km<sup>2</sup>) and the population is amount of people within 100m x 100m grid cells.

175

In order to discriminate the residential typology for both built-up and population, the Corine Land Cover (CLC 2018) code 1.111 ([continuous urban fabric](#)) and 2.112 ([discontinuous urban fabric](#)) is used as the artificial explanatory layer.

## 2.2 Single hazard hotspots analysis

The study uses a hotspot analysis in order to identify clusters (concentrations) of regions — LAUs, -with assets ([or elements at risk](#)) [exposed](#) to single-hazard. The chosen approach enables the recognition of spatial patterns and trends which are not immediately apparent in raw data and which exhibit underlying spatial processes at work that are not the result of random processes (Getis and Ord., 1992). We argue that these spatial patterns (hotspots) once combined across multiple-hazards will describe the statistically significant multi-hazard [potential exposure](#)-of regions.

180

Various methods for combining single hazard data are considered in literature, including classifications and index developments. For more information on this topic, the reader can refer to Kappes et al., (2012).

185

For this study, the Gi\*(d) statistic is used for local spatial autocorrelation analysis using the python-based Exploratory Spatial Data Analysis (ESDA)<sup>6</sup> package. The method describes the spatial autocorrelation as Z-score (standard deviations), p-value (probability), and confidence level (significance) for each feature (each LAU region). Very high (positive) or very low (negative) Z-scores, associated with very small p-values (e.g. values of  $-p < 0.1$ ), describe spatial clusters as cold spots and respectively hotspots with high significance level. In the field of disaster risk reduction and management, identifying both cold spots and hotspots is crucial for allocating resources efficiently. In the present study the hot spots refers to areas or regions with higher susceptibility of risk from multi-hazard while the cold spots can be considered less prone to multi-hazard risks.

190

<sup>6</sup> <https://pysal.org/esda/>



195 **-Conceptualization of Spatial Relationship.** A known characteristic is that the statistics we are interested in (high Z-scores, low  $p$ -values) are placed in the tails of the distribution and therefore are susceptible to noise and spatial outliers. Moreover, the skewness of a distribution can bias the statistics (Cousineau D.,2010). These are important to consider as the resulting distribution areas of the single hazard clusters needs to be homogeneous in order to be significantly combined in a multi-hazard spatial cluster through meta-analysis (Hak et al., 2016).

200 To ensure reliable results, the study addresses noise and outliers through a spatial weights matrix. This matrix defines neighboring regions and we use the k-Nearest Neighbour (Fix and Hodges, 1951, Cover and Hart 1967) algorithm which is based on the proximity ( $k$ ) information in order to represent the spatial relationship between regions (LAUs). We have selected this method over contiguity based weights, since the k-nearest neighbour weights displays no ‘‘island’’ problem (isolated polygons that do not share any boundaries with other polygons), and every region has at least one neighbour. More  
205 information on the factors which affect clustering performance can be found in Zhao, M., et al. 2016, on the merits of a weighted matrix.

The study also considers the optimization of spatial autocorrelation/clustering across single hazard exposures by selecting the optimal neighborhood size ( $k$ ) in the k-Nearest Neighbor (k-NN) algorithm (we present it in Supplementary material section 2).

### 210 **2.3 Meta-Analysis: Identifying Regions with Multi-Hazard Potential driven by exposure**

The study adopts a meta-analysis approach to identify regions with multi-hazard potential. This involves combining probabilities ( $Z$ -scores and  $p$ -values) from independent hotspots. From the hotspot analysis of different hazards exposure, the same region can show statistically significant positive clustering (hotspot), statistically significant negative clustering (coldspot) as well as statistically non-significant clustering. By the combined outcome of these individual tests that  
215 sometimes differ and contradict each other, we measure the multi-hazard potential at regional level. Meta-analysis serves as a viable solution for addressing the challenge of seemingly conflicting evidence in research (Hak et al., 2016; Borenstein et al., 2009). Notably, it serves as a potent tool for conducting robust significance tests (Hak et al., 2016). Consequently, meta-analysis also proves instrumental in resolving the issue of ‘‘insignificant results.’’ In the context of our study, meta-analysis serves as a mechanism for synthesizing findings from various clustering analyses. Furthermore, by elucidating the statistical  
220 significance of the common estimation, it furnishes an objective ‘‘statistical proof’’ of the potential for multi-hazard clustering in our particular case.

Many  $p$ -values or  $Z$ -scores combining methods are used in meta-analysis to aggregate summary statistics. Most used methods are the following:

- i. Fisher method (Fisher, 1932) based on  $p$ -value to test the significance of the aggregations;
- 225 ii. Lancaster’s method (Lancaster, 1961) is a generalization of Fisher’s test by assigning different weights;
- iii. Stouffer’s method (Souffer, 1949) based on  $Z$ -transform test,
- iv. Lipták’s method (Lipták, 1958) which is Stouffer’s method with weights, known as weighted  $Z$ -test;

- v. the binomial test (Wilkinson, 1951) which counts the number of  $p$ -values that are below a threshold  $\alpha$
- vi. the truncated P-value methods (Zaykin et al., 2002) which adds up  $p$ -values that fall below a threshold  $\alpha$

230 For a good overview and comparison of these methods please refer to Whitlock, 2005; Zaykin, 2011; Chen, 2011. Meta-analysis have a widespread use due to their applicability, primarily in psychology, biology and medicine (McFarland, V, L., 2015). Within the field of disaster risk management, meta-analysis have been used mainly to assign the macroeconomy of disasters (van Bergeijk et. Al., 2015).

235 We chose to use the Stouffer's method (Z-transform test), without weighting, applied on the two-tailed distribution of the single clusters:

$$Z_s = \frac{\sum_{i=1}^k Z_i}{\sqrt{k}}$$

240 The sum of Z-scores ( $Z_i$ ), divided by the square root of the number of tests,  $k$ , provides a test of the cumulative evidence on the common null hypothesis (Whitlock, 2005).

Generally the Z-transform test converts the one-tailed  $p$ -values, from each of  $k$  independent tests into standard normal deviates  $Z_i$ . A common approach in meta-analysis is to sum the Z-scores across studies, weighting them appropriately using the sample sizes. On considering two-tailed method please see Whitlock (2005); Yoon et al., (2021) and on advantages and disadvantages of using the unweighted version of this method please see Becker, B.J. 1994. The z-transform test was  
 245 performed in python using the *scipy.stats*.<sup>7</sup>


### 3. Results

We identify the regions (LAUs) in Europe **at risk exposed** to multi-hazards by combining the Z-scores and  $p$ -values across the hotspots of single hazard exposure (i.e. population and built-up) computed on absolute and relative (%) aggregations. In fig. 3, we map these regions and further we consider for a statistical overview the regions with more than 1 hazard exposure (Hz  
 250 > 1) and confidence level set at 90% ( $p$ -value < 0.10 and positive Z-score > 0). In the Supplementary material (fig. S25) we present also the map with all hazard types identified at the level of LAU and depicted by the analysis done on the relative population exposure (the analysis performed on other assets types will presents a different spatial distribution of the hazard types).

255

---

<sup>7</sup> <https://docs.scipy.org/doc/scipy/tutorial/stats.html>



260 **Figure 3. Regions (LAUs) at-risk exposed to multi-hazards identified by the meta-data analysis performed on a.) absolute population exposure , b.) relative (%) population exposure, c.) absolute residential built-up exposure and d.) relative (%) residential built-up exposure**

The identification of these regions yielded disparate outcomes contingent upon the specific exposure types scrutinized within our analysis, namely, population density or residential built-up areas. Moreover, the choice of aggregation method, whether relative (expressed as a percentage) or absolute (in terms of the number of individuals or square kilometres of residential built-up areas at-risk exposed), introduced variations in both the quantity and spatial arrangement of regions identified as susceptible to multi-hazard events. The difference in multi-hazard exposure among regions, when considering absolute versus relative aggregation, is influenced by clustering algorithm sensitivity to distance (computed by the  $k$ -parameter) and similarity measures, where absolute aggregation accentuates variance and is susceptible to outliers, while relative aggregation smooths dominance of extreme values, potentially overlooking high-exposure areas within densely populated regions. Also Aa higher number of regions at the European level were identified as susceptible to multi-hazard risks when

265  
270

considering population-based criteria, as opposed to residential built-up exposures (see Figure 4). Furthermore, there is a significant difference between the amount of regions being at-risk-exposed to multi-hazards identified on absolute (12% for population and 10.6% for residential built-up) compared with the relative aggregation (21% - population and 13.6% - residential built-up).

275




Figure 4. Local administrative units (as % from the total in Europe) identified as being prone to multi-hazards ~~risk~~-based on different indicators -(population and residential built-up) and aggregation types (relative and absolute)

280

In order to simplify the interpretation of the results and clearly present the potential of the methodology used, we further focus only on the regions at-risk-exposed to multi-hazards identified by the relative (%) population.

### 3.1. Regions (LAUs) with significant multi-hazard potential


285

Based on population exposure we found 26 058 administrative regions, LAUs (fig. 5) prone to multi-hazards ~~risk~~ in Europe with high significance level (regions with > 90 % confidence interval and number of hazards >1). Most of these regions (20 912) are described statistically as hotspots with highest confidence, 99%, and in only 6 regions in Europe all of the hazard considered for this analysis are present (5 in Italy and 1 in Croatia) (fig. 5 c). These are mountainous and coastal regions.

290

Regions prone to multiple hazards represents 21.4% of the local administrative units of Europe and around 87 mil. people (18.8 % of Europe population) (fig. 5 c and d). In figure 5 d, we show that almost half of the population is at-risk-exposed to more than 3 hazards. Most of these regions are found in France 6956 LAUs, Italy 4627 LAUs, Slovenia 3802, Bulgaria 1876, Spain 1779, Germany and Romania (around 1000 LAUs each). Almost a quarter of the population at-risk-is exposed-is

in Italy (21.4 mil) and together with the Netherlands (10.1 mil), France (9.5 mil), Spain and Germany (7.1 mil each) they total more than 55% of population at riskexposed to multi-hazards (fig. 5 b and 5 d).



295

300

**Figure 5.** Regions (LAUs) with population at riskexposed to multi-hazards by significance level (a.); Sum of population at riskexposed to multi-hazards assessed at NUTS3 (only hotspots regions with > 90 % confidence interval) (b.); Number of administrative areas at riskexposed to multi-hazards by confidence interval and number of hazards (c.); Population at riskexposed to multi-hazards by confidence interval and number of hazards (d.).

We present a statistical overview of these regions identified as being at riskexposed to multi-hazards, looking at their spatial distribution and their population at riskexposed considering (see 3.1):

- i. various level of economic development (high, middle high, middle low and low income regions - LAUs)
- ii. urbanisation level: rural and urban (according to URAU audit 2018 definitions across European LAUs)

- 305      *iii.* identifying metropolitan<sup>8</sup> areas ~~at risk~~exposed to multi-hazards  
*iv.* and comparing city centres (city cores – C ) and Functional urban areas (FUA) levels in a metropolitan area

(*i.*) In fig. 6 we present the results per income group and degree of urbanisation at European level (fig. 6, a., c.) and by countries (fig. 6 b, d) .

310 From fig. 6 a., about 36% (9496) of the administrative regions (LAUs) identified as having population ~~at risk~~exposed to multi-hazards are low income regions and together with the low middle income they sum up to 67% . High income regions represent 10% of the LAUs and high middle income regions 23%. However, the groups of high and high middle income administrative regions total around 50% (43.4 mil) of the population ~~at risk~~exposed to multi-hazards (fig. 6. c).

315 In fig. 6, b., based on income group and degree of urbanisation, we present the top countries with administrative regions (LAUs) identified as being ~~at risk~~exposed to multi-hazards.

Based on the income groups, most of the high income administrative regions ~~at risk~~exposed to multi-hazards are in Switzerland (30,9 %), Italy (19.1%) France (16.7%) and Austria, Germany, the Netherlands ( each >5% ) while the low income administrative regions are mostly found in the southern and eastern Europe in Slovenia (31.6%), Bulgaria (19.8%), Romania (10.4%), Hungary (8.9%) and in Italy and Portugal (each > 5%).

320 In fig. 6, d., most of the low income population ~~at risk~~exposed to multi-hazards are concentrated in Romania (23%), Italy, Hungary, Poland and Bulgaria (each > 10%) while the high income population ~~at risk~~exposed to multi-hazards is found in the Netherlands (33%), Germany, Italy and Austria (each >10%).

(*ii.*) Also, from fig. 6, a., the number of administrative areas (LAUs) that are characterised as urban area (based on URAU 2018 definition and on correspondence with LAUs) is much smaller than the number of rural administrative areas  
325 (respectively 26.3% or 6585 versus 73.7% or 19200).

Nevertheless, the urban population ~~at risk~~exposed to multi-hazards total 54% (46.8 mil) compared with the rural administrative areas 46% (40.1 mil) (fig. 6. c.).

330 Based on the urbanisation degree, 15 countries in Europe have a higher share of population ~~at risk~~exposed to multi-hazards in rural areas compared to urban areas: Sweden, Norway (100%) or Croatia, Cyprus, Portugal, Slovakia (between 70%-90%) and Hungary, Spain, Belgium, Slovenia, Romania, Switzerland, (between 50%-70%). In the rest of the countries like the Netherlands, Austria (> 80%), Poland, Germany, Greece, (60%-80%), Ireland, United Kingdom, France, Denmark, Czech Republic, Bulgaria (50%-60%) the share of population ~~at risk~~exposed to multi-hazards in urban areas is higher compared to rural area.

335 This indicates that people are more ~~at exposed to risk to~~ multi-hazards if they live in regions with higher GDP and more densely populated (high and high middle income and urban areas, these are 12% of the administrative regions in Europe) compared with people living in regions with lower GDP and less populated (low and middle low income and rural

---

<sup>8</sup> The metropolitan areas' according to URAU 2018 definitions and represented here as composed by: core city, Functional Urban Area, Greater city and Trans-national Functional Urban Area (codes: C, F, K, T)

areas, these are 54% of the administrative regions in Europe). Also considering the degree of urbanisation only, people are more at risk/exposed to multi-hazards either if they live in high income urban areas (compared with low income urban areas) or low income rural areas (compared with high income rural areas) (fig. 6. c.).

340





Figure 6. Number of administrative areas (LAUs) with population at risk/exposed to multi-hazards by income level and urbanization level (a. – Europe wide, b. - the 15 highest ranked countries) (upper part); Population at risk/exposed to multi-hazards by income level and urbanization level (c. – Europe wide, d. - the 15 highest ranked countries) (lower part)

345

From fig. 7 exploring the differences between various income classes we find that as countries and their regions get richer they get more exposure to multi-hazard risk. After they reach a higher level of income (in the middle income category), the population at risk/exposed from multi-hazard decreases towards the high income. This can suggest that low income countries have the major part of the population at risk/exposed in the rural areas compared to the high income countries where most of the population at risk/exposed is in densely populated urban area and only a quarter from the population at risk/exposed (25%) live in the rural area. The peak in the countries with regions in the middle income category could suggest a balance between the high number of urban areas (the largest across various income classes) and the rural areas with high densities in population.




**Figure 7. Population at risk/exposed per income level. The markers represent countries' population at risk/exposed of multi-hazard split by income level. The blue line links the 75th quantile of the income classes**

(iii.) Using the Urban Audit 2018 definition and based on correspondence with LAUs we have identified 46% of the urban/metropolitan areas in Europe (442 of a total of 952) have population at risk/exposed to multi-hazards. These urban areas, totalling 46.8 mil people, are mostly high and middle high income (62.4%). The high income urban areas are mostly found in the Nederland (28), UK (23), Germany (20), France (9) and Italy (9) while the low income (110 at European level) are found in Romania (17), Poland (15), Hungary (13), Czech Republic (11) and Bulgaria (16) and others (in Supplementary fig.S23 and table S6)

(iv.) In fig. 8 a). and b). we further explore the distribution of population at risk/exposed to multi-hazards within the urban areas comparing the categories: cities (or city cores/centers - C) and larger urbanized zones (commuting zone/Functional Urban - F). We show that from this local perspective, the population at risk/exposed to multi-hazards is



governed either by urban population densities or the expansion of urban land. The commuting zones (FUA) are more **at riskexposed** to multi-hazards compared with the city centers: 58% (in 257 out of 442) compared with the city centers. However, 57% of the population **at riskexposed** from the metropolitan areas in Europe live in City Centers. This would suggest that the more population density in the city centers, the more **at riskexposed** is the metropolitan area. This positive relationships is depicted in fig. 8 b. but is particularly weak in the case of high income metropolitan areas, as shown by the almost flat fitting (red) line, and stronger for the middle high and low income. This shows that going towards the richer metropolitan area the risk increases due to the expansion of the urban area (into the functional urban areas) and diversely, going towards less rich metropolitan areas, due to the densities increase. This is confirmed, with some exceptions (the Netherlands, Austria, Island), by the high income Nordic and Western countries metropolitan areas where higher proportion of population **at riskexposed** is found in the functional urban areas compared with the city centers: Denmark, Luxembourg (100%), Finland, Belgium, Switzerland (between 60%-80%) and Ireland, Italy, Germany and UK (between 50%-60%). Contrarily, in France, Spain and Portugal, most of population **at riskexposed** (between 50%-60%) is concentrated in city centres of the middle income metropolitan areas which are also the most populated. For the Eastern European lower income countries, the population **at riskexposed** to multi-hazards is greater in the city centres compared with the functional urban areas: Latvia, Romania, Poland, (> 70%), Bulgaria, Slovenia, Slovakia, Hungary, Czech Republic (between 60%-70%) (Supplementary fig. S24). However, it is evident that the intended comparison could be better explained through complex urban processes such as changing patterns of residential-choice behaviour due to socio-economic growth which we do not address in this work.



**Figure 8. Population at riskexposed to multi-hazards at the level of Metropolitan area; a). European countries' population at riskexposed within Metropolitan categories: City Centres (C) and Functional urban area (F). The lower and upper whiskers represent, respectively, the lowest 5% and the highest 95% of the calculated population at riskexposed to multi-hazards for each metropolitan category; b). Linear relation between population at riskexposed and total population assessed as difference from FUA of the City Centres. A flatter fitted line indicates a weaker or less pronounced relationship between the population at**

395


riskexposed and the total population. In this case, changes in the total population have a relatively smaller impact on the population at riskexposed. It suggests that the income category of the region may not strongly influence the risk factors within the population. A *less flat fitted line*, on the other hand, indicates a stronger relationship between the population at riskexposed and the total population, with changes in the total population having a more significant impact on the population at riskexposed.

#### 4. Validation

400

The validation proposed is based on- Spearman correlation analysis of the population at riskexposed from multi-hazard with 2 empirical datasets as independent variable: the DRMKC RDH recorded data on fatalities from past events and the count of events with fatalities (for the period 1980-2019), for common hazards: coastal floods, earthquakes, river floods, landslides, subsidence and wildfires. The input data, both the population at riskexposed to multi-hazards and the empirical data are brought to a common geographical scale, the NUTS3 and metric (Z-scores and p-values of clusters). We use the same methodological approach explained in this study in order to arrive to single hazard (clusters) hotspots. The single hazard hotspots of empirical data (fatalities and event count) and population at riskexposed to multi-hazards are combined through meta-analysis in order to arrive to a multi-hazard hotspots, of fatalities, event count and of population at riskexposed scaled at NUTS3 level (fig.9). Finally hot/cold-spots regions of the 2 independent variables (fatalities and event count) are compared with the population at riskexposed from multi-hazard.

405



410

**Figure 9. Identified hot /cold -spots regions (NUTS3) with a.) population at riskexposed to multi hazard; b.) fatalities from multi-hazard, c.) number of events with fatalities; -used in Spearman correlation analysis for the validation purpose**

By using the correlation coefficient analysis we tried to capture the strength of the relation between the two paired datasets, numerically.

415


We focused on a non-parametric test, the Spearman correlation analysis, because it does not assume that the data is from a specific distribution and is computed on ranks and so depicts monotonic relationships. We choose it as a neutral way of assessing the general central tendency (median) among the pairs of variables at NUTS3. As interpretation, the Spearman

shows the degree by which two variables tend to change in the same direction. Therefore, variables with high correlation increase and decrease simultaneously, while variables with low absolute correlation rarely increase and decrease together.

420

425

430



**Figure 10 Spearman correlation between the multi-hazard clusters' size (Z-scores) of population at risk exposed with the empirical fatalities from past events (lefts) and events count (right )**

435

The results presented in fig. 10 refers to the correlation coefficients between paired population at risk exposed with the: a.) amount of fatalities (absolute) and b.) count of events of the empirical data.

440


We find a rather inconclusive relationship between the multi-hazard risk data and the empirical data, if we consider all regions for all significance levels. The scatterplot suggests a positive correlation between the variables but their increasing monotonic relationship is weak ( $r=0.37$  with fatalities and  $r=0.25$  with the event count) .

However, if we consider only the regions with higher significance ( $p<0.01$ ,  $p<0.05$ ,  $p<0.10$ ) we notice a stronger correlation (table 2 and fig. 11). This means that going towards more significant clustering (hot/cold-spots), the independent variables used for the validation tend to follow better the changes in value of the population at risk exposed to multi-hazards.

445

**Table 2. Spearman correlation coefficient between the empirical data (fatalities and count of past events) and the population at risk exposed from multi-hazard for regions (NUTS3) with different significance levels.**

Variables	$p\_value<0.01$	$p\_value<0.05$	$p\_value<0.10$	All regions
Fatalities absolute	0.59	0.51	0.46	0.37
Count events	0.30	0.40	0.35	0.25



450 Figure 11. Regions (NUTS ) ~~at risk~~exposed to multi-hazards identified with high significance,  $p < 0.01$  (left), and  $p < 0.10$  (right) as  
 455 hotspots/cold-spots and their correlation coefficient (Spearman  $r$ ) with independent variables: a.) empirical data – fatalities, b.)  
 empirical data – count of events .

Therefore, more significant the multi-hazard clustering, stronger is the relationship with the independent variables. The  
 monotonic relationship is strong–  $r = 0.59$  with the fatalities as independent variable for the regions with the highest  
 455 significance  $p < 0.01$  while the for the event count the strongest correlation ( $r = 0.40$ ) is reached for the regions with the  
 significance  $p < 0.05$ . This makes the recorded data on fatalities a better explanatory variable for the clustered population ~~at~~  
~~risk~~exposed to multi-hazards.

## 5. Discussions

460 The identification of exposure or risk on the DRMKC RDH platform is done generally from relating an asset to a hazard. There is also the possibility to relate an asset to multiple hazards and have a multi-hazard assessment (of exposure or risk ) on the single asset. This latter situation is the central aspect of the analysis presented in this study which considers the relation of a single asset (e.g. population or the residential built-up) to multiple hazards: landslide, coastal flood, river flood, earthquake, wildfires and subsidence. Starting from this initial setting of the analysis specific characteristics and limitations  
465 need to be presented.

First, we show that the proposed methodology allows for the detection of the regions at-riskexposed to multi-hazards, differently, as function of the typology of the assets. This is important as it directly reveals relationships between assets types and threats, valuable for the identification of the disaster risk management pathways in multi-hazard assessment (Ward, P. J., et al., 2022).

470 Furthermore, we argue for an approach that identifies the regions (local administrative units) prone to multi-hazard with high level of significance. We adopt a meta-analysis approach, combining single hazard hotspots which seeks to solve the problem of “insignificant results” and provides an objective “statistical proof” of the multi-hazard potential of a region. We support these results through a validation process which considers empirical data as explanatory variables. We show that more significant is the multi-hazard clustering, stronger is the correlation relationship with the independent variables.

475 With this study we also show that the proposed methodology allows for detecting changing patterns of the population being at-riskexposed from multi-hazard by considering the socio-economic dimension. Our findings are in line with previous studies which present an increase in risk to multi-hazard from low income countries towards higher income countries, and then a decrease as countries’ income is the highest (Koks et al., 2019). We have also evidenced the highly urbanized regions (urban area) as a space of risk (Hansjurgens et al., 2008) compared with the rural administrative units from  
480 multi-hazard occurrence. Furthermore, we show the potential of this methodological approach in detecting the risk to multi-hazard associated with complex socio-economic urban processes. We indicate that high density of population is a good explanatory variables for the increase in risk of the metropolitan areas. However this situation is particularly different in the case of high income metropolitan areas where more at-riskexposed to multi-hazards are the population living in the (less densely populated) functional urban area.

~~We also exemplify the applicability of the multi hazard methodological approach within the multi hazard interaction theoretical framework developed by Gill and Malamud (2014). We demonstrate it is scalable down to local and regional level and we present it as one future step in support of DRM decision making at national and subnational level.~~

Whilst we believe that the disaster risk management for multi-hazard assessment is brought forward by the ability of the proposed approach to identify regions (LAUs) being at-riskexposed to multi-hazards with high significance, several  
490 shortcomings are identified.

Most important shortcoming is that the presented case study does not consider the vulnerability for the assessment of the assets (population and residential built-up) ~~at risk~~exposed to multi-hazards. The multi-hazard potential of regions is measured in this study by means of exposure (or assets ~~at risk~~exposed). Nevertheless, the overall analytical approach is detecting significant patterns of multi-hazard potential across regions, revealing spatially explicit clusters in a heterogeneous groups of data and thereby setting the ~~bas~~ies for more precise and focused analysis.

Furthermore the clusters are identified at the level of areal dimension (represented by LAUs). The areal dimension approach excludes a detailed level of study that could more accurately examine the spatial coincidence of multiple hazards at localized levels. Also, by subdividing the exposure data at the level of areal dimension which are heterogenous in size (see 2.1.1.) will introduce underestimations or overestimations of the clusters especially when the clustering analysis is based on neighbouring relations defined by distance. However, we identified the optimal  $k$  value (dynamic for any relation hazard-asset) in order to reduce the susceptibility to noise and outliers used in the clustering analysis .

A way of improving the results accuracy and a direction for future research includes the revision of the meta-analysis (based on the Stouffer's method), used in this analysis. The choice is whether to use the weighted or unweighted versions of the Z-transform test for Stouffer method when combining the single hazards hotspots into multi-hazard hotspots. There are arguments in the statistical literature (Whitlock, M.C. 2005) that favour the weighted Z-approach especially when there is variation in the sample size across studies/clusters (e.g. the number of regions depending on the exposure type) as it is the case in our study. However the weighted or unweighted version of this test is actually an open question in meta-analysis (Becker, 1994).

## 6 Conclusions

To our knowledge, this is the first study that uses spatial patterns (clusters/hotspots) and meta-analysis in order to identify the regions at European level ~~at risk~~ exposed to multi-hazards. The methodology presented in this study provides multi-hazard enhanced insights, valuable for the identification of the disaster risk management pathways in multi-hazard risk assessments. The findings point out the socio-economic dimension as a determinant factor for the spatial variability and the risk potential of the local administrative units to multi-hazard. We show that the high density of population is a good explanatory variable for the identification of the regions ~~exposed~~at risk to multi-hazards but the economic aspect is the main driver that controls the risk status at local level: within rural and urban areas and in complex socio-economic urban structure. We also exemplify the applicability of the multi-hazard methodological approach within a multi-hazard interaction theoretical- framework and we demonstrate that is scalable to local and regional level and replicable.

-By identifying local administrative units with high level of significance as being ~~exposed~~at risk to multi-hazards we also narrow the- uncertainty around the major challenges related with multi-hazard studies: identification of the regions prone to multi-hazard; ~~and~~ quantification of multi-hazard ~~exposure, impact and characterization of the relation between the multiple~~

hazards. Future research should aim to address identified limitations by incorporating vulnerability assessments into multi-hazard analyses and the consideration of a multi-hazard interaction framework, for improved accuracy and reliability of multi-hazard hotspots identification.

525

The outcome of this study brings forward an useful methodological input that is made available for use through the Risk Data Hub platform. The objective of this study is to support national authorities on addressing the multi-hazard approach in the National Risk Assessments preparation.

530

535

540

545

## References

- Aldenderfer, M., S. and Blashfield, R. K., *Cluster Analysis (Quantitative Applications in the Social Sciences)* (Newbury Park, CA: SAGE Publications, Inc), 1984
- 550 Baker, F.B., Hubert, L.J., “*Measuring the Power of Hierarchical Cluster Analysis.*” *Journal of the American Statistical Association*, 70(349), 31–38,1975
- Bates, P. D., Horritt, M. S., and Fewtrell, T. J.: *A simple inertial formulation of the shallow water equations for efficient two-dimensional flood inundation modelling*, *J. Hydrol.*, 387, 33–45, 2010.
- 555 Becker, B.J. 1994. *Combining significance levels. In: The Handbook of Research Synthesis* (H. Cooper & L. V. Hedges, eds), pp. 15–230. Russell Sage Foundation, New York.
- Birkmann J, Cardona OM, Carreño ML, Barbat AH, Pelling M, Schneiderbauer S, Kienberger S, Keiler M, Alexander D, 560 Zeil P, Welle T (2013) *Framing vulnerability, risk and societal responses: the MOVE framework*. *Nat Hazards* 67(2):193–211.
- Borenstein, M., Hedges, L. V., Higgins, J. P. T. and Rothstein, H. R. (2009) *Front Matter, in Introduction to Meta-Analysis*, John Wiley & Sons, Ltd, Chichester, UK. doi: 10.1002/9780470743386.
- 565 Cabinet Office (2011) *Keeping the country running: natural hazards and infrastructure. A guide to improving the resilience of critical infrastructure and essential services*. Cabinet Office, London. [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/61342/natural-hazards-infrastructure.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/61342/natural-hazards-infrastructure.pdf). Accessed 22 Mar 2015
- 570 [Camus, P., Haigh, I.D., Nasr, A.A., Wahl, T., Darby, S.E., Nicholls, R.J., 2021. Regional analysis of multivariate compound coastal flooding potential around Europe and environs: sensitivity analysis and spatial patterns. Nat. Hazard Earth Sys. 21.](#)
- Cardona, O.D., M.K. van Aalst, J. Birkmann, M. Fordham, G. McGregor, R. Perez, R.S. Pulwarty, E.L.F. Schipper, and B.T. Sinh, 2012: Determinants of risk: exposure and vulnerability. In: *Managing the Risks of Extreme Events and Disasters to 575 AdvanceClimate Change Adaptation* [Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.D. Mastrandrea, K.J. Mach,G.-K. Plattner, S.K. Allen, M. Tignor, and P.M. Midgley (eds.)]. *A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC)*. Cambridge University Press, Cambridge, UK, and New York, NY, USA, pp. 65-108.



- 580 Charlton, M., Kemp, K. K.,: *Encyclopaedia of Geographic Information Science*, SAGE Publications, Inc. (2008), pp. 289-  
290
- Chen, Z. 2011. *Is the weighted z-test the best method for combining probabilities from independent tests?*, *Journal of Evolutionary Biology* 24: 926–930.
- 585 Claassen, J. N., Ward, P. J., Daniell, J., Koks, E. E., Tiggeloven, T., & de Ruiter, M. C. (2023). A new method to compile global multi-hazard event sets. *Scientific Reports*, 13(1), 1–14. <https://doi.org/10.1038/s41598-023-40400-5>
- Cousineau, D. and Chartier, S. (2010) Outliers Detection and Treatment: A Review. *International Journal of Psychological*  
590 *Research*, 3, 58-67.
- Cover, T. M., Hart, P. E. , "*Nearest Neighbor Pattern Classification*," *IEEE Trans. Inform. Theory*, vol. IT-13, pp. 21- 27, 1967.
- 595 [Dallo, I., Stauffacher, M., Marti, M., \(2020\): What defines the success of maps and additional information on a multi-hazard platform?. International Journal of Disaster Risk Reduction, Volume 49, 2020,101761, ISSN 2212-4209, https://doi.org/10.1016/j.ijdr.2020.101761](https://doi.org/10.1016/j.ijdr.2020.101761)
- 600 [De Angeli, S., Malamud, B. D., Rossi, L., Taylor, F. E., Trasforini, E., Rudari, R., A multi-hazard framework for spatial-temporal impact analysis, \(2022\), International Journal of Disaster Risk Reduction, Volume 73, 102829.](https://doi.org/10.1016/j.ijdr.2022.102829)
- 605 [De Fino, M., Tavolare, R., Bernardini, G., Quagliarini, E., Fatiguso, F., \(2023\) Boosting urban community resilience to multi-hazard scenarios in open spaces: A virtual reality – serious game training prototype for heat wave protection and earthquake response, Sustainable Cities and Society, Volume 99, 104847, ISSN 2210-6707, https://doi.org/10.1016/j.scs.2023.104847.](https://doi.org/10.1016/j.scs.2023.104847)
- De Pippo, T., Donadio, C., Pennetta, M., Petrosino, C., Terlizzi, F., and Valente, A.: *Coastal hazard assessment and mapping in Northern Campania, Italy*, *Geomorphology*, 97, 451- 466, doi:10.1016/j.geomorph.2007.08.015, 2008.
- 610 Dilley, M., Chen, R. S., Deichmann, U., Lerner-Lam, A. L., and Arnold, M.: *Natural Disaster Hotspots, a global risk analysis*, the World Bank, p. 112, p. 132, 2005.

Dunford R, Harrison PA, Jager J, Rounsevell MDA, Tinch R (2014) *Exploring climate change vulnerability across sectors and scenarios using indicators of impacts and coping capacity*. Clim Chang. doi:10.1007/s10584-014-1162-8

615

[Fekete, A., Rufat, S., \(2023\), Should everyone in need be treated equally? A European survey of expert judgment on social vulnerability to floods and pandemics to validate multi-hazard vulnerability factors, International Journal of Disaster Risk Reduction, Volume 85, 103527, ISSN 2212-4209, https://doi.org/10.1016/j.ijdr.2023.103527.](https://doi.org/10.1016/j.ijdr.2023.103527)

620 Fotheringham, A.S. and Wong, D.W.S. (1991) *The modifiable areal unit problem in multivariate statistical analysis, Environment and Planning A*, 23: 1025-1044.

Fisher, R. 1932. *Statistical methods for research workers*. Oliver and Boyd, Edinburgh.

625 Fix, E. & Hodges, J.L., 1951: "Nonparametric Discrimination: Consistency Properties", Randolph Field, Texas, Project 21-49-004, Report No. 4.

Gallina, V., Torresan, S., Critto, A., Sperotto, A., Glade, T., Marcomini, A., 2016. *A review of multi-risk methodologies for natural hazards: consequences and challenges for a climate change impact assessment*. J. Environ. Manage. 168, 123-132,

630 <https://doi.org/10.1016/j.jenvman.2015.11.011>.

Gaillard, J. C., & Mercer, J. (2013). From knowledge to action: Bridging gaps in disaster risk reduction. *Progress in Human Geography*, 37(1), 93–114. <https://doi.org/10.1177/0309132512446717>.

635 [Gentile, R., Cremen, G., Galasso, C., Jenkins, L. T., Manandhar, V., Mentese, E. Y., Guragain, R., McCloskey, J., \(2022\), Scoring, selecting, and developing physical impact models for multi-hazard risk assessment, International Journal of Disaster Risk Reduction, Volume 82, 103365, ISSN 2212-4209, https://doi.org/10.1016/j.ijdr.2022.103365.](https://doi.org/10.1016/j.ijdr.2022.103365)

640 [Ghanbari, M., Arabi, M., Kao, S.-C., Obeysekera, J., & Sweet, W. \(2021\). Climate change and changes in compound coastal-riverine flooding hazard along the U.S. coasts. Earth's Future, 9, e2021EF002055. https://doi.org/10.1029/2021EF002055.](https://doi.org/10.1029/2021EF002055)

Gill, J. C. and Malamud, B. D.: *Reviewing and visualizing the interactions of natural hazards*, Rev. Geophys., 52, 680–722, 2014.

645

Gill, J. C. and Malamud, B. D (2016).: Hazard interactions and interaction networks (cascades) within multi-hazard methodologies. Earth Syst. Dynam., 7, 659–679, <https://doi.org/10.5194/esd-7-659-2016>, 2016.

650 Gill, J. C., Malamud, B. D., Barillas, E. M., and Guerra Noriega, A.: Construction of regional multi-hazard interaction frameworks, with an application to Guatemala, *Nat. Hazards Earth Syst. Sci.*, 20, 149–180, <https://doi.org/10.5194/nhess-20-149-2020>, 2020.

Getis, A. and Ord, J. K. (1992), *The Analysis of Spatial Association by Use of Distance Statistics*. *Geographical Analysis*, 24: 189–206. doi:10.1111/j.1538-4632.1992.tb00261.x

655

G., T, Kappes, M., S., Frigerio S, Malet J.,P. (2012) *Multi-hazard exposure analyses with multirisk—a platform for user-friendly analysis*. 12th Congress INTERPRAEVENT, Grenoble, France, pp 487–495

660 Granger, K., Jones, T., Leiba, M., & Scott, G. (1999). *Community Risk in Cairns: A Multi-Hazard Risk Assessment*. Canberra: Commonwealth of Australia Publishing.

Greiving, S., Fleischhauer, M., and Lückenkötter, J.: *A methodology for an integrated risk assessment of spatially relevant hazards*, *J. Environ. Plann. Man.*, 49, 1–19, 2006.

665 Grünthal, G., Thieken, A. H., Schwarz, J., Radtke, K. S., Smolka, A., and Merz, B.: *Comparative risk assessment for the city of Cologne-storms, floods, earthquake*, *Nat. Hazards*, 38, 21–44, 2006.

Hak, T., Van Rhee, H. J., & Suurmond, R. (2016). *How to interpret results of meta-analysis*. (Version 1.3) Rotterdam, The Netherlands: Erasmus Rotterdam Institute of Management. [www.irim.eur.nl/researchsupport/meta-essentials/downloads](http://www.irim.eur.nl/researchsupport/meta-essentials/downloads)

670

Hewitt, K. and Burton, I. (1971) *The Hazardousness of a Place: A Regional Ecology of Damage Events*, University of Toronto Press.

675 Han, J., Wu, S., and Wang, H.: *Preliminary study on geological hazard chains*, *Earth Science Frontiers*, 14, 11–20, doi:10.1016/S1872-5791(08)60001-9, 2007.

IPCC (2012) *Managing the risks of extreme events and disasters to advance climate change adaptation. A special report of working groups I and II of the intergovernmental panel on climate change*. Cambridge University Press, Cambridge.

- 680 Jelinski, D.E. and Wu, J. 1996: *The modifiable areal unit problem and implications for landscape ecology*. Landscape Ecology 11, 129–40.
- Kappes, M. S., Keiler, M., and Glade, T.: *From single- to multi-hazard risk analyses: a concept addressing emerging challenges*, in: *Mountain Risks: Bringing Science to Society*, edited by: Malet, J. P., Glade, T., and Casagli, N., CERG  
685 Editions, Strasbourg, France, 351–356, 2010.
- Kappes M.S., Gruber, K., Frigerio, S., Bell, R., Keiler, M., Glade, T. 2012. *The multirisk platform: the technical concept and application of a regional-scale multihazard exposure analysis tool*. *Geomorphology* 151–152:139–155. doi:10.1016/j.geomorph.2012.01.024
- 690
- Kappes, M. S., Keiler, M., von Elverfeldt, K., and Glade, T.: *Challenges of analyzing multi-hazard risk: a review*, *Nat. Hazards*, 64,1925–1958, 2012
- Kwan, M-P.:. *The Uncertain Geographic Context Problem*, *Annals of the Association of American*  
695 *Geographers*, volume:102/5, pg.958-968, Routledge, doi:10.1080/00045608.2012.687349,2012.
- .
- Lee, R., White, C. J., Adnan, M. S. G., Douglas, J., Mahecha, M. D., O’Loughlin, F. E., Patelli, E., Ramos, A. M., Roberts, M. J., Martius, O., Tubaldi, E., van den Hurk, B., Ward, P. J., & Zscheischler, J. (2024). Reclassifying historical disasters: From single to multi-hazards. *Science of the Total Environment*, 912(October 2023), 169120.  
700 <https://doi.org/10.1016/j.scitotenv.2023.169120>
- Lipták, T. 1958. *On the combination of independent tests*. *Magyar Tud. Akad. Mat. Kutato Int. Közl3*: 171–196
- Liu, B., Siu, Y. L., and Mitchell, G.: Hazard interaction analysis for multi-hazard risk assessment: a systematic classification based on hazard-forming environment, *Nat. Hazards Earth Syst. Sci.*, 16, 629–642, <https://doi.org/10.5194/nhess-16-629-2016>, 2016  
705
- Marzocchi, W., Mastellone, M., Di Ruocco, A., Novelli, P., Romeo,E., and Gasparini, P.: Principles of Multi-Risk Assessment: Interactions Amongst Natural and Man-Induced Risks, European Commission, Directorate-General for  
710 Research, Environment Directorate, Luxembourg, 72 pp., 2009
- Marzocchi, W., Garcia-Aristizabal, A., Gasparini, P., Mastellone,M., and Di Ruocco, A.: *Basic principles of multi-risk assessment: a case study in Italy*, *Nat. Hazards*, 62, 551–573, 2012.

- 715 Mahendra, R. S., Mohanty, P. C., Bisoyi, H., Kumar, T. S., and Nayak, S.: *Assessment and management of coastal multi-hazard vulnerability along the Cuddalore–Villupuram, east coast of India using geospatial techniques*, *Ocean Coast. Manage.*, 54, 302–311, doi:10.1016/j.ocecoaman.2010.12.008, 2011.
- Lancaster, H. :*The combination of probabilities: An application of orthonormal functions*. *Aust. J. Stat.* 3, 20–33 (1961).
- 720 Openshaw, S. (1984) : *The Modifiable Areal Unit Problem*, CATMOG 38, GeoBooks, Norwich.
- Peter A. G. van Bergeijk, Sara L., 2015. "Macroeconomics of Natural Disasters: Strengths and Weaknesses of Meta-Analysis Versus Review of Literature," *Risk Analysis*, John Wiley & Sons, vol. 35(6), pages 1050-1072, June.
- 725 Preston, B. , D., Abbs, B., Beveridge, C., Brooke, Gorddard R, Hunt G, Justus M, Kinrade P, Macadam L, Measham T G,McInnes K, Morrison C, O'Grady J, Smith T F and Withycombe G 2007 : *Spatial approaches for assessing vulnerability and consequences in climate change assessments Modelling & Simulation*, Society of Australia & New Zealand 261.
- Sadegh, M., Moftakhari, H., Gupta, H. V., Ragno, E., Mazdiyasn, O., Sanders, B., Matthew, R., & AghaKouchak, A. (2018). Multihazard scenarios for analysis of compound extreme events. *Geophysical Research Letters*, 45, 5470–5480.
- 730 [https:// doi.org/10.1029/2018GL077317](https://doi.org/10.1029/2018GL077317)
- Schneider, S. H., Semenov, S., Patwardhan, A., Burton, I., Magadza, C. H.D., Oppenheimer, M., Pittock, A. B., Rahman, A., Smith, J.B., Suarez, A., and Yamin, F. (2007), *Assessing Key Vulnerabilities and the Risk from Climate Change*. In: *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (M .L. Parry, O.F. Canziani, J.P. Palutikof, P.J. van der Linden and C.E. Hanson, Eds.). Cambridge University Press, Cambridge, UK.
- 735 *Report of the Intergovernmental Panel on Climate Change* (M .L. Parry, O.F. Canziani, J.P. Palutikof, P.J. van der Linden and C.E. Hanson, Eds.). Cambridge University Press, Cambridge, UK.
- Stillwell, J., Daras, K., Bell, M. and Lomax, N. (2014) *The IMAGE studio: a tool for internal migration analysis and modelling*, *Applied Spatial Analysis and Policy*, 7(1): 5-23
- 740
- Tarvainen, T., Jarva, J., and Greiving, S.: *Spatial pattern of hazards and hazard interactions in Europe*, in: *Natural and Technological Hazards and Risks Affecting the Spatial Development of European Regions*, edited by: Schmidt-Thomé, P., Geological Survey Of Finland, Espoo, Finland, 42, 83–91, 2006
- 745 Theodoridis S, Koutroubas K (2008). *Pattern Recognition*. 4th edition. Academic Press.
- Tibshirani R, Walther G, Hastie T (2001). "Estimating the Number of Clusters in a Data Set Via the Gap Statistic." *Journal of the Royal Statistical Society B*, 63(2), 411–423.

- 750 van Westen, C. J., Montoya, L., Boerboom, L., & Badilla Coto, E. (2002). *Multi-hazard risk assessment using GIS in urban areas: a case study for the city of Turrialba*, Costa Rica. 120-136.
- van Westen C., Kappes M.S., Luna B.Q., Frigerio S., Glade T., Malet JP. (2014): *Medium-Scale Multi-hazard Risk Assessment of Gravitational Processes*. In: Van Asch T., Corominas J., Greiving S., Malet JP., Sterlacchini S. (eds)
- 755 *Mountain Risks: From Prediction to Management and Governance*. Advances in Natural and Technological Hazards Research, vol 34. Springer, Dordrecht
- Veerbeek W and Husson H 2013: *Vulnerability to Climate Change: Appraisal of a vulnerability assessment method in a policy context*. KfC report number 98/2013 (Unesco-IHE OR/MST/177)
- 760 Ward, P. J., Daniell, J., Duncan, M., Dunne, A., Hananel, C., Hochrainer-Stigler, S., Tijssen, A., Torresan, S., Ciurean, R., Gill, J. C., Sillmann, J., Couasnon, A., Koks, E., Padrón-Fumero, N., Tatman, S., Tronstad Lund, M., Adesiyun, A., Aerts, J. C. J. H., Alabaster, A., Bulder, B., Campillo Torres, C., Critto, A., Hernández-Martín, R., Machado, M., Mysiak, J., Orth, R., Palomino Antolín, I., Petrescu, E.-C., Reichstein, M., Tiggeloven, T., Van Loon, A. F., Vuong Pham, H., and de Ruiter, M.
- 765 C.: Invited perspectives: A research agenda towards disaster risk management pathways in multi-(hazard-)risk assessment, *Nat. Hazards Earth Syst. Sci.*, 22, 1487–1497, <https://doi.org/10.5194/nhess-22-1487-2022>, 2022.
- Ward, P.J., de Moel,H., Aerts, J.C.J.H., *How are flood risk estimates affected by the choice of return-periods?* *Natural Hazards and Earth System Science*, 11 (2011), pp. 3181-3195, 10.5194/nhess-11-3181-2011
- 770 Wipulanusat, W., Nakrod, S., Prabnarong, P. (2009): *Multi - hazard risk assessment using GIS and RS applications: a case study of Pak Phanang basin*. *Walailak Journal of Science and Technology* 6, 109 - 125.
- Whitlock, M.C. 2005: *Combining probability from independent tests: the weighted Z-method is superior to Fisher's approach*. *Journal of Evolutionary Biology* 18: 1368–1373.
- 775 Stouffer, S., DeVinney, L. & Suchmen, E. 1949: *The American soldier: Adjustment during army life*, vol. 1. Princeton University Press Princeton, US.
- 780 Zaykin, D., Zhivotovsky, L., Westfall, P. & Weir, B. 2002: *Truncated product method for combining P-values*. *Genetic Epidemiology* 22: 170–185.

Zaykin D. V. 2011: *Optimally weighted Z-test is a powerful method for combining probabilities in meta-analysis*. Journal of evolutionary biology, 24(8), 1836–1841. <https://doi.org/10.1111/j.1420-9101.2011.02297.x>

785

Zhao, M., Jingchao C., 2016: *Improvement and Comparison of Weighted k Nearest Neighbors Classifiers for Model Selection*. Journal of Software Engineering, 10: 109-118.

.

Ming Zhao and Jingchao Chen, 2016. *Improvement and Comparison of Weighted k Nearest Neighbors Classifiers for Model Selection*. Journal of Software Engineering, 10: 109-118.

790

Yoon, S., Baik, B., Park, T. et al. : *Powerful p-value combination methods to detect incomplete association*. Sci Rep 11, 6980 (2021). <https://doi.org/10.1038/s41598-021-86465-y>