

Answer to Referee 2

Although the subject of the study is interesting, there are many points that need to be adequately addressed.

OK.

1) The English in the text is not fluent, resulting in misunderstandings in several sections. It requires thorough proofreading by a native English speaker.

The language has been improved.

2) Throughout the text, the authors appear to use the term "seismic risk" to refer to the "precursor" of large events. It is important to note that seismic risk is defined as the product of seismic hazard and vulnerability. Seismic hazard refers to the natural phenomena generated by an earthquake, while seismic risk is the probability of human loss or damage to the built environment when exposed to a seismic hazard.

OK, we follow this precision and appropriate changes have been done through the text.

3) In the Introduction, the authors state: "The aim of the present analysis is to find patterns in the data sequence that could lead to understanding the following aspects of the process: a) entropic activity of the zone or sub-zone that can be an indicator of seismic risk; b) parameters serving as indicators of seismic risk; c) behavior of the data sequence during the earthquake and immediate aftershocks; d) recovery to the 'normal' or previous seismic activity following the aftershock period." However, these patterns or analyses seem poorly discussed in the text.

These points are now discussed in more details in Section 3. In particular, we now stress those aspects that could be considered precursors of important seismic activity. A paragraph dedicated to this aspect is now part of the conclusions.

4) The description of data selection in Section 2.1 appears to be quite complicated. Initially, the authors extract earthquakes with a maximum depth of 100 km, then they re-select those with a maximum depth of 70 km. Why not select events with a maximum depth of 70 km from the beginning? Furthermore, there is another potentially more critical issue: the authors first apply the GR law to select all events (among those with a maximum depth of 100 km) with a magnitude larger than the completeness magnitude, then from this subset, they extract events with a maximum depth of 70 km. I believe the selection process should be different: first, they should select events with a maximum depth of 70 km, then apply the GR law to this dataset and consider only the events with a magnitude larger than the completeness magnitude.

Yes, you are absolutely right. Several extractions were done during the preparation of this paper which caused the confusion. But now we go directly to the facts: what we use is an extraction with 19549 seisms which are used to get the GR plot (Fig. 2). From here we filter the data with magnitudes equal and over magnitude 2.1 and depth less or equal to 70 km, ending with a final catalog of 13871 events.

5) The estimation of b-value is missing.

Yes, in the present version the parameters of the adjustment to Gutenberg-Richter law, namely, $a=5.70$ and $b=0.73$, are explicitly given in the text and in Fig. 2.

6) Lines 72-73 “To the left, we appreciate how the activity of this region continuously increased irregularly since the beginning of the century”: this is not “activity” (generally indicated by the number of events in a specified timescale), but the magnitude or the “energy”.

This sentence was changed according to your comment.

7) Lines 75-77: “At this point, we can follow two complementary ways to analyze these data through entropy: a) magnitude sequence, and b) sequence of intervals between consecutive seisms. We have chosen the former since entropy is more easily associated with energy, even since the early thermodynamics courses. However, we will include some results using the latter in the Appendix”. Actually, this is quite a peculiar way to indicate their preference for analyzing magnitude instead of interevent times. Moreover, they mention that interevent times will be analyzed in the Appendix, which is missing!

Yes, we decided now to leave out any discussion concerning the interevent times. We think this point needs to be studied under a broader scope, including different geographical areas, to compare it with the magnitude analysis. Some early results we have indicate that entropy on interevent catalogs can have advantages over the magnitude analysis, but this needs to be established properly. In a previous version, Fig. 13 was in an Appendix whose heading was removed but the reference to it remained. Sorry for that. The section on time intervals and previous figure 13 were removed from the present version.

8) Line 78: “In Fig. 3, the depiction presents the magnitude of each of the 15 011 seisms at the time of their production.” Fig. 3 should be Fig. 4. And, please, rephrase: “Fig. 3 shows the time distribution of the analysed magnitudes”.

Yes, this was fixed. Figs. 3 and 4 were wrongly addressed in the previous version.

9) Lines 82-84: The sentence is really complicated. The authors can simply write “We calculated the Tsallis entropy and mutability in a moving window of size W , shifting by one event through the entire catalog and associate the calculated values to the time of the last event in the window”.

OK.

10) Line 90: “states” are simply “magnitudes”.

Done

11) Line 94-98: why so complicated way to calculate the average magnitude? Just simply say that within each window you calculate the average magnitude.

OK.

12) I don't understand why the authors mention Shannon entropy (eq. 4) if it is never used in the study! While I understand the desire to shorten the theoretical treatment, the way the authors have done it makes the text very difficult to comprehend. It would be better to include all the details that allow the reader to understand how to calculate q .

There are two different ideas in this comment. First, a referee for a previous paper insisted that we should mention Shannon entropy as another source of treating the data, so we included previous Eq. (4) to show the connection to our approach. Now we removed Eq. (4) and left just a comment (not a numbered equation) in the section devoted to Mutability. Second, since we had explicitly provided the calculation of q in previous articles (Posadas and Sotolongo-Costa 2023 and Posadas et al. 2023.), we considered unnecessary to repeat the entire procedure here. However, we have included the most relevant part of the treatment in Section 2.2 guiding the interested reader to the appropriate literature.

13) The same issue applies to the description of mutability. Mathematical details are required. Simply writing "the weight in bytes of the map created by wzip of the time window at position n " leaves the reader unsure about what "weight," "wzip," and "map" refer to.

Since this concept is rather new, we have included details using a portion of the catalog defined here to illustrate the procedure and the definitions of the terms you mention. Although similar examples were included in previous papers, it is perhaps helpful to remind them with a close example. The subsection 2.3, "Mutability" contains all this material in the present version.

14) Lines 119-123 are repeated in lines 131- 134.

Yes, this was fixed.

15) Line 152: "Fig. 5 reports the Tsallis' entropy calculated for windows $W = 512$ and $W = 1024$ ", Line 155: "Fig. 5 presents the average value of the Tsallis entropy over windows with $W=512$ and $W=1024$ ": is it the average value of Tsallis entropy or not??

Sorry, there is nothing like instant values in our calculation. Results always correspond to average values depending on the initial time and the span of the time window. Thank you for noticing this misunderstanding.

16) Lines 153-154: "The left-hand side presents the variations along the 24 years of this study, while the right-hand side zooms on the last 4 years, where most of the important activity shows up." This is repeated few lines below.

Repetition was eliminated.

17) Line 158: “it can be noticed that at the precise moment the large earthquakes are produced, the Tsallis entropy suddenly decreases”: better to write “The Tsallis entropy is characterized by a decrease before the occurrence of a large shock”

Yes, we followed your suggestion.

18) Line 161: “to detect risk in advance” please see my comment in the point2)

Yes, this was fixed.

19) Line 161: “poor statistics”??

We rephrased this expression.

20) Line 162: “The average magnitude tends to be minimized before a large earthquake”. Where is the figure showing this behavior of mean magnitude?

This comment was eliminated.

21) Lines 165-175: the authors try to explain on a physical basis the behavior of Tsallis entropy, but honestly, I cannot follow their argument. They talk about 4 degrees latitude, 6 degrees longitude, stalagmites, stalactites, compressible gas...I would suggest to delete all this explanation, which, furthermore, lacks of adequate literature to be based on.

OK, it was deleted.

22) In Line 158 “at the precise moment the large earthquakes are produced, the Tsallis entropy suddenly decreases.”; in line 179 “Mutability decreases previous to a large earthquake because small seisms of restricted magnitude dominate the sequence”, but in line 188 for $W=256$ “simultaneous increases of Tsallis entropy and decreases of mutability as the large earthquake approaches” the behavior of the two quantities is opposing. So, I should argue that the Tsallis and mutability have a correlated behavior for $W=512$ and 1024 and anticorrelated for $W=256$???. A DEEP explanation is NECESSARY.

Yes, you are right. We are comparing two instances: previous to the earthquake (precursors) and at the moment of the earthquake and soon afterwards (aftershock). They became entangled. We have rewritten the entire paragraph.

23) Line 205: “We choose to present the time in number of days “ but Figs. 8-12 shows years.

Everything is in terms of years now.

24) Line 217: “Then, near day 7500,” so, when?

Now we refer to the middle of year 2020 instead.

25) Section 3.5 shows results only for $W=256$. At least for sub-zoness A and D, the results should also be presented for $W=512$ (due to what observed in my previous point 22).

Yes, we followed your suggestion. It works well for subzone D, but not so well for subzone A. However, it serves to justify our discussion regarding to a minimum of records to do the statistics.

26) Fig. 13 should present also the results for $W=1024$.

The whole discussion on inter-event times was deleted, so Fig. 13 is no longer in the paper.

27) A further figure is necessary showing the time variation of the Tsallis entropy S_q for the interevent times.

Previous answer applies to this comment as well.

Let us say that we really appreciate your time and important comments, corrections and suggestions. Thank you so much.

2021 Alaska Earthquake: entropy approach to its precursors and aftershock regimes.

Eugenio Vogel^{1,2,3}, Denisse Pastén⁴, Gonzalo Saravia⁵, Michel Aguilera⁶, and Antonio Posadas^{7,8}

¹Departamento de Física, Universidad de La Frontera, Casilla 54-D, Temuco, Chile

²Center for the Development of Nanoscience and Nanotechnology (CEDENNA), 9170124 Santiago, Chile

³Facultad de Ingeniería, Universidad Central de Chile, Santiago 8330601, Chile

⁴Departamento de Física, Facultad de Ciencias, Universidad de Chile, Santiago, Chile

⁵Los Eucaliptus 1189, 4812537 Temuco, Chile

⁶Universidad Técnica Federico Santa María, Valparaíso, Chile.

⁷Departamento de Química y Física, Universidad de Almería, 04120 Almería, Spain.

⁸Instituto Andaluz de Geofísica, Campus Universitario de Cartuja, Universidad de Granada, 18071 Granada, Spain.

Correspondence: Denisse Pastén denisse.pasten.g@gmail.com

Abstract. We have conducted an entropy analysis in Alaska, a seismic-rich region in a subduction zone that exhibits a nontrivial behavior: the subduction arc alters the seismic activity from the eastern zone to the western zone, demonstrating a decrease in activity along the subduction. We analyze this zone through the Tsallis entropy and the mutability (or dynamic entropy) for the first time. Considering 13 870 seismic events after appropriate filtering, we analyzed a data set for the selected Alaska zone between 2000 and 2023. We have found agreement between the results for the two entropies. We have followed the value of the q parameter of the Tsallis entropy (S_q) finding values between 1.70 and 1.85, in concordance with values found in other seismic regions of the planet. The values of S_q decrease slightly over time but show a broad increase before the major earthquakes. Just opposite to Tsallis entropy, mutability shows a tendency to decrease previous to the major earthquakes. We used the simpler mutability method to further analyze this zone upon dividing the region into four sub-zones. The results show how mutability can identify the seismic activity in each zone. This study shows how an entropy approach can shed light on understanding the seismicity in subduction zones.

1 Introduction

The seismic background in Alaska has been a source of questions and studies for the last decades. Particularly, the subduction of the Pacific Plate under the North American Plate produces the Aleutian trench zone, which runs almost parallel to the arc of islands running on the Southern part of Alaska, pointing to Asia towards the west. The Aleutian arc has an extension of approximately 3000 km, from the Gulf of Alaska (east) to the Kamchatka Peninsula (west) (USGS, 2024). Like other subduction zones on the planet, it is a geologically active area in both underground seismicity and surface volcanic eruptions.

So, the Alaska-Aleutian Region has a rich history of large earthquakes which we briefly summarize next: Shumagin Island in 1938 (M_w 8.2); Aleutian Islands in 1946 (M_w 8.6); Queen Charlotte Islands in 1949 (M_w 8.1); Lituya Bay in 1958 (M_w 8.2); Prince William Sound in 1964 (M_w 9.2) which is the second largest earthquake worldwide since there are reliable registers;

Rat and Near Island segment in 1965 (M_w 8.7) (Qu et al., 2022). In recent years, at least four large earthquakes have occurred in this zone: the Simeonof earthquake in July 2020 (M_w 7.8); the Alaska Peninsula earthquake in October 2020 (M_w 7.6); the South Alaska Peninsula in July 2021 (M_w 8.2), almost exactly one year after the previous and the late aftershock of 2023 (M_w 7.2). This high seismic activity in the Alaska Peninsula in the last three years deserves dedicated attention, which is one
25 main purpose of this article. Researchers have studied the Alaska subduction zone from different points of view for several years (Biswas et al., 1986; Doser and Rodriguez, 2011; Smith and Tape, 2019; Daly et al., 2021; Qu et al., 2022). In this present article, we will present an analytical and numeric approach based on the variations of two complementary forms of entropy, applied to catalogs of magnitudes.

The Pacific Plate is moving in a northwest direction at average rates of 60 mm/year in the east, and 76 mm/year in the west.
30 The rich activity of this zone lies in the different tectonic sources of its seismicity (Martin-Short et al., 2018). On one hand, the seismicity of the central and the eastern portions of the arc are both greater than the one of the western portion. In the latter, the seismic activity is shallow, and the volcanic activity decreases, in comparison with the central and eastern zones of the Aleutians. At least 12 large earthquakes have occurred in the eastern and central areas of the arc in the last century, with magnitudes greater than M_w 7.5.

35 As has been shown, the analysis of seismicity through information theory has deepened our understanding of this system out of equilibrium. Studies based on entropy have been able to follow the time evolution of seismicity and they are especially useful in subduction tectonic zones (Sigalotti et al., 2023; Skordas et al., 2020; Varotsos et al., 2018; Vallianatos et al., 2015; Telesca, 2010, 2011; Vogel et al., 2017; Posadas et al., 2022, 2023; Pasten et al., 2023). There are different ways to define and use entropy. In this study, we will use two specific approaches: Tsallis entropy and mutability (a form of dynamical entropy).
40 To study the time evolution of the seismic data, we will use an enumeration of the events; this method can be compared to the well-known concept of natural time (Varotsos et al., 2011, 2019).

The primary aim of the present analysis is to find patterns in the data sequence that could lead to understanding the following aspects of the process: a) entropic activity of the zone or subzone that can be an indicator of seismic hazard; b) parameters serving as indicators of seismic hazard; c) behavior of the data sequence during the earthquake and immediate aftershocks; d)
45 recovery to the “normal” or previous seismic activity following the aftershock period.

The next section devotes itself to methods, starting with the extraction of data from the USGS catalog and filtering mechanisms. Then, the researchers quickly review Tsallis entropy and mutability, primarily referring to previous publications to avoid repetition. Section 3 is where the authors focus on results and discussions, while they dedicate Section 4 to conclusions.

2 Methodology

50 2.1 Data

We make use of the catalog of the United States Geological Survey (USGS) on the website <https://earthquake.usgs.gov/earthquakes/search/>. The process starts by defining the geographical area of interest, which, in our case, is determined by the

geographic coordinates: $53.5^{\circ} \text{ (N)} \leq \text{Latitude} \leq 57.5^{\circ} \text{ (N)}$ and $161.0^{\circ} \text{ W} \leq \text{Longitude} \leq 155.0^{\circ} \text{ W}$. In this way, we include the most important recent earthquakes in Alaska.

55 Next, we need to specify the period of observation. Since we want to study signs of previous activity, we will extend it from January 1, 2000, to December 31, 2023. The original extraction of data yielded 19 549 seisms, set that was used to do the Gutenberg Richter analysis (details below). From this last analysis, we set a minimum seismic magnitude of 2.1; to concentrate on seismicity near the planet's surface (mainly subduction) we set a limit at a depth of 70 km, which also coincides with a previous study in a different region (Posadas et al., 2023). After applying this filtering, we got a catalog of 13 870 seisms that
60 we will use for the analysis and to draw figures from 3 to 7. The analysis by subzones has its own subsets of seisms defined in subsection 3.4.




Figure 1. Illustration of the seisms with epicenters within the volume between 53.5° N and 57.5° N , 155° W and 161° W and up to 70 km deep. The largest stars correspond to the 4 earthquakes listed in Table 1 and they are labeled A, B, C, and D, following the order of occurrence. Other symbols correspond to seisms of increasing magnitude according to the following color code: $2.0 \leq M_w \leq 3.0$ yellow; $3.1 \leq M_w \leq 4.0$ cyan; $4.1 \leq M_w \leq 5.0$ blue; $5.1 \leq M_w \leq 6.0$ orange; $6.1 \leq M_w \leq 9.0$ red.

| Date | M_w | Latitude | Longitude | Depth (km) | Label |
|------------|-------|----------|-----------|------------|-------|
| 2020.07.22 | 7.8 | 55.07N | 158.60W | 28.0 | A |
| 2020.10.19 | 7.6 | 54.60N | 159.63W | 28.4 | B |
| 2021.07.29 | 8.2 | 55.36N | 157.89W | 35.0 | C |
| 2023.07.16 | 7.2 | 54.39N | 160.76W | 25.0 | D |

Table 1. Data for the four main seisms within the zone of interest.

Fig. 1 shows the geographical area defined by this search, representing the magnitude of the seisms with the following color codes: $2.0 \leq M_w \leq 3.0$ yellow; $3.1 \leq M_w \leq 4.0$ cyan; $4.1 \leq M_w \leq 5.0$ blue; $5.1 \leq M_w \leq 6.0$ orange; $6.1 \leq M_w \leq 9.0$ red. Large red stars denote the four largest seisms tabulated in Table 1, identified by the letters A, B, C, and D, following the order of happening.

Fig. 2 reports the Gutenberg Richter analysis for the original extraction of 19 549 records. Following the principle of maximum curvature, we can pick a magnitude of $M_w 2.1$ as the minimum magnitude compatible with a distribution characterized by a linear decrease in this diagram. The linear regression is represented by a straight line with $a = 5.70$ (interception with the ordinate axis), and $b = 0.73$ (slope with a negative sign in the expression). Following Eq. (4) we get $q = 1.73$. In addition, we set at 70 km the maximum depth to concentrate on the seisms nearer to the surface, then we are left with 13 870 seisms for the global analysis.




Figure 2. Left) Figure 2. Left) Gutenberg Richter plot for the 19 549 seisms initially considered in this paper; threshold magnitude from the maximum curvature method is $M = 2.1$. Blue circles represent the number of events in each magnitude bin and orange circles are cumulative frequency-magnitude distributions for all seismic events. A linear fit over the linear part of the blue circles distribution provides the correlation coefficients a and b , which are given in the equation as an inset. An overall estimate of the b value is 0.73. Right) Depth frequency for the 19 549 original seisms, with a red vertical line at 70 km to show the maximum depth to be considered in the rest of this paper.

We present the time distribution of the 13 870 seisms in Fig. 3. To the left, we appreciate how the energy of this region continuously increased irregularly since the beginning of the century. To the right, the lack of activity prior to the largest seisms is more evident upon zooming to the last 4 years, which is one of the precursors we will bring out of the analysis. At first glance, we notice the irregularity of the sequence, as we can expect from this phenomenon which is largely stochastic. In addition, the decrease in activity preceding the large earthquakes reflects the accumulation of energy to be released with the incoming seisms of larger magnitudes.




Figure 3. Left) Magnitude of the sequence of 13 870 along the years 2000-2023. Right) Detail of the years 2020-2023 where the seisms A, B, C, and D are clearly appreciated by their magnitudes over 7.0. Noteworthy is the decrease in seismic activity just before these large earthquakes.

We calculated the Tsallis entropy and mutability in a mobile window of size W , shifting by one event through the entire catalog, and associate the calculated values to the time of the last event in the time window W .

80 2.2 Tsallis Entropy

Within each window of W registers, we find a distribution of data where f_i corresponds to the frequency of magnitude M_i . Then, we can easily calculate the probability p_i of obtaining the value M_i as:

$$p_i = \frac{f_i}{W} . \tag{1}$$

It immediately follows that:

$$85 \sum_{i=1}^{\Omega} p_i = 1 , \tag{2}$$

adding over all the Ω accessible microstates within the observation period.

From here on we make use of the fragment-asperity model developed by Sotolongo-Costa and Posadas since 2004 (Sotolongo-Costa and Posadas, 2004) and continued to recent extensions of the model (Posadas et al., 2023; Pasten et al., 2023). We take it from Eq. (1) of this last reference, namely:

$$90 \quad \log(N(> M)) = \log(N) + \frac{2-q}{1-q} \log(1 + a(q-1)(2-q)^{\frac{1-q}{q-2}} 10^{2M}), \quad (3)$$

where the left-hand side is given as the upper function in the Gutenberg-Richter law, N is the abscissa there, q is the Tsallis entropic index to be determined, a is the proportionality between the seismic energy and the size of the fragments, which can be left as an adjusting parameter. Following Sarlis et al. 2010 (Sarlis et al., 2010), Telesca et al. 2010 (Telesca, 2010) a relationship between b value and q parameter can be found. Following (Posadas et al., 2023; Pasten et al., 2023) we can write:

$$95 \quad b = 2 \frac{2-q}{q-1}, \quad (4)$$

which can be compared to the linear descent of the Gutenberg-Richter law. From here, we can obtain the b value and then q .

Different approaches have been followed to get the non-extensive parameters, and the interested reader can find details in recent literature (Posadas et al., 2023; Flores-Márquez et al., 2024). In addition, combining these techniques with natural time analysis has provided a deep insight into the changes in entropy previous to a large earthquake. The early work of Sarlis, Skordas, and Varotsos (Sarlis et al., 2010) concentrated on the seismicity of California and Japan. This work was continued to report precursors one day before the Tohoku undersea mega-thrust earthquake of 11 March 2011 (M_w 9.0) tsunami lasting 6 minutes, which also caused a large tsunami Varotsos et al. (2023a). This method was extended to the seismology of Japan, Mexico, and California (Varotsos et al., 2023b). Moreover, the work by Telesca and collaborators has also inspired the present work (Telesca, 2010). This brief review shows that non-extensive entropy analysis can be very important in determining seismic hazards all over the world.

Once q is determined, the Tsallis entropy calculation is straightforward:

$$S_q = \frac{1}{q-1} \left(1 - \sum_{i=1}^{\Omega} p_i^q \right) \quad (5)$$

Boltzmann-Gibb's entropy is recovered in the limit $q \rightarrow 1$ but for most seismic zones $q > 1.0$ (interested reader is referred to Appendix A in Flores-Márquez et al., (2024) (Flores-Márquez et al., 2024)).

110 2.3 Mutability

Let us consider a segment of $W = 24$ consecutive magnitudes corresponding to the beginning of the year 2023 within the rectangle already defined for Alaska. They are listed in the second column of Table 2, while the first column labels the sequence.

To form the third column, go to the original vector file (second column) and retrieve the first record (2.4 in this case), write it down in the first slot of the third column, followed by the distance to the origin of this file (0 in this case). ii) Go then to the next (second) record (2.3) and verify if this has already an assigned slot: if not, it writes this magnitude value in the third column followed by its distance to the origin; if yes, it increases the corresponding slot to the right as explained below. iii)

| i | Magn | Map | f_i | $p_i = f_i/W$ |
|----|------|---------------|-------|---------------|
| 1 | 2.4 | 2.4 0 16 | 2 | 2/24 = 0.083 |
| 2 | 2.3 | 2.3 1 13 6 | 3 | 3/24 = 0.125 |
| 3 | 2.6 | 2.6 2 11 | 2 | 2/24 = 0.083 |
| 4 | 2.2 | 2.2 3,2 3 16 | 4 | 4/24=0.167 |
| 5 | 2.2 | 2.1 5,2 2,5 5 | 8 | 8/24 = 0.333 |
| 6 | 2.1 | 2.7 15 7 | 2 | 2/24 = 0.083 |
| 7 | 2.1 | 3.0 16 | 1 | 1/24=0.042 |
| 8 | 2.2 | 3.7 18 | 1 | 1/24=0.042 |
| 9 | 2.1 | 2.5 19 | 1 | 1/24=0.042 |
| 10 | 2.1 | | | |
| 11 | 2.1 | | | |
| 12 | 2.1 | | | |
| 13 | 2.1 | | | |
| 14 | 2.6 | | | |
| 15 | 2.3 | | | |
| 16 | 2.7 | | | |
| 17 | 3.0 | | | |
| 18 | 2.1 | | | |
| 19 | 3.7 | | | |
| 20 | 2.5 | | | |
| 21 | 2.3 | | | |
| 22 | 2.4 | | | |
| 23 | 2.7 | | | |
| 24 | 2.2 | | | |

Table 2. Example of the way wzip constructs the map storing the compressed file for the first 24 seisms of the year 2023. Sequential numbers are listed in column 1; the corresponding measured magnitudes for these seisms is given in column 2, whose vector file has a weight W in bytes; Column 3 stores the map created by wzip as explained in the text, weighting W^* bytes; The frequency of these magnitudes is listed in column 4, leading to the normalized probability of finding this magnitude in this sample given in column 5.

Continue with the next record in the second column (2.6 in this example); if it is new, it writes it down on the next slot in the third column: if not (as in this example) it goes to the next slot in the third column followed by 2, which is its distance to the origin. iv) It repeats the process for the next register (2.2) followed by its distance to the origin (3), but there is a novelty: since this value repeats itself forming a string of 2 consecutive values, then it adds a comma and the number of consecutive repetition: 2 in this case. v) Then, the program follows to find the record 2.4, which is already stored in the first slot of the third column. It goes there and writes 5 as the distance to the previous equivalent record. vi) Then it is 2.1, the next record, which is new, 6 positions away from the origin. vii) The next record is 2.2, which is already known, with at a distance 3 from to the previous last appearance. viii) Then the already known 2.1 register comes again, two positions after the last show, and repeats itself 5 consecutive times (2,5 to the right). And it continues in this way, as the reader can verify with the rest of the file.

An appropriate algorithm can readily count the frequency f_i for the i -th value in the fourth column of Table 2. Following Eq. 1, we can calculate the probability p_i of each accessible value by dividing f_i by W . With this information,

we could calculate Shannon entropy $H = -\sum p_i \ln(p_i)$. However, we will not go in that direction in the present article to concentrate on the seismological activity described by Tsallis entropy and mutability.

130 The third column is the recognized or compressed file, whose weight is w^* . Then the mutability ζ is given by:

$$\zeta = \frac{w^*}{w}, \quad (6)$$

where w is the weight of the original file in the second column.

The more repetitive and the sooner these repetitions occur, the lower the value of ζ . A succession of quite different registers occupies distinct lines producing an extensive file, increasing w^* and hence increasing also the value of ζ .

135 We recently provided all technical details concerning this process in (Pasten et al., 2023), so we will not repeat additional details here. Let us just remind that mutability reaches its minimum value for repetition of information (magnitude of the seisms in the present case). Before a major earthquake occurs, the subduction mechanism nearly halts, allowing only small advancements between the plates. These small advancements result in medium to low magnitudes that are very similar to each other, enabling better compression and a lower value of mutability. When the rocks finally collapse, they produce a large quake
140 with a high magnitude, followed by a dispersion of magnitudes that differ significantly from each other. This sudden change in magnitudes causes the mutability to rise abruptly, resembling upward needles in the mutability diagrams below.

3 Results and Discussion

3.1 Results for Tsallis entropy

From equation 5, we realize that Tsallis entropy is determined by the value of q got from the data distribution. Fig. 4 presents
145 the variations of q with time for the region under study, for $W = 512$ along the 24 years. Values span a range between 1.70 and 1.85. The texture of the curves varies according to the window size, and the ranges decrease slightly with larger windows. Fluctuations around 1.75 (most of the values in Fig. 5) agree well with previously calculated q values for other earthquakes in other regions of the world (Sotolongo-Costa and Posadas, 2004; Silva et al., 2006; Telesca and Chen, 2010; Telesca, 2011; Darooneh and Mehri, 2010; Valverde-Esparza et al., 2012; Varotsos et al., 2023a, b).

150 Sotolongo-Costa introduced a method (which we essentially follow here) and reported q values in the range 1.60 to 1.64 for Southern Spain and 1.65 for California (Sotolongo-Costa and Posadas, 2004). From there, this method and its variations have been applied to various regions of the world. Thus, we find the work of Silva et al. (Silva et al., 2006) reporting q values of 1.60, 1.63, and 1.71 for data from Brazil, the USA, and Turkey, respectively. Then, Dahoornet and Mehri (Darooneh and Mehri, 2010) reported values of 1.78 and 1.81 for seismic areas in Iran and California, respectively. Luciano Telesca studied
155 the L'Aquila seismic region by non-extensive entropy and reported q values of 1.48 in the early period increasing to 1.74 and 1.70 later on (Telesca, 2010). A similar analysis was then done for the seismicity of Taiwan reporting $q = 1.685$ (Telesca and Chen, 2010). Also, the value of $q = 1.66$ was found for Japan in (Sarlis et al., 2010). For the seismicity of California, a variation of these methods yielded $q = 1.54$ (Telesca, 2011). The seismicity of Mexico was also explored by nonextensive methods, and q values of 1.7, 1.69, 1.63, and 1.64 were reported for different areas (Valverde-Esparza et al., 2012).




Figure 4. Variations of the Tsallis q value along the period of study for mobile overlapping windows of $W = 512$ consecutive events.

160 The differences among the q values in the brief review of the previous paragraph can be because of both variations in the method and the different geological characteristics of the different zones picked for this sampling. These facts clearly establish that: 1) q is much larger than 1.0, requiring non-extensive treatments, and 2) The ranges reported by Fig. 4 are of the same order of magnitude as other active seismic zones.

165 Upon reaching the period of stronger seismic activity (after 2020 in Fig. 4) large variations of the q value occur. Prior to this period, q values oscillate with periods of 2 to 4 years over a baseline (not drawn) which is slightly descending. This can be a precursor behavior, showing that the zone is increasing the chance of a large earthquake in the future.

Fig. 5 reports the average Tsallis' entropy calculated for windows of $W = 512$ and $W = 1024$, along with the 13 870 seisms covered by this report. The left-hand side presents the variations along the 24 years of this study, while the right-hand side zooms in the last 6 years, where most of the important activity shows up.

170 The relevant seisms A, B, C, and D given in Table 1 are also marked in Fig. 5 by the symbols given in the upper part of the plot. Regardless of the time windows, a decrease in Tsallis entropy is observed with the occurrence of an enormous shock. The larger window ($W = 1024$) keeps most of the characteristics of the function, but it has two disadvantages: i) It shows less texture than the one with $W = 512$, and ii) it shows results after 5 years of gathering data to reach the 1024 seisms; such delay is not desirable to detect hazard in advance. So we will prefer $W = 512$ when possible, accepting $W = 256$ where the number
175 of records is small.




Figure 5. Left) Tsallis' entropy for the seismic sequence from 2000 to 2023 using mobile windows of $W = 512$ events (black) and 1024 events (red). On the top line, the largest seisms A, B, C, and D of Table 1 are clearly indicated. Noteworthy is the slight long-run increase in entropy during the years prior to the large earthquakes. Right) Zoom on the recent years of the same data.

3.2 Results for mutability

180 Fig. 6 reports the variations of mutability for the same windows of 512 and 1024 consecutive events used with Tsallis entropy plotted in the previous figure. The left-hand side reports the mutability for the entire time range from 2000 to 2023, while the right-hand side zooms on the last 4 years, which is where the large earthquakes concentrate. Mutability decreases previous to a large earthquake because small seisms of restricted magnitude dominate the sequence. The general slope of decreasing the mutability value is also clear for the $W = 1024$ window, with the disadvantage of the delay in time between cause and effect. The absolute minimum mutability after the $M_w 7.6$ earthquake, whose aftershock regime was suppressed, is notorious. This is almost an announcement of the 8.2 earthquake, which shows here as an isolated “needle” pointing upwards on the right-hand side of the mutability function. So the decrease in the mutability value can also be a precursor of large earthquakes.




Figure 6. Left) Mutability on magnitude for the seismic sequence from 2000 to 2023 using mobile windows of $W = 512$ events (black) and 1024 events (red). On the top line, the largest seisms A, B, C, and D of Table 1 are clearly indicated. Noteworthy is the long run decrease in mutability during the years previous to the large earthquakes. Right) Zoom on the recent years of the same data.

185 3.3 Tsallis entropy and mutability

The agreement between Tsallis entropy and mutability is clearly reflected in Fig. 7. This brief window of $W = 256$ events is noisier than previous ones, which underlines the coincidences between these two functions. Despite being calculated by completely different algebraic procedures, the information content embedded in the distribution of magnitude values leads to simultaneous increases of Tsallis entropy and decreases of mutability as the large earthquake approaches. Downwards needles from Tsallis entropy coincide completely with upwards needles from mutability. The response of both functions to the $M_w 7.2$ is weak, which is noteworthy.

This agreement was even more significant with the seismic activity a few days previous to the $M_w 8.1$ earthquake in front of Iquique (Chile) in the year 2014 (see Fig. 7 of Ref. (Pasten et al., 2023)). Both mutability and Tsallis entropy revealed an incoming seismic activity, increasing from days to minutes before the large quake. In the present case, we did not observe this effect for the recent Alaska earthquakes. For each of the 4 events reported in Table 1, we found a sudden increase (decrease) in mutability (Tsallis entropy) with no previous announcement. Obviously, the underground for these two zones can be completely different, and no generalizations are possible. But even for these 4 seisms reported in Table 1 each earthquake can reflect different underground dynamics. This is an excellent opportunity to investigate this point for 4 seisms close in geography and time. A careful look at Fig. 7 reveals that mutability shows more texture than Tsallis entropy for the same data. For this reason, we carry on with calculations of mutability only since these results have all the information that the time series can yield.




Figure 7. Tsallis entropy and mutability on magnitude along the years 2018-2023, for overlapping mobile windows of 256 events.

3.4 Subzones

We begin by defining 4 non-overlapping subzones on the map given in Fig. 1. Each subzone experienced an earthquake with a magnitude of over 7.0 near its center. It included a large cluster of smaller seismic events around it from January 1, 2020, to December 31, 2023, and up to a depth of 70 km. Table 3 summarizes the geographic definitions of the 4 subzones.

| Seism | M_w | Date | Latitude | Longitude | # events |
|-------|-------|------------|----------------|------------------|----------|
| A | 7.8 | 2020.07.22 | 54.0N to 55.0N | 160.5W to 161.0W | 929 |
| B | 7.6 | 2020.10.19 | 54.5N to 55.5N | 157.3W to 158.2W | 4021 |
| C | 8.2 | 2021.07.29 | 54.0N to 55.0N | 159.5W to 160.0W | 629 |
| D | 7.2 | 2023.07.16 | 50.0N to 50.5N | 158.0W to 158.5W | 908 |

Table 3. Definition of the subzones studied separately by means of mutability. The first column gives the label of the main seism; the second and third columns give its magnitude and its date, respectively. The fourth column gives the range in latitudes, while the fifth column gives the range in longitudes. The sixth column gives the number of seisms for each subzone.

205 The first important difference among the subzones is the total number of seisms collected over the 24 years for each one of them. They go from 629 for subzone C (containing the largest M_w 8.2 earthquake!) to 4 021 for the subzone B. Analyses based on $W = 1024$ are not possible, while those based on $W = 516$ are nearly meaningless. So, we will use $W = 256$ for the subzone analysis.




Figure 8. Magnitude of seismic activity in subzones A, B, C, and D for the last 8 years. Open diamonds mark the 4 important seisms characterizing each of the subzones at the top of the diagram.

The time distribution of the earthquakes shows apparent differences in their dynamics. Fig. 8 shows the evolution of the 210 magnitudes for the years 2016-2023. All subzones are quiet before the triggering event of 2020.07.22, except for subzone D (to some extent). Each subzone will be now characterized separately.

Subzone A shows nearly no seisms previous to the M_w 7.8 earthquake of 2020.07.22 and the magnitudes of the few previous 215 events are less than 4.0. Around mid-year 2020, the M_w 7.8 earthquake occurred with no precursor, followed by an usual aftershock regime of seisms at short intervals and decreasing magnitude. However, subzones B or D do not reflect any of this activity. But just a few days after this triggering event (July 22, say), a swarm of quakes with magnitudes around 5.0 occurred in the northeast sector of subzone D (Blue circles and orange triangles in Fig. 1). The upper part of Fig. 8 reflects this activity. It is not clear if the swarm in Subzone D is a consequence of the major earthquake in Subzone A, but it is a possibility.

Subzone B had weak previous activity, but the M_w 7.6 earthquake of 2020.10.19 unleashed a continuous aftershock regime 220 lasting until the last days of our recollection of data. Fig. 1 shows a “vertical” cluster of blue circles and orange triangles near latitude -159.5, showing this. The North-South activity spans all magnitudes from 2.1 to approximately 5.0. A recent example of the latter is the 2023.10.09 M_w 5.0 earthquake, just very close to the epicenter of earthquake B.

Subzone C had very weak activity before previous seisms; it did not show any response to neighboring seism A, but it follows the general aftershock regime generated by the 7.6 seism of 2020.10.19 in subzone B. Then, the subzone lost activity

to almost none, just prior to the M_w 8.2 earthquake that generated a modest aftershock activity here and in the neighboring A
225 subzone to some extent. Currently, the activity of sub zone C is like that of sub zone A.

Subzone D had recovered 2021 its quiet behavior with almost no seismic activity until the M_w 7.2 seism came in 2023, generating aftershock activity only here. These seisms are shown by blue circles and orange triangles to the East of the meridian at 160.5°W in Fig. 1. Comparison to other subzones in Fig. 8 shows only a possible connection to neighboring subzone B, near the corner 55°N and 160.5°W where a swarm of activity can be seen as blue circles and orange triangles in Fig. 8.

230 The magnitude sequences of the four subzones (Fig. 8) share the following common features: irregular time distribution of the seismic events; sparse time distribution of the seisms before the main earthquake of the subzone; nearly halting of the seismic sequence a few weeks before the main earthquake; abundance of seisms of different magnitudes after the main quakes (aftershock regime); little or no influence of large earthquakes in the sequence of other subzones.

They present large differences also: different number of total seisms; different distribution of the seisms with time; subzone
235 C presents almost no activity previous to the large earthquakes, while subzone D presents varieties of intensity activity previous to the largest earthquake in the subzone; subzone B exhibits an intense aftershock regime with seisms of magnitude over 4.0, while subzone A show much lower activity with few quakes reaching magnitude 3.0.

Some of the previous features can be useful to define precursors of a future earthquake in a determined area. However, the main message is that premonitory activity tends to be associated to lack of important seisms immediately before the main
240 quake, a signature that can be lost if the zone or subzone is too large and other activity foci cover up this lack of seisms.

All the previously described diversity in the seismic behavior is an indication of large differences in the subduction processes. Subzones only 100 km away (or even less) show a completely different evolution, although they are wrapped under the common process of plate subduction. We will now examine this process from the optics of the entropy encoded in the sequence of data.

3.5 Mutability analysis for the subzones

245 In the plots of Figs. 9, 10 11 and 12, we mark with a red filled circle the time of the M_w 8.2 earthquake, regardless of the zone. This is to realize that even the largest seism of this catalog does not have a direct influence on subzones nearby.

Fig. 9 shows an approximately constant initial mutability near a value 0.62 evidencing a calm period. Then a descent of mutability begins during 2018, reaching a minimum value of 0.60 to suddenly jump to higher levels on the day of the 7.8 seism of 2020. The empty green triangle shows the position of this A earthquake in coincidence with the upward needle shown by
250 the mutability function. The aftershock regime produces a chain of seisms with a variety of magnitudes that cannot be easily compressed by w/zip keeping high the values of mutability until it finally decreases at the beginning of 2021. However, during 2021, a lower tiny second needle in the mutability is appreciated: this is because of the M_w 8.2 earthquake in the neighboring region C. From there, it goes through a true minimum from which it recovers remaining at lower levels than before years 2016 and 2017. This tendency toward low values could be a precursor for future activity in this subzone.

255 The details in the discussion of the previous figure will be not explicitly repeated for the other three figures, since the cases are similar. Mutability levels cannot be strictly compared among subzones because they reflect the dynamics of the underground subduction in each place.




Figure 9. Mutability on magnitude for the 929 seisms of subzone A. An empty green triangle identifies the time for the largest seism of this subzone, while a solid red circle marks the time of the largest seism of the whole zone under study.

Highly surprising is subzone B: despite having the largest amount of seisms among the 4 subzones, it has almost no activity prior to the M_w 7.6 earthquake marked with an empty green triangle during the second half of year 2020 in Fig. 10. The initial calculation, making no modifications, yields a mutability of 0.61, which subsequently shows a strong maximum corresponding to the seism B, only to decrease later because of the aftershock regime. The mutability function clearly marks the needle coinciding with the 7.6 earthquake B, and then it gradually decreases to 0.53 in the typical oscillatory pattern of aftershock regimes. Its recovery is far from being complete, remaining at levels near 0.58, which is even lower than present values for subzone A. No significant activity can be associated here to the earthquake B despite its large magnitude.




Figure 10. Mutability on magnitude for the 4021 seisms of subzone B. An empty green triangle identifies the time for the largest seism of this subzone, while a solid red circle marks the time of the largest seism of the whole zone under study.

265 Subzone C has so few seismic events that the curve begins approximately 4 months before the largest M_w 8.2 earthquake. The needle of the mutability function coincides with the red circle marking the time of seism C. The mutability function goes down, remaining low at values near 0.615. It is hard to say anything else here because of the small amount of data.




Figure 11. Mutability on magnitude for the 629 seisms of subzone C. A solid red circle marks the time for the largest seism of this subzone, which is also the largest seism of the whole zone.

Subzone D shows the expected needle at the position of the M_w 7.2 earthquake marked with an empty green triangle towards the far right of this plot. However, we find an unexpected broad maximum during the second half of the year 2020. We had to go back to the time series to discover that this activity corresponds to a swarm of seisms with magnitudes between M_w 5.5 and M_w 5.8 beginning in this subzone on July 2020 and during for a few months. The present level of the mutability function is 0.62, similar to subzones A, and C. There is no activity associated to the largest M_w 8.2 earthquake marked with a red solid circle.




Figure 12. Mutability on magnitude for the 908 seisms of sub zone D. An empty green triangle identifies the time for the largest seisms of this subzone, while a solid red circle marks the time of the largest seisms of the whole zone under study.

Despite the differences among sub zones, we can notice the sensitivity of mutability to detect the seismic activity in the time sequence. Although a premonitory protocol is still far away, we can recognize the decrease in the mutability value as one element to investigate further as a precursor.

4 Conclusions

We have analyzed the seismicity in the Alaska subduction zone measured during the last 24 years through non-extensive thermodynamics using two entropies: Tsallis entropy and mutability (or dynamical entropy). We have followed the time evolution of the seismic activity in the eastern and central zones of the Aleutian arc between the years 2000 and 2023, mainly focusing on four large earthquakes that occurred in that zone during the last years of the period under study.

We have used an entropic approach to deal with data produced by the seismic activity grouped in a sequential series of W consecutive events. Such non-overlapping time windows are analyzed to produce results on Tsallis entropy $S_q(t)$, which can be summarized as follows. The dynamical values for $q(t)$ in the definition of the Tsallis entropy vary between 1.70 and 1.85,

285 which agrees with ranges of values reported in the literature in other subduction areas (Telesca, 2010, 2011; Silva et al., 2006; Telesca and Chen, 2010; Darooneh and Mehri, 2010; Valverde-Esparza et al., 2012).

Tsallis entropy increases before seismic activity and collapses to minimal values at the instant of the mega-thrust. This is a manifestation of the set of magnitude values brought in by the aftershock regime.

290 Mutability on magnitude values has precisely the opposite behavior than the one shown by mutability as presented in Fig. 7. Thus, mutability minimizes before an incoming large earthquake since the magnitudes of nearby seisms remain small because of the clogging in the plate subduction. This accumulates stresses until a rupture occurs, producing chains of interrelated events (like a domino effect), which are described by non-additive entropies like the Tsallis entropy.

The definition of small sectors within the area of study leads to 4 subzones with similarities and differences. Thus, in all subzones, the frequency of inter events halted before a large earthquake ($M_w > 7.0$), but the level of activity is large (like in 295 region D) or low (like in region B). So, even within a geographically restricted zone, the subduction process is different for processes separated just for a few dozen km.

The production of a large earthquake, like the M_w 8.2 in subzone C, can occur in a quiet zone, without previous activity and with no previous sign. This requires permanent monitoring of the seismic activity of Alaska to find indicators of premonitory activity.

300 Several possible precursors were emphasized in previous discussions. The low density of seisms prior to important earthquakes was stated in relation to Fig. 3. The evolution of q , the non extensive parameter, is also a possible precursor as its value oscillates with a general tendency to lower values prior to a large earthquake. Tsallis entropy S_q oscillates, tending slightly towards larger values. Simultaneously, the values of mutability decrease prior to a large earthquake. This last observation was confirmed during the subzone analysis. These variations occur over decades. It is highly probable that these observations are 305 not independent, but there is only a minor extra effort to calculate them all. Eventually, after applying them to different geographical areas and along different times of observation, a couple of them can serve as a long-term parameters warning of a possible earthquake in the coming years.

Finer precursors could be possible for relatively small subzones. However, in advance one does not know where to draw this small zone. Then the only possibility is to monitor several or many subzones, eventually overlapping among them, to find 310 out a couple of them showing an evolution similar to those presented in Fig. 8 just before a large quake. This can be further confirmed by the measurement of the mutability of the seismic sequence in that area.

The behavior of this subduction zone is not comparable to the previous analysis of the subduction zone between the Nazca Plate and the South-American Plate, where premonitory behaviors were detected. In the present case, we did not find a behavior that warns about an earthquake coming. However, both Tsallis entropy and mutability show the state in which the study areas 315 have been left, warning that the central area of the Aleutian arc is still very active.

Data availability. <https://earthquake.usgs.gov/earthquakes/search/>

Author contributions. The authors are contributed equally to this manuscript.

Competing interests. The authors declare no competing interest.

Acknowledgements. Partial support from the following two Chilean sources is acknowledged: Fondecyt under contract 1230055, Financiamiento Basal para Centros Científicos y Tecnológicos de Excelencia (Chile) through the Center for Development of Nanoscience and Nanotechnology (CEDENNA) under contract AFB220001. This research has been partially supported by the Agencia Estatal de Investigación (grant no. PID2021-124701NBC21 y C22); the Universidad de Almería (grant no. FEDER/UAL Project UAL2020-RNM-B1980); the Consejería de Universidad, Investigación e Innovación, Junta de Andalucía (grant no. RNM104). PPITUAL, Junta de Andalucía-FEDER 2021-2027. Programa: 54.A. A.P., D.P. and E.E.V. have been partially funded by the Spanish Project LEARNIG PID2022-143083NB-I00 by the Agencia Estatal de Investigación.

References

- Biswas, N.N. and Pujol, J., Tytgat, G., and Dean, K.: Synthesis of seismicity studies for western Alaska., *Tectonophysics*, 131, 369–392, [https://doi.org/doi.org/10.1016/0040-1951\(86\)90183-6](https://doi.org/doi.org/10.1016/0040-1951(86)90183-6), 1986.
- Daly, K. A., Abers, G. A., Mann, M. E., Roecker, S., and Christensen, D. H.: Subduction of an oceanic plateau across southcentral Alaska: High-resolution seismicity., *Journal of Geophysical Research: Solid Earth*, 126, e2021JB022809, <https://doi.org/doi.org/10.1029/2021JB022809>, 2021.
- Daroonch, A. H. and Mehri, A.: A nonextensive modification of the Gutenberg-Richter law: q-stretched exponential form, *Physica A: Statistical Mechanics and its Applications*, 389, 509–514, <https://doi.org/https://doi.org/10.1016/j.physa.2009.10.006>, 2010.
- Doser, D. I. and Rodriguez, H.: A seismotectonic study of the Southeastern Alaska Region, *Tectonophysics*, 497, 105–113, <https://doi.org/https://doi.org/10.1016/j.tecto.2010.10.019>, 2011.
- Flores-Márquez, E., Ramírez-Rojas, A., and Sigalotti, L.: Non-Extensive Statistical Analysis of Seismicity on the West Coastline of Mexico., *Fractal Fract.*, 8, 306, <https://doi.org/doi.org/10.3390/fractalfract8060306>, 2024.
- Martin-Short, R., Allen, R., Bastow, I. D., Porritt, R. W., and Miller, M. S.: Seismic imaging of the Alaska subduction zone: Implications for slab geometry and volcanism., *Geochemistry, Geophysics, Geosystems*, 19, 4541–4560, <https://doi.org/doi.org/10.1029/2018GC007962>, 2018.
- Pasten, D., Vogel, E. E., Saravia, G., Posadas, A., and Sotolongo, O.: Tsallis Entropy and Mutability to Characterize Seismic Sequences: The Case of 2007-2014 Northern Chile Earthquakes, *Entropy*, 25, <https://doi.org/10.3390/e25101417>, 2023.
- Posadas, A., Morales, J., Ibáñez, J., and Posadas, A.: Shaking earth: Non-linear seismic processes and the second law of thermodynamics: A case study from Canterbury (New Zealand) earthquakes., *Chaos, Solitons and Fractals. Nonlinear Science, and Nonequilibrium and Complex Phenomena*, 151, 111 243, <https://doi.org/https://doi.org/10.1016/j.chaos.2021.111243>, 2022.
- Posadas, A., Pasten, D., Vogel, E. E., and Saravia, G.: Earthquake hazard characterization by using entropy: application to northern Chilean earthquakes, *Natural Hazards and Earth System Sciences*, 23, 1911–1920, <https://doi.org/10.5194/nhess-23-1911-2023>, 2023.
- Qu, R., Ji, Y., Zhu, W., Zhao, Y., and Zhu, Y.: Fast and Slow Earthquakes in Alaska: Implications from a Three-Dimensional Thermal Regime and Slab Metamorphism, *Applied Sciences*, 12, <https://doi.org/10.3390/app122111139>, 2022.
- Sarlis, N., Skordas, E., and Varotsos, P.: Nonextensivity and natural time: The case of seismicity., *Phys Rev E*, 82, 021 110, <https://doi.org/https://doi.org/10.1103/PhysRevE.82.021110>, 2010.
- Sigalotti, L., Ramírez-Rojas, A., and Vargas, C.: Tsallis q-Statistics in Seismology., *Entropy*, 25, 408, <https://doi.org/https://doi.org/10.3390/e25030408>, 2023.
- Silva, R., França, G. S., Vilar, C. S., and Alcaniz, J. S.: Nonextensive models for earthquakes, *Phys. Rev. E*, 73, 026 102, <https://doi.org/10.1103/PhysRevE.73.026102>, 2006.
- Skordas, E., Sarlis, N., and Varotsos, P.: Precursory variations of Tsallis non-extensive statistical mechanics entropic index associated with the M9 Tohoku earthquake in 2011., *Eur. Phys. J. Spec. Top.*, 229, 851–859, <https://doi.org/https://doi.org/10.1140/epjst/e2020-900218-x>, 2020.
- Smith, K. and Tape, C.: Seismic noise in central Alaska and influences from rivers, wind, and sedimentary basins., *Journal of Geophysical Research: Solid Earth*, 124, 11,678–11,704, <https://doi.org/doi.org/10.1029/2019JB017695>, 2019.
- Sotolongo-Costa, O. and Posadas, A.: Fragment-Asperity Interaction Model for Earthquakes, *Phys. Rev. Lett.*, 92, 048 501, <https://doi.org/10.1103/PhysRevLett.92.048501>, 2004.

- Telesca, L.: Analysis of Italian seismicity by using a nonextensive approach, *Tectonophysics*, 494, 155–62, <https://doi.org/https://doi.org/10.1016/j.tecto.2010.09.012>, 2010.
- 365 Telesca, L.: Tsallis-based nonextensive analysis of the southern California seismicity, *Entropy*, 13, 1267–80, <https://doi.org/https://doi.org/10.3390/e13071267>, 2011.
- Telesca, L. and Chen, C.-C.: Nonextensive analysis of crustal seismicity in Taiwan, *Natural Hazards and Earth System Sciences*, 10, 1293–1297, <https://doi.org/10.5194/nhess-10-1293-2010>, 2010.
- USGS: M 8.6 - 1946 Aleutian Islands (Unimak Island), Alaska Earthquake, https://earthquake.usgs.gov/earthquakes/eventpage/official19460401122901_30/region-info, 2024.
- 370 Vallianatos, F., Michas, G., and Papadakis, G.: A description of seismicity based on non-extensive statistical physics: a review. In: D'Amico S, editor. *Earthquakes and their impact on society.*, Cham: Springer Natural Hazards, pp. 1–41, <https://doi.org/https://dx.doi.org/10.1007/978-3-319-21753-6>, 2015.
- Valverde-Esparza, S., Ramirez-Rojas, A., Flores-Márquez, E. L., and Telesca, L.: Non-extensivity analysis of seismicity within four subduction regions in Mexico., *Acta Geophys.*, 60, 833–845, <https://doi.org/doi.org/10.2478/s11600-012-0012-1>, 2012.
- Varotsos, P., Sarlis, N., and Skordas, E.: Tsallis Entropy Index q and the Complexity Measure of Seismicity in Natural Time under Time Reversal before the M9 Tohoku Earthquake in 2011, *Entropy*, 20, 757–73, <https://doi.org/https://doi.org/10.3390/e20100757>, 2018.
- Varotsos, P., Sarlis, N., Skordas, E., Nagao, T., and Kamogawa, M.: Natural time analysis together with non-extensive statistical mechanics shorten the time window of the impending 2011 Tohoku M9 earthquake in Japan, *Communications in Nonlinear Science and Numerical Simulation*, 125, 107 370, <https://doi.org/https://doi.org/10.1016/j.cnsns.2023.107370>, 2023a.
- 380 Varotsos, P., Sarlis, N., Skordas, E., Nagao, T., Kamogawa, M., Flores-Márquez, E., Ramírez-Rojas, A., and Perez-Oregon, J.: Improving the Estimation of the Occurrence Time of an Impending Major Earthquake Using the Entropy Change of Seismicity in Natural Time Analysis., *Geosciences*, 13, 222, <https://doi.org/https://doi.org/10.3390/geosciences13080222>, 2023b.
- Varotsos, P. A., Sarlis, N., Skordas, E. S., Uyeda, S., and Kamogawa, M.: Natural time analysis of critical phenomena, *P. Natl. Acad. Sci. USA*, 108, 11 361–11 364, 2011.
- 385 Varotsos, P. A., Sarlis, N. V., and Skordas, E. S.: Phenomena preceding major earthquakes interconnected through a physical model, *Ann. Geophys.*, 37, 315 – 324, 2019.
- Vogel, E., Saravia, G., and Ramirez-Pastor, A.: Phase transitions in a system of long rods on two-dimensional lattices by means of information theory, *Physical Review E*, 96, 062 133, 2017.