

What can we learn about multi-hazard impacts from global disaster records?

Wiebke S. Jäger¹, Marleen C. de Ruiter¹, Timothy Tiggeloven¹, Philip J. Ward^{1,2}

¹Institute for Environmental Studies (IVM), Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

5 ²Deltares, Delft, The Netherlands

Correspondence to: Wiebke S. Jäger (w.s.jaeger@vu.nl)

Abstract. Recent studies have reported more extreme, compounding impacts from multi-hazards than from single hazards owing to complex interrelationships of hazard, exposure and vulnerability. However, our current understanding of multi-hazard impacts is primarily based on case studies of individual events. To complement this, we examine the disaster records of the global emergency events database EM-DAT from 2000 to 2018. We develop an algorithm to identify multi-hazard events using the information on associated hazards as well as spatiotemporal relationships between disaster records. We find that 35% of events and 61% of hazards are multi-hazard when considering a spatial overlap of at least 50% and a time lag of at most 3 months. Overall, the multi-hazards accounted for 78% of the total damages, 83% of the total people affected and 69% of the total deaths. We also statistically compare the impacts of hazards pairs, single hazards and combinations of two single hazards. The analysis suggests distinct patterns of compounding impacts, which vary depending on hazard and impact type. We conceptualize four archetypes (“the whole is greater than the sum of its parts”, “the whole equals the sum of its parts”, “one part determines the whole”, and “the whole and the parts are limited by total impact”) to describe these patterns and to guide the integration of multi-hazard interrelationships into risk assessments. All archetypes have in common that hazard pairs have at least as much impact as single hazards and as combinations of two single hazards. Nonetheless, the uncertainties and limitations encountered highlight that future research should focus on improving data on multi-hazards and their impacts.

1 Introduction

In the past decade, multiple studies have reported disproportionate impact amplifications during multi-hazard or compound events (de Ruiter et al., 2020; Gill & Malamud, 2016; Zscheischler et al., 2018). Such events are characterised by hazardous conditions overlapping in space or time or both. Examples are: the 2018 Osaka earthquake followed by flooding and landslides (de Ruiter et al., 2020); concurrent heatwaves in major breadbasket regions (Kornhuber et al., 2020); and floods from spatially and temporally coinciding pluvial, fluvial and coastal drivers (Eilander et al., 2023).

Impact amplifications during multi-hazard events can arise from several different elements of disaster risk that interrelate with each other (De Angeli et al., 2022). These interrelationships can be on the hazard, exposure as well as vulnerability level and include feedback and dynamic processes. Throughout the article, we follow the UNDRR (2017) definitions for risk, hazard, exposure and vulnerability. Moreover, we use the term “multi-hazard impact” for impact generated from multiple hazards and accounting for all interrelationships on any level following Ward et al. (2022). Table 1 provides an overview of the key definitions used in this article.

35

It is widely recognized that disregarding such interrelationships can lead to an over- or underestimation of risk (De Angeli et al., 2022; de Ruiter & van Loon, 2022; Hillier et al., 2020; Kappes et al., 2012; Leonard et al., 2014; Terzi et al., 2019; Ward et al., 2022; Zscheischler & Seneviratne, 2017). It can also lead to ineffective or even harmful risk reduction strategies as measures to decrease the risk of one hazard may lead to increase of risk of another hazard (de Ruiter et al., 2020; Hurk et al., 2023; Ward et al., 2022). This means that multi-hazard impact cannot simply be modelled by adding up the impacts from single hazard models, which is known as multilayer single-hazard approach (Zschau, 2017). Instead, interrelationships of the risk elements should be considered in risk modelling, as highlighted in the UN's Sendai Framework (UNDRR, 2017) and reflected in the IPCC's AR6 cycle (IPCC, 2023).

So far, most multi-hazard research has focused on hazard-hazard interrelationships, and several classification systems have been proposed (De Angeli et al., 2022; Gill & Malamud, 2014; Liu et al., 2016; Tilloy et al., 2019; van Westen & Greiving, 2017; Zscheischler et al., 2020). Though the terms used in these classifications differ, they describe similar and overlapping concepts including statistical (in)dependence between hazards, spatiotemporal relationships, amplifications of magnitude, and triggering relationships. Methodological reviews and guidelines for quantifying interrelationships have also been published (Bevacqua et al., 2021; Tilloy et al., 2019). Hazard-exposure and hazard-vulnerability interrelationships have been researched less extensively, but a number of types have already been identified. For instance, changes in exposure can arise due to migration and evacuation (Tierolf et al., 2023) or due to losses and damages from a previous hazard that are not yet recovered (De Angeli et al., 2022). Furthermore, de Ruiter & van Loon (2022) identified and discuss key types of dynamics of vulnerability, such as the effects of an earlier hazard on the vulnerability at the time of a second hazard. It has also been identified that a combined load from multiple hazards can cause higher damage than the summed damages of the separate hazards (Li et al., 2012; Zuccaro et al., 2008).

To our knowledge, few studies exist that consider the interrelationships on all levels together to investigate their overall effect on impact. One example is the comprehensive modelling framework developed by De Angeli et al. (2022), which integrates interrelationships on all levels to assess multi-hazard impact on the built environment and illustrates this by a case study with a hypothetical combined seismic and flood scenario. Another example is the multivariate linear regression analysis by Budimir et al. (2014), which shows that past earthquake-and-landslide events were associated with more

fatalities than earthquakes alone when considering several independent covariates representing hazard, exposure and vulnerability elements. However, data limitations prevented the authors from assessing whether the hazard pair is associated with more fatalities than the sum of the constituent hazards or, in other words, whether impact amplifications arise.

In general, the way in which impact data are currently collected and stored makes it difficult to study and understand compounding impacts in a multi-hazard or compound event context. Issues range from missing data and biases, which affect the reliability of the data (e.g., Gall et al., 2009), to the single-hazard focus and limited spatiotemporal information of many well-known impact databases, such as HANZE for floods (Paprotny et al., 2018), the NOAA natural hazards data for tsunamis, earthquakes and volcanic eruptions (NOAA, n.d.) or DESINVENTAR (UNDRR, n.d.) for various types of hazards. The single-hazard focus necessitates the investigation and linkage of possible multi-hazard conditions, but this is hindered by the limited spatiotemporal information. Finally, impacts are being observed and stored on an event level and additional (statistical) methods are needed to attribute them to individual hazard components (Budimir et al., 2014).

Recently, Lee et al. (2024) have shown that the information on main and associated disasters in the emergency events database EM-DAT (Delforge et al., 2023) can be used to classify the disaster records into different types of multi-hazard events. However, hazards occurring simultaneously or in close succession at the same location have been reported in separate disaster records in multiple instances. The Guatemala 2010 volcanic eruption and tropical cyclone (Gill & Malamud, 2014) and the tropical cyclones Idai and Kenneth hitting Mozambique in 2019 (de Ruiter & van Loon, 2022) are two examples of hazards that are recognized as multi-hazard events in the scientific literature but reported as separate records in EM-DAT.

New possibilities for leveraging EM-DAT for multi-hazard analyses arise from the recently developed GDIS dataset of geocoded disaster locations (Rosvold & Buhaug, 2021) as well as MYRIAD-HESA, an algorithm for identifying multi-hazard events on the basis of spatiotemporal overlaps (Claassen et al., 2023). In this article, we make use of these possibilities to reexamine the disaster records in EM-DAT. Our aim is to gain a better understanding of compounding impacts of multi-hazards for different types of hazards and impacts.

We identify multi-hazard events following the approach by Lee et al. (2024) in combination with the GDIS dataset and a MYRIAD-HESA-inspired algorithm to account for spatiotemporal overlaps of disaster records. We focus on events with a (partial) spatial overlap and their immediate impacts rather than so-called systemic or complex impacts which can also arise from spatially distinct but temporally coinciding events due to global and sectoral interconnectedness (Hochrainer-Stigler et al., 2020; Simpson et al., 2021). We extract and derive impacts of hazard pairs and impacts of single hazards for different hazard types and perform a statistical analysis to compare impacts of hazard pairs, single hazards and combinations of two

single hazards. Based on the identified differences and similarities, we distinguish four “archetypes” of compounding impacts, which can guide the integration of multi-hazard interrelationships into risk assessments.

Table 1 Definitions of terms used in this article

Term	Definition	Source
Risk	A combination of hazard, exposure and impact vulnerability as illustrated by the conceptual equation: Hazard x Exposure x Vulnerability	(UNDRR, 2017)
Hazard	A process, phenomenon or human activity that may cause loss of life, injury or other health impacts, property damage, social and economic disruption, or environmental degradation.	(UNDRR, 2017)
Exposure	The situation of people, infrastructure, housing, production capacities and other tangible human assets located in hazard-prone areas.	(UNDRR, 2017)
Vulnerability	The conditions determined by physical, social, economic, and environmental factors or processes which increase the susceptibility of an individual, a community, assets, or systems to the impacts of hazards.	(UNDRR, 2017)
Multi-hazard	The selection of multiple major hazards that the country faces and the specific contexts where specific hazards may occur over time simultaneously, cascadingly or cumulatively over time, and taking into account interrelated effects.	(UNDRR, 2017)
Multi-hazard impact /risk	Impact / risk generated from multiple hazards as well the interrelationships between these hazards and on the vulnerability and exposure level.	(Ward et al., 2022)

100 **2 Data**

This study uses the international disaster database EM-DAT (Delforge et al., 2023), which contains information on natural hazards and their impacts together with the global dataset of geocoded disaster locations GDIS (Rosvold & Buhaug, 2021), which contains geospatial footprints of the impact areas.

2.1 EM-DAT

105 EM-DAT is, to our knowledge, the only publicly available data source with global coverage for disaster events that includes multiple hazard types and quantitative information on socio-economic impacts. The database records events with substantial impact that are related to natural as well as technological hazards from 1900 – present. Substantial impact is defined as an event that resulted in either at least ten deaths, at least 100 people affected, a call for international assistance or an emergency declaration. Each entry corresponds to a disaster event at country level. Events that span multiple countries are reported
110 separately for each country, as opposed to being recorded as a single physical disaster event. For example, the 2004 Indian Ocean earthquake and tsunami is reported in 12 individual disaster records, including records in several Asian and African countries.

Each disaster record in EM-DAT contains mandatory and optional fields. The mandatory fields relevant to this study are the unique event identifier, the country, the continent, the start year, and the disaster type. We also use the optional fields, although data are frequently missing. Relevant optional fields are: the disaster subtype; a first and second associated disaster, which represent subsequent or co-occurring hazards that may have contributed to the disaster impact; the start date and end date; and several human and economic impact variables.

In terms of impact, we consider the number of people affected, number of deaths, and damages. Throughout the following sections we will use the term “impact” to refer to these three quantities. Their definitions are:

- Number of people affected: number injured, number affected and number homeless, where number affected are the people needing immediate assistance due to the disaster. If only the number of families affected or houses damaged are reported, the figure is multiplied by the average family size for the affected area.
- Number of deaths: confirmed fatalities directly imputed to the disaster plus missing people whose whereabouts since the disaster are unknown and so they are presumed dead based on official figures.
- Damages: refers to total economic damage in US \$ adjusted for inflation.

While EM-DAT is being widely used in disaster risk science (Jones et al., 2022), it has well-known issues related to reporting biases (Gall et al., 2009) as well as the general reliability and accuracy of the impact data (Guha-Sapir & Below, 2002; Moriyama et al., 2018; Panwar & Sen, 2020). The biases are due to having entire records missing rather than fields missing within records (Gall et al., 2009). These include time bias, hazard-related bias, threshold bias, accounting bias, geographic bias, as well as systemic bias. We exclude data from before the year 2000 to minimize time bias as recommended by the maintainers of EM-DAT (Delforge et al., 2023). However, the other bias types remain, posing a limitation to this study. For example, heatwaves are known to be underreported in EM-DAT (Brimicombe et al., 2021; Harrington & Otto, 2020).

Guidelines for handling biases and missingness are still lacking in disaster risk science. Approaches for missingness differ across studies. Deletion, augmentation and imputation, or a combination of these, are most common for studies using EM-DAT as a primary or secondary data source (Jones et al., 2023). Deletion is simpler, but deemed inferior to augmentation and imputation because it poses a higher risk of introducing bias especially when data are not missing at random (Nakagawa & Freckleton, 2008). However, bias can be introduced by augmentation and imputation as well, if the dataset used to develop those methods is already biased due to the missing cases.

We use two approaches for dealing with missing data. First, we use a deletion approach for distributions or statistics of hazard impacts. The approach, called ‘available case analysis’, utilizes only the observed data points for each variable.

Because variables with few observations are less likely to be representative of the various possible underlying conditions in terms of hazard intensity, vulnerability and exposure than variables with many observations, we only conduct in-depth analyses for variables with at least 50 observations. Second, we use an imputation approach for total aggregate results that involve sums. Here, we assume missing values to be zeros. This is currently the standard approach in the literature though it inevitably leads to an underestimation of total impacts (Jones et al., 2023; Lee et al., 2024). After preprocessing to handle biases and missingness, we obtained a data subset containing 5868 disaster records of which 74% have one hazard, 22% have two hazards, and 4% have three hazards. This corresponds to a total of 7605 hazards ($5868 \times 74\% \times 1 + 5868 \times 22\% \times 2 + 5868 \times 4\% \times 3 = 7605$).

2.2 GDIS

GDIS is an open-source extension to EM-DAT and provides geographical approximations for main geophysical, meteorological, hydrological and climatological disaster types from 1960 – 2018 (Rosvold and Buhaug, 2021). It includes spatial geometries for floods, storms, earthquakes, volcanic activity, extreme temperatures, landslides, and droughts, however not for wildfires. Overall, GDIS provides impact zones for almost 90% of these types of records.

The spatial geometries in GDIS correspond to administrative areas, as contained in the Global Administrative Areas database (GADM, n.d.). The geometries are derived from EM-DAT's country field or optional "Location" field, which lists the name(s) of the affected administrative area(s), or "Latitude" and "Longitude" fields, which provide coordinates for the location. Most locations can be described on the spatial resolution of administrative level 1 (typical state/province/region). The highest resolution corresponds to level 3 (district/commune/village) and the lowest resolution corresponds to level 0 (country). However, as hazards are unlikely to affect the precise area of an administrative region, the spatial geometries have to be regarded as crude approximates of the impact zones.

3 Method

Our method has two main parts and is outlined in Figure 1. In the first part we re-classify the disaster records into single-hazard and multi-hazard events. In the second part we analyse the impacts.

The first part involves three steps. First, we preprocess the EM-DAT disaster records by geo-referencing them using GDIS (Section 3.1.1). Second, we identify spatiotemporally overlapping disaster record pairs (Section 3.1.2). Third, we derive the single-hazard and multi-hazard events (Section 3.1.3).

The second part also involves three steps. First, we assess the share of multi-hazards and their impacts in global disasters (Section 3.2.1). Second, we identify and statistically compare impacts of hazard pairs, single hazards and combinations of

two single hazards (Section 3.2.2). Third, we identify and conceptualize patterns of compounding impacts from multi-
180 hazards in four distinct archetypes (Section 3.2.3).




Figure 1 The two main parts and sub-steps of the methodology

3.1 Re-classifying disaster records into single-hazard and multi-hazard events

185 In this section, we describe the three steps to identify single-hazard and multi-hazard events based on the EM-DAT disaster records. The python code of the algorithm can be found on GitHub (link to be added upon publication).

3.1.1 Preprocessing and georeferencing disaster records

We preprocess and geo-reference disaster records using EM-DAT and GDIS data. Given the properties of, and guidelines for, the datasets, we only include disaster records that fall within the period 2000 – 2018 and belong to one of the seven disaster types listed in the second column of .

EM-DAT uses a hierarchical classification system with types and subtypes for the main disasters. The associated disasters do not follow the main classification system of EM-DAT but appear to correspond to either the disaster type or the disaster subtype. For consistency we map them to a disaster type. If the associated disasters cannot be mapped to one of the seven disaster types, we exclude the record from the analysis.

In the remainder of the paper we use the term hazard types instead of disaster type to be in line with terminology of the disaster risk field (PreventionWeb, 2023). We use nine different hazard types that capture different combinations of disaster type and subtype as well as the associated disasters. EM-DAT records that contain other hazard types are excluded for this analysis. We use the same terms for the hazard types as (Claassen et al., 2023); they are given in the first column of .

Finally, we link the GDIS geometries to the EM-DAT data set via the unique disaster event identifier that is present in both datasets.

Table 2 Hazard types used in this article versus terms used in EM-DAT

<i>Terms used in this article</i>	<i>Terms used in EM-DAT</i>		
Hazard types	<i>Disaster type</i>	<i>Disaster subtype</i>	<i>Associated disaster</i>
Earthquake (eq)	Earthquake	Ground movement	Earthquake
Tsunami (ts)		Tsunami	Tsunami/tidal wave,
Volcanic eruption (vo)	Volcanic activity	Ashfall, lahar, pyroclastic flow, lava flow	Volcanic activity
Landslide (ls)	Landslide	Landslide, rockfall, mudslide, avalanche (snow, debris, mudflow, rock)	Slide (land, mud, snow, rock), avalanche (snow, debris)
Coldwave (cw)	Extreme temperature	Cold wave	Cold wave
Heatwave (hw)		Heat wave	Heat wave
Extreme wind (ew)	Storm	Convective storm, tropical cyclone, extra-tropical storm	Storm
Flood (fl)	Flood	Coastal flood, riverine flood, flash flood	Flood
Drought (dr)	Drought	All (Drought)	Drought

3.1.2 Identifying spatiotemporally overlapping disaster record pairs

- 210 We identify spatiotemporal overlaps between pairs of disaster records using their spatial geometries and dates. We explain the algorithm with the illustration in Figure 2. This example has five disaster records A – E. Figure 2a shows the relevant information for the algorithm as obtained from EM-DAT. Each disaster record has a start date as well as a main hazard type and optionally one or two associated hazard types. End dates are often missing. The algorithm works as follows:
1. We create a list of all possible pairwise combinations of disaster records per country. We focus on pairs within a single
215 country, because this is how disasters are recorded in EM-DAT. Alternatively, we could have merged the records of different countries when they correspond to the same physical disaster event. This would give a better estimation of the number of events from a physical perspective. However, our main goal is to assess and compare differences in impact from single-hazards and multi-hazards focussing on spatiotemporal overlaps. We reason that considering the records of different countries individually enables us to better separate areas that are affected by multiple hazards from areas that
220 are affected by a single hazard. Suppose disaster records A – E are in one country, then all possible pairs would be “A, B”, “A, C”, “A, D”, “A, E”, “B, C”, “B, D”, “B, E” and “D, E”.
 2. We assess the spatial overlap for each of the pairs from step 1. We calculate the intersecting area between the records as well as the fractions of the intersecting areas compared to the areas of the individual events. We refer to the higher value of the two fractions as the intersection percentage and use a minimum value as criterion to define spatially overlapping
225 disaster records. We reason that the greater the intersecting area of two footprints, the more likely it is that the actual disaster impact zones overlap. We use a threshold of 50% and perform a sensitivity analysis (0%, 25%, 50%, 75%, 100%)¹ on this choice. Given the spatial geometries in Figure 2b, the spatially overlapping pairs would be “A, B”, “A, C” and “B, D”.
 3. We assess the temporal overlap for each of the pairs from step 1. We calculate the time difference between the start
230 dates of the pair. We use a maximum time lag as a criterion to define temporally overlapping disaster records. We use a time lag of 3 months and perform a sensitivity analysis on this choice (0 months, 1 month, 3 months, 6 months, 12 months)². Figure 2c depicts the time lags. Suppose all the times between disasters (Δt_{21} , Δt_{32} , Δt_{43} , Δt_{54}) are 1 month, the temporally overlapping pairs using a 3 month time-lag would be “A, B”, “A, C”, “A, D”, “B, C”, “B, D”, “B, E” and “D, E”.
 4. We identify all spatiotemporally overlapping disaster record pairs based on the previous assessments of spatial and
235 temporal overlap. In the example, these are “A, B”, “A, C” and “B, D”.

¹ The criterion is \geq for all spatial overlap values, except for the 0% value. In this case the criterion is $>$.

² The criterion is \leq for all time lag values.

3.1.3 Deriving events

We derive single-hazard events and multi-hazard events by identifying all disaster records that have potentially contributed to the reported impact through direct or indirect spatiotemporal overlaps. We use an iterative algorithm on the previously identified overlapping disaster record pairs. We explain the algorithm expanding the previous example in Figure 2d:

- For each disaster record, we find all pairs of spatiotemporally overlapping disaster records that include this disaster record. The other record in the pair is considered to be a contributing disaster record, if it is preceding in time. For example, if D is the disaster record of interest, then B is a contributing disaster record.
- If the contributing disaster record has in turn another contributing disaster record, we add that one as well, thus considering indirect contributions. Here, A is contributing disaster record to B. Hence, we add A as contributing disaster record to D as well. Adding indirectly contributing disaster records is a recursive process. For the example the recursive process stops here, because A has no further contributing disaster records.
- An event consists of the disaster record of interest and all contributing disaster records ordered in time (A, B, D) as well as the hazards included in those disaster records (A1, B1, D1, D2). Events which are fully included in another event are marked as duplicates and not used in part of the analysis. They have been events at the time their impact was recorded but evolved to include additional hazards later on. For example, “A1, B1” is fully included in “A1, B1, D1, D2”, which is marked by parenthesis in Figure 2e. Note that the same hazard can be part of multiple events. For example, hazard “A1” is part of event “A1, C1, C2, C3” as well as “A1, B1, D1, D2”

Figure 2e illustrates how the terms hazard, disaster record and event relate to each other.




Figure 2 Example with five disaster records A, B, C, D and E. a) Relevant fields obtained from EM-DAT; b) Spatial geometries of disaster records; c) Start dates of disaster records; d) Disaster records, their contributing disaster records and corresponding

260 events. Parentheses indicate events which are fully included in another event. e) Illustration of how the terms hazard, disaster record and event relate to each other.

3.2 Analysing impacts

In this section, we describe the three steps for analysing and comparing impacts of single-hazards and impacts of multi-hazards. The python code of the algorithm can be found on GitHub (link to be added upon publication).

3.2.1 Assessing the share of multi-hazards and their impacts

265 We follow a number of steps to assess the share of multi-hazards and their impacts in global disasters, as illustrated by the flowchart in the box of step 3.2.1 in Figure 1. Starting point are the derived single- and multi-hazards events with duplicates removed. First, we determine the total number of events. The ones that contain one hazards are single-hazard events. The ones that contain more than one hazard are multi-hazard events. Then, we determine the share of multi-hazard events in the total number of events. We also determine the share of hazards that occur in multi-hazard events. We determine this by
270 subtracting the share of single-hazards from 100% to avoid double counting due to hazards being part of more than one multi-hazard event. The share of single hazards is given by the number of single-hazard events divided by the total number of hazards in the disaster records and converted to a percentage. Similarly, we determine the share of impacts caused by multi-hazards: we subtract the share of impacts caused by single hazards from 100%, where the share of impacts caused by single hazards is the sum of impacts caused by single hazards divided by the total impacts of all disaster records and
275 converted to a percentage.

3.2.2 Identifying and comparing impacts of hazard pairs, single hazards and combinations of two single hazards

We create a dataset of the human and socioeconomic impacts of single hazards and of hazard pairs. We focus on damages, number of people affected, and number of deaths. We create the dataset by selecting all events (including the duplicates)
280 consisting of one or two hazards. These would be “A1”, “A1, B1” and “E1, E2” in the example of Figure 2d. If the event consists of one hazard, we record a single hazard impact. If an event consists of two hazards, we record a hazard pair impact. Note that the hazards of a hazard pair can belong to one disaster record or two disaster records. If they belong to two disaster records, we sum their impacts to obtain the total of both hazards.

285 We examine distributions and compare means to analyse impacts of hazard pairs, single hazards and combinations of two single hazards for different hazard and impact types. We have the distribution of impacts for hazard pairs and for single hazards from the previous step but not for the combinations of two single hazards. For the latter, we can only derive the mean by summing the mean impacts of the underlying single hazards. To compare the means, we construct confidence

intervals (CIs) with a percentile bootstrap (N=10,000). If the CIs do not overlap, we conclude that the difference in impacts
290 is statistically significant. Otherwise, we conclude that the difference is statistically not significant.

3.2.3 Identifying and conceptualizing patterns of compounding impacts

We identify different patterns of compounding impacts based on the detected differences and similarities in impacts of
hazard pairs, single hazards and combinations of two single hazards. We conceptualize these patterns and call them
archetypes, inspired by the field of system dynamics, which uses the term to describe common dynamics that recur in many
295 different settings (Senge, 1990).

4. Results

We first show the share of events that are multi-hazard events and their impacts (step 3.2.1). Then, we show the results of the
statistical comparison of impacts of hazard pairs, single hazards and combinations of two single hazards (step 3.2.2). Finally,
we describe distinct patterns of compounding impacts (step 3.2.3). Additional results on intermediate outputs of steps 3.1.1
300 and 3.1.2 can be found in sections A.1 and A.2 of the Appendix.

4.1 Share of multi-hazards events and their impacts

The datasets of identified (multi-)hazard events for different criteria can be found at Zenodo (link to be added upon
publication). We find that a higher share of events and higher share of hazards are multi-hazard than Lee et al. (2024).
According to their approach, there are 5868 events of which 26% are multi-hazard events and include 43% of the hazards.
305 The multi-hazard events caused 57% of the total damages, 40% of the total people affected and 49% of the total deaths
globally.

The higher shares that we find are due to spatiotemporal overlaps of the disaster records in EM-DAT. Figure 3a-f show the
results for different assumptions of spatiotemporal overlap. Figure 3a shows the total number of identified events, Figure 3b
310 shows the share of events that are multi-hazard events, Figure 3c shows the share of hazards that are associated with multi-
hazard events and Figures 3d-f show the shares of total damages, number of people affected and number of deaths caused by
the share of hazards associated with multi-hazard events.

The lower the criterion for minimum spatial overlap and the higher the criterion for maximum time lag, the lower the
315 number of events and the higher the share of multi-hazards and their impacts. For example, compared to Lee et al. (2024),
the number of events decreases to 4636 and the share of multi-hazard events increases to 35% when assuming a spatial
overlap of at least 50% and a time lag of at most 3 months. The identified multi-hazard events include 218 different hazard
combinations and consist of up to 32 individual hazards from 5 different hazard types rather than at most 3 different hazards

and hazard types in the disaster records. Furthermore, the share of hazards associated with multi-hazard events increases by almost 50% to 61%. Together these hazards caused 78% of the total damages, 83% of the total people affected and 69% of the total deaths globally.




Figure 3 (a) Number of events, (b) share of multi-hazard events, (c) share of hazards associated with multi-hazard events as well as their share in (d) total damages, (e) total people affected and (f) total deaths when accounting for spatiotemporal overlaps between the disaster records in EM-DAT.

4.2 Comparison of impacts of hazard pairs, single hazards and combinations of two single hazards

We present the results for a spatial overlap of at least 50% and a time lag of at most 90 days. Figure 4 shows the boxplots of the distributions of impacts of hazard pairs and single hazards as well as the mean values with 95% confidence interval (CI) for hazard pairs, single hazards and combinations of two single hazards. We only show impact types and hazard types with sample sizes $N \geq 50$ in an attempt to capture the broad range of underlying hazard intensity, exposure and vulnerability

conditions from which the impacts arise. There are eight cases in terms of combination of impact and hazard types that fulfil this criterion. For extreme winds and floods, sufficient data are available for damages, number of deaths, and number of people affected (first column of Figure 4). For floods and landslides, as well as consecutive floods, sufficient data are available for number of deaths and number of people affected (second and third column of Figure 4). For earthquake and landslides, sufficient data are available for people affected (fourth column of Figure 4). Sample sizes are reported in Table C1 and Table C2 in the Appendix.

For all variables (impacts of single hazards and of hazard pairs), the mean value is higher than the 75%-quantile; for 9 of the 16 variables the mean value is higher than the 95%-quantile. Thus, the majority of impacts are clustered towards the lower end of the impact range while a few very high data points pull the mean upwards. Consequently, the uncertainties about the mean are large compared to the spread of the distributions: For 10 of the 16 variables, the uncertainty about the mean is larger than the 75% inter-quantile range.

The first four columns of the matrix in Table 3 list the results of the comparison of the mean impacts of hazard pairs and single hazards. In three cases, the impacts of the hazard pair are significantly higher than those of both single hazards (damages for extreme winds and floods, number of deaths for consecutive floods, and number of people affected for consecutive floods). Also, in three cases, the impacts of the hazard pair are significantly higher than those of one but not the other of the single hazards (number of deaths for floods and landslides, number of people affected for floods and landslides, and number of people affected for earthquakes and landslides). Finally, in two cases, the average impacts of the hazard pair are not significantly different than those of either of the single hazards (number of deaths for extreme winds and floods and number of people affected for extreme winds and floods). In no case is the average impact of a hazard pair significantly lower than those of either or both single hazards.

The last four columns of the matrix in Table 3 list the results of the comparison of mean impacts of hazard pairs and combinations of two single hazards. In two cases, the impacts of the hazard pair are significantly higher than the combinations of two single hazards (number of deaths of consecutive floods and number of people affected of consecutive floods). In all other cases, no statistical difference is detected. In no case is the average impact of a hazard pair significantly lower than those of the combined impacts of the two single hazards.




Figure 4 Boxplots of impact data for single hazards as well as hazard pairs (,) for different impact types and hazard types as well as mean values and their bootstrap 95% CI for single hazards, hazard pairs (,) and the combined impact of two single hazards (+). The rows show different impact metrics. The columns show different hazard types (ew – extreme wind, fl – flood, ls – landslide, eq – earthquake). Only combinations of impact and hazard type with N > 50 are shown.

365

370 **Table 3 Statistically significant differences in mean impacts of hazard pairs, single hazards and two single hazard combined (ew – extreme wind, fl – flood, ls – landslide, eq – earthquake). Pairs are denoted by a ‘,’ and combinations of two single hazards are denoted by a ‘+’. A ‘>’ indicates that the impact of the variable in the row is higher than of the variable in the column and a ‘=’ denotes no difference.**

		ew	fl	eq	ls	ew+fl	fl+fl	fl+ls	eq+ls
Damages	ew,fl	>	>			=			
Number of deaths		=	=			=			
Number of people affected		=	=			=			
Damages	fl,fl								
Number of deaths			>				>		
Number of people affected			>				>		
Damages	fl,ls								
Number of deaths			>		=			=	
Number of people affected			=		>			=	
Damages	eq,ls								
Number of deaths									
Number of people affected				=	>				=

375 **4.3 Archetypes of compounding impacts**

We identify four distinct patterns of compounding impacts based on the differences and similarities in mean impacts of hazard pairs, single hazards and combinations of two single hazards (Table 4). In the first archetype, the impact of the hazard pair is higher than the impacts of both underlying single hazards as well as their combined impact. We see this pattern in number of deaths and number of people affected of flood-flood pairs. In the second archetype, the impact of the hazard pair is higher than the impacts of both underlying single hazards, but not different from their combined impact. We see this pattern in damages of extreme wind – flood pairs. In the third archetype, the impact of hazard pair is higher than the impacts of one of the underlying single hazards but neither different from the impact of the other or from their combined impact. We see this pattern in number of deaths and number of people affected of flood – landslide pairs as well as number of people affected in earthquake – landslide pairs. In the fourth archetype, the impact of a hazard pair is neither different from the impact of any of the underlying single hazards nor from their combined impact.

Table 4 Four archetypes describing distinct patterns of compounding impacts of hazard pairs. “haz1,haz2” denotes a hazard pair; “haz1” and “haz2” denote the underlying single hazards; and “haz1+haz2” denotes the underlying single hazards combined. A ‘>’ indicates that the impact of the variable in the row is higher than of the variable in the column and a ‘=’ denotes no difference.

		haz1	haz2	haz1+haz2
Archetype 1	haz1,haz2	>	>	>
“The whole is greater than the sum of its parts”				
Archetype 2		>	>	=
“The whole equals the sum of its parts”				
Archetype 3	haz1,haz2	>	=	=
“One part determines the whole”				
Archetype 4	haz1,haz2	=	=	=
“The whole and the parts are limited by total impact”				

5 Discussion

395 The aim of this study was to gain understanding of multi-hazards and their compounding impacts by analysing the emergency events database EM-DAT.

We found that the number of events and number of hazards that are considered multi-hazard are likely higher than recorded in EM-DAT and identified by Lee et al. (2024). This is due to spatiotemporal overlaps between the disaster records as multi-
400 hazard events may involve multiple disaster records. However, there remains substantial uncertainty in the identification of multi-hazard events in EM-DAT.

On one hand, the uncertainty is due to the limited spatiotemporal information in the data. The resolution of spatial footprints in GDIS varies from local administrative units to country level and varies in size introducing uncertainty regarding the actual
405 overlap of hazard-exposed areas (see Appendix A.2 for examples and a discussion). In addition, the individual hazards within a disaster record are associated with the same footprint, even though their footprints may have very different extents. Similarly, the temporal information in EM-DAT is crude. As end dates are partially missing, we used start dates and a time lag, which is a crude approximation of the actual time between hazards. The temporal information used here is also provided on disaster record level, but not on the level of individual hazards. Finally, we still lack understanding on how much time lag
410 and overlap should be considered, although some suggestions and sensitivity analysis is provided by Claassen et al. (2023). De Ruiter et al. (2020) suggest that hazards should be analysed together if direct impacts of a subsequent hazard spatially overlap before recovery from a previous hazard is considered to be completed, but information on recovery duration is limited.

415 On the other hand, the increasing trend in the reporting of associated disasters in EM-DAT (Lee et al., 2024) suggest that impacts may have been, and still are being, assigned to a single main hazard even though additional hazards have occurred leading to an underestimation of multi-hazards. We also encountered several similar cases during our analysis. An example is tropical cyclone Grace, which followed the 7.2-magnitude earthquake in Haiti in August 2021 (Daniels, 2021) but is not reported in EM-DAT. Despite these uncertainties in the data and method, multi-hazards are likely underreported in EM-DAT.

We also found that the reported multi-hazards contribute to a disproportionately high share of total impacts globally compared to single hazards, which is in agreements with the results of Lee et al. (2024). When statistically comparing impacts of hazard pairs, single hazards as well as combinations of two single hazards, the results differ per impact and hazard type, suggesting that there are different patterns of compounding impacts. Nonetheless, there are commonalities across all cases: The average impact of a hazard pair is as high or higher than the average impact of a single hazard and as high or higher than the combined average impacts of the two underlying single hazards, while the opposite was not found. This suggests that multi-hazard interactions leading to increased impact tend to outweigh multi-hazard interactions leading to decreased impact. Again, the results need to be treated with care because of large uncertainties in the impact data but also because of known biases and limitations of EM-DAT.

In cases where there is a significant difference, this could indeed point to an actual difference in impacts. However, it may also be caused by biases such as systematic double counting of impacts of consecutive disasters or geographical biases. For example, two consecutive earthquakes in Iceland have the exact same amount of reported total damages (reported under disaster numbers 00-0076-ISL and 2000-0335-ISL), suggesting potential double counting when adding them up. The fact that we came across these inconsistencies by chance, suggests that there are many more.

In cases where there is no significant difference found between two impact variables, this could either mean that there is indeed no difference between those variables or that there is a difference but not sufficient evidence in the dataset to detect it. Several factors contribute to the uncertainties. Firstly, a handful of extremely high data points are pulling the mean impact value up from the bulk of data points clustered at the lower end of the distribution. In addition, the sample sizes of impact data are small per hazard pair type, which is due to the many different event types as well as missing impact data in EM-DAT. While we considered 9 hazard types, we could only analyse impacts for four hazard types and four hazard pair types when requiring a sample size of at least $N=50$. Especially, impacts for extreme temperatures and droughts are missing. For these types, the complexity and difficulty to assess impacts is well known (Wilhite et al., 2007). Lastly, the hazards occurred under diverse conditions in terms of hazard intensity, exposure and vulnerability, which can cause a wide range of impacts. Ideally, these factors would be controlled for in the analysis, as done for example by Budimir et al. (2014). However, EM-DAT does not contain sufficient information to do so.

450 Finally, we could identify and conceptualize four distinct patterns of compounding impacts. While we acknowledge the substantial uncertainty in the association of each case to a particular archetype in our results, there are possible explanations and real world cases that support the existence of such patterns in general:

1. Archetype 1 (“The whole is greater than the sum of its parts”): A pattern where the impact of a hazard pair is significantly higher than the impact of two hazards combined would arise when both hazards in the pair exacerbate each other’s impacts if they co-occur simultaneously or consecutively. Real world examples of such a pattern are a previous flood that increases vulnerability leading to more impacts from a second flood (de Ruiter et al. 2020), a previous flood that intensifies a second flood due to already saturated soils and thus leading to higher impacts (Berghuijs et al., 2019), or a progressive increase of a building’s physical vulnerability due to multiple loads (Zuccaro et al., 2008).
2. Archetype 2 (“The whole equals the sum of its parts”): A pattern where the impact of a hazard pair is comparable to the combined impact of two single hazards would emerge when the hazards do not significantly affect each other’s impact. This could, for example, be the case for damages to the built environment caused by extreme wind and a flood, because they have different damage causing mechanisms. Floods tend to affect the interior of buildings and the lower floors, whereas extreme winds tend to damage the exterior of buildings and, in particular, the roof (Amini & Memari, 2020).
3. Archetype 3 (“One part determines the whole”): A third pattern would emerge when one hazard is so impactful that, in comparison, the contribution of other hazard is negligible, possibly combined with a “the whole and the parts are limited by total impact”-effect (Archetype 4). This could, for example, be the case for the number of people affected by flood – landslide pairs and earthquake – landslide pairs. Floods and earthquakes usually occur on larger spatial scales than landslides and trigger landslides within the already affected area so that the landslide will not add to the number of affected people, even though it may add to the severity in which they are affected.
4. Archetype 4 (“The whole and the parts are limited by total impact”): Another pattern could emerge when one hazard causes an ultimate impact to an exposed element, such as total loss for a building or death for a person, or when the impact metric only reports that an element has been affected but not to what degree. In both cases, a second hazard acting on the same elements cannot increase the value of the impact metric anymore. This could potentially be the case for the number of people affected by extreme wind and flood pairs when the same area is hit by both hazards.

6 Conclusion

By accounting for spatiotemporal overlaps between disaster records in EM-DAT, we found that almost 50% more hazards occurred in a multi-hazard context than previously reported, although substantial uncertainty remains in the identification of multi-hazard events. Despite the uncertainty, the identified multi-hazard events provide promising case studies for investigating impact-relevant spatiotemporal relations between hazards and their role in compounding impacts. Overall, multi-hazards have caused disproportionately high impacts in global disasters, but it appears that there are different patterns of compounding impacts which depend on the impact and hazard type. We conceptualized four distinct archetypes to capture the encountered patterns. In all archetypes, hazard pairs tend to have at least as much impact than single hazards or combinations of two single hazards but never less impact. This suggests that multi-hazard interactions leading to increased impact tend to outweigh multi-hazard interactions leading to decreased impact.

7 Recommendations

We propose the further investigation and development of archetypes to capture the different patterns. Such archetypes could help decide on the level of complexity to take into account in risk assessments and risk management for a region of interest if relevant hazard types and impact metrics are known. For some types of hazard and impact, modelling the impact of one dominant hazard may yield a reasonable approximation of multi-hazard impact, while in other cases modelling single-hazards impacts separately and adding them up may yield a reasonable approximation, while yet in other cases, it may be important to take into account interaction effects leading to either increased or decreased impacts compared to a simple sum of individual impacts. However, further research using more reliable data sources is needed to confirm these archetypes, to validate them for use in forward-looking risk assessments, to explore potentially additional forms of compounding impacts, and to expand them for additional hazard types and impact metrics.

Future research should also be directed at improving the reporting of multi-hazards and their impacts. In the long run, standardized reporting procedures should be established and implemented to ensure that multi-hazards and their impacts are documented sufficiently and consistently. This is essential for aligning research efforts and advancing the disaster risk field, as well as for translating findings into effective policy recommendations for risk assessment and reduction.

In the short-term, we recommend to improve and support the existing information in EM-DAT. To start with, a quality control of the impact data that solely focuses on the most disastrous records could already improve overall reliability because these records dominate any statistical data analysis that is based on mean values or total values. Data science techniques could be explored to automate the identification of errors and inconsistencies. Another key area to improve the usability of EM-DAT would be to develop high resolution data sets of (multi-)hazard intensities as well as exposure and vulnerability that can readily be linked to the disaster records. On one hand, this would enable a deeper analysis of multi-hazard

occurrence and impacts because factors determining the context in which the hazards occurred can be controlled for. On the other hand, this could help identify as yet unrecorded impactful events and reduce reporting bias. Finally, consistent and standardized methods to estimate impacts are needed to better enable comparison across countries and even events. Quality of impact data could also be improved by leveraging new impact data sets that are becoming available with novel methods. These data sets should also be made linkable to EM-DAT to enable cross-validation and increase the sample size, which in turn could enable an analysis of (multi-)hazard types that had to be excluded from the statistical analysis in this study.

Appendices

Appendix A An Exploratory Data Analysis of the Joint EM-DAT and GDIS Data Set

This section relates to method step 3.1.1. The data set of geo-referenced disaster events covers the period 2000 – 2018 and the nine hazard types listed in . It contains 5,868 disaster records. Table A1 shows the availability of data in the optional fields in these records. In case of associated disasters, we assume that empty fields means that no other hazards have taken place. In all other cases, we assume that data are missing. The temporal information is most complete. All events have a start year and start month as well as an end year. All events other than droughts also have an end month. The exact day is missing more frequently.

Table A1 Count of events and the data availability of key variables in the geo-refenced data set

	Event count	Start date	End date	Damages	Number of deaths	Number of people affected	Geospatial footprint
fl	2782	100%	92%	30%	72%	88%	91%
ew	1629	100%	97%	52%	74%	73%	77%
cw	198	100%	47%	8%	83%	29%	85%
dr	188	96%	4%	40%	4%	60%	80%
hw	118	100%	48%	13%	84%	31%	79%
ls	353	100%	98%	12%	96%	64%	91%
eq	480	100%	100%	42%	64%	97%	96%
vo	93	100%	90%	15%	13%	89%	82%
ts	27	100%	100%	70%	100%	85%	93%

The availability of impact data depends on the hazard type as well as impact type and ranges from 4% for number of deaths due to droughts to 100% for number of deaths due to tsunamis (Table A1). Human impact is available more than total damages. Availability also fluctuates across the years and continents. For total deaths it ranges from 64% in 2004 to 78% in 2007 and from 40% in Oceania to 79% in Asia. For total affected it ranges from 73% in 2010 and 2018 to 87% in 2017 and

from 55% in Europe to 90% in Africa. For damage is ranges from 22% in 2006 to 48% in 2013, and from 14% in Africa to 44% in Oceania.

535

The availability of spatial footprints also differs per hazard type, year and continent, but less so than the impact variables. It ranges from 77% for extreme wind to 96% of earthquakes, from 71% in 2018 to 93% in 2006 and from 77% in Europe to 89% in the Americas. Overall, we could associate 87% (5090/5668) of all events with a spatial footprint which is in line with the 89% reported by the developers of GDIS (Rosvold & Buhaug, 2021).

540

These results suggests that data for impact and geospatial footprint are not missing at random in our extracted data set which poses a risk of bias in the subsequent analysis. This is in line with the findings of Jones et al. (2022) who identified the year the disaster occurred, income-classification of the affected country and hazard types as significant predictors of missingness for human and economic impact variables in a formal statistical analysis of the entire EM-DAT data set.

545 **Appendix B Pairs of Spatiotemporally Overlapping Disaster Records**

This section relates to method step 3.1.2. Out of the 5,868 disaster records, 5,090 events have spatial footprints. These can be grouped into 12,951,505 unique combinations of two events. 107,406 pairs have spatial overlap.

Figure B1 shows a histogram of the intersection percentage. Notable is the high number of event pairs with 0% overlap and with 100% when rounded to 2 decimals. The high number of events 0% overlap is likely caused by rounding errors for events that impacted adjacent administrative areas and these event pairs are considered to not be overlapping. Figure B2a shows an example. The high number of events with 100% overlap is also likely due to the fact that the resolution is on administrative boundary level: As soon as events are within the same administrative district they fully overlap, whereas only large scale impact events that affected multiple administrative districts can partially overlap. Figure B2b-c show examples with different overlap percentages.

555




Figure B1 Histogram of the intersection percentage of the 107406 EM-DAT events with spatial overlap

560 There is uncertainty on whether or not the actual impact zones overlap for all pairs of intersecting events, because the spatial
footprints are an approximation on the level of administrative regions and the events are unlikely to have affected the entire
region (Rosvold & Buhaug, 2021). This uncertainty could potentially reduce by considering the combination of the scale of
the natural hazards (e.g., landslides are local events while heat waves and cold waves are regional or national level events),
the extent of the damage (e.g., higher damages and fatalities are likely to stem from larger impact zones), and the
565 administrative level of the footprint (e.g., a footprint consisting of multiple district level polygons which have been joint to a
greater area is more likely to represent the actual impact area than a footprint consisting of a single country-level polygon).




Figure B2 Example event pairs with spatial overlap. Individual event impact zones are plotted in blue and yellow. The overlapping impact zone is plotted in green.

For example, in Figure B2b, event 2014-0009-USA is a cold wave and event 2014-0317-USA is a convective storm associated with a cold wave. As these are larger scale weather phenomena the overlapping administrative zone is likely to reflect the actual overlapping impact zone. However, in Figure B2c, event 2006-0128-USA is storm associated with a flood and event 2008-0173-USA is a riverine flood, and, in Figure B2d, event 2006-0598-USA is a riverine flood associated with heavy rain and event 2006-0744-USA is a storm. In these two cases, additional data on impact extent, or by proxy hazard extent, would be required to confirm actual overlap.

Table B1 shows the number of pairs of overlapping events for different spatial and temporal criteria. As expected, the number of pairs of overlapping events is lowest when requiring a high intersection percentage and a low time lag as overlap criteria.

Table B1 Number of pairs of overlapping events using different spatial and temporal overlap criteria.

Time lag \ Intersection	>0%	>=50%	>=100%
0 days	31	21	17

1 month	1,339	758	480
3 months	3,575	1,917	1,164
6 months	5,865	3,023	1,798
12 months	11,630	6,059	3,631
Any	107,406	54,712	33,245

Appendix C Sample size of impact data for single hazards and hazard pairs

585 This section presents sample size for the data used in Sect. 4.2. Table C1 shows the sample sizes for single hazards for a spatial overlap of at least 50% and a time lag of maximum 91 days. Table C2 shows the sample sizes for hazard pairs for a spatial overlap of at least 50% and a time lag of maximum 91 days.

590 **Table C1 Sample sizes of impact data for single hazards for a minimum spatial overlap of 50% and a time lag of maximum 91 days (ew – extreme wind, fl – flood, ls – landslide, eq – earthquake, dr – drought, cw – cold wave, hw – heat wave, vo – volcanic activity, ts – tsunami)**

Hazard	Damages	Number of deaths	Number of people affected
fl	428	1102	1479
ew	354	527	524
eq	109	171	281
dr	52	4	91
ls	21	230	147
cw	12	146	50
vo	10	6	65
ts	5	6	4
hw	3	64	23

Table C2 Sample size of impact data for hazard pairs for a minimum spatial overlap of 50% and a maximum time lag 91 days (ew – extreme wind, fl – flood, ls – landslide, eq – earthquake, dr – drought, cw – cold wave, hw – heat wave, vo – volcanic activity, ts – tsunami)

Hazard 1	Hazard 2	Damages	Number of deaths	Number of people affected
fl	ls	178	417	439
ew	fl	133	205	220
eq	ls	28	44	56
ew	ew	16	26	23
fl	fl	15	67	87

ew	ls	15	28	31
fl	ew	12	23	24
ew	cw	11	22	11
ts	ts	10	15	13
hw	dr	7	9	2
ls	fl	6	23	22
eq	eq	5	8	15
dr	hw	5		3
eq	ts	4	9	11
ls	ls	3	12	9
eq	fl	3	5	5
fl	dr	2	1	5
fl	cw	2	5	4
dr	fl	1	1	6
fl	eq	1	4	5
vo	eq	1		4
fl	ts	1	2	3
eq	ew	1	2	2
eq	dr	1		2
ls	eq	1	1	1
dr	cw	1	1	1
vo	ts	1	1	1
dr	eq	1		1
ew	dr	1		1
vo	ls		1	3
hw	ew		5	2
fl	hw		3	2
eq	hw		1	2
hw	fl		4	1
cw	cw		3	1
ew	hw		3	1
vo	fl		1	1
hw	ls		1	1
ls	dr			1
cw	eq			1
eq	cw			1

dr	ew			1
vo	vo			1
cw	fl		1	
cw	hw		1	
ls	ts		1	
cw	ls		1	87
ls	cw		1	31

595

Code Availability

The code to develop multi-hazard event data sets as well as to perform the statistical analysis of impacts has been publicly released on GitHub. (Link to be added upon publication).

Data Availability

600 The multi-hazard event dataset compiled during the is study, is openly available on Zenodo. (Link to be added upon publication).

Author Contribution

MCdR and PJW conceived the study. All co-authors contributed to the development and design of the methodology. WSJ performed the analysis, interpreted the results and prepared the paper, with contributions from all co-authors.

605 **Competing interests**

Some authors are members of the editorial board of NHESS.

Acknowledgements

610 WSJ, TT, MCdR and PJW received support from the MYRIAD-EU project, which received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No [101003276](#). MCdR also received support from the Netherlands Organisation for Scientific Research (NOW) (VENI; grant no. [VI.Veni.222.169](#)). This work used the Dutch national e-infrastructure with the support of the SURF Cooperative using Grant No. EINF-4493.

References

- Amini, Mehrshad, and Ali M.Amini, M., & Memari, A. M. (2020). Review of Literature on Performance of Coastal Residential Buildings under Hurricane Conditions and Lessons Learned. *Journal of Performance of Constructed Facilities*, 34(6), 04020102. [https://doi.org/10.1061/\(ASCE\)CF.1943-5509.0001509](https://doi.org/10.1061/(ASCE)CF.1943-5509.0001509)
- Berghuijs, W. R., Harrigan, S., Molnar, P., Slater, L. J., & Kirchner, J. W. (2019). The Relative Importance of Different Flood-Generating Mechanisms Across Europe. *Water Resources Research*, 55(6), 4582–4593. <https://doi.org/10.1029/2019WR024841>
- Bevacqua, E., De Michele, C., Manning, C., Couasnon, A., Ribeiro, A. F. S., Ramos, A. M., Vignotto, E., Bastos, A., Blesić, S., Durante, F., Hillier, J., Oliveira, S. C., Pinto, J. G., Ragno, E., Rivoire, P., Saunders, K., van der Wiel, K., Wu, W., Zhang, T., & Zscheischler, J. (2021). Guidelines for Studying Diverse Types of Compound Weather and Climate Events. *Earth's Future*, 9(11), e2021EF002340. <https://doi.org/10.1029/2021EF002340>
- Brimicombe, C., Di Napoli, C., Cornforth, R., Pappenberger, F., Petty, C., & Cloke, H. L. (2021). Borderless Heat Hazards With Bordered Impacts. *Earth's Future*, 9(9), e2021EF002064. <https://doi.org/10.1029/2021EF002064>
- Budimir, M. E. A., Atkinson, P. M., & Lewis, H. G. (2014). Earthquake-and-landslide events are associated with more fatalities than earthquakes alone. *Natural Hazards*, 72(2), 895–914. <https://doi.org/10.1007/s11069-014-1044-4>
- Claassen, J. N., Ward, P. J., Daniell, J., Koks, E. E., Tiggeloven, T., & de Ruiter, M. C. (2023). A new method to compile global multi-hazard event sets. *Scientific Reports*, 13(1), 13808. <https://doi.org/10.1038/s41598-023-40400-5>
- Daniels, J. P. (2021, August 17). Tropical Storm Grace's heavy rains pour misery on Haiti earthquake survivors. *The Guardian*. <https://www.theguardian.com/global-development/2021/aug/17/haiti-tropical-storm-grace-earthquake-rain>
- De Angeli, S., Malamud, B. D., Rossi, L., Taylor, F. E., Trasforini, E., & Rudari, R. (2022). A multi-hazard framework for spatial-temporal impact analysis. *International Journal of Disaster Risk Reduction*, 73, 102829. <https://doi.org/10.1016/j.ijdr.2022.102829>

- 635 de Ruiter, M. C., Couasnon, A., van den Homberg, M. J. C., Daniell, J. E., Gill, J. C., & Ward, P. J. (2020). Why We Can No Longer Ignore Consecutive Disasters. *Earth's Future*, 8(3), e2019EF001425. <https://doi.org/10.1029/2019EF001425>
- de Ruiter, M. C., & van Loon, A. F. (2022). The challenges of dynamic vulnerability and how to assess it. *iScience*, 25(8). <https://doi.org/10.1016/j.isci.2022.104720>
- 640 Delforge, D., Wathelet, V., Below, R., Sofia, C. L., Tonnelier, M., Loenhout, J. van, & Speybroeck, N. (2023). *EM-DAT: The Emergency Events Database*. <https://doi.org/10.21203/rs.3.rs-3807553/v1>
- Eilander, D., Couasnon, A., Leijnse, T., Ikeuchi, H., Yamazaki, D., Muis, S., Dullaart, J., Haag, A., Winsemius, H. C., & Ward, P. J. (2023). A globally applicable framework for compound flood hazard modeling. *Natural Hazards and Earth System Sciences*, 23(2), 823–846. <https://doi.org/10.5194/nhess-23-823-2023>
- 645 GADM. (n.d.). Retrieved July 5, 2024, from <https://gadm.org/data.html>
- Gall, M., Borden, K. A., & Cutter, S. L. (2009). *When Do Losses Count?* <https://doi.org/10.1175/2008BAMS2721.1>
- Gill, J. C., & Malamud, B. D. (2014). Reviewing and visualizing the interactions of natural hazards. *Reviews of Geophysics*, 52(4), 680–722. <https://doi.org/10.1002/2013RG000445>
- Gill, J. C., & Malamud, B. D. (2016). Hazard interactions and interaction networks (cascades) within multi-hazard methodologies. *Earth System Dynamics*, 7(3), 659–679. <https://doi.org/10.5194/esd-7-659-2016>
- 650 Guha-Sapir, D., & Below, R. (2002). The quality and accuracy of disaster data: A comparative analyse of 3 global data sets. *CRED Work. Pap*, 1–18.
- Harrington, L. J., & Otto, F. E. L. (2020). Reconciling theory with the reality of African heatwaves. *Nature Climate Change*, 10(9), 796–798. <https://doi.org/10.1038/s41558-020-0851-8>
- 655 Hillier, J. K., Matthews, T., Wilby, R. L., & Murphy, C. (2020). Multi-hazard dependencies can increase or decrease risk. *Nature Climate Change*, 10(7), 595–598. <https://doi.org/10.1038/s41558-020-0832-y>
- Hochrainer-Stigler, S., Colon, C., Boza, G., Poledna, S., Rovenskaya, E., & Dieckmann, U. (2020). Enhancing resilience of systems to individual and systemic risk: Steps toward an integrative framework. *International Journal of Disaster Risk Reduction*, 51, 101868. <https://doi.org/10.1016/j.ijdr.2020.101868>

- 660 Hurk, B. J. J. M. van den, White, C. J., Ramos, A. M., Ward, P. J., Martius, O., Olbert, I., Roscoe, K., Goulart, H. M. D., &
Zscheischler, J. (2023). Consideration of compound drivers and impacts in the disaster risk reduction cycle.
iScience, 26(3). <https://doi.org/10.1016/j.isci.2023.106030>
- IPCC. (2023). *Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment
Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]*.
665 <https://dx.doi.org/10.59327/IPCC/AR6-9789291691647>
- Jones, R. L., Guha-Sapir, D., & Tubeuf, S. (2022). Human and economic impacts of natural disasters: Can we trust the
global data? *Scientific Data*, 9(1), Article 1. <https://doi.org/10.1038/s41597-022-01667-x>
- Jones, R. L., Kharb, A., & Tubeuf, S. (2023). The untold story of missing data in disaster research: A systematic review of
the empirical literature utilising the Emergency Events Database (EM-DAT). *Environmental Research Letters*,
670 18(10), 103006. <https://doi.org/10.1088/1748-9326/acfd42>
- Kappes, M. S., Keiler, M., von Elverfeldt, K., & Glade, T. (2012). Challenges of analyzing multi-hazard risk: A review.
Natural Hazards, 64(2), 1925–1958. <https://doi.org/10.1007/s11069-012-0294-2>
- Kornhuber, K., Coumou, D., Vogel, E., Lesk, C., Donges, J. F., Lehmann, J., & Horton, R. M. (2020). Amplified Rossby
waves enhance risk of concurrent heatwaves in major breadbasket regions. *Nature Climate Change*, 10(1), 48–53.
675 <https://doi.org/10.1038/s41558-019-0637-z>
- Lee, R., White, C. J., Adnan, M. S. G., Douglas, J., Mahecha, M. D., O’Loughlin, F. E., Patelli, E., Ramos, A. M., Roberts,
M. J., Martius, O., Tubaldi, E., van den Hurk, B., Ward, P. J., & Zscheischler, J. (2024). Reclassifying historical
disasters: From single to multi-hazards. *Science of The Total Environment*, 912, 169120.
<https://doi.org/10.1016/j.scitotenv.2023.169120>
- 680 Leonard, M., Westra, S., Phatak, A., Lambert, M., van den Hurk, B., McInnes, K., Risbey, J., Schuster, S., Jakob, D., &
Stafford-Smith, M. (2014). A compound event framework for understanding extreme impacts. *WIREs Climate
Change*, 5(1), 113–128. <https://doi.org/10.1002/wcc.252>
- Li, Y., Ahuja, A., & Padgett, J. E. (2012). Review of Methods to Assess, Design for, and Mitigate Multiple Hazards. *Journal
of Performance of Constructed Facilities*, 26(1), 104–117. [https://doi.org/10.1061/\(ASCE\)CF.1943-5509.0000279](https://doi.org/10.1061/(ASCE)CF.1943-5509.0000279)

- 685 Liu, B., Siu, Y. L., & Mitchell, G. (2016). Hazard interaction analysis for multi-hazard risk assessment: A systematic classification based on hazard-forming environment. *Natural Hazards and Earth System Sciences*, 16(2), 629–642. <https://doi.org/10.5194/nhess-16-629-2016>
- Moriyama, K., Sasaki, D., & Ono, Y. (2018). Comparison of Global Databases for Disaster Loss and Damage Data. *Journal of Disaster Research*, 13(6), 1007–1014. <https://doi.org/10.20965/jdr.2018.p1007>
- 690 Nakagawa, S., & Freckleton, R. P. (2008). Missing inaction: The dangers of ignoring missing data. *Trends in Ecology & Evolution*, 23(11), 592–596. <https://doi.org/10.1016/j.tree.2008.06.014>
- NOAA. (n.d.). *Natural Hazards Data*. U.S. Department of Commerce. Retrieved January 27, 2025, from <https://www.ngdc.noaa.gov/hazard/>
- Panwar, V., & Sen, S. (2020). Disaster Damage Records of EM-DAT and DesInventar: A Systematic Comparison. *Economics of Disasters and Climate Change*, 4(2), 295–317. <https://doi.org/10.1007/s41885-019-00052-0>
- 695 Paprotny, D., Morales-Nápoles, O., & Jonkman, S. N. (2018). HANZE: A pan-European database of exposure to natural hazards and damaging historical floods since 1870. *Earth System Science Data*, 10(1), 565–581. <https://doi.org/10.5194/essd-10-565-2018>
- PreventionWeb. (2023, September 18). *Hazard Information Profiles (HIPs)*. [https://www.preventionweb.net/drr-](https://www.preventionweb.net/drr-glossary/hips)
- 700 [glossary/hips](https://www.preventionweb.net/drr-glossary/hips)
- Rosvold, E. L., & Buhaug, H. (2021). GDIS, a global dataset of geocoded disaster locations. *Scientific Data*, 8(1), Article 1. <https://doi.org/10.1038/s41597-021-00846-6>
- Senge, P. (1990). *The Fifth Discipline*. [https://www.managementboek.nl/boek/9781905211203/the-fifth-discipline-peter-](https://www.managementboek.nl/boek/9781905211203/the-fifth-discipline-peter-senge)
- senge
- 705 Simpson, N. P., Mach, K. J., Constable, A., Hess, J., Hogarth, R., Howden, M., Lawrence, J., Lempert, R. J., Muccione, V., Mackey, B., New, M. G., O'Neill, B., Otto, F., Pörtner, H.-O., Reisinger, A., Roberts, D., Schmidt, D. N., Seneviratne, S., Strongin, S., ... Trisos, C. H. (2021). A framework for complex climate change risk assessment. *One Earth*, 4(4), 489–501. <https://doi.org/10.1016/j.oneear.2021.03.005>

- Terzi, S., Torresan, S., Schneiderbauer, S., Critto, A., Zebisch, M., & Marcomini, A. (2019). Multi-risk assessment in
710 mountain regions: A review of modelling approaches for climate change adaptation. *Journal of Environmental
Management*, 232, 759–771. <https://doi.org/10.1016/j.jenvman.2018.11.100>
- Tierolf, L., Haer, T., Botzen, W. J. W., de Bruijn, J. A., Ton, M. J., Reimann, L., & Aerts, J. C. J. H. (2023). A coupled
agent-based model for France for simulating adaptation and migration decisions under future coastal flood risk.
Scientific Reports, 13(1), 4176. <https://doi.org/10.1038/s41598-023-31351-y>
- 715 Tilloy, A., Malamud, B. D., Winter, H., & Joly-Laugel, A. (2019). A review of quantification methodologies for multi-
hazard interrelationships. *Earth-Science Reviews*, 196, 102881. <https://doi.org/10.1016/j.earscirev.2019.102881>
- UNDRR. (n.d.). *United Nations DesInventar Open Source Initiative—Official Website*. Retrieved January 27, 2025, from
<https://www.desinventar.net/index.html>
- UNDRR. (2017). *Sendai Framework Terminology on Disaster Risk Reduction*. [http://www.undrr.org/drr-](http://www.undrr.org/drr-glossary/terminology)
720 [glossary/terminology](http://www.undrr.org/drr-glossary/terminology)
- van Westen, C. J., & Greiving, S. (2017). Multi-hazard risk assessment and decision making. *Environmental Hazards
Methodologies for Risk Assessment and Management*, 31.
- Ward, P. J., Daniell, J., Duncan, M., Dunne, A., Hananel, C., Hochrainer-Stigler, S., Tijssen, A., Torresan, S., Ciurean, R.,
Gill, J. C., Sillmann, J., Couasnon, A., Koks, E., Padrón-Fumero, N., Tatman, S., Tronstad Lund, M., Adesiyun, A.,
725 Aerts, J. C. J. H., Alabaster, A., ... de Ruiter, M. C. (2022). Invited perspectives: A research agenda towards
disaster risk management pathways in multi-(hazard-)risk assessment. *Natural Hazards and Earth System Sciences*,
22(4), 1487–1497. <https://doi.org/10.5194/nhess-22-1487-2022>
- Wilhite, D. A., Svoboda, M. D., & Hayes, M. J. (2007). Understanding the complex impacts of drought: A key to enhancing
drought mitigation and preparedness. *Water Resources Management*, 21(5), 763–774.
730 <https://doi.org/10.1007/s11269-006-9076-5>
- Zschau, J. (2017). *Where are we with multihazards, multirisks assessment capacities?*
https://drmkc.jrc.ec.europa.eu/portals/0/Knowledge/ScienceforDRM/ch02/ch02_subch0205.pdf

- Zscheischler, J., Martius, O., Westra, S., Bevacqua, E., Raymond, C., Horton, R. M., van den Hurk, B., AghaKouchak, A., Jézéquel, A., Mahecha, M. D., Maraun, D., Ramos, A. M., Ridder, N. N., Thiery, W., & Vignotto, E. (2020). A
735 typology of compound weather and climate events. *Nature Reviews Earth & Environment*, 1(7), 333–347.
<https://doi.org/10.1038/s43017-020-0060-z>
- Zscheischler, J., & Seneviratne, S. I. (2017). Dependence of drivers affects risks associated with compound events. *Science
Advances*, 3(6), e1700263. <https://doi.org/10.1126/sciadv.1700263>
- Zscheischler, J., Westra, S., van den Hurk, B. J. J. M., Seneviratne, S. I., Ward, P. J., Pitman, A., AghaKouchak, A., Bresch,
740 D. N., Leonard, M., Wahl, T., & Zhang, X. (2018). Future climate risk from compound events. *Nature Climate
Change*, 8(6), 469–477. <https://doi.org/10.1038/s41558-018-0156-3>
- Zuccaro, G., Cacace, F., Spence, R. J. S., & Baxter, P. J. (2008). Impact of explosive eruption scenarios at Vesuvius. *Journal
of Volcanology and Geothermal Research*, 178(3), 416–453. <https://doi.org/10.1016/j.jvolgeores.2008.01.005>

745