

# Brief Communication: Drivers of the recent warming of the Mediterranean Sea, and its implications for hail risk

Stephen Cusack<sup>1</sup>, Tyler Cox<sup>2</sup>

<sup>1</sup>Stormwise Ltd, Luton, LU4 9DU, United Kingdom

<sup>2</sup>Inigo Limited, London, EC3A 5AY, United Kingdom

Correspondence to: Stephen Cusack ([stephen.cusack@stormwise.co.uk](mailto:stephen.cusack@stormwise.co.uk))

## Abstract

The Mediterranean Sea has warmed by about 2 K since the early 1980s, and ~~past research has established how~~ this heating intensifies ~~damage from~~ moisture-driven perils such as hailstorms and floods in continental Europe ~~via the basic thermodynamic effect~~. This study uses the DAMIP (Detection and Attribution Model Intercomparison Project) set of climate model experiments to ~~identify~~explore the drivers of the recent Mediterranean warming. The models simulate the observed multidecadal variations of Mediterranean Sea temperatures in the modern ~~industrial~~ period accurately, and indicate anthropogenic aerosol forcing was largely responsible for the cooler period from about 1900 to the late 1970s, while rising greenhouse gases are the main cause of warming waters since then. Next, we reviewed ~~the quantitative impact of this heating on hailstorm risk~~trends in damaging hail, and found ~~hail damages in higher risk countries~~they have been increasing by around 2% per year ~~in key parts of mainland Europe~~ since 1980. The rising risk fits with the established mechanism whereby warmer waters moisten low-level air ~~and intensify thunderstorms leading to more severe thunderstorm hazards, though the exact relation between a warming Mediterranean and hail loss growth remains uncertain~~. Anthropogenic forcings will continue warming the Mediterranean for the next couple of decades at least, ~~indicating~~suggesting further increases to hail damages in ~~central and southern~~high-risk parts of Europe.

## 1 Introduction

Climate impacts from anthropogenic forcings have regional variations, caused by factors such as ~~local~~ dynamical feedbacks and spatial inhomogeneities of some drivers such as aerosols (e.g. Seneviratne et al., 2021). The Mediterranean region has been a notable climate hotspot over the past few decades (e.g. MedECC, 2020; Ali et al., 2022). For example, sea surface temperatures warmed by 0.41 K/decade over the 1982-2023 period according to the Copernicus Marine Service (Roquet et al., 2016; Mulet et al., 2018) which is almost double the rate of the global oceans. Further, its warming is projected to continue outpacing the global mean in the future (e.g. Lionello and Scarascia, 2018).

30 One of the main effects of a warming sea is the humidification of the lower levels of the atmosphere. The warming sea surface is accompanied by similar increases in temperature of the overlying air, increasing its water-holding capacity, and as a result more water is evaporated into the atmosphere. In general, greater amounts of low-level water vapour in the atmosphere act to raise the severity of weather perils such as heavy precipitation events (reviewed in Seneviratne et al., 2021) and large hail (e.g. Raupach et al., 2021; Chen and Dai, 2023). More specifically to central and southern parts of mainland Europe, moisture  
35 ~~amounts~~ from the Mediterranean Sea ~~have been~~was a key ingredient in many of the most ~~damaging~~extreme floods (e.g. James et al., 2004; Volosciuk et al., 2016; Krug et al., 2022; Tradowsky et al., 2023) and hailstorms ~~((~~(e.g. Heimann and Kurz, 1985; Kunz et al., 2018; Piper et al., 2019; Kunz et al., 2020; Kopp et al., 2023) in recent decades.

Given how research has established low-level Mediterranean air masses as an important component of past major weather disasters, and that these air masses are moistening due to sea surface warming~~heating~~, it would be prudent~~useful~~ to learn more  
40 about the Mediterranean Sea warming trend, and its consequences on severe weather risk. The twin aims of this study are to gain insights into the drivers of the warming Mediterranean in recent decades, and to measure the trends in damaging hail over the same period. Section 2 contains a description of the data and processing used in this study, then in section 3 we analyse model simulations and identify the key drivers of multidecadal variations in Mediterranean temperatures over the industrial period. Section 4 begins with a review of how Mediterranean Sea temperatures affect the occurrence of damaging hail over  
45 the higher-risk parts of Europe, then we present estimates of trends in damaging hail over recent decades from previous studies, and a new analysis of insurance loss data. Conclusions are presented in Section 5.

## 2 Data and methods

### 2.1 Mediterranean temperatures

~~The main aim of this study is to use modelling results from~~This study uses data from climate model experiments performed in  
50 the DAMIP (Detection and Attribution Model Intercomparison Project; Gillett et al., 2016) sub-project of ~~CMIP6~~ for information on how sea surface temperatures in the Mediterranean respond to different forcings of the climate system. Section 2 contains a description of the data and processing used in this study, followed by an analysis in Section 3 which identifies the key drivers of Mediterranean heat changes over the industrial period based on DAMIP results. A secondary aim was to quantify the link from Mediterranean Sea warming to its impacts on hailstorm risk in Europe, using hazard and insurance loss  
55 information from recent decades. This investigation is presented in Section 4. A summary of the main findings is presented in Section 5.

2 Data and methods

2.1 Mediterranean temperatures

This study uses data from the DAMIP sub-project, which is one of 23 sub-projects that formed part of the sixth version of the Coupled Model Intercomparison Project (CMIP6; Eyring et al., 2016). DAMIP was designed to investigate the impacts of various external forcings on global and regional climate. Its experiments explore the modern industrial period from 1850 to 2014, and its Tier 1 model simulations consist of setting one type of forcing to suitable values for the historical period, with all others fixed at pre-industrial values. Initial conditions are taken from pre-industrial control runs (from the main CMIP6 model experiments), and Tier 1 historical forcings are split into three distinct types: natural forcings (solar and volcanic, hereafter Nat), greenhouse gases (GHG), and anthropogenic aerosols (Aero). We analyse results from six different modelling centres providing monthly-mean near-surface temperature diagnostics (variable ‘tas’) for at least five different ensemble members, for all three forcing tests. In addition, the Historical experiments with all forcings (Hist) performed as part of the central CMIP project are also analysed. Table 1 summarises the simulations analysed in this study. More information on the climate models is provided in CMIP (2025).

Table 1: summary details of the DAMIP climate model simulations.

Model	Reference ID in-text	No. of ensemble members	Release year	Atmosphere resolution (km)	Ocean resolution (km)
CNRM CM6-1	1	40	2017	250	100
CanESM5	2	45	2010	500	100
GISS E2-1-G	3	5	2010	250	100
HadGEM3-GC31-LL	4	45	2016	250	100
MIROC6	5	40	2017	250	100
MPI ESM1-2 LR	6	45	2017	250	250

Observed sea surface temperatures (SSTs) are taken from the Hadley Centre Sea Ice and Sea Surface Temperature data set (HadISST; Rayner et al., 2003), a global dataset of monthly sea surface temperature values at 1° x 1° resolution from January 1870 and continually updated to the present day. These SST data are used to assess the validity of near-surface air temperature variations in DAMIP simulations. Anomalies in both quantities are very similar at the large spatial and temporal scales analysed in this study. For example, Rubino et al. (2020) described their close correspondence at interdecadal timescales (their Figure 4), and the two quantities are often used interchangeably in long climate reconstructions (e.g. Morice et al., 2012).

85 **Table 1: summary details of the DAMIP climate model simulations.**

<u>Model</u>	<u>Reference ID in text</u>	<u>No. of ensemble members</u>	<u>Release year</u>	<u>Atmosphere resolution (km)</u>	<u>Ocean resolution (km)</u>
<u>CNRM-CM6-1</u>	<u>1</u>	<u>10</u>	<u>2017</u>	<u>250</u>	<u>100</u>
<u>CanESM5</u>	<u>2</u>	<u>15</u>	<u>2019</u>	<u>500</u>	<u>100</u>
<u>GISS-E2-1-G</u>	<u>3</u>	<u>5</u>	<u>2019</u>	<u>250</u>	<u>100</u>
<u>HadGEM3-GC31-LI</u>	<u>4</u>	<u>15</u>	<u>2016</u>	<u>250</u>	<u>100</u>
<u>MIROC6</u>	<u>5</u>	<u>10</u>	<u>2017</u>	<u>250</u>	<u>100</u>
<u>MPI-ESM1-2-LR</u>	<u>6</u>	<u>15</u>	<u>2017</u>	<u>250</u>	<u>250</u>

Both observed and modelled data are processed similarly. Monthly mean values are initially area-averaged over a region of the Mediterranean Sea shown in Figure 1, then combined to form an extended summer half-year average (May-October) corresponding to the annual peak of Mediterranean influence on flood and hail risk, in order to focus on the most relevant seasonal warming trends (Garcia-Monteiro et al., 2022). Anomalies were defined using the climate from the common baseline period of 1870-2014 for observed and modelled values. The climate of the Hist simulations in this common 1870-2014 period has been used to define all model anomalies, for consistency with how observed anomalies are defined. Finally, a second-order low-pass Butterworth filter (Butterworth, 1930) with a 5-year cutoff was applied to all timeseries to reduce large amounts of interannual noise obscuring longer-term trends. The 5-year cutoff was chosen to retain potential signals from Nat forcings such as 11-year solar cycles and major volcanoes, as well as the slower changes from GHG and Aero.




Figure 1: map of a part of Europe, with shaded area denoting the Mediterranean Sea region used in the analysis.

2.2 European hail losses


National insurance associations often monitor losses to their market, and some of their records extend to the past few decades. Three publicly available historical records of hail losses are explored in Section 4 for guidance on trends in the risk from this weather peril. These data are now described.

- 105
- The German Insurance Association (GDV) issue annual losses indexed to 2023 inventory and prices using data on numbers of insured vehicles, and the cost of automobile parts and repair from the Federal Statistical Office of Germany (GDV, 2024). The second dataset comprises annual hail losses to all insured buildings in France, based on information from France Assureurs (2023). Their annual data consists of the ratio of all insured risks making a hail claim (a frequency ratio) and the average hail claim size indexed to 2022 values using the standard FFB Index, which is based on costs for various elements of a standard
- 110
- apartment in Paris and specifically designed to index insurance policies (FFB, 2025). We have computed total industry losses by multiplying these two quantities together, then scaling them by the total number of industry risks (N). The total industry loss in 2022 was used to define N, then we apply this value of N to all other years to obtain annual industry losses for all years based on the number of risks in 2022. This approach ensures indexation captures both the growth in claim severity, and how claim numbers increase with changes in number of insured risks.
- 115
- The third national loss dataset contains annual hail loss costs in Switzerland for the 1980 to 2023 period for all buildings in the 19 Swiss cantons covered by public insurance, from VKG (2022, 2024). The loss cost is defined as the ratio of the total cost of repair to the total exposure value. It is a useful loss metric in insurance, because total exposure value in the denominator is usually defined to reflect the same growths in insurance claim size and number which are present in the numerator, hence socio-economic trends are absent from the ratio: loss costs are indexed (aka normalised) by design.
- 120
- Trends in losses over the past few decades are derived from a regression of log(loss quantity) with year, providing best-fit estimates of the growth in units of % per year.

3 Results

3.1 Past changes in Mediterranean SST

Figure 2 displays timeseries of SST anomalies in both the global (60°S to 60°N) and Mediterranean regions for summer half-  
125 years in 1940 to 2024. The Mediterranean region cooled from roughly 1950 to 1980, then warmed rapidly since then. Changes in the global mean have been more muted, with anomalies near zero from 1940 to the mid-1970s, and a more gradual warming since then. Figure 2 also contains the best-fitting linear trend for SST in these two areas from 1980 to 2024, and it was found that average SSTs over each area have been warming at rates which are significantly different from zero at the 1% level (p-values are below 1e-10 in both cases). However, the most notable feature is the summertime Mediterranean SSTs trending  
130 upward at more than three times the global-mean warming rate since 1980. The Mediterranean Sea has been an oceanic hotspot over the past four decades.





135 **Figure 2: timeseries of global (black) and Mediterranean (red) SST anomalies in the summer half-year (May to October), with**  
 140 **associated best-fitting linear trends in the 1980-2024 period. Data are from HadISST, anomalies are computed with respect to the**  
 145 **1870 to 2014 period, and global data are from 60°S to 60°N.**

### 3.2 Drivers of past changes in DAMIP experiments

140 The causes of the more rapid warming of the Mediterranean Sea are now explored using DAMIP modelling results. Figure 3a  
 shows the timeseries of Mediterranean temperature anomalies over the extended historical period for observed and multi-  
 model ensemble means for the Historical and three DAMIP single-forcing experiments. Earlier multidecadal variations in  
observations, consisting of minima around 1910 and the late 1970s, and a local maximum from about 1930 to the early 1960s,  
are replicated in Hist, albeit with reduced amplitude. To our knowledge, there are no published analyses on the relative roles  
 145 of internal climate variability and external forcing toward the observed multidecadal variations in the Mediterranean basin  
prior to 1980. However, these earlier fluctuations are similar to those in the North Atlantic sector analysed by Booth et al.  
(2012), which they found to be caused by both anthropogenic and volcanic aerosol forcing, in addition to internal climate  
variability. The observed warming since about 1980 is of greater concern since it has four times larger amplitude than the mid-  
20<sup>th</sup> century peak. Notably, the multimodel ensemble mean simulates this recent rapid warming accurately. This comparison  
 150 to observed variations bolsters confidence in the fidelity of CMIP6-DAMIP climate model simulations of surface temperature  
anomalies in the Mediterranean, particularly its dominant feature of rapid warming in recent decades. We now use the validated  
model ensemble to assess the contribution of individual forcings to the total signal.  
 The multi-model ensemble mean signal from all forcings (Hist — solid red line) is similar to observed values (solid black line)  
 over the whole period. Specifically, the modelled rate of Mediterranean warming since the 1970s closely follows HadISST  
 155 data, and the earlier multidecadal oscillation in observations, consisting of minima around 1910 and the late 1970s, and a local

maximum from about 1930 to the early 1960s, is replicated in Hist, albeit with reduced amplitude. Control simulations with forcings fixed at preindustrial levels (part of the main CMIP6 protocol) have an interannual standard deviation ( $\sigma$ ) of 0.38 K, hence a 70 member mean is expected to have  $\sigma=0.046$  K (from the Central Limit Theorem), therefore the simulated multidecadal oscillation in Hist with amplitude of 0.5 K is very unlikely to be caused solely by internal climate variability. Research in nearby regions also find a role for external forcing in the multidecadal cycle over the first seven decades of the 20<sup>th</sup> century, such as Booth et al. (2012) in the broader North Atlantic, and Aizawa et al. (2022) in the Arctic. Having established the validity of CMIP6 climate models to simulate temperature anomalies in the Mediterranean, we now discuss the contribution of individual forcings to the total signal.





**Figure 3: Timeseries of Mediterranean temperature anomalies. (a) observed (solid black line) and multi-model mean anomalies for the various forcing experiments, with major volcanic eruptions indicated by the long red tick marks on the time axis. (b) DAMIP multi-model mean (solid lines), and the mean of each of the six models (various symbols), for the Aero (blue) and GHG (gold) experiments.**

The impact of Nat forcing (green dotted line in Figure 3a) on Mediterranean temperatures is simulated to be near-zero over the long-term. However, volcanic eruptions exert influence on temperature anomalies at timescales shorter than about 10 years in DAMIP simulations. Six of the largest climate-changing eruptions in the simulation period are indicated as long red tick-marks on the x-axis (Krakatoa 1883; Santa Maria 1902; Novarupta 1912; Mount Agung 1963; El Chichon 1982; Pinatubo 1991), and are associated with anomalies of up to -0.5 K for a few years following the eruption. Such a size of perturbation is smaller than the recent GHG and Aero forcings, and of much shorter duration, hence DAMIP model results indicate Nat forcing is a minor consideration in the modern industrial period.

GHG (gold dashed line) acts to warm the Mediterranean throughout the period. The magnitude of its forcing was small up to around 1920, and has grown continuously since then. DAMIP models indicate GHG had caused almost 2 K of warming by the end of the simulations in 2014, and caused almost three-quarters of the total modelled warming from 1975 to 2014.

185 The Mediterranean cooled in response to Aero forcing (blue dash-dotted line in Figure 3a). This forcing dominated total climate forcing up to 1920, and outweighed GHG forcing for much of the 20th century. Its peak impact on the Mediterranean was reached in the late 1970s, then stabilised for a short time before reducing in magnitude from about 1990 onwards, in line with aerosol emissions (e.g. Lund et al., 2019). The net warming from 1990 to 2014 due to reduced aerosol burdens caused almost one-quarter of the total modelled warming in this recent period.

190 Finally, we note how the sum of responses to individual forcings are consistent with Hist signals (not shown), suggesting Mediterranean warming is largely explained as a linear response to external forcings. Confidence in these DAMIP signals was also assessed using the inter-model spread. Figure 3b contains mean signals for each of the six climate models, for the two main forcings of GHG and Aero. All six models agree on the respective sign of the two forcings, and signal amplitudes are similar too. We conclude that GHG and Aero forcings of Mediterranean temperatures are a robust feature of DAMIP models

195 and that these two forcings are the primary drivers of recent Mediterranean Warming.

Formatted: Space After: 1 line, Line spacing: single

4 Trends in hail risk

200 In this section, we review past research showing a link between the Mediterranean Sea and damaging hail events in the higher risk parts of Europe, and how rising low-level moisture amounts are implicated in increasing trends of hail in these areas, thus connecting the warming Mediterranean to rising hail risk. Next, we provide quantitative estimates of trends in damaging hail, based on both published results and a new analysis of hail losses provided by national insurance bodies, to quantify how the warming Mediterranean may be impacting this peril.

205 It is difficult to quantify the fraction of all hail damage across the continent which involves a Mediterranean moisture source, since hail damage records are spatially and temporally incomplete. However, past research has linked many of the most damaging events in Europe to low-level Mediterranean air masses (e.g. 4.1 Multidecadal connection from Mediterranean temperatures to hail risk)

210 It is difficult to precisely measure the impacts that Mediterranean temperatures have on hail risk at multidecadal scales, mainly due to the incompleteness of hail damage records both spatially and temporally, though their strong connection has been established. Basic thermodynamic principles indicate a warming sea surface is accompanied by similar increases in temperature of the overlying air, increasing its water-holding capacity, and as a result increasing the low-level moisture in the atmosphere. Further, the low-level moisture is an important ingredient for severe hail events. Indeed, previous studies of extreme damaging hail events in Europe reveal how their low-level moisture is drawn from the Mediterranean basin (e.g.

Heimann and Kurz, 1985; Kunz et al., 2018; Piper et al., 2019; Kunz et al., 2020; Kopp et al., 2023). ~~Meanwhile,~~ Kunz et al. (2020) performed a more comprehensive analysis of hailstorms in their study area (Germany, France, Belgium, Luxembourg) and noted that the typical case involves a trough drawing moist Mediterranean air northwards over Europe on its eastern flank, while the increased shear and lifting processes from the trough produce conditions conducive for organised thunderstorm development. Other sources of low-level moisture may also cause damaging hail in some parts of Europe, such as the Cantabrian Sea (e.g. de Pablo Dávila et al., 2021) and the Black Sea (e.g. Piper et al., 2019), ~~but it though the Mediterranean is clear that the~~ largest and warmest body of water in the region, ~~the Mediterranean Sea, is hence very likely~~ the main source of the high dewpoints in the damaging hail events in the higher risk ~~areas of~~ central and ~~southern parts of western mainland~~ Europe.

A number of studies ~~identify increases in~~ have expanded on the connection between hail risk, low-level moisture as being the ~~and a warming Mediterranean to show that low-level moisture increases have been a main causedriver of rising hailstorm~~ trends in damaging hail in high-risk across parts of Europe over the past few decades. Low-level moistening increases the severity of hail in, via two ways. First pathways. The first pathway concerns moisture ~~drives driving~~ convective instability, hence greater amounts of water vapour will tend to intensify thunderstorms and produce more severe hail (e.g. Kunz et al. 2009; Mohr and Kunz, 2013; Púčik et al., 2017; Taszarek et al., 2021; Wilhelm et al., 2024). Second, Schemm et al. (2017) found more frequent strong fronts ~~hazard and losses. A number of studies identify rising amounts of low-level moisture as the~~ main cause of upward trends in damaging hail over the past few decades, in different parts of the higher-risk zone in central and western Europe: for example in France (Dessens, 1995), Germany (Kunz et al. 2009; Mohr and Kunz, 2013), Switzerland (Wilhelm et al., 2024), northern Italy (Battaglioli et al., 2023) and across the higher-risk areas of central and western Europe (e.g. Rädler et al., 2018). Further, multiple climate model projections suggest rising low-level moisture will continue to drive upward trends in damaging hail in the future (Rädler et al., 2019). The second pathway relating moisture to damaging hail concerns the rising frequency of extremely strong fronts. Schemm et al. (2017) found very strong fronts became more frequent over the bulk of Europe from 1979 to 2014 due to rising humidity, and Kunz et al. (2020) described how hail associated with fronts were more damaging, due to larger hailstones and longer hail swathes. Severe hail

The occurrence of individual severe hail events depends on other quantities such as deep layer shear, steep mid-level lapse rates and the low-level cap, and ~~Taszarek et al. (2021) show any long-term changes to these quantities could also modulate hail risk. Past studies have investigated these other ingredients have quite and find relatively steady, or even slightly inhibiting recently, implying trends (e.g. Mohr and Kunz, 2013; Rädler et al., 2018; more fully documented in Taszarek et al., 2021). These results support increasing~~ low-level moisture amounts ~~areas~~ the ~~dominant main~~ driver of upward trends in hail risk over the past few decades.

The full picture of trends in hail risk at all European locations will require consideration of other moisture source such as the Black Sea, and other environmental ingredients such as the warmer, drier air that inhibits convective initiation, and future research into their contributions would be valuable. ~~At the present time, Our current knowledge suggests~~ it is ~~clear~~ very likely that a warming Mediterranean is ~~a primary the main~~ contributor to the recent trends in damaging hail across the higher-risk in

key parts of Europe. We now seek to quantify the trend in hail risk in the higher-risk areas, for guidance on the impacts of the how a warming Mediterranean may be impacting this peril.

#### 4.2 Measured trends in hail risk

Specific measures of recent trends in hail damage fall into two broad types: those reflecting hailstone occurrence, and those based on observed hail damages. The former type has the benefit of more homogeneous hazard data over time, but are often based on models of hail occurrence at a location given the atmosphere conditions, with uncertainty as to whether the models accurately capture the relative roles of all hailstorm ingredients over the past few decades. In contrast, evaluations based on damages reflect hail occurrence at the ground, however, there is some uncertainty in estimated losses at the time of older events, and in how the losses are indexed to represent the total damages if the events occurred in the present day. Given these uncertainties, we will consider estimates of trends from both methods.

Hazard-based studies generally find that large hail has become increasingly common across most of Europe since the 1970s. For instance, extensive hailpad networks indicate a trend toward more intense hail in both southern France over the 1989-2009 period (Berthet et al., 2011), and the central-eastern Alpine area of Italy over the 1975-2009 period (Eccel et al., 2012). A second type of hazard-based study provides more spatially complete and longer hail reconstructions, based on weather ingredients from reanalyses and calibrated to observations such as damaging hail on the ground (e.g. Mohr and Kunz, 2013; Rädler et al., 2018; Battaglioli et al., 2023). The hail model developed by Rädler et al. (2018) indicated the annual amount of hail  $\geq 2$  cm increased at around 2.4% p.a. over 1979-2015 in a region covering Germany and the Alps, and up by 1.4% p.a. over the greater area of western and central Europe. More recently, Battaglioli et al. (2023) described a model which had been calibrated to observed lightning and large hail reports, then estimated linear trends in damaging hail occurrence from 1950 to 2021, and found it had been rising generally across Europe, with relatively bigger more positive changes for larger hail. They highlighted how hailstones  $\geq 5$  cm are now three times more likely than in the 1950s in northern Italy, though the trend was not uniform throughout the whole period, with most of the increase occurring since the 1980s, hence linear trends since 1950 are likely to underestimate trends over the past four decades. Some insurance industry research used models similar to those of Rädler et al. (2018) and Battaglioli et al. (2023): Stormwise Ltd. (2024) studied linear trends over 1960-2023 while Partner Re (2024) examined 1950-2022, and both report hazard driving increases in annual hail risk of around 1 to 1.5% p.a. over much of mainland Europe. The trends from Rädler et al. (2018) are higher than others, however, they base and likely due to their trends on representing the more recent 1979-2015 period hence more synchronised aligned with the period of Mediterranean warming, which began around since about 1980.

There are various sources of property losses due to hail and they all also show an upward trend in hail risk over recent decades. Many in the insurance industry get their impression of rising hail severity in Europe from publicly issued industry loss estimates. There have been some notable events in the past few years, with 4.5 billion USD of insured losses from severe thunderstorms in Europe during summer 2021 (Swiss Re, 2021), then in 2022, French property and automobiles suffered five billion euros of insured loss due to hail (France Assureurs, 2023), while in 2023, hail storms caused over five billion euros of

insured loss, mostly in Italy (Swiss Re, 2024). These loss severities were rarer in earlier times: ~~while~~ storms Andreas (2013) and Ela (2014) caused multi-billion euro insured losses, mainly to Germany and France respectively (Swiss Re, 2014 and 2015) and were the first major events since the multi-billion loss in the Munich hailstorm of 1984 (e.g. ~~PueikPúčík~~ et al., 2019). While the sudden spate of multi-billion euro events is impactful, it ~~is viewed as~~ provides relatively weak evidence of rising hail risk, due to significant errors in estimates of damages from older storms and statistical uncertainty from small sample sizes.

Consideration of other data sources and analysis provides a more robust view on hail loss trends. Early work included a study by Dessens (1995) which found an upward trend in crop-hail insurance losses in France after 1980, while Kunz et al. (2009) examined damage to buildings in southwest Germany over the period 1986-2004, and found the number of severe hail days had increased significantly, by around 3.5% per year.

We supplement earlier damage studies with a new analysis of annual hail losses published by national insurance associations, which were described in subsection 2.2. These industry bodies often monitor losses to their market, and some of their records extend to the past few decades. Figures 4a-c show timeseries of hail losses in Germany, France and Switzerland respectively. The timeseries of annual losses in the 1973-2023 period for automobiles in Germany due to the combined wind and hail perils (the vast majority of which are due to hail) contains an annual growth rate of 3.1% over the past 50 years, while more modern data since 1980 grow by 2.2% per year, and both trends are significantly different from zero at the 1% level ~~based on a standard two-sided t-test (used in all significance testing of trends)~~. Figure 4b presents annual hail losses to all insured buildings in France, containing a +5.8% annual growth rate of hail losses from 1990 to 2022, which drops to +4.7% p.a. when excluding the final year with its huge hail losses. Both of these trends are different from zero at the 1% level. Finally, Figure 4c shows annual hail loss costs in Switzerland for the 1980 to 2023 period for all buildings in the 19 Swiss cantons covered by public insurance, and the best-fitting trend is 2.76% per year over the past 44 years, and *not* different from zero at the 1% level (p-value = 0.058076).

In summary, the datasets shown in Figure 4a-c indicate losses have been trending upward at 2 to 5% per year over the past few decades, and consistently faster than the 1 to 2% range suggested by the above-mentioned hazard-based studies. Losses from national bodies are based on accurate information from surveys of reported losses combined with careful indexation methods using data widely regarded as suitable for its purpose, and represent the industry's best estimates of loss growth. However, hazard-based estimates use hail models based on our best understanding of atmospheric processes, and calibrated to observed hail on the ground, and they represent the best estimates from meteorology. The cause of their different growth rates is unclear, and future investigation into how fast hail damage has been increasing would be useful. In the meantime, we suggest the best estimate of rising hail risk to property is around 2% per year for mainland Europe.




Figure 4: timeseries of national hail damages for three countries. Insurance industry hailstorm losses are shown for (a) automobiles in Germany (GDV, 2024), (b) buildings in France (France Assureurs, 2023), and (c) buildings in Switzerland (VKG, 2022, 2024).

In practice, a 2% per year trend in hail risk may accumulate ~~and to~~ harm ~~the insurance industry companies~~. Their ~~view of risk~~ pricing is often ~~based on models~~ calibrated to past experience ~~of hazard or loss, and they will have lower hence may lag behind~~ current levels of risk ~~than the present day due to the hail trends~~. As an ~~For~~ example, if a ~~model view of risk~~ is calibrated to a time period with a midpoint of 10 years ago, and the risk has been increasing at 2% per year, then this translates to a  $1.02^{10} = 21.9\%$  underestimate of the present-day risk. This is material in the context of larger companies typically aiming for annual profits of around 5 to 10%.




Figure 4: timeseries of national hail damages for three countries. Insurance industry hailstorm losses are shown for (a) automobiles in Germany (CDV, 2024), (b) buildings in France (France Assureurs, 2023), and (c) buildings in Switzerland (VKG, 2022, 2024).

## 5 Conclusions

The Mediterranean Sea has been warming at three times the rate of global mean SSTs over the past 45 years (Figure 2), and its drivers of this rapid heating were investigated using results from the CMIP6-DAMIP experiments. The DAMIP climate models were found to simulate most of reproduce the observed variations in observed Mediterranean temperatures at multidecadal timescales since 1870, especially the rapid heating in recent decades, building confidence in the realism fidelity of these models multimodel mean estimates. They indicate anthropogenic forcings from greenhouse gases (GHG) and aerosols (Aero) have had the greatest influence on multidecadal changes in Mediterranean temperatures. Anthropogenic aerosols dominated up to the late 1970s to produce a slightly cool period on average, then clean-air acts reduced their impacts while GHG forcing continued to grow increase, resulting in a rapid warming of the Mediterranean of about 0.54 K / decade since 1980. Rising GHG rises amounts were responsible for about three-quarters 75% of this warming over the past four decades, with the remaining portion mostly due to declining aerosol burdens. Major volcanoes modify Mediterranean SSTs, though their peak temporary cooling of 0.5 K over a few years is minor compared to anthropogenic forcings.

The temperature trends in the Mediterranean Sea are a concern for hail risk in central and southern Europe. It is known that the Mediterranean is the main source of the high moisture amounts that drive severe are a key ingredient of the damaging hailstorms in the higher risk parts of Europe, and its warming fits with the increases in low-level moisture which are driving hail trends since about 1980. Recent trends in hail damages over these higher risk parts of Europe were reviewed. We

345 ~~synthesized various studies and datasets~~ to measure ~~the impacts due to Mediterranean warming trends in European hail risk~~  
~~since 1980~~. Two different types of studies produce a similar sign of trend, but different rates of growth in risk over recent  
decades. First, models of hail based on environmental conditions and calibrated to observed occurrences of larger hail indicate  
~~risk damage~~ across mainland Europe has been rising by 1 to 2% per year since the mid-20<sup>th</sup> century ~~due to ongoing warming~~,  
and the trend since 1980 ~~may be~~ nearer the upper end of this range. Second, trends in estimates of hail damage to property,  
indexed to the present-day, show annual hailstorm losses are increasing by 2 to 5% in some key countries (Germany, France  
and Switzerland). ~~above exposure-driven growth~~. Both methods have strengths and weaknesses, and future work on resolving  
350 their different estimates would be useful. We suggest a best estimate of 2% per year growth in hail damage based on current  
evidence. Uncertainty is large, though large-scale trends in the higher-risk ~~area~~ ~~areas of western and central Europe~~ are positive  
from almost all studies.

Impacts of rising trends in ~~damaging~~ hail are material to society. For example, if an estimate of hail risk was based on a climate  
centred on 10 years ago, then a 2% per year growth accumulates to a 22% underestimate of the damages from a peril which  
355 causes billions of euros of losses to Europe each year.

~~While Past research establishes how the warming Mediterranean played a dominant role in upward trends of damaging hail~~  
~~over the past few decades, though~~ further work is required to precisely measure ~~how much of the increase in hail risk in central~~  
~~and southern Europe is due to the warming Mediterranean, it is clear~~ the relation. ~~In this study, we find that it has played an~~  
~~important role, and will continue to do so given~~ the current trajectory of anthropogenic forcings. ~~We indicate sustained~~  
360 ~~warming of the Mediterranean, and we~~ recommend including ~~hail~~ ~~upward~~ trends in ~~our~~ views of ~~risk~~ ~~hail losses across high-~~  
~~risk parts of mainland Europe.~~

## Acknowledgements

The authors are grateful to the reviewers and the public commentator for ~~their~~ feedback which improved this manuscript. ~~SC~~  
365 ~~is very grateful, and~~ to Inigo Limited for funding part of this research.

## Code/Data availability

HadISST data are available at <https://www.metoffice.gov.uk/hadobs/hadisst/> (accessed on 24<sup>th</sup> October 2024). CMIP6-DAMIP  
climate model results were downloaded from the Earth System Grid Federation (ESGF): <https://aims2.llnl.gov/search/cmip6/>  
(accessed July-October 2024).

370 **Author Contribution**

SC designed the tests and did the analysis. SC and TC prepared the manuscript.

**Competing Interests**

The authors declare no competing interest.

375 **References**

~~Aizawa T., Oshima N. and Yukimoto S.: Contributions of anthropogenic aerosol forcing and multidecadal internal variability to mid-20th century Arctic cooling—CMIP6/DAMIP multimodel analysis. *Geophys. Res. Lett.*, 49, e2021GL097093, <https://doi.org/10.1029/2021GL097093>, 2022.~~

Ali, E., W. Cramer, J. Carnicer, E. Georgopoulou, N.J.M. Hilmi, G. Le Cozannet, and P. Lionello, 2022: Cross-Chapter Paper 4: Mediterranean Region. In: *Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegria, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 2233–2272, doi:10.1017/9781009325844.021.

380 Battaglioli F., Groenemeijer P., Púčik T., Taszarek M., Ulbrich U. and Rust H.: Modeled Multidecadal Trends of Lightning and (Very) Large Hail in Europe and North America (1950–2021), *J. Appl. Meteorol. Clim.*, 62, 1627–1653, <https://doi.org/10.1175/JAMC-D-22-0195.1>, 2023.

Berthet C., Dessens J. and Sanchez, J. L.: Regional and yearly variations of hail frequency and intensity in France. *Atmos. Res.*, 100, 391–400, <https://doi.org/10.1016/j.atmosres.2010.10.008>, 2011.

390 Booth B. B. B., Dunstone N. J., Halloran P. R., Andrews T. and Bellouin, N.: Aerosols implicated as a prime driver of twentieth-century North Atlantic climate variability. *Nature*, 484(7393), 228–232. <https://doi.org/10.1038/nature10946>, 2012.

Butterworth S., “On the Theory of Filter Amplifiers,” *Experimental Wireless and the Wireless Engineer*, Vol. 7, pp. 536-541, [https://www.changpuak.ch/electronics/downloads/On\\_the\\_Theory\\_of\\_Filter\\_Amplifiers.pdf](https://www.changpuak.ch/electronics/downloads/On_the_Theory_of_Filter_Amplifiers.pdf), 1930,

Chen, J., & Dai, A. (2023). The atmosphere has become increasingly unstable during 1979–2020 over the Northern Hemisphere. *Geophysical Research Letters*, 50, e2023GL106125. <https://doi.org/10.1029/2023GL106125>.

395 CMIP: CMIP Model and Experiment Documentation, <https://wcrp-cmip.org/cmip-model-and-experiment-documentation/>, 2025.

de Pablo Dávila, F., Soriano, L.J.R., Alonso, C.J., García, M.M. and Martín, J.R.: Synoptic patterns of severe hailstorm events in Spain. *Atmospheric Research*, 250, 105397. <https://doi.org/10.1016/j.atmosres.2020.105397>, 2021.

400 Dessens J.: Severe convective weather in the context of a nighttime global warming. *Geophys. Res. Lett.*, 22, 1241–1244, <https://doi.org/10.1029/95GL00952>, 1995.

Eccel E., Cau P., Riemann-Campe K. and Biasioli F.: Quantitative hail monitoring in an alpine area: 35-year climatology and links with atmospheric variables. *Int. J. Climatol.*, 32, 503–517, <https://doi.org/10.1002/joc.2291>, 2012.

Eyring V., Bony S., Meehl G. A., Senior C. A., Stevens B., Stouffer R. J. and Taylor K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model Dev.*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.

405 FFB: FFB Construction Cost Index, [https://www.ouutils.ffbatiment.fr/federation-francaise-du-batiment/le-batiment-et-vous/en\\_chiffres/indices-index/Chiffres\\_Index\\_FFB\\_Construction.html](https://www.ouutils.ffbatiment.fr/federation-francaise-du-batiment/le-batiment-et-vous/en_chiffres/indices-index/Chiffres_Index_FFB_Construction.html), 2025

France Assureurs.: L'Assurance des Événements Naturels en 2022, Fédération Française de l'Assurance, <https://www.franceassureurs.fr/wp-content/uploads/assurance-des-evenements-naturels-en-2022.pdf>, 2023.

410 García-Monteiro S., Sobrino J., Julien Y., Soria G., and Skokovic D.: Surface Temperature trends in the Mediterranean Sea from MODIS data during years 2003–2019. *Regional Studies in Marine Science*, 49:102086, <https://doi.org/10.1016/j.rsma.2021.102086>, 2022.

GDV: Data service for the natural hazards report, <https://www.gdv.de/gdv/statistik/datenservice-zum-naturgefahrenreport>, 2024.

415 Gillett N. P., Shiogama H., Funke B., Hegerl G., Knutti R., Matthes K., Santer B. D., Stone D. and Tebaldi C.: The Detection and Attribution Model Intercomparison Project (DAMIP v1.0) contribution to CMIP6, *Geosci. Model Dev.*, 9, 3685–3697, <https://doi.org/10.5194/gmd-9-3685-2016>, 2016.

Heimann D., and Kurz M.: The Munich hailstorm of July 12, 1984. A discussion of the synoptic situation. *Beitr. Phys. Atmos.* 58: 528–544, <https://elib.dlr.de/64721/1/85-heimann.pdf>, 1985

420 James P., Stohl A., Spichtinger N., Eckhardt S., and Forster C.: Climatological aspects of the extreme European rainfall of August 2002 and a trajectory method for estimating the associated evaporative source regions, *Nat. Hazards Earth Syst. Sci.*, 4, 733–746, <https://doi.org/10.5194/nhess-4-733-2004>, 2004.

Kopp J., Schröer K., Schwierz C., Hering A., Germann U. and Martius O.: The summer 2021 Switzerland hailstorms: weather situation, major impacts and unique observational data. *Weather*, 78: 184-191. <https://doi.org/10.1002/wea.4306>, 2023.

425 Krug, A., Aemisegger, F., Sprenger, M., and Ahrens, B.: Moisture sources of heavy precipitation in Central Europe in synoptic situations with Vb-cyclones, *Clim. Dynam.*, 59, 3227–3245, <https://doi.org/10.1007/s00382-022-06256-7>, 2022.

Kunz M., Sander J. and Kottmeier C.: Recent trends of thunderstorm and hailstorm frequency and their relation to atmospheric characteristics in southwest Germany. *Int. J. Climatol.*, 29, 2283–2297, <https://doi.org/10.1002/joc.1865>, 2009.

430 Kunz, M., Blahak, U., Handwerker, J., Schmidberger, M., Punge, H. J., Mohr, S., Fluck, E., and Bedka, K. M.: The severe hailstorm in southwest Germany on 28 July 2013: characteristics, impacts and meteorological conditions, *Q. J. Roy. Meteorol. Soc.*, 144, 231–250, <https://doi.org/10.1002/qj.3197>, 2018.

Formatted: French (France)

- Kunz M., Wandel J., Fluck E., Baumstark S., Mohr S. and Schemm S.: Ambient conditions prevailing during hail events in central Europe, *Nat. Hazard Earth Sys.*, 20, 1867–1887, <https://doi.org/10.5194/nhess-20-1867-2020>, 2020.
- 435 Lionello, P. and L. Scarascia, 2018: The relation between climate change in the Mediterranean region and global warming. *Reg. Environ. Change*, 18(5), 1481–1493, <https://doi.org/10.1007/s10113-018-1290-1>.
- Lund M. T., Myhre G. and Samset B. H.: Anthropogenic aerosol forcing under the Shared Socioeconomic Pathways, *Atmos. Chem. Phys.*, 19, 13827–13839, <https://doi.org/10.5194/acp-19-13827-2019>, 2019.
- MedECC (2020) Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the
- 440 Future. First Mediterranean Assessment Report [Cramer, W., Guiot, J., Marini, K. (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, 632pp. ISBN: 978-2-9577416-0-1 / DOI: 10.5281/zenodo.7224821
- Mohr S. and Kunz M.: Recent trends and variabilities of convective parameters relevant for hail events in Germany and Europe. *Atmos. Res.*, 123, 211–228, <https://doi.org/10.1016/j.atmosres.2012.05.016>, 2013.
- ~~Morice, C. P., Kennedy, J. J., Rayner, N. A., and Jones, P. D.: Quantifying uncertainties in global and regional temperature~~
- 445 ~~change using an ensemble of observational estimates: The HadCRUT4 data set, *J. Geophys. Res.*, 117, D08101, <https://doi.org/10.1029/2011JD017187>, 2012.~~
- Mulet, S., Buongiorno Nardelli, B., Good, S., Pisano, A., Greiner, E., Monier, M., Autret, E., Axell, L., Boberg, F., Ciliberti, S., Drévillon, M., Droghei, R., Embury, O., Gourrion, J., Høyer, J., Juza, M., Kennedy, J., Lemieux-Dudon, B., Peneva, E., Reid, R., Simoncelli, S., Storto, A., Tinker, J., Von Schuckmann, K., Wakelin, S. L., 2018. Ocean temperature and salinity.
- 450 In: Copernicus Marine Service Ocean State Report, Issue 2, Journal of Operational Oceanography, 11:sup1, s5–s13, DOI: 10.1080/1755876X.2018.1489208
- Partner Re: The Contribution of Climate Change to Europe's Increasing Hail Losses, <https://www.partnerre.com/perspectives/the-contribution-of-climate-change-to-europes-increasing-hail-losses/>, 2024
- Piper, D. A., Kunz, M., Allen, J. T., and Mohr, S.: Investigation of the temporal variability of thunderstorms in Central and
- 455 Western Europe and the relation to large-scale flow and teleconnection patterns, *Q. J. Roy. Meteor. Soc.*, 145, 3644–3666, <https://doi.org/10.1002/qj.3647>, 2019.
- ~~Púčik, T., Groenemeijer, P., Rädler, A. T., Tijssen, L., Nikulin, G., Prein, A. F., van Meijgaard, E., Fealy, R., Jacob, D., and~~
- ~~Teichmann, C.: Future changes in European severe convection environments in a regional climate model ensemble, *J. Climate*, 30, 6771–6794, <https://doi.org/10.1175/JCLI-D-16-0777.1>, 2017.~~
- 460 ~~Púčik, T.,~~ Castellano C., Groenemeijer P., Kuhne T., Rädler A. T., Antonescu B. and Faust E.: Large hail incidence and its economic and societal impacts across Europe. *Mon. Weather Rev.*, 147, 3901–3916, <https://doi.org/10.1175/MWR-D-19-0204.1>, 2019.
- Rädler A. T., Groenemeijer P., Faust E. and Sausen R.: Detecting severe weather trends using an additive regressive convective hazard model (AR-CHaMo). *J. Appl. Meteorol. Clim.*, 57, 569–587, <https://doi.org/10.1175/JAMC-D-17-0132.1>, 2018.

465 [Rädler, A. T., Groenemeijer, P. H., Faust, E., Sausen, R., and Púcik, T.: Frequency of severe thunderstorms across Europe expected to increase in the 21st century due to rising instability, npj Clim. Atmos. Sci., 2, 30, <https://doi.org/10.1038/s41612-019-0083-7>, 2019.](#)

Raupach T. H., Martius O., Allen J. T., Kunz M., Lasher-Trapp S., Mohr S., Rasmussen K. L., Trapp R. J. and Zhang Q. (2021): The effects of climate change on hailstorms, Nat. Rev. Earth Environ., 2, 213–226, <https://doi.org/10.1038/s43017-020-00133-9>, 2021.

470 Rayner N. A., Parker D. E., Horton E. B., Folland C. K., Alexander L. V., Rowell D. P., Kent E. C. and Kaplan A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, J. Geophys. Res., 108, 4407, <https://doi.org/10.1029/2002JD002670>, 2003.

Roquet, H., Pisano, A., Embury, O., 2016. Sea surface temperature. In: von Schuckmann et al. 2016, The Copernicus Marine Environment Monitoring Service Ocean State Report, Jour. Operational Ocean., vol. 9, suppl. 2. doi:10.1080/1755876X.2016.1273446.

475 [Rubino, A., Zanchettin, D., De Rovere, F., McPhaden, M. J.: On the interchangeability of sea-surface and near-surface air temperature anomalies in climatologies, Sci. Rep. 10, 7433, <https://doi.org/10.1038/s41598-020-64167-1>, 2020.](#)

Schemm, S., Sprenger M., Martius O., Wernli H., and Zimmer, M.: Increase in the number of extremely strong fronts over Europe? A study based on ERA-Interim reanalysis (1979–2014), Geophys. Res. Lett., 44, 553–561, <https://doi.org/10.1002/2016GL071451>, 2017.

480 Seneviratne S., Zhang X., Adnan M., Badi W., Dereczynski C., Luca A. D., Ghosh S., Iskandar I., Kossin J., Lewis S., Otto F., Pinto I., Satoh M., Vicente-Serrano S., Wehner M., and Zhou, B.: Weather and Climate Extreme Events in a Changing Climate. In Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, <https://doi.org/10.1017/9781009157896.013>, 2021.

Stormwise Ltd.: Climate Change impacts on hail, <https://stormwise.co.uk/climate-change-and-hail>, 2024.

Swiss Re: Natural catastrophes and man-made disasters in 2013, Swiss Re Institute, Sigma No.1/2014, <https://www.swissre.com/institute/research/sigma-research/sigma-2014-01.html>, 2014.

Swiss Re: Natural catastrophes and man-made disasters in 2014, Swiss Re Institute, Sigma No.2/2015, <https://www.swissre.com/institute/research/sigma-research/sigma-2015-02.html>, 2015.

490 Swiss Re: Global insured catastrophe losses rise to USD 112 billion in 2021, the fourth highest on record, Swiss Re Institute estimates, <https://www.swissre.com/media/press-release/nr-20211214-sigma-full-year-2021-preliminary-natcat-loss-estimates.html>, 2021.

Swiss Re: Natural catastrophes in 2023: gearing up for today's and tomorrow's weather risks, Swiss Re Institute, Sigma No.1/2024, <https://www.swissre.com/dam/jcr:c9385357-6b86-486a-9ad8-78679037c10e/2024-03-sigma1-natural-catastrophes.pdf>, 2024.

Formatted: Portuguese (Portugal)

Formatted: Portuguese (Portugal)

Formatted: Portuguese (Portugal)

Taszarek M., Allen J. T., Brooks H. E., Pilguy N. and Czernecki B.: Differing trends in United States and European severe thunderstorm environments in a warming climate. *B. Am. Meteorol. Soc.*, 102, E296–E322, <https://doi.org/10.1175/BAMS-D-20-0004.1>, 2021.

- 500 Tradowsky, J. S., Philip, S. Y., Kreienkamp, F., Kew, S. F., Lorenz, P., Arrighi, J., Bettmann, T., Caluwaerts, S., Chan, S. C., De Cruz, L., De Vries, H., Demuth, N., Ferrone, A., Fischer, E. M., Fowler, H. J., Goergen, K., Heinrich, D., Henrichs, Y., Kaspar, F., Lenderink, G., Nilson, E., Otto, F. E. L., Ragone, F., Seneviratne, S. I., Singh, R. K., Skålevåg, A., Termonia, P., Thalheimer, L., Van Aalst, M., Van Den Bergh, J., Van De Vyver, H., Vannitsem, S., Van Oldenborgh, G. J., Van Schaeybroeck, B., Vautard, R., Vonk, D., and Wanders, N.: Attribution of the heavy rainfall events leading to severe flooding  
505 in Western Europe during July 2021, *Clim. Change*, 176, 90, <https://doi.org/10.1007/s10584-023-03502-7>, 2023.

VKG: Analyse langfristiger Gebäudeschadendaten, published by Vereinigung Kantonalen Gebäudeversicherungen VKG, [https://cms.vkg.ch/media/at1kb01p/vkf\\_analyse-schadendaten\\_de.pdf](https://cms.vkg.ch/media/at1kb01p/vkf_analyse-schadendaten_de.pdf), 2022.

VKG: Elementar: Übersicht 2004-2023 – Elementarschäden an Gebäuden (20-Jahresvergleich) [https://cms.vkg.ch/media/tutg0vgg/elementarschaeden-an-gebaeuden-2004-2023\\_de.pdf](https://cms.vkg.ch/media/tutg0vgg/elementarschaeden-an-gebaeuden-2004-2023_de.pdf), 2024.

- 510 Volosciuk, C., Maraun, D., Semenov, V. et al. Rising Mediterranean Sea Surface Temperatures Amplify Extreme Summer Precipitation in Central Europe. *Sci Rep* 6, 32450 (2016). <https://doi.org/10.1038/srep32450>.

Wilhelm, L., Schwierz, C., Schröer, K., Taszarek, M., and Martius, O. (2024). Reconstructing hail days in Switzerland with statistical models (1959–2022), *Nat. Hazards Earth Syst. Sci.*, 24, 3869–3894, <https://doi.org/10.5194/nhess-24-3869-2024>.