

**Cities near volcanoes: Which cities are most exposed to volcanic hazards?**

Formatted: Font: Times New Roman

Elinor S. Meredith<sup>1,2\*</sup>, Rui Xue Natalie Teng<sup>1</sup>, Susanna F. Jenkins<sup>1</sup>, Josh L. Hayes<sup>3</sup>, Sébastien Biass<sup>4</sup>, Heather Handley<sup>2,5</sup>

**Affiliations:**

<sup>1</sup> Earth Observatory of Singapore, Asian School of the Environment, Nanyang Technological University, Singapore, 639754

<sup>2</sup> Department of Applied Earth Sciences, ITC Faculty, University of Twente, Enschede, The Netherlands

<sup>3</sup> GNS Science, P.O. Box 30368, Lower Hutt, 5040, New Zealand

<sup>4</sup> Department of Earth Sciences, University of Geneva, 13, Rue des Maraîchers, CH-1205 Geneva, Switzerland

<sup>5</sup> School of Earth, Atmosphere and Environment, Monash University, Clayton, Australia

\* Contact: [e.s.meredith@utwente.nl](mailto:e.s.meredith@utwente.nl)

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

## 1. Abstract

Cities near volcanoes expose dense concentrations of people, buildings, and infrastructure to volcanic hazards. Identifying cities globally that are exposed to volcanic hazards helps guide local risk assessment for better land-use planning and hazard mitigation. Previous city exposure approaches have used the city centroid to represent an entire city, and to assess population exposure and proximity to volcanoes. But cities can cover large areas and populations may not be equally distributed within their bounds, meaning that a centroid may not accurately capture the true exposure. In this study, we suggest a new framework to rank global city exposure to volcanic hazards. We assessed global city exposure to volcanoes in the Global Volcanism Program database that are active in the Holocene by analysing populations located within 10, 30, and 100 km of volcanoes. These distances are commonly used in volcanic hazard exposure assessment. City margins and populations were obtained from the Global Human Settlement (GHS) Model datasets. We ranked 1,406,133 cities based on the number of people exposed at different distances from volcanoes, the distance of the city margin from the nearest volcano, and by the number of nearby volcanoes. Notably, 50% of people living within 100 km of a volcano are in cities. We highlight Jakarta, Bandung, and San Salvador, as scoring highly across these rankings. Bandung, Indonesia ranks highest overall with over 8 million people exposed within 30 km of up to 12 volcanoes. South-east Asia has the highest number of exposed city populations (~162 million). Jakarta (~38 million), Tokyo (~30 million), and Manila (~24 million) having the largest number of people within 100 km. Central America has the highest proportion of its city population exposed, with Quezaltepeque and San Salvador exposed to the most volcanoes (n=23). Additionally, we ranked the 1,283,264 Holocene volcanoes by the city populations exposed within 10, 30, and 100 km, the number of nearby cities, and distance to nearest city. Tangkuban Parahu, Tampomas, and San Pablo Volcanic Field, and Tampomas score highly across these rankings. Notably, Gede-Pangrango (~48 million), Laguna Caldera, Tangkuban Parahu (~8 million), and Nejapa-Miraflores (~0.8 million) volcanoes have the largest city populations within 100, 30, and 10 km, respectively. We developed a web app to visualise all the cities with over 100,000 people exposed. This study provides a global perspective on city exposure to volcanic hazards, identifying critical areas for future research and mitigation efforts.

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

## 2. Introduction

48  
 49 As of 2023, more than half (57 %) of the world's population reside in cities (World Bank, 2023).  
 50 These dense urban clusters of buildings, infrastructure, and populations are particularly vulnerable  
 51 to natural hazards (Degg, 1992; Godschalk, 2003), as urban residents are heavily reliant on city  
 52 infrastructure (UNDP, 2021). Such susceptibility exposes cities to high potential losses and  
 53 cascading systematic impacts that can affect the wider region, country, or world (Thouret, 1999;  
 54 Chester et al., 2000; Heiken, 2013; Mani et al., 2021). Recent rapid urbanisation into hazardous  
 55 areas escalates the threat to cities (Pelling, 2012; Freire et al., 2019; Iglesias et al., 2021), driving  
 56 increasing disaster impacts globally (Gu, 2019). Identification of the most exposed cities and  
 57 analysis of the spatio-temporal patterns of urban hazard exposure is crucial for guiding effective  
 58 land-use planning and mitigation efforts. This focus will help prioritise cities that need focussed  
 59 attention for sustainable development and improved preparedness and resilience against future  
 60 disasters (Ariyanti et al., 2020).

61  
 62 Cities situated near volcanoes face a variety of direct threats from volcanic hazards resulting from  
 63 eruptions of various intensities. Historically, volcanic flows have destroyed whole cities; for  
 64 example, pyroclastic density currents (PDCs) emplaced within 10 km of their volcano destroyed  
 65 Herculaneum, Italy, in 79 CE (Volcanic Explosivity Index, VEI 5), Saint Pierre, Martinique, in  
 66 1902 CE (VEI 4), and Plymouth, Montserrat, in 1997 CE (VEI 3). Some cities are repeatedly  
 67 impacted, such as Goma, Democratic Republic of Congo (DRC), less than 30 km from the volcano,  
 68 which was partially inundated by lava flows in 1977 CE, 2002 CE, and 2021 CE. Lahars (volcanic  
 69 mudflows) destroyed the city of Armero, Colombia, approximately 50 km from the Nevado del  
 70 Ruiz volcano in 1985 CE (VEI 3), and the city of Lumajang, Indonesia, 35 km from Semeru  
 71 volcano, in 1909 CE (VEI 2). Some cities are built on old lahar deposits, suggesting they are likely  
 72 to be impacted again (e.g., Arequipa, Peru; ~~Mexico City, Mexico~~; San Salvador, El Salvador). The  
 73 more widely dispersed hazard of tephra falls has destroyed the city of Akrotiri, Greece, around  
 74 1600 BCE (VEI 7), and disrupted and damaged the cities of Kagoshima, Japan in 1914 CE (VEI  
 75 4), Anchorage, USA in 1989 CE (VEI 3), and Angeles City, the Philippines in 1991 CE (VEI 5).

76  
 77 In order to reduce the risks faced by cities situated close to volcanoes, we first must identify which  
 78 cities are most exposed to volcanic hazards. Some past studies have used a localised approach,

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

79 whereby they identify exposed urban areas close to case study volcanoes (e.g., Thouret et al., 2001;  
80 Sandri et al., 2014; Strader et al., 2015; Magill & Blong, 2005; Alberico et al., 2011; Delgado  
81 Granados & Jenkins, 2015; Torres et al., ~~2022~~2023). However, a more systematic regional or  
82 global approach based on the location of all cities would reveal which cities are most exposed. For  
83 example, taking a regional multi-volcano approach, Jenkins et al. (2018) evaluated the potential  
84 impact of tephra fall on 16 major cities in the Asia-Pacific region, highlighting Tokyo, Japan;  
85 Jakarta, Indonesia; and Manila, Philippines as being most exposed. Ranking cities or volcanoes by  
86 city population exposure would allow us to identify global exposure hotspots which may be missed  
87 by a more localised approach.

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

88  
89 Whilst global city analyses have been conducted for other hazards such as coastal flooding  
90 (Hanson et al., 2011), earthquakes (Bilham, 2009), and multi-hazards (Degg, 1992; Brecht et al.,  
91 2013; Gu, 2019), for volcanic hazards, reports or studies often focus on growth rates or total  
92 population numbers of case study cities, based on the location of city centroids close to volcanoes.  
93 For example, in 1989 CE, the UN estimated that of the top 50 fastest growing cities, only four  
94 were exposed to volcanic hazards (UN United Nations, 1989). Three years later, it was estimated  
95 that 10 of the world's most populated cities were located within 30 km of an active volcano (Degg,  
96 1992), Pelling (2012) and Blaikie (~~1994~~2004) highlight Jakarta as a major city at risk from volcanic  
97 activity. Heiken (2013) identified 67 cities (with populations greater than 100,000), home to a total  
98 of ~116 million people, located on or near ~~to~~ active volcanoes. Other studies considered the city's  
99 distance and direction from volcanoes. Donovan and Oppenheimer (2014) list 49 cities within 100  
100 km of a volcano with recognised Quaternary activity and Erfurt-Cooper (2014) provided a list of  
101 25 cities in close proximity to active volcanoes. Chester et al. (2000) and subsequently Auker et  
102 al. (2013) plotted polar charts with examples of highly populated cities in relation to the distance  
103 and direction of the city centroid from nearby volcanoes, within 200 km and 50 km of volcanoes,  
104 respectively. Brown et al. (2015) identified seven capital cities globally within 10 km of volcanoes,  
105 37 within 30 km, and 69 within 100 km. A systematic approach quantifying all global city  
106 populations at specific distances from volcanoes would better capture the variable distribution of  
107 populations across cities and exposure to volcanoes.

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

109 Key cities identified as exposed to volcanic hazards can then be targeted for more localised  
110 research and mitigation efforts. For example, Jenkins et al. (2022) ranked volcanoes in South-East  
111 Asia by population and physical exposure, and proposed that populations near Guntur volcano,  
112 Indonesia, are highly exposed and thus require further study. For identified highly exposed cities,  
113 combining high resolution exposure with hazard maps or footprints can identify key areas in a city  
114 on which to focus future mitigation, such as structural adaptations (e.g., to tephra fall in  
115 Kagoshima: Durand, 2001) and/or land-use planning efforts (e.g., Nieto-Torres et al., 2021;  
116 Strader et al., 2015; Thouret et al., 2001).

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

117  
118 In this study, we propose a new framework to quantify and rank global cities exposed to volcanoes  
119 situated within varying proximity thresholds (10, 30, and 100 km), based on three variables:  
120 population exposed; distance to nearest volcano; and the total count of volcanoes each city is  
121 exposed to. Leveraging high-resolution updated population data allows for assessment of the  
122 spatial variation of population exposure within each city. Instead of looking at the total population  
123 of the city as presented in past studies, this approach more accurately identifies the proportion of  
124 the city population within each distance threshold. Furthermore, our framework also ranks  
125 volcanoes by city exposure based on their proximity to city populations, considering the total count  
126 of cities and their respective populations within the specified distances from volcanoes.

## 128 2.1 Measuring population exposure around volcanoes

129  
130 To quantify global population exposure around volcanoes, studies typically count the number of  
131 people residing within concentric radial buffers of certain sizes around the volcanic vent (e.g.,  
132 Small & Naumann, 2001; Ewert, 2007; Aspinall et al., 2011; Brown et al., 2017; Freire et al., 2019;  
133 Nieto-Torres et al., 2021; Guimarães et al., 2021). For example, Freire et al. (2019) calculated total  
134 populations living within 10, 30, 50, and 100-km radial buffers around active volcanoes between  
135 1975 and 2015 CE. The concentric radial buffer size approach allows for a conservative estimate  
136 of exposure at each volcano so that the cities can be compared and ranked, especially useful for  
137 volcanoes without historical eruptions or high-resolution topography data (Biass et al., 2024). The  
138 radius sizes of buffers around the volcanic vent used to calculate exposure are determined by the  
139 maximum distances of primary volcanic hazards and the extents of potential Volcano Explosivity

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Index (VEI) events. For instance, the emplacement of the majority of primary hazards in a VEI <4 eruption are within a 10-km radius around the volcano, including extents of ballistic projectiles and most dome collapse pyroclastic density currents (PDCs) (Biass et al., 2024). A 30-km radius generally represents the extent of the majority of VEI <5 eruption hazard extents, while a 100-km radius signifies the extent of the majority of VEI <6 eruption hazard extents. However, larger eruptions, secondary hazards, or eruptions occurring from fields or fissures beyond the vent might extend beyond these maximum hazard distances. For example, shown by modelling of a VEI 5 scenario in Biass et al. (2024), tephra loads exceeding  $1 \text{ kmkg/m}^2$  could extend as far as 600 km away.

These exposure counts can be assigned to specific indices for comparison across volcanoes. For example, the Volcanic Population Index (VPI) estimates the number of people living within 5 and 10 km of volcanoes (Ewert & Harpel, 2004). Aspinall et al. (2011) developed this methodology to assess populations within 10, 30, and 100 km of volcanoes. To calculate the Population Exposure Index (PEI), the population counts are weighted according to evidence on historical distributions of fatalities within a given distance from the vent and each volcano is assigned to one of seven PEI indices (Brown et al., 2015). On a local or regional scale, past studies combined these population counts with physical exposure (e.g., Etna volcano, Italy: Del Negro et al., 2022; Rainier volcano, USA: Wood & Soulard, 2009) and/or hazard factors to understand localised volcanic threat (e.g., Ewert et al., 2018; Mangan et al., 2018; Nieto-Torres et al., 2021). Nieto-Torres et al. (2021) developed a volcanic risk index for Central America, considering 41 different factors related to hazards and exposure, assessing population risks within distances of 5, 10, 30, and 100 km from volcanoes. Researchers used various buffer sizes in ongoing efforts to accurately assess the risks that volcanic hazards pose to total populations. No studies conducted exposure analysis based on population exposure by urban type, such as city populations. In this study, we quantify and rank global cities exposed to volcanoes situated within 10, 30, and 100 km of volcanoes.

### 3. Methods

We developed a framework designed to quantify, rank, and assess city exposure to volcanoes (see Fig. 1 and Fig. 2). In this study, *exposed populations* refer to populations in cities within 100 km (unless the distance is specified) from at least one volcano active in the Holocene, as explained in

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman


Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman


Formatted: Font: Times New Roman

Section 2.1. Below we detail how we prepared the city outlines, extracted populations, and ranked the cities and volcanoes. The R code used to generate the results of this study is provided at: <https://github.com/vharg/VolcCities>. The ~~Fig.s~~figures presented in this study focus on the top 10 cities or volcanoes. We developed a web app to comprehensively present results of this study, including the population exposure and maps of city polygons for all the cities with over 100,000 people exposed, which is available at: [https://nteng.shinyapps.io/vharg.github.io/ExposureOfCitiesshiny\\_VolcCities/](https://nteng.shinyapps.io/vharg.github.io/ExposureOfCitiesshiny_VolcCities/).



**Fig. 1:** A flowchart of our framework to assess city exposure to volcanoes. Input datasets (yellow rectangles) are fed into the GIS process (blue rectangles) to collate city polygons (left orange rectangle) and attribute data, clipped by radial buffers (central orange rectangle), into the exposed population polygon dataset (right orange rectangle). ~~For input datasets from the Global Human Settlement~~ For the 2020 urban centres, the GHS-UCDB (2024) dataset was used, and the spatially joined centroids of this dataset were used for names. GHS\_DUC (2023) Name\_2 column was used for city names if they were not available. The cities were ranked by the population size, the distance to nearest volcano, and the maximum number of volcanoes exposed to, and the average of these was presented as the composite rank. The volcanoes chosen are those active in the Holocene epoch. GVP stands for the Global Volcanism Program and GHS stands for Global Human Settlement. The code used in this paper is provided at: <https://github.com/vharg/VolcCities>

192 e



193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

**Fig. 2:** A schematic of our framework for assessing city exposure to volcanoes. The framework includes the following steps: 1) obtaining city polygons for 2020 ~~CE~~ from the GHS-UCDB (2024) urban centre polygons dataset, 2) joining city and country names from GHS-UCDB (2024) centroids, otherwise GHS\_DUC polygons were used, and Natural Earth (2023) was used for subregion names, 3) conducting distance analysis by clipping city polygons within 10, 30, and 100-km buffers from volcanoes, and analysing the number of volcanoes by counting overlaps of 100-km volcano buffers across city polygons, and 4) extracting the population of city polygons within volcano buffers using a population raster. Cities were ranked by population size and the maximum number of exposed volcanoes. Volcanoes included are those active in the Holocene, as per the Global Volcanism Program (GVP). The code used in this paper is provided at: <https://github.com/vharg/VolcCities>

### 3.1 Preparing the exposed city population polygon dataset

We obtained city area outlines from the 2024 ~~CE~~ release of the Global Human Settlement Layer - Settlement Model (GHS-UCDB; Marí Rivero et al., 2024). This dataset is in the World Mollweide (EPSG:54009) projection and we did not reproject it. This dataset identifies the *urban centres*, classified as having a population density of  $\geq 1500$  inhabitants/km<sup>2</sup> and a total population size of at least 50,000 people (Marí Rivero et al., 2024). These city areas will be referred to from here, as *city polygons*. We identified the centroid ~~co-ordinates~~ *coordinates* and the areas of the *city polygons*.

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman



214 We selected only those polygons that had all or part of their area within 100 km of a volcano  
 215 resulting in 1,458 ~~city~~ polygons. Volcano locations in this study were based on coordinates from  
 216 the 1,283 ~~264~~ Holocene Volcano List from the Volcanoes of the World (VOTW) ~~v4.11.1v~~ 5.2.7  
 217 database (Global Volcanism Program, 2024). ~~Co-ordinates~~ Coordinates of the Holocene volcanoes  
 218 in the database are positioned either at the summit for volcanoes with a distinct primary edifice, or  
 219 close to known vents. For volcanic fields with multiple vents, the database presents the most  
 220 prominent or active vent, most recently erupting vent, or centre of the volcanic field, depending  
 221 on information availability. ~~Only the 958 volcanoes that are located within 100 km of a city polygon~~  
 222 ~~are used within this study~~ Using a single point to represent distributed volcanism introduces  
 223 uncertainty, as the precise location of future eruptions is unknown.  
 224  
 225  
 226 The *city polygons* generated from the raster lacked attribute data, therefore, we spatially joined  
 227 each polygon with the city and country names of the 2025 ~~CE~~ projected city polygons from the  
 228 2024-~~CE~~ release of the Global Human Settlement Layer - Settlement Model, Urban Centre  
 229 Database (GHS-UCDB; Marí Rivero et al., 2024). The spatial join was based on the overlap of the  
 230 centroids with the *city polygons*. For the 30 cities that did not have city names (as they are not in  
 231 the 2025 ~~CE~~ dataset), we spatially joined the *city polygons* to the 2023-~~CE~~ release of the GHS  
 232 Degree of Urbanisation Classification (GHS-DUC; Schiavina et al., 2023a) polygons dataset and  
 233 selected the name from the Name\_2 column, and for one case, the Name\_1 column as Name\_2  
 234 was empty. For the 35 cities that did not have any country names, these were added from the  
 235 spatially joined GHS\_DUC dataset. The spatial join was based on the location of the *city polygons*'  
 236 centroids. For the subregions, we spatially joined the *city polygons* to the Natural Earth (2023)  
 237 country polygons and selected the subregions, based on the location of the nearest polygon. This  
 238 dataset was chosen as there were more subregions included than the GHS-UCDB dataset. We then  
 239 grouped the *city polygons* by city name in each country, resulting in a dataset of 1,406 ~~133~~ cities.  
 240  
 241 We made other calculations for the other columns in the dataset shown in Table 1, we included the  
 242 *city polygon* centroid ~~co-ordinates~~ coordinates based on the largest polygon, calculated the *city*  
 243 *polygon* area (km<sup>2</sup>), and measured the shortest distance (km) of the *city polygon* margin that lies  
 244 closest to the nearest volcano. To calculate the total population of the city, we extracted the 2020

**Formatted:** Font: Times New Roman, Not Italic

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

**Formatted:** Font: Times New Roman

population from Global Human Settlement Layer - Population (GHS-POP; Schiavina et al., 2023b) 100-m population rasters within each of the *city polygons* using the *exactextract* function in R (Levine, 2022). This population dataset is in the World Mollweide (EPSG:54009) projection and we did not reproject it.

**Table 1:** The results collated in this study are set out in a dataset with the columns named as in this table. One row represents one city.

Column Name		
Total city polygon	Name	
	Centroid <del>eo-ordinates</del> coordinates	Latitude, Longitude
	Country	
	Continent	
	Subregion	
	Total city area (km <sup>2</sup> )	
	Total city population	
	Nearest volcano	
	Distance to nearest volcano (km)	
	Rank by distance	
	Maximum number of volcanoes exposed by	
	Composite rank	
Results by distance	Rank (by <100 km)	
	Population by distance	<10 km, <30 km, <100 km, >100 km (not exposed), 30 – 10 km, 100 – 30 km
Results by number of volcanoes	Rank (by max <100km)	
	Population by number of volcanoes <100 km	1, 2-5, 6-10, 11-20, 20+
	Population by number of volcanoes <30 km	1, 2-5, 6-10, 11-20, 20+
	Population by number of volcanoes <10 km	0, 1, 2-5, 6-10, 11-20, 20+

### 3.2 Ranking cities

Using the volcano location ~~eo-ordinates~~coordinates as the centroids we created 10, 30, and 100-km radial buffers around each volcano, as these are commonly used to assess volcanic exposure based on typical maximum distances of primary volcanic hazards (Ewert, 2007; Brown et al., 2017; Biass et al., 2024). From here, these will be referred to as *volcano buffers*.

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

We clipped the *city polygons* by each of the three 10, 30, and 100-km radial *volcano buffers* and merged these together to create a map of the city classified by distance to a volcano. This means that parts of the city, or specific proportions, fall into one of the distance categories: <10 km, 10–30 km, 30–100 km, or >100 km from a volcano. We extracted the population using the method shown in 3.1 and where the volcano buffer partially covers a population raster pixel, this function extracts the population number based on the proportion of pixel covered. We then merged and pivoted the dataset so that it resulted in the population in each *volcano buffer* for each city (Table 1). For each *volcano buffer* we ranked the cities by the total populations exposed.

We also ranked cities by how many volcanoes each city's population is exposed to. We overlaid the 100-km *volcano buffers* and calculated the number of buffers that were in contact with each *city polygon*. By intersecting the overlaid buffers by the *city polygons*, we created a map where the areas within the city are classified by the number of volcanoes it is exposed to. From this map, we then extracted the population, so that, for each city, the numbers of people exposed to numbers of volcanoes is known (Table 1). This was repeated for the 30-km and 10-km *volcano buffers*. In separate columns we entered the maximum number of volcanoes that the city is exposed to (within 100 km) and ranked the cities by this amount. This means that the maximum number of volcanoes the city is exposed to relates to the total number of 100-km *volcano buffers* in contact with the *city polygon*.

Finally, we calculated a composite ranking for each city, summing the three rankings of distance to nearest volcano, population <100 km, and number of volcanoes <100 km, and dividing by three to create a final ranking of cities. This assumes the same weighting for each ranking.

### 3.3 Ranking volcanoes

In a separate dataset shown in Table 2, for each volcano, we recorded the volcano name, name, vent ~~eo ordinates~~coordinates, and country from the Global Volcanism Program (2023) Holocene Volcano List. We also calculated the distance of the vent location to the nearest *city polygon*. For each *volcano buffer* of each volcano, we calculated the total populations located within each

Formatted: Font: Times New Roman

*volcano buffer* and *city polygons* using the extraction method detailed in 3.1. We then ranked the volcanoes by the total city populations within 100 km of the volcano. We also calculated the total number of *city polygons* within each *volcano buffer* for each volcano and ranked the volcanoes by the total number of cities within 100 km of the volcano. Finally, we calculated a composite ranking for each volcano, summing the three rankings of total population <100 km, number of cities <100 km, and distance to nearest city, and dividing by three to create a final ranking of volcanoes. This assumes the same weighting for each ranking.

**Table 2:** The results collated in this study are set out in a dataset with the columns named as in this table. One row represents one city.

Column Name		
Volcano	Volcano Name	
	Vent	Latitude, Longitude
	<del>ordinates</del> coordinates	
	Country	
	Nearest city	
	Distance to nearest city (km)	
Results by city population	Composite rank	
	Rank (by <100 km)	
	City population by distance	<100 km, <30 km, <10 km, 100 – 30 km, 30 – 10 km
Results by number of cities	Rank (by max <100 km)	
	Number of cities	
		<100 km, <30 km, <10 km, 100 – 30 km, 30 – 10 km

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

#### 4. Results

We present our results in two parts, firstly we present the rankings for cities and volcanoes within each of the 10, 30, and 100-km *volcano buffers*, and secondly, we present regional trends in the city exposure. Cities were ranked based on: a) the total ~~exposed~~city population ~~exposed~~ to one or more volcanoes, b) the number of volcanoes the city is exposed to, and c) the distance to nearest volcano. Volcanoes were ranked based on: a) the total exposed city population, b) the number of cities exposed, and c) the distance to the nearest city. Here, we define an exposed city population as the population of cities located within 100 km of a volcano. In the following section, we explore the quantification and ranking of these city populations and volcanoes, detailing spatial trends in

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, Not Italic

Formatted: Font: Times New Roman

city exposure. We also explore regional trends in city exposures. The results for all our exposure analyses are presented in Section 9.

#### 4.1 Quantifying and ranking city exposure to volcanoes

Globally, 1,406,133 cities have some proportion of their populations living within 100 km of at least one volcano active in the Holocene (Table 3). Within these cities, ~430,431 million people are exposed within 100 km of volcanoes (Table 3), representing ~50.4% of the total population exposed within 100 km (n= 852,989,097) and ~422.1% of the total population of cities globally (n=3,511,560,764/2,022,199,789). For each city, the exposed population varies from four people in Hamamatsu, Japan, to ~38.1 million people in Jakarta.

**Table 3:** The total number of cities with some proportion of their area within the three volcano buffer distances of at least one volcano, and city populations exposed to volcanoes within each volcano buffer size used in our study.

Distance from nearest volcano	Number of cities exposed	City populations exposed
10	9994	13,756,890/182,876
30	381	428,065,560/118,852,034
100	1,406,133	429,145,006/430,696,763

The greatest number of cities and populations exposed are in Indonesia (see Fig. A3 in the Appendix). The top five cities with the most people living within 100 km from at least one volcano are Jakarta, Indonesia; Tokyo, Japan; Manila, Philippines; Mexico City, Mexico; and Seoul, South Korea (Fig. 4). For these cities, almost the entire populations are exposed to at least one volcano (Jakarta: Salak, Perbakti-Gagak; Manila: ~~Laguna Caldera~~San Pablo Volcanic Field, Taal; Mexico City: Iztaccihuatl, Chichinautzin Volcanic Field, Popocatepetl), except for Tokyo and Seoul where parts of the cities are not exposed to any volcanoes (farther than 100 km; see Fig. A3 in the Appendix). These top five cities represent ~28 % (~121 million) of the total exposed city populations (~429,431 million), with over 677,000 people within 10 km from a volcano. For the other exposed cities, exposed populations decrease gradually as rank increases (Fig. 4; Fig. A2 in

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, Not Highlight

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

the Appendix). When combining rankings of number of people within 100 km, number of volcanoes within 100 km, and distance to nearest volcano, Bandung is the highest ranked city (Fig. 3).

As the spatial distribution of populations varies across cities, we found that some of the highly ranked cities have proportions of their populations located more than 100 km from volcanoes (e.g., Tokyo, Japan; Seoul, South Korea; Tehran, Iran), whilst others have high proportions of their populations located very close to volcanoes. Notably, for the 30 km *volcano buffer*, the entire populations of Bandung, Indonesia (n=8,443,660) and Quito, Ecuador (n=2,435,784) reside within 30 km of at least one volcano. The top five cities ranked by people living within 100 km of volcanoes have relatively low proportions of their populations within 30 km (~18.1%), with most (~82.78%) living between 30 and 100 km. As a result, for rankings at smaller *volcano buffer* distances, these cities drop down to lower rankings (Fig. 4). However, Jakarta and Bandung feature in the top ten ranked cities for exposed populations within all three *volcano buffer* distances and have the highest number of people exposed to at least one volcano within 100 km (n=38,050,484 and 46,494,144) and 30 km (n=8,443,660 and 15,456,793). The top two ranked cities for the 10-km *volcano buffer*: Naples, Italy, and Managua, Nicaragua, combined have a smaller total population than Jakarta; however, they have 40% and 66% of their city populations living within 10 km of volcanoes, respectively, compared to 1% of Jakarta (Fig. 4). Thus, cities may have low total populations but have high proportions of their populations living in close proximity to nearby volcanoes.

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

City Rankings						Volcano Rankings					
Rank	City Name	Population exposed rank	Number of volcanoes rank	Distance to nearest volcano rank	Average of ranks	Rank	Volcano Name	Population exposed rank	Number of cities exposed rank	Distance to nearest city	Average of ranks
1	Bandung, Indonesia	8	30	11	16.33	1	Tangkuban Parahu, Indonesia	4	22	11	12.33
2	Jakarta, Indonesia	1	40	16	19	2	Tampomas, Indonesia	16	9	27	17.33
3	San Salvador, El Salvador	39	1	25	21.67	3	San Pablo V.F., Philippines	8	42	4	18
4	Quito, Ecuador	28	14	29	23.67	4	Guntur, Indonesia	19	11	26	18.67
5	Managua, Nicaragua	50	30	1	27	5	Talagabodas, Indonesia	23	6	34	21
6	Guatemala City, Guatemala	24	13	66	34.33	6	Ciremai, Indonesia	18	4	42	21.33
7	Tokyo, Japan	2	56	65	41	7	Penanggungan, Indonesia	27	9	28	21.33
8	Tarogong, Indonesia	47	56	28	43.67	8	Patuha, Indonesia	10	42	15	22.33
9	Surabaya, Indonesia	9	104	23	45.33	9	Sundoro, Indonesia	33	5	33	23.67
10	Tasikmalaya, Indonesia	25	75	39	46.33	10	Galunggung, Indonesia	24	11	49	28
11	Kuningan, Indonesia	85	75	37	65.67	11	Papandayan, Indonesia	25	27	35	29
12	Bishoftu, Ethiopia	154	56	3	71	12	Kendang, Indonesia	22	27	39	29.33
13	Malang, Indonesia	34	104	78	72	13	Arjuno-Welirang, Indonesia	29	11	52	30.67
14	Adama, Ethiopia	100	40	88	76	14	Bishoftu V.F., Ethiopia	50	51	3	34.67
15	Cirebon, Indonesia	40	75	129	81.33	15	Sumbing, Indonesia	30	11	65	35.33

**Fig. 3: Top 2015 composite rankings of city exposure to volcanic hazards and their score within the different ranks. For cities these are based on three different ranks: a) the total city population exposed to one or more volcanoes within 100 km of volcanoes, b) the number of volcanoes the city is exposed to, c) the distance to nearest volcano. For volcanoes, these rankings are based on three different ranks: a) the total exposed city population within 100 km, b) the number of cities exposed, c) the distance of the volcano to the nearest city. The cells are coloured by the ranking position within the column, with darker red as a higher ranking. The cities and volcanoes are ordered by the average of three rankings (composite rank shown in the Rank column rounded to the nearest integer). V.F. stands for Volcanic Field and V.C. stands for Volcanic Complex.**


Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman



**Fig. 4:** Horizontal stacked bar charts of city population (left) and proportion (right) of each city classified by distance of the city margin to the nearest volcano (within <10, 10 - 30, and 30 - 100-km distances from volcanoes), and ranked by 100, 30, and 10-km radial buffers. Each bar represents a city exposed to volcanoes, and the x axis limits (left) reflect the population range for that volcano buffer.

Fig. 4 shows city rankings of populations by distances to their nearest volcanoes, with 9994 cities having their city margins within 10 km from a volcano. However, ~6367% of cities with populations within 100 km of volcanoes (n=698615) are exposed to more than one volcano. Fig. 5 presents the top 20 ranking of cities by exposed populations, classified by the number of volcanoes the city is exposed to. The Fig. shows that portions of the populations of Tokyo, Japan; Seoul, South Korea; and Nagoya, Japan, are not exposed to any volcanoes, but these cities still rank within the top 20 exposed cities (Fig. 5). Interestingly, not all the largest exposed populations are exposed to multiple volcanoes, such as Seoul, South Korea; Tehran, Iran; Barcelona, Spain; Antananarivo, Madagascar; and Athens, Greece, which have populations exposed to only one volcano. The top four cities with exposed populations have much larger populations compared to the others, with millions more residents. In contrast, the differences in exposed populations among the remaining

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman



390 cities exhibit a more gradual increase, illustrated in Fig. 5 and for mid-sized cities (populations  
391 under 1 million) in the Appendix (Fig. A4).

392  
393 Most of the cities exposed to multiple volcanoes are in Central America, ~~with~~ populations in  
394 Quezaltepeque, El Salvador, is exposed to the highest number of volcanoes (n=23), and San  
395 Salvador, El Salvador is the city with >1 million people exposed to the highest number of  
396 volcanoes (n=~~22~~23) (Fig. 6). Although San Salvador is ranked 39<sup>th</sup> by population within 100 km,  
397 it is third in the composite ranking behind Bandung and Jakarta, due to its high ranking of the  
398 number of volcanoes (Fig. 3). Despite this proximity of cities to multiple volcanoes, there is the  
399 same ranking of the top 10 cities when ranked by the people exposed to the nearest volcano, and  
400 those exposed to at least one volcano. Bandung ranks highly for both ~~the~~ maximum number of  
401 volcanoes to which the city is exposed, and the number of people exposed, with exposed  
402 populations within 100 km of between 9 and 12 volcanoes, with the most, ~4 million, exposed to  
403 10 volcanoes (Fig. 6).




Formatted: Font: Times New Roman

Formatted: Font: Times New Roman


Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman



**Fig. 5:** Stacked bar charts of city population exposure for the top 20 cities coloured by the number of volcanoes the populations are exposed to, within A) 100-km, B) 30-km, and C) 10-km volcano buffers.



**Fig. 6:** Heatmap for the top 10 cities with total populations of over 1 million ranked by exposure to number of volcanoes, showing the city population exposed to increasing numbers of volcanoes for A) 100-km, B) 30-km, and C) 10-km volcano buffers.

#### 4.2 Ranking volcanoes by city exposure

Almost 47.46% of Holocene volcanoes (n=596,578) have city populations located within 100 km of their assumed vents. Indonesia dominates the rankings for city populations within 100 km of volcanoes. Five, seven, six, eight, and three of the top 10 volcanoes within the 100-km, 30-km and 10-km buffers, respectively, are in Indonesia. Among the 596,578 volcanoes close to city populations, Gede-Pangrango, Salak, and Pebakti-Gagak volcanoes, all located in Indonesia, rank highest in terms of city population exposure within 100 km (Fig. 7), including the entire population of Jakarta for Salak and Gede-Pangrango, shown in the Appendix (Fig. A3). For the top 10 volcanoes with cities within 100 km, up to 35.9% of the population live within 30 km, with most living between 30 and 100 km (Fig. 7). Notably, Gede-Pangrango volcano has the most city

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

populations living within 100 km and Guntur volcano hosts the largest population within a 10-km distance in Indonesia, reaching populations in Bandung and Jakarta. Outside of Indonesia, volcanoes Nejapa-Miraflores, Nicaragua, and the Auckland Volcanic Field have almost 50 % of city populations nearby living within 10 km of the volcanoes. In fact, Auckland Volcanic Field and San Salvador have almost the entire city population exposure within 30 km of the volcanoes and the entire country of El Salvador is within 30 km of a volcano (Fig. 9).

In addition to distances, it is also important to consider the number of cities that are close to volcanoes. Ethiopia is noted for its high number ( $n=40896$ ) of cities close to volcanoes, and the greatest number of cities within the country ( $n=2318$ ) have the Northern Lake Abaya Volcanic Field, Ethiopia, as their nearest volcano, as detailed in the Appendix (Fig. A1). Tangkuban Parahu volcano ranks highly across our rankings of the number of people within 100 km, the number of cities, and the distance to the nearest city (Fig. 3).


Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman



**Fig. 7:** Horizontal stacked bar charts of city population (left) and proportion (right) around each volcano, classified by distance of the city margin to the nearest volcano (within <10, 10 - 30, and 30 - 100-km distances from volcanoes), and ranked by 100, 30, and 10-km radial buffers. Each

Formatted: Font: Times New Roman

bar represents a volcano, and the x axis limits (left) reflect the population range for that volcano buffer.

#### 4.4 Regional trends in city exposure to volcanoes

The largest amount of people living in cities exposed to at least one volcano are in South-East Asia, with ~~6,450,774~~; ~~62,503,491~~; ~~5,847,751~~; ~~52,621,515~~; and ~~462,357,787~~ ~~160,870,875~~ people living within 10, 30, and 100 km of a volcano, respectively (Fig. 8A). However, the highest proportion of city populations exposed to volcanoes are in Central America (Fig. 8B) where approximately ~~66%~~, ~~34.57 %~~, ~~27 %~~ and ~~43 %~~ of city populations are located within 100 km, 30 km, and 10 km, respectively, of at least one volcano. In Australia and New Zealand, and Europe, subregions there are the least number of people exposed, with ~~2,018,698~~ ~~1,660,474~~ and ~~22,942,510~~ ~~77,280~~, respectively, within 100 km. City populations in the subregions of Australia and New Zealand, and Europe, are also exposed to fewer volcanoes than the other continents, with more than 10 % of the exposed populations exposed to one or two volcanoes (Fig. 8B). However, 4 % of the total city population of Australia and New Zealand lies within 10 km of at least one volcano, the highest proportion of any subregion (Fig. 8).

Indonesia stands out globally for having the greatest number of cities and residents located in proximity to volcanoes, with the largest populations within 10, 30, and 100 km of at least one volcano (Fig. 9). Japan ranked second for the 100-km *volcano buffer* but ranks ~~fifth~~ ~~sixth~~ and ~~fourteenth~~ ~~thirteenth~~ for the 30-km and 10-km *volcano buffers* as a high proportion of the city population are living within 30 and 100 km of volcanoes. This is also the case for South Korea, Iran, and Chile, with high numbers of people living within 30 and 100 km of volcanoes compared to <30 km. Meanwhile, the Philippines, ranked third for the 100-km buffer, has a high proportion of its city populations residing within all three distances, meaning that it ranks highly for all three buffer sizes. For the ~~32~~ ~~33 %~~ of city populations exposed in East Africa (Fig. 8), these are dominated by Ethiopia and Yemen (Fig. 9), where the cities are not reflected in the top city rankings of Fig. 4 as the city populations are split across multiple smaller cities. Ethiopia has almost 100 cities exposed, and Yemen has ~~~almost~~ ~~60~~ cities exposed to volcanoes (see Fig. A1 in the Appendix). Remarkably, El Salvador presents a unique case for the top 10 ranked cities, where

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, Not Bold

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman


Formatted: Font: Times New Roman

Formatted: Font: Times New Roman


Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

the entirety of its population lives within 30 km of at least one volcano. Both Nicaragua and El Salvador have over 40 % of their city populations living within 10 km of at least one volcano.



**Fig. 8:** The exposed city population (A) and proportion of total city population (B) split by subregion and classified by the distance of the population to nearest volcano. The subregions are highlighted on the bottom right schematic map showing an approximate location of where the cities are located. Note that cities in French Polynesia are included in Melanesia and Polynesia, and Réunion, Martinique, and Mayotte Islands, and Kamchatka, are included in Europe.



**Fig. 9:** Horizontal stacked bar charts of city population (left) and proportion (right) around each volcano, grouped by country, classified by distance of the city margin to the nearest volcano (within <10, 10 - 30, and 30 - 100-km distances from volcanoes), and ranked by 100, 30, and 10-km radial buffers. Each bar represents a country, and the x axis limits (left) reflect the population range for that volcano buffer .

## 4 Discussion

### 4.1 City Rankings

Our methodology offers a ranking of cities and revised population metrics and distance to the nearest volcano, advancing the polar charts of past studies showing the distance, direction, and population of example cities (Auker et al., 2013; Chester et al., 2000). We updated city populations: for instance, while Chester et al. (2000) reported Manila had a population of 7.94 million and Tokyo had a population of 25 million, our 2020 ~~CE~~ data was ~ 16 million and ~ 5 million higher, respectively. In this study, ~~557~~575 cities exposed to volcanoes have populations greater than 100,000 people, a substantial increase from the 67 cities documented by Heiken


Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

(2013). We identify Jakarta, Tokyo, and Manila, as having the highest exposure within 100 km of volcanoes. These large cities have high proportions of people living between 30 and 100 km from the nearest volcano and population spread across the three *volcano buffers* (Fig. 4; Table 4), highlighting the potential for variable impacts. Jenkins et al. (2018) also identified these three cities as having the greatest tephra fall hazard and risk in Asia, when wind conditions, eruption characteristics, and tephra transport were accounted for. Jakarta is highlighted as the largest city exposed to volcanoes in Pelling (2012) and Blaikie (1994). However, Heiken (2013) identifies Tokyo, Manila, and Mexico City as the only megacities exposed to volcanoes. Jakarta is not selected as an example by Heiken (2013), Auker et al. (2013), or Chester et al. (2000). This may be due to potential differences in criteria used in assessing exposure.



**Fig. 10:** Polar chart to show the top 10 cities with the most exposed city populations within 100 km of the nearest volcano. The city centroids (left) and nearest city margin to the volcano (right) are plotted as diamonds relative to the nearest volcano (centre of the plots), and the diamond size refers to the size of the population exposed. The centroid of Seoul is located outside of the 100-km radial buffer, but the population within the 100 km is high enough to rank the city ~~into~~ the top 10 exposed.

Identifying an exposed city by its centroid alone is not an ideal representation of city exposure as it does not capture the spatial extent (sprawl) of a city, or changes in population density across the city. City margins are closer to volcanoes than the centroids (Fig. 10). Some cities have higher population densities closer to volcanoes despite having greater areas of the city farther away (Table 4), suggesting that cities can have higher population densities in the city peripheries away from

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman



the city centre. For example, the city centroid of Seoul is located more than 100 km from Mount Baekdu (Changbaishan), however approximately half of Seoul's city population ( $n=11,705,767$ ), and some of the most densely populated parts of the city ( $\sim 11,419,362$  people per  $\text{km}^2$ ) are located within the 100-km volcano buffer (Fig. 10), with the closest margin of the city 64 km from the volcano. This is likely due to residential zones being situated farther from the city centre, in areas where population density may increase as cities expand. These close city margins place dense populations very close to potential volcanic hazards. Therefore, our approach better captures the spatial sprawl and variation in population across the cities, which are overlooked by the centroid approach.

**Table 4:** Cities with populations across three different buffer distances of volcanoes: 10, 10 – 30, and 30 – 100 km, with city population exposed, area of city polygon, and the averaged population density within each buffer distance. The population, area, and densities in bold font are those higher than the other two buffer distances. The cities are in order of population density < 10 km of volcanoes. Numbers are rounded to the nearest integer.

City Name	Population			Area ( $\text{km}^2$ )			Population Density ( $/\text{km}^2$ )		
	10 km	10 - 30 km	30 - 100 km	10 km	10 - 30 km	30 - 100 km	10 km	30 - 100 km	30 - 100 km
Taipei, Taiwan	457,807	<b>6,633,853</b>	2,181,179	64	512	374	7,116	<b>12,965</b>	5,836
Jakarta, Indonesia	453,914	6,559,219	<b>31,037,351</b>	110	1,123	<b>3,258</b>	4,142	5,838	<b>9,520</b>
Surabaya, Indonesia	387,688	2,927,379	<b>3,471,151</b>	102	<b>769</b>	429	3,805	3,806	<b>8,088</b>
Manila, Philippines	220,952	1,399,747	<b>22,753,383</b>	48	215	<b>2,227</b>	4,636	<b>6,499</b>	629
Tasikmalaya, Indonesia	167,908	<b>2,182,277</b>	306,365	70	<b>403</b>	49	2,393	5,420	<b>6,246</b>
Guatemala City, Guatemala	23,943	<b>2,020,662</b>	694,688	6	<b>326</b>	120	3,705	<b>6,192</b>	5,776

Rome, Italy	16,698	<b>2,160,308</b>	279,120	5	<b>399</b>	63	3,206	<b>5,418</b>	4,430
Mexico City, Mexico	2,373	5,987,736	<b>11,607,299</b>	2	733	<b>1,391</b>	1,052	8,170	<b>8,348</b>
Tokyo, Japan	21	558,703	<b>29,275,558</b>	<1	170	<b>4,229</b>	290	3,284	<b>6,924</b>

In this research we focussed on identifying cities with high populations exposed to volcanic hazards. However, we emphasise the importance of assessing both the total numbers of exposure as well as the proportions of the city (Fig. 4B), as less populated and smaller cities may have their entire population exposed. While these cities may not rank highly in terms of total population, smaller cities may be more susceptible to complete destruction (Pelling, 2012), whereas an eruption affecting part of a larger city may allow for greater capacity to recover. Limited resources and less infrastructure in smaller cities may also magnify the impacts of eruptions. Thus, eruptions affecting an entire city potentially cause challenges in ~~ter~~terms of evacuation, continuity, and recovery. An eruption that affects multiple neighbouring cities can compound this effect. If the eruption affects multiple small cities instead of one large city, it may result in a high total number of exposed individuals dispersed across multiple cities. This scenario can strain emergency response efforts and limit the options for evacuation and recovery. We accounted for this in our composite ranking by including ranks by the number of nearby cities (Fig. 3).

Conversely, an eruption affecting part of a large populous city can have implications for the entire city, neighbouring cities, the country and potentially globally. In large urban areas, the interdependence of infrastructure systems, such as transportation and utilities, means that even minor damage can lead to widespread functional disruptions, affecting populations beyond the immediate impacted area (Pelling, 2012; Heiken, 2013; Weir et al., 2024). For example, lahars triggered following the 1991 Pinatubo eruption in the Philippines (VEI 5) damaged highways to the North of Metro Manila, resulting in cascading impacts that disrupted access to the city (Solway, 1994; Pelling, 2012). These indirect impacts can extend beyond infrastructure disruption, affecting supply chains, labour markets, and public health systems, with potential for the effects of an eruption to impact regionally or globally, particularly for those cities that are central to financial systems (Pelling, 2012). Thus, these large cities that have a small proportion exposed may be

Formatted: Font: Times New Roman, Highlight

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, 11 pt, English (United Kingdom)

Formatted: Font: Times New Roman, Highlight

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

ranked low in our analysis in terms of exposure, but this could mask potential widespread indirect impacts across the broader city (Mossler, 1996; Pelling, 2012). Understanding these wider effects is crucial for assessing the full scope of urban vulnerability to volcanic hazards. Further localised assessments and systematic risk evaluations are recommended to capture a more complete range of potential impacts (Mossler, 1996; Pelling, 2012).

#### 4.2 Volcano Rankings

We highlight key volcanoes such as Salak, Gede-Pangrango, and Perbakti-Gagak with the most people exposed within cities nearby, as well as Tangkuban Parahu and Tampomas, Indonesia, and San Pablo Volcanic Field, Philippines, that rank highly in our composite ranking. Other studies highlight volcanoes in Indonesia as having high levels of exposure. For example, in our study, Gede-Pangrango was ranked ~~second~~first and seventh for populations living within 100 km and 30 km of volcanoes, respectively. Gede-Pangrango was also ranked by Small and Naumann (2001) as the most populous volcano globally and by Jenkins et al. (2022), as the fifth highest volcano for exposure in Indonesia and the Philippines when considering both populations and physical assets threatened by VEI 3-5 eruptions. Jenkins et al. (2022) also identified Guntur volcano as a volcano in need of further study due to its resulting exposure and the lack of localised hazard or exposure assessments. Our findings support this, as Guntur volcano ranks highest in Indonesia for city populations within 10 km of the volcano and ranks fourth in the composite ranking. Our findings highlight key hotspot volcanoes for further localised exposure and hazard assessments.

Large city populations within 10 km of volcanoes, a distance reached by destructive proximal hazards, can be affected by a smaller or moderate eruption at one of these volcanoes, which may have greater direct impacts than a larger eruption where populations are located farther away. A large proportion of these close populations are residing near volcanoes in Central and South America (Fig. 4; Fig. 9). Escobar et al. (2007) found that a high proportion of city populations in Central America are living close to the most dangerous volcanoes in the region, something supported by our study (Fig. 7). Ewert and Harpel (2004) highlight Central America as having potentially significant exposure to volcanic hazards, with 2.7 million people (data from 2004 CE) within 10 km of

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, Highlight

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, Highlight

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

volcanoes living in both urban and rural areas. In our study, using 2020 ~~CE~~ city population data for the same buffer distance, we find that a similar number (2.6 million) of people now live just within ~~the~~ urban (city) areas. The high proportions of urban populations close to volcanoes in Central and South America (Fig. 9) is perhaps linked to the colonial past of the region, whereby investments into agriculture in the early to mid-20<sup>th</sup> Century resulted in a growth of agricultural settlements through time (Swyngedouw, 2006). Nejapa-Miraflores volcano, Nicaragua is one example of where almost 1 million people, or ~~60~~45 % of the exposed population, are located within 10 km of the volcano. These populations are distributed over multiple cities, adding complexity to potential eruption impacts and response strategies.

#### 4.4 Future Research Directions

In this study, we define cities as those areas classified in the GHS-UCDB dataset as *urban centres*, which excludes suburbs and regions of lower population densities. For example, the remaining ~~~25~~46 % (n=~~336~~686) of volcanoes that do not have city populations within 100 km, may still be situated near suburban or rural populations, or have tourists, not assessed in this study. ~~Through time, these~~ The delineation between urban, peri-urban, and rural areas is not always clear (Chen et al., 2018; Sorensen and Labbé, 2020). Urban centres can sprawl into surrounding peri-urban and rural regions, and through time, urban centres may merge together and densify. Therefore, total city populations may slightly differ when using other datasets. If suburban and rural populations were also considered, the ranking of volcanoes would likely change, as these areas may have higher exposure to volcanic hazards. Past volcanic eruptions such as 1982 El Chichon eruption, Mexico, have primarily affected rural communities rather than city centres. Our approach could be expanded or used to explore more than just cities, by assessing changing land-use patterns, transient populations, or other using different population densities datasets (e.g., LandScan ambient population dataset: Lebakula et al., 2024), or gradations of population density around volcanoes using different buffer distances. We can also explore past trends in city population exposure through time, and future projections, to quantify rates of exposure change. Mapping the urban sprawl of these cities allows us to identify key areas around volcanoes to focus on future mitigation efforts and land-use planning. Further research is needed to understand how the interconnectedness of cities and suburbs influences impact.

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, Highlight

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, English (United Kingdom)

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, English (United Kingdom)

632 Additionally, our 2020-~~CE~~ dataset includes 137 cities exposed to volcanoes located in other  
633 countries, underscoring the need for research in cross-border cooperation and planning to mitigate  
634 the impacts of eruptions with transboundary effects (Donovan & Oppenheimer, 2019).

Formatted: Font: Times New Roman  
Formatted: Font: Times New Roman  
Formatted: Font: Times New Roman  
Formatted: Font: Times New Roman

637 We used a comparative approach to quantify the hazard by using the distance to the nearest volcano  
638 and the number of nearby volcanoes, which could be explored further. For example, the volcano  
639 buffers used in this study relate to the average maximum distances of primary hazards. However,  
640 our use of a 100-km radial buffer does not account for far-reaching volcanic flows or tephra falls  
641 that may reach beyond 100 km (Biass et al., 2024) or cascading hazards that may extend beyond  
642 this distance, such as tsunamis. Future studies could explore these cascading and widespread  
643 impacts, in the key cities identified in this study, to capture potential losses.

644 The selection of the location of the volcanic crater as point ~~co-ordinates~~coordinates provided by  
645 the Global Volcanism Program (2023), does not account for the uncertainty in the next eruption  
646 site, for example ~~at~~within volcanic fields or rift zones. ~~Future work could classify volcanoes by last~~  
647 ~~eruption, VEI range, or tectonic setting, to better understand the specific types of potential hazard~~  
648 ~~they pose to nearby cities.~~ Additionally, research could explore alternative methods to the  
649 traditional volcano buffers, considering approaches that account for the spatial variability of  
650 ~~volcanic fields and shield volcanoes. There is also necessity in localised exposure and risk~~  
651 ~~assessment~~distributed volcanism, such as those used in Nieto-Torres et al. (2021), and of shield  
652 ~~volcanoes. Future work could classify volcanoes by probability of eruption, last eruption, VEI~~  
653 ~~range, or tectonic setting, to better understand the specific types of potential hazard they pose to~~  
654 ~~nearby cities. For example, the presence of multiple volcanoes within 100 km, as in San Salvador,~~  
655 ~~does not directly equate to higher hazard, as eruption probabilities vary. While our approach~~  
656 ~~identifies cities with high volcanic exposure, localised hazard assessments are essential for a more~~  
657 ~~precise evaluation of the threat~~ (Biass et al., 2024; Jenkins et al., 2022; Diefenbach et al., 2015).  
658 Future research can incorporate local topography, seasonal weather patterns, eruption frequencies,  
659 and hazard probabilities as well as exposures beyond population, and the vulnerability of these to  
660 the hazards to understand the specific risks faced by individual cities.  
661

Formatted: Font: Times New Roman, Highlight  
Formatted: Font: Times New Roman  
Formatted: Font: Times New Roman  
Formatted: Font: Times New Roman  
Formatted: Font: Times New Roman  
Formatted: Font: Times New Roman  
Formatted: Font: Times New Roman  
Formatted: Font: Times New Roman  
Formatted: Font: Times New Roman

## 5 Conclusions

Cities close to volcanoes are at high risk of volcanic hazards. ~~In our study, we~~ We present a framework to quantify and rank global city population exposure to volcanoes by extracting and calculating 2020 CE population counts from city polygons. We evaluated the exposure of 1,106 cities within 100 km of 596 Holocene volcanoes, measuring exposure within 10, 30, and 100 km radial buffers, reflecting maximum distances for various eruption VEIs. Although topography and weather can affect hazard footprints (Biaass et al., 2024), our method using concentric radial buffers conservatively estimates and compares city exposure. ~~These buffer sizes may overestimate exposure, but we preferred this approach to avoid underestimating it. Our results are provided at Section 9 and as a web application for visualisation of all city exposures~~ ([https://vharg.github.io/shiny\\_VolcCities/](https://vharg.github.io/shiny_VolcCities/)). The rankings ~~can~~ of 1,133 cities within 100 km of 578 Holocene volcanoes provide a foundation to identify areas for future detailed and localised exposure or risk assessments, especially for cities with limited past hazard or exposure data. Ranking cities by exposure also helps identify key locations for future research and land-use planning. ~~Our results are provided at Section 9 and as a web application for visualisation of all city exposures~~ (<https://nteng.shinyapps.io/ExposureOfCities/>). ~~Using this framework, we ranked cities by exposed population, number of nearby volcanoes, and distance to the nearest volcano.~~ We highlight Jakarta, Bandung, and San Salvador as scoring highly across these rankings. Jakarta, Bandung, and Naples have the largest city populations within 100, 30, and 10 km, respectively, of at least one volcano. San Salvador, Guatemala City, and Managua are cities of over 1 million people that have the largest number of people exposed to the largest number of volcanoes within 100, 30 and 10 km respectively with 22, 523, 6, and 2 volcanoes, respectively. We also ranked volcanoes in three ways: by number of exposed city populations, by the number of nearby cities, and the distance of the nearest city. We highlight Tangkuban Parahu, Tampomas, and San Pablo Volcanic Field, and Tampomas as scoring highly across these rankings. Gede-Pangrango, ~~Laguna Caldera~~ Tangkuban Parahu, and Nejapa-Miraflores volcanoes have the largest number of city populations within 100, 30, and 10 km respectively. These rankings reveal hotspot cities with high populations exposed to multiple volcanoes; for example, ~8.5 million people in Bandung are exposed within 100 km to 12 volcanoes.

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, English (United Kingdom)

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Globally, 50% of people exposed to volcanoes (within 100 km) live in cities. The size, number, and distance of cities near volcanoes, or spread of population density across cities, create different challenges regarding exposure to volcanoes. For example, Jakarta has high population density between 30 – 100 km of volcanoes, while 2339 smaller cities in Ethiopia are exposed to a single volcano: Northern Lake Abaya Volcanic Field. Other cities, such as Auckland, are located on volcanic fields. Some countries are highly exposed to volcanic hazards. For example, all cities in El Salvador are located within 30 km of a volcano and in the Philippines, 90-80% of cities have some part of the city located within 100 km of a volcano. Understanding these diverse exposures is crucial for developing effective risk management strategies tailored to the specific needs of each country.

~~This research highlights key exposed cities at risk of volcanic hazards ranked in different ways.~~ These findings can inform decision-making and further research around volcanoes. This work prompts more localised studies that overlay these exposures with probabilistic hazard maps which can enhance our understanding of the dynamic risks surrounding volcanoes.

## 8 Appendix A




Fig. A1: Horizontal stacked bar charts of number of cities with populations within each buffer distance (left) and proportion (right) around each volcano, grouped by country, classified by distance of the city margin to the nearest volcano (within <10, 10 - 30, and 30 - 100-km distances from volcanoes), and ranked by 100, 30, and 10-km radial buffers. If the city is spread across multiple buffers, the closest to the volcano is selected. Each bar represents a country, and the x axis limits (left) reflect the population range for that volcano buffer.




Fig. A2: Top 20 cities with less than one million exposed city populations below one million, coloured by the number of volcanoes for which the population is exposed by city lies within a 100 km radius.






Fig. A3: City populations exposed to individual volcanoes within 100 km, ranked by the dominant city. Bars are coloured by the dominant city exposed population. *V.F. stands for Volcanic Field, V.G. stands for Volcanic Group, and V.C. stands for Volcanic Complex.*

**Formatted:** Font: Times New Roman, Italic

**Formatted:** Font: Times New Roman, Italic

**Formatted:** Font: Times New Roman




Fig. A4: Top forty countries with a total area of  $>2,000 \text{ km}^2$ , ranked in order of their total land area exposed to volcanic hazards, and coloured by the percentage of area. Is stands for islands and Fr stands for French. *Land area from Natural Earth (2023)*.

#### 8 Code Availability

The R code used in this manuscript is available at: <https://github.com/vharg/VolcCities>.

#### 9 Data Availability

The web application presenting results for all cities is available at

<https://nteng.shinyapps.io/ExposureOfCities/> [https://vharg.github.io/shiny\\_VolcCities/](https://vharg.github.io/shiny_VolcCities/).

Ranking results of city exposure to volcanic hazards and volcanoes by exposed populations is available at <https://researchdata.ntu.edu.sg/privateurl.xhtml?token=65c3712c-461b-4d95-b499-f2eebb5ab30d>.

#### 10 Author Contribution

The project was conceptualised by ESM, RXNT, SJF and SB. The methodology was developed by ESM and RXNT with support from SFJ, JLH, and SB. Visualisations were made by ESM and the web app was developed by RXNT. The original draft was written by ESM. All authors contributed to editing and reviewing the manuscript.

Formatted: Font: Times New Roman, Italic

Formatted: Font: Times New Roman, Italic

Formatted: Font: Times New Roman, Italic

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

750 11 Competing Interests

751 The authors declare that they have no conflict of interest.

752

753 12 Acknowledgements

754 We kindly thank the helpful reviews from Amiel Nieto Torres and Luis E. Lara that enhanced the  
755 manuscript. We thank editor Giovanni Macedonio for overseeing the review process. This research  
756 was partly supported by the Ministry of Education, Singapore, under its MOE Academic Research  
757 Fund Tier 3 InVEST project (Award MOE-MOET32021-0002). This work comprises EOS  
758 contribution number 631. SFJ acknowledges financial support from the AXA Joint Research  
759 Initiative. We acknowledge support for this work from the Centre for Disaster Resilience, ITC  
760 Faculty, University of Twente.

761

762 13 References

763

764 Alberico, I., Petrosino, P., & Lirer, L. (2011). Volcanic hazard and risk assessment in a multi-  
765 source volcanic area: The example of Napoli city (Southern Italy). Nat. Hazards Earth Syst.  
766 Sci., 11(4), 1057–1070. <https://doi.org/10.5194/nhess-11-1057-2011>

767 Ariyanti, V., Gaafar, T., De La Sala, S., Edelenbos, J., & Scholten, P. (2020). Towards liveable  
768 volcanic cities: A look at the governance of lahars in Yogyakarta, Indonesia, and  
769 Latacunga, Ecuador. *Cities*, 107, 102893. <https://doi.org/10.1016/j.cities.2020.102893>

770 Aspinall, W., Auker, M., Hincks, T., Mahony, S., Nadim, F., Pooley, J., Sparks, R. S. J., & Syre,  
771 E. (2011). Volcano hazard and exposure in GFDRR priority countries and risk mitigation  
772 measures—GFDRR Volcano Risk Study.

773 Auker, M. R., Sparks, R. S. J., Siebert, L., Croweller, H. S., & Ewert, J. (2013). A statistical  
774 analysis of the global historical volcanic fatalities record. *J. Appl. Volcanol.*, 2(1), 2.  
775 <https://doi.org/10.1186/2191-5040-2-2>

776 Biass, S., Jenkins, S. F., Hayes, J. L., Williams, G. T., Meredith, E. S., Tennant, E., Yang, Q.,  
777 Lerner, G. A., Burgos, V., Syarifuddin, M., & Verolino, A. (2024). How well do concentric  
778 radii approximate population exposure to volcanic hazards? *Bull. Volcanol.*, 86(1).  
779 <https://doi.org/10.1007/s00445-023-01686-5>

780 Bilham, R. (2009). The seismic future of cities. *Bull. Earthq. Eng.*, 7(4), 839–887.  
781 <https://doi.org/10.1007/s10518-009-9147-0>

782 Blaikie, P., Cannon, T., Davis, I., & Wisner, B. (2004). *At risk: Natural hazards, people's*  
783 *vulnerability and disasters*. Routledge.

784 Brecht, H., Deichmann, U., & Wang, H. G. (2013). A global urban risk index. [World Bank  
785 Working Paper]. <http://econ.worldbank.org>

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, No underline,  
Font color: Auto

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

786 Brown, S. K., Jenkins, S. F., Sparks, R. S. J., Odbert, H., & Auken, M. R. (2017). Volcanic fatalities  
787 database: Analysis of volcanic threat with distance and victim classification. *J. Appl.*  
788 *Volcanol.*, 6(1), 2. <https://doi.org/10.1186/s13617-017-0067-4>.

789 Chen, C., LeGates, R., Zhao, M., & Fang, C. (2018). The changing rural-urban divide in China's  
790 megacities. *Cities*, 81, 81-90. <https://doi.org/10.1016/j.cities.2018.03.017>

791 Chester, D. K., Degg, M., Duncan, A. M., & Guest, J. E. (2000). The increasing exposure of cities  
792 to the effects of volcanic eruptions: A global survey. *Environ. Hazards*, 2(3), 89–103.  
793 <https://doi.org/10.3763/ehaz.2000.0214>.

794 Degg, M. (1992). Natural disasters: Recent trends and future prospects. *Geography*, 77(3), 198–  
795 209.

796 Del Negro, C., Cappello, A., Bilotta, G., Ganci, G., Hérault, A., & Zago, V. (2020). Living at the  
797 edge of an active volcano: Risk from lava flows on Mt. Etna. *Bull. Geol. Soc. Am.*, 132(7–  
798 8), 1615–1625. <https://doi.org/10.1130/B35290.1>.

799 Delgado Granados, H., & Jenkins, S. F. (2015). Extreme volcanic risks 1: Mexico City. In J.  
800 Shroder (Ed.), *Volcanic hazards, risks and disasters* (pp. 315–354). Elsevier.

801 Diefenbach, A. K., Wood, N. J., & Ewert, J. W. (2015). Variations in community exposure to lahar  
802 hazards from multiple volcanoes in Washington State (USA). *J. Appl. Volcanol.*, 4(1), 4.  
803 <https://doi.org/10.1186/s13617-015-0024-z>.

804 Donovan, A., & Oppenheimer, C. (2014). Extreme volcanism: disaster risk and societal  
805 implications. *Extreme natural hazards: Disaster risks and societal implications*, 29-46.

806 Donovan, A., & Oppenheimer, C. (2019). Volcanoes on borders: A scientific and (geo)political  
807 challenge. *Bull. Volcanol.*, 81(5), 28. <https://doi.org/10.1007/s00445-019-1291-z>

808 Durand, M. (2001). Impacts of, and responses to ashfall in Kagoshima from Sakurajima volcano:  
809 Lessons for New Zealand. *Inst. Geol. Nucl. Sci. Ltd., New Zealand*.

810 Erfurt-Cooper, P. (2014). Volcanic tourist destinations. Springer. [https://doi.org/10.1007/978-3-](https://doi.org/10.1007/978-3-642-16191-0)  
811 [642-16191-0](https://doi.org/10.1007/978-3-642-16191-0)

812 Escobar, R., Alvarado, G., Soto, G., Navarro, M., Escobar, D., Pullinger, C., & Bonis, S. (2007).  
813 Volcanic activity, hazards, and monitoring. In *Central America* (pp. 837–880). Taylor &  
814 Francis. <https://doi.org/10.1201/9780203947043.ch38>.

815 Ewert, J. W. (2007). System for ranking relative threats of US volcanoes. *Nat. Hazards Rev.*, 8(4),  
816 112–124.

817 Ewert, J. W., Diefenbach, A. K., & Ramsey, D. W. (2018). 2018 Update to the U.S. Geological  
818 Survey National Volcanic Threat Assessment. *Sci. Invest. Rep.*, 2018-5140.

819 Ewert, J. W., & Harpel, C. J. (2004). In harm's way: Population and volcanic risk. *Geotimes*, 49(4),  
820 14–17.

821 Freire, S., Florczyk, A. J., Pesaresi, M., & Sliuzas, R. (2019). An improved global analysis of  
822 population distribution in proximity to active volcanoes, 1975–2015. *ISPRS Int. J. Geo-*  
823 *Inf.*, 8(8), 341. <https://doi.org/10.3390/ijgi8080341>.

824 Global Volcanism Program. (2024). Volcanoes of the World (v. 5.2.0; 6 Jun 2024). Smithsonian  
825 Institution. <https://doi.org/10.5479/si.GVP.VOTW5-2024.5.2>.

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, 11 pt, English  
(United Kingdom)

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

- Godschalk, D. R. (2003). Urban hazard mitigation: Creating resilient cities. *Nat. Hazards Rev.*, 4(3), 136–143. <https://doi.org/10.1061/ASCE1527-698820034:3136>
- Gu, D. (2019). Population exposure and vulnerability to natural disasters for world's cities. United Nations Populations Division Working Paper. [www.unpopulation.org](http://www.unpopulation.org).
- Guimarães, L. F., Nieto-Torres, A., Bonadonna, C., & Frischknecht, C. (2021). A new inclusive Volcanic Risk Ranking, Part 2: Application to Latin America. *Frontiers in Earth Science*, 9, 936. <https://doi.org/10.3389/feart.2021.757742>
- Hanson, S., Nicholls, R., Ranger, N., Hallegatte, S., Corfee-Morlot, J., Herweijer, C., & Chateau, J. (2011). A global ranking of port cities with high exposure to climate extremes. *Clim. Change*, 104(1), 89–111. <https://doi.org/10.1007/s10584-010-9977-4>
- Heiken, G. (2013). *Dangerous neighbors: Volcanoes and cities*. Cambridge Univ. Press.
- Iglesias, V., Braswell, A. E., Rossi, M. W., Joseph, M. B., McShane, C., Cattau, M., Koontz, M. J., McGlinchy, J., Nagy, R. C., Balch, J., Leyk, S., & Travis, W. R. (2021). Risky development: Increasing exposure to natural hazards in the United States. *Earth's Future*, 9(7).
- Jenkins, S. F., Biass, S., Williams, G. T., Hayes, J. L., Tennant, E., Yang, Q., Burgos, V., Meredith, E. S., Lerner, G. A., Syarifuddin, M., & Verolino, A. (2022). Evaluating and ranking Southeast Asia's exposure to explosive volcanic hazards. *Nat. Hazards Earth Syst. Sci.*, 22(4), 1233–1265. <https://doi.org/10.5194/nhess-22-1233-2022>
- Jenkins, S. F., Magill, C. R., & Blong, R. J. (2018). Evaluating relative tephra fall hazard and risk in the Asia-Pacific region. *Geosphere*, 14(2), 492–509. <https://doi.org/10.1130/GES01549.1>
- Lebakula, V., Epting, J., Moehl, J., Stipek, C., Adams, D., Reith, A., Kaufman, J., Gonzales, J., Reynolds, B., Basford, S., Martin, A., Buck, W., Faxon, A., Cunningham, A., Roy, A., Barbose, Z., Massaro, J., Walters, S., Woody, C., ... Urban, M. (2024). *LandScan Silver Edition [Data set]*. Oak Ridge National Laboratory. <https://doi.org/10.48690/1531770>
- Levine, D. J. (2022). exactextractr: Fast Extraction from Raster Datasets using Polygons (R package version 0.8.3). Available at: <https://CRAN.R-project.org/package=exactextractr> [Accessed: January 2024].
- Loughlin, S. C., Sparks, S., Brown, S. K., Jenkins, S. F., & Vye-Brown, C. (2015). *Global volcanic hazards and risk*. Global Volcanic Hazards and Risk. Cambridge University Press. <https://doi.org/10.1017/CBO9781316276273>
- Magill, C., & Blong, R. (2005). Volcanic risk ranking for Auckland, New Zealand. II: Hazard consequences and risk calculation. *Bull. Volcanol.*, 67(4), 340–349. <https://doi.org/10.1007/s00445-004-0375-5>
- Mani, L., Tzachor, A., & Cole, P. (2021). Global catastrophic risk from lower magnitude volcanic eruptions. *Nat. Commun.*, 12(1). <https://doi.org/10.1038/s41467-021-25021-8>
- Mangan, B. M., Ball, J., Wood, N., Jones, J. L., Peters, J., Abdollahian, N., Dinitz, L., Blankenheim, S., Fenton, J., & Pridmore, C. (2018). SIR 2018–5159 ver. 1.1: California's Exposure to Volcanic Hazards.

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

- Mari Rivero, I., Melchiorri, M., Florio, P., Schiavina, M., Krasnodębska, K., Politis, P., Uhl, J., Pesaresi, M., Maffeni, L., Sulis, P., Crippa, M., Guizzardi, D., Pisoni, E., Belis, C., Oom, D., Branco, A., Mwaniki, D., Kochulem, E., Githira, D., Carioli, A., Ehrlich, D., Tommasi, P., Kemper, T., & Dijkstra, L. (2024). Joint Research Centre (JRC) [Dataset]. <https://doi.org/10.2905/1A338BE6-7EAF-480C-9664-3A8ADE88CBCE>
- Mossler, M. (1996). Environmental hazard analysis and small island states: rethinking academic approaches. *Geogr. Zeitsch.*, 1, 86–93.
- Natural Earth. (2023). Admin 0 – Countries. 1:10m, 1:50m, and 1:110m scales. Available at: <https://www.naturalearthdata.com/downloads/10m-cultural-vectors/10m-admin-0-countries/> [Accessed: January 2024].
- Newhall, C., Self, S., & Robock, A. (2018). Anticipating future Volcanic Explosivity Index (VEI) 7 eruptions and their chilling impacts. *Geosphere*, 14(2), 572–603. <https://doi.org/10.1130/GES01513.1>
- Nieto-Torres, A., Guimarães, L. F., Bonadonna, C., & Frischknecht, C. (2021). A New Inclusive Volcanic Risk Ranking, Part 1: Methodology. *Front. Earth Sci.*, 9. <https://doi.org/10.3389/feart.2021.697451>
- Pelling, M. (2012). *The Vulnerability of Cities: Natural Disasters and Social Resilience*.
- Pesaresi, M., Florczyk, A., Schiavina, M., Melchiorri, M., Maffeni, L. (2019). GHS settlement grid, updated and refined REGIO model 2014 in application to GHS-BUILT R2018A and GHS-POP R2019A, multitemporal (1975-1990-2000-2015), R2019A. European Commission, Joint Research Centre (JRC). DOI: 10.2905/42E8BE89-54FF-464E-BE7B-BF9E64DA5218 PID: <http://data.europa.eu/89h/42e8be89-54ff-464e-be7b-bf9e64da5218>
- Sandri, L., Thouret, J. C., Constantinescu, R., Biass, S., & Tonini, R. (2014). Long-term multi-hazard assessment for El Misti volcano (Peru). *Bull. Volcanol.*, 76(2), 1–26. <https://doi.org/10.1007/s00445-013-0771-9>
- Schiavina, M., Melchiorri, M., & Freire, S. (2023a). GHS-DUC R2023A - GHS Degree of Urbanisation Classification: Application of the Degree of Urbanisation methodology (stage II) to GADM 4.1 layer, multitemporal (1975–2030) [Data set]. European Commission, Joint Research Centre (JRC). <https://doi.org/10.2905/DC0EB21D-472C-4F5A-8846-823C50836305>
- Schiavina, M., Freire, S., Carioli, A., & MacManus, K. (2023b). GHS-POP R2023A - GHS population grid multitemporal (1975-2030). In European Commission, Joint Research Centre (JRC) [Dataset] doi: 10.2905/2FF68A52-5B5B-4A22-8F40-C41DA8332CFE PID: <http://data.europa.eu/89h/2ff68a52-5b5b-4a22-8f40-c41da8332cfe>
- Schiavina, M., Melchiorri, M., & Freire, S. (2021). GHS-DUC R2019A - GHS Degree of Urbanisation Classification (2015, 2000, 1990, 1975), R2019A - OBSOLETE RELEASE. In European Commission, Joint Research Centre (JRC) [Dataset] doi: 10.2905/ED8E8E11-62C3-4895-A7B9-5EF851F112ED PID: <http://data.europa.eu/89h/ed8e8e11-62c3-4895-a7b9-5ef851f112ed>

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, 11 pt

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

- Small, C., & Naumann, T. (2001). The global distribution of human population and recent volcanism. *Environ. Hazards*, 3, <http://www.LDEO.columbia.edu/Bsmall/PopVol.html>.
- Solway, L. (1994). Urban developments and megacities: vulnerability to natural disasters. *Disaster Manag.-Redhill*, 6, 160.
- Sorensen, A., & Labbé, D. (2020). *Megacities, megacity-regions, and the endgame of urbanization. In Handbook of megacities and megacity-regions (pp. 1-19). Edward Elgar Publishing.*
- Strader, S. M., Ashley, W., & Walker, J. (2015). Changes in volcanic hazard exposure in the Northwest USA from 1940 to 2100. *Nat. Hazards*, 77(2), 1365–1392. <https://doi.org/10.1007/s11069-015-1658-1>.
- Swyngedouw, E. (2006). Human Development Report 2006 Power, Water and Money: Exploring the Nexus United Nations Development Programme Background Paper Power, Water and Money: Exploring the Nexus.
- Thouret, J.-C. (1999). Urban hazards and risks; consequences of earthquakes and volcanic eruptions: an introduction. *GeoJournal*, 49(2).
- Thouret, J.-C., Finizola, A., Fornari, M., Legeley-Padovani, A., Suni, J., & Frechen, M. (2001). Geology of El Misti volcano near the city of Arequipa, Peru. *Geology*, 113(12). [https://doi.org/10.1130/0016-7606\(2001\)1132.0.CO;2](https://doi.org/10.1130/0016-7606(2001)1132.0.CO;2).
- Torres, A. N., Del Pozzo, A. L. M., Gropelli, G., & Viera, M. D. C. J. (2023). Risk scenarios for a future eruption in the Chichinautzin monogenetic volcanic field, South México City. *Journal of Volcanology and Geothermal Research*, 433, 107733. <https://doi.org/10.1016/j.jvolgeores.2022.107733>
- UNDP. (2021). Urban Risk Management and Resilience Strategy. United Nations Development Programme. [www.undp.org](http://www.undp.org).
- United Nations (1989) "Prospects of world urbanization, 1988", *Population Studies No. 112, (ST/ESA/SER.A/112)*, New York: United Nations, Department of International Economic and Social Affairs
- Weir, A. M. (2024). Quantifying systemic vulnerability of interdependent critical infrastructure networks: A case study for volcanic hazards. <https://orcid.org/0000-0001-7449-016X>.
- Wood, N., & Soular, C. (2009). Variations in population exposure and sensitivity to lahar hazards from Mount Rainier, Washington. *J. Volcanol. Geotherm. Res.*, 188(4), 367–378. <https://doi.org/10.1016/j.jvolgeores.2009.09.019>.
- World Bank. (2023). Urban Development. Washington, DC: World Bank. Available at: <https://data.worldbank.org/topic/urban-development> [Accessed: November 21, 2024].

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, 11 pt

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, 11 pt

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman