

# Demonstrating the use of UNSEEN climate data for hydrological applications: case studies for extreme floods and droughts in England

Kay, Alison L<sup>1</sup>, Dunstone, Nick<sup>2</sup>, Kay, Gillian<sup>2</sup>, Bell, Victoria A<sup>1</sup>, Hannaford, Jamie<sup>1</sup>

<sup>1</sup>UK Centre for Ecology & Hydrology, Wallingford, UK, OX10 8BB

<sup>2</sup>Met Office, Exeter, UK, EX1 3PB

*Correspondence to:* A Kay ([alkay@ceh.ac.uk](mailto:alkay@ceh.ac.uk))

**Abstract.** Meteorological and hydrological hazards present challenges to people and ecosystems worldwide, but the limited length of observational data means that the possible extreme range is not fully understood. Here, a large ensemble of climate model data is combined with a simple grid-based hydrological model, to assess unprecedented but plausible hydrological extremes in the current climate across England. Two case studies are selected—dry (Summer 2022) and wet (Autumn 2023)—with the hydrological model initialised from known conditions then run forward for several months using the large climate ensemble. The modelling chain provides a large set of plausible events including extremes outside the range from use of observed data, with the lowest flows around 28% lower on average for the Summer 2022 drought study and the highest flows around 42% higher on average for the Autumn 2023 flood study. The temporal evolution and spatial dependence of extremes is investigated, including the potential time-scale of recovery of flows to normal and the chance of persistent extremes. Being able to plan for such events could help improve the resilience of water supply systems to drought, and improve flood risk management and incident response.

**Keywords.** River flows, extreme events, floods, droughts, hydrological modelling

## 1 Introduction

Meteorological and hydrological hazards—storms, floods and droughts—present challenges to people, infrastructure and ecosystems globally (Beevers et al. 2022; Mokhtari et al. 2023). The relatively limited period of available observational data means that the possible range of extremes of such events is often not fully understood. For example, Thompson et al. (2017) show that in SE England there is a 7% chance of exceeding the current observed rainfall record in at least one month in any given winter, with a 34% chance of breaking a regional record somewhere in England and Wales. Similarly, Kent et al. (2022) investigate plausible summer rainfall extremes, showing an approximately 1% chance per year of exceeding current daily rainfall records in the UK in the current climate. Chan et al. (2023b) show an approximately 9% chance of a summer month with lower rainfall than the observed driest summer in SE England in the current climate. When such unprecedented events do inevitably occur, they can lead to very severe impacts (Bertola et al. 2023). This is not just because of their unprecedented magnitude but due to inherent unpreparedness, given that water supply systems, flood infrastructure, and related risk management strategies are typically adapted to historical ranges of variability (Kjeldsen and Prosdocimi 2016).

One possible way to assess how events may unfold is the use of a so-called ‘storyline’ approach; a storyline is typically defined as a “physically self-consistent unfolding of past events, or of plausible future events or pathways” (Shepherd et al. 2018). This approach has been advocated in the context of future climate change, as a

1 way of circumventing the deep uncertainties associated with future climate projections by placing more emphasis  
2 on driving factors and plausibility than probability (Shepherd et al. 2018). In this context it has links to so-called  
3 H++ scenarios, which are plausible but high-end scenarios of climate change (Reynard et al. 2017). But storylines  
4 can equally be applied in the context of past events by developing plausible counterfactuals, i.e. alternative ways  
5 that those events could have unfolded even in the current climate (Sillmann et al. 2021). For example, Chan et al.  
6 (2022) developed storylines for the UK drought of 2010-2012, by applying changes to the observed event based  
7 on i) antecedent conditions (applying progressively drier conditions), ii) temporal sequencing (adding a dry winter  
8 before or after the observed event), and iii) climate change. The results showed the importance of hydrological  
9 initial conditions, and the vulnerability of catchments in Britain to a ‘third dry winter’. Such studies can aid  
10 preparedness by enabling planning for events similar to, but more extreme than, known events (with known  
11 responses and impacts). However, the development of storylines in this way requires expert judgment on  
12 plausibility, and on the factors important to the development of a particular event. Possible spatial factors may  
13 also be neglected; for example Chan et al. (2022) apply the same storylines for catchments across Britain, treating  
14 catchments essentially independently.

15 The use of large ensembles of climate data can reduce the need for expert judgment and enable spatially consistent  
16 analyses and estimation of likelihoods. The extreme winter rainfall study of Thompson et al. (2017) was based on  
17 a large ensemble of high-resolution initialised global climate simulations (termed ‘UNSEEN’, UNprecedented  
18 Simulated Extremes using ENsembles), thus “directly sampling more extreme cases than the available  
19 observations, allowing the identification of unprecedented rainfall events to assess their likelihood in the real  
20 world”. Statistical modelling could also be used to estimate the probability of unprecedented rainfall from  
21 observations, and this is increasingly done in practice in UK water resources management using stochastic  
22 simulation (e.g. Dawkins et al. 2022). However, the use of a dynamical model is judged to better preserve physical  
23 plausibility and spatial dependence (Thompson et al. 2017). Data from either could be used to drive hydrological  
24 models, to enable subsequent assessment of potentially unprecedented hydrological extremes, but the likely better  
25 representation of spatial structures in dynamical models is important if large or multiple (not independent)  
26 catchments are being considered. Chan et al. (2023a) used a large ensemble of seasonal global model hindcast  
27 data to drive catchment-based hydrological models for 16 catchments (plus a groundwater model for 10 boreholes)  
28 in the Anglian region of England, and used the summer 2022 drought as a case study to explore plausible storylines  
29 of development into 2023. Brunner and Slater (2022) show that pooling reforecast ensemble members of European  
30 river flows enables more robust estimates of very extreme flood events (those occurring less than twice in 100  
31 years), with reduced uncertainty bounds compared to observation-based estimates.

32 Here, an expanded version of the UNSEEN ensemble of Thompson et al. (2017) is used in combination with a  
33 simple grid-based hydrological model for Great Britain, to assess unprecedented but plausible hydrological  
34 extremes in the current climate. For hydrological modelling, the antecedent conditions (e.g. water stored in the  
35 soil and groundwater) are an important factor in subsequent river flows. Thus two case studies are selected, one  
36 very dry (Summer 2022, associated with a major national-scale drought) and one very wet (Autumn 2023,  
37 associated with persistent and large-scale flooding), with the hydrological model initialised from known  
38 conditions at the start of each case study, then run forward for a number of months using the large ensemble of  
39 UNSEEN climate data. The case studies are used to illustrate the potential of the approach to provide  
40 unprecedented but plausible temporal and spatial hydrological extremes.

## 1 2 Data and methods

### 2 2.1 The UNSEEN climate datasets


3 The UNSEEN dataset used in this study comes from two hindcast ensembles of the Met Office Decadal Prediction  
4 System version 3 (DePreSys3; Dunstone et al. 2016), which is based on the Hadley Centre global coupled model  
5 (GCM) HadGEM3-GC3 (Williams et al. 2018). The model has an atmospheric grid resolution of ~60 km in the  
6 midlatitudes, and an ocean resolution of 0.25°. The first ensemble is initialised every 1st November and the second  
7 is initialised every 1st May, and each has forty ensemble members (realisations) but they are run for different  
8 lengths of time:

- 9 • 16-month periods starting in each November from 1959 up to 2021;
- 10 • 11-month periods starting in each May from 1960 up to 2022.

11 Differences between ensemble members are solely due to natural variability; i.e. there are no changes to GCM  
12 structure or parameterisation. Monthly rainfall rates (mm/day) are provided on a lat-long grid, for a region  
13 covering most of the UK (Figure 1a). The data for each realisation in an n-month period are considered dependent.

14 The GCM rainfall data are processed via the two steps below.

- 15 1. *Bias-corrected using simple monthly factors.* Biases in the temporal and spatial patterns of the GCM rainfall  
16 data are corrected as simply as possible, using monthly 60km grids of correction factors. These are derived  
17 for each month of the Nov-initialised and May-initialised data separately (i.e. there are 16 resulting grids of  
18 correction factors for the Nov-initialised runs and 11 for the May-initialised runs), by comparing the GCM  
19 monthly mean precipitation across all initialisation-years and realisations against monthly mean observed  
20 precipitation from HadUK-Grid 1km data for 1961-2020 (Met Office et al. 2021) averaged up to the GCM  
21 grid. The grids of correction factors are then smoothed using a 3x3 grid around each cell, with a weight of  
22 1/2 for the centre cell and 1/16 for each of the 8 surrounding cells (unless any surrounding cell has missing  
23 data, in which case its weight is added to that of the centre cell). A similar bias correction is derived for RCM  
24 precipitation data by Kay (2021). Maps of the correction factors are shown in Supp. Figures S1 and S2; these  
25 show that the GCM rainfall is typically too low (correction factor > 1) in Aug/Sep, but is too high in other  
26 months in some parts of the country.
- 27 2. *Downscaled to the 1km GB national grid.* The hydrological model (Section 2.2) requires 1km inputs, which  
28 are derived from the ~60km lat-lon data by identifying the GCM grid box to use for each 1km grid box (Figure  
29 1a) and distributing non-uniformly over the 1km grid boxes within each GCM grid box by using ratios of  
30 1km to GCM grid box-mean standard average annual rainfall (SAAR) (Bell et al. 2007; Kay et al. 2023b).  
31 The SAAR ratios are shown in Figure 1b; these show greater variation in ratios (and therefore in the  
32 downscaled rainfall data) within 60km grid boxes to the north/west, which are typically hillier, with much  
33 less variation in ratios within grid boxes in the flatter south/east. The ratios thus principally indicate  
34 topographic effects on spatial rainfall distribution, including rain-shadow effects.



1

2 **Figure 1 a) Identifying GCM grid boxes on the GB national grid. b) The SAAR ratios used for downscaling GCM**  
 3 **rainfall to the 1km grid. c) The 17 UK Hydrological Outlook regions, with the eight regions used here in the left-hand**  
 4 **list (green) and the rest in the right-hand list (blue).**

5 **2.2 The hydrological model**

6 The Grid-to-Grid (G2G) is a grid-based runoff-production and routing model, operating on a 1km grid at a 15-  
 7 minute time-step across Great Britain (Bell et al. 2009). It is used for operational flood forecasting for England,  
 8 Wales and Scotland (Price et al. 2012; Cranston et al. 2012), and has been used to estimate the potential future  
 9 impacts of climate change on river flows across Britain (e.g. Kay et al. 2023a). However, the short time-step of

1 the model (required for stability of the routing scheme given the 1km grid scale) means that runs take a not  
2 insignificant amount of time. For a seasonal forecasting application, where a relatively large number of runs were  
3 required using coarse (temporal and spatial) climate data, a simple monthly Water Balance Model (WBM) was  
4 developed, based upon G2G (Bell et al. 2013, 2017).

5 The WBM uses data from long historical runs of G2G (1km grids of the long-term means of monthly actual  
6 evaporation (AE), flow, and subsurface water storage) as well as information on the network of flow paths used  
7 by G2G, and is initialised using an estimate of subsurface water storage on a 1km grid across GB, also taken from  
8 G2G for the required date. The WBM forms one component of the UK Hydrological Outlook (UKHO;  
9 Prudhomme et al. 2017), where it is initialised using a G2G estimate of subsurface water storage derived using  
10 the most recent observations of rainfall and potential evaporation (PE), and then run forward driven by an  
11 ensemble of Met Office rainfall forecasts for 1- and 3-months ahead, to provide forecasts of regional mean river  
12 flows. The results are combined with those from a number of other forecasting approaches to produce a monthly  
13 hydrological outlook for the UK (hydoutuk.net, ukho.ceh.ac.uk).

14 Bell et al. (2017) provide an assessment of the performance of the WBM compared to G2G, driving both with  
15 observed 5km gridded rainfall data (1962-2010) and initialising the WBM from G2G at the start of each month.  
16 Regional means of standardised 1km river flows for 17 regions across GB (those in Figure 1b), for 1- and 3-  
17 months ahead, show correlations of over 0.8 in all cases.

### 18 **2.3 Applying the climate data – two case studies**

19 As described in Section 2.2, the WBM needs to be initialised using an estimate of subsurface water storage on a  
20 1km grid across GB, which is generally taken from a run of G2G driven by observed rainfall and PE data. Here,  
21 the WBM is initialised for two case study events;

- 22 1. Summer 2022 drought, and
- 23 2. Autumn 2023 flood.

24 For each case study, the G2G model is run with daily observed data up to the end of the previous month, to produce  
25 the initial conditions used by the monthly WBM. The observed data consists of 1km HadUK-Grid precipitation  
26 (Met Office et al. 2021) and MORECS PE (Hough and Jones 1997). Then the WBM is run forwards for a number  
27 of months using both UNSEEN ensembles for all realisations and all initialisation-years (1961-2022), regardless  
28 of the year in which the UNSEEN climate data was initialised (hereafter ‘WBM UNSEEN’, Table 1).

29 For each case study, the WBM is also run forwards for the same period using all years of historical HadUK-Grid  
30 1km precipitation data (1961-2022). not just the year corresponding with the case study (hereafter ‘WBM Obs’).  
31 This is analogous to the ‘Ensemble Streamflow Prediction’ method used as part of the UKHO (Harrigan et al.  
32 2018). where an ensemble of historical sequences of daily weather data is used to drive a lumped hydrological  
33 model from given initial conditions, for catchments across the UK.

34 The use of UNSEEN (and observed) data from all years allows the widest examination of possible subsequent  
35 rainfall pathways following summer 2022 and autumn 2023. The relatively long lead-times of the UNSEEN data  
36 used here (Table 1), and hence the relatively weak predictable UK rainfall signals, results in a large dataset with  
37 a wide range of possible outcomes including well-sampled dry (after summer 2022) and wet (after autumn 2023)

extreme tails. The WBM Obs ensemble provides a ‘benchmark’ that enables an assessment of the added-value of the WBM UNSEEN ensemble.

For droughts, which typically evolve relatively slowly due to accumulated rainfall deficits, the monthly time-step of the modelling chain is likely sufficient. Floods can develop and recede much more quickly, so a finer time-step would ideally be required (particularly for smaller/flashier catchments), but the monthly system can still give an indication of flood potential; this is discussed further in Section 4.2.

**Table 1** The time-periods covered by the Nov- and May-initialised UNSEEN climate ensembles, and their use in each case study. The grey boxes indicate UNSEEN data months regarded as spin-up. The green vertical lines indicate the WBM initialisation for each case study (end-July 2022 for the summer 2022 drought study and end-October 2023 for the Autumn 2023 flood study).

| Year                        | 1    |      | 2    |      |      |      |      |      |       |      |      |       | 3    |      |      |      |      |
|-----------------------------|------|------|------|------|------|------|------|------|-------|------|------|-------|------|------|------|------|------|
| Month                       | N    | D    | J    | F    | M    | A    | M    | J    | J     | A    | S    | O     | N    | D    | J    | F    | M    |
| <b>UNSEEN ensembles:</b>    |      |      |      |      |      |      |      |      |       |      |      |       |      |      |      |      |      |
| Nov-initialised             | Grey | Blue | Blue | Blue | Blue | Blue | Blue | Blue | Blue  | Blue | Blue | Blue  | Blue | Blue | Blue | Blue | Blue |
| May-initialised             |      |      |      |      |      |      | Grey | Blue | Blue  | Blue | Blue | Blue  | Blue | Blue | Blue | Blue | Blue |
| <b>Use in Case studies:</b> |      |      |      |      |      |      |      |      |       |      |      |       |      |      |      |      |      |
| <b>Summer 2022 Drought:</b> |      |      |      |      |      |      |      |      |       |      |      |       |      |      |      |      |      |
| Nov-initialised             |      |      |      |      |      |      |      |      | Green | Blue | Blue | Blue  | Blue | Blue | Blue | Blue | Blue |
| May-initialised             |      |      |      |      |      |      |      |      |       | Blue | Blue | Blue  | Blue | Blue | Blue | Blue | Blue |
| <b>Autumn 2023 Flood:</b>   |      |      |      |      |      |      |      |      |       |      |      |       |      |      |      |      |      |
| Nov-initialised             |      |      |      |      |      |      |      |      |       |      |      | Green | Blue | Blue | Blue | Blue | Blue |
| May-initialised             |      |      |      |      |      |      |      |      |       |      |      |       | Blue | Blue | Blue | Blue | Blue |

### 2.3.1 Summer 2022 drought

July 2022 was exceptionally warm and dry in the UK, with widespread record-breaking maximum temperatures and extremely low rainfall across England (<70% of average for the majority of the country, but <10% over large parts of the south/east including Anglia) (Barker et al. 2024, Hannaford et al. 2022). The preceding four months had also been dry (<90% of average across the whole country, with <70% in large areas, particularly to the south/east) (Sefton et al. 2022), leading to river flows and reservoirs reaching exceptionally low levels across much of England, particularly in catchments to the south, and the introduction of temporary use bans by several water companies in southern England (Barker et al. 2024, Hannaford et al. 2022).

We here ask the question ‘how much worse might the situation have become’? To do this, the WBM is initialised from G2G sub-surface conditions from end-July 2022 and run to end-Feb 2023 (7 months), with every UNSEEN initialisation-year and realisation, and with every observed data year. This gives 4960 WBM UNSEEN runs for each month (2 ensembles \* 62 initialisation-years \* 40 realisations), vs only 62 historical sequences for WBM Obs, hence enabling a more robust assessment of rare low extremes.

### 2.3.2 Autumn 2023 flood

October 2023 was exceptionally wet across most of England, with rainfall totals exceeding 150% of average widely (apart from the southwest and northwest) and exceeding 250% in some areas (particularly to the north/east)

1 (Hannaford et al. 2023). September had also been relatively wet (exceeding 110% of average across much of the  
2 country, apart from the far southeast and some limited pockets elsewhere, and exceeding 150% in some areas to  
3 the northwest, northeast and southwest) and, while August was near-average, July also saw exceptional rainfall  
4 across the country, leading to notable rainfall accumulations over this three-month period (Sefton et al. 2023a).  
5 This led to river flows reaching notably high levels across much of England, with elevated flood risk in many  
6 areas.

7 We again ask the question ‘how much worse might the situation become’? In a similar way to the Summer 2022  
8 analysis, the WBM is initialised from G2G sub-surface conditions from end-Oct 2023 and run to end-Feb 2024  
9 (4 months), with every UNSEEN initialisation-year and realisation, and with every observed data year. This again  
10 gives 4960 WBM UNSEEN runs for each month, vs only 62 for WBM Obs, hence enabling a more robust  
11 assessment of rare high extremes.

## 12 **2.4 Flow analyses**

13 The WBM provides monthly mean river flows on a 1km grid, but in the UKHO these are typically standardised  
14 (by dividing by long-term mean flow from a set of observation-driven runs) and the standardised flows are  
15 displayed on a 1km grid or averaged across 17 regions (Figure 1c). This WBM approach was originally developed  
16 to provide an indication of the relative magnitude of monthly and regional mean river flows across GB using  
17 spatially-coarse GloSEA5 rainfall forecasts (Bell et al. 2017). A very similar approach is used here; although the  
18 DePreSys3 hindcasts have an improved spatial resolution, they are still relatively coarse, and focusing on regional  
19 mean flows greatly simplifies subsequent analyses and plotting. The results focus on eight regions of England;  
20 NW England, Northumbria, Severn-Trent, Yorkshire, Thames, Anglian, Wessex, SE England (Figure 1c).

### 21 **2.4.1 Fidelity tests**

22 Fidelity tests on WBM flow estimates are performed in a similar way to that applied for the UNSEEN rainfall  
23 data (Thompson et al. 2017 Fig. 2). That is, for each case study and region,

- 24 • the WBM UNSEEN (May- and Nov-initialised) simulated flows are resampled 1000 times, with each sample  
25 randomly selected from the 40\*2 available realisations for the year, producing a time-series of the same length  
26 as the WBM Obs time-series;
- 27 • for each resample, the mean, standard deviation, skewness and kurtosis statistics are calculated;
- 28 • the same statistics are calculated for the WBM Obs simulated flows;
- 29 • if, for all four statistics, the WBM Obs flow value sits within the 2.5%-97.5% range of the values from the  
30 1000 resamples of the WBM UNSEEN simulations, then the WBM UNSEEN simulations are considered to  
31 have passed the test (i.e. the distributions of WBM Obs and WBM UNSEEN flow values for the region and  
32 month are considered indistinguishable). If the test is only failed for one of the four statistics, then this is  
33 noted.

34 Passing the fidelity tests gives confidence in the ability of the WBM and UNSEEN GCM precipitation to simulate  
35 appropriate distributions of monthly mean river flows in a region, given the WBM initialisation for each case  
36 study.



## 1    **2.4.2    Extreme flows**

2    For each case study, the monthly time-series of regional mean flows from the full set of years and realisations of  
3    WBM UNSEEN are plotted as the median and range (5<sup>th</sup>-95<sup>th</sup> percentiles, and overall min and max). The Nov-  
4    and May-initialised ensembles are combined together. For comparison, the median and range from WBM Obs are  
5    also shown.

6    For historical context, the time-series of monthly regional mean flows for the previous two years are shown for  
7    each region, along with long-term mean flow ranges derived for the preceding years (from 1963). These are  
8    derived from runs of the WBM driven by observed data for 1963-2023, with initialisation (using G2G data) at the  
9    start of every month (hereafter ‘WBM Obs 1m’). For each region, long-term mean flow ranges are derived by  
10    extracting the 5<sup>th</sup>, 13<sup>th</sup>, 28<sup>th</sup>, 72<sup>th</sup>, 87<sup>th</sup> and 95<sup>th</sup> percentiles for each month, along with overall min and max. These  
11    percentiles are chosen to match those defining the seven classes used in the UKHO: ‘Exceptionally low’ (<5<sup>th</sup>),  
12    ‘Notably low’ (5<sup>th</sup>-13<sup>th</sup>), ‘Below normal’ (13<sup>th</sup>-28<sup>th</sup>), ‘Normal’ (28<sup>th</sup>-72<sup>th</sup>), ‘Above normal’ (72<sup>th</sup>-87<sup>th</sup>), ‘Notably  
13    high’ (87<sup>th</sup>-95<sup>th</sup>), ‘Exceptionally high’ (>95<sup>th</sup>) ([ukho.ceh.ac.uk](http://ukho.ceh.ac.uk)).

14    The ‘WBM Obs 1m’ run described above is not directly comparable to the ‘WBM UNSEEN’ and ‘WBM Obs’  
15    ensemble runs, since the former is re-initialised from G2G for every month whereas the latter are only initialised  
16    at the start of each case study then run forward for the  $n$ -months of each study (where  $n$  is 7 for the Summer 2022  
17    drought and 4 for the Autumn 2023 flood). To demonstrate any difference this may make, a further run is done  
18    using observed driving data for each case study but only initialising at the start then running forward for  $n$  months  
19    (hereafter ‘WBM Obs n-m’).

## 20    **2.4.3    Temporal and spatial variation in extreme flows**

21    The UNSEEN-derived extreme flows are assessed, for each case study, by selecting and plotting the ensemble  
22    member giving the most extreme flows for each month of the simulation, and investigating how these vary  
23    temporally (for consecutive months) and spatially (for neighbouring regions). The UNSEEN ensemble member  
24    is defined by the ensemble (Nov or May), the initialisation-year, and the realisation number (out of 40); see  
25    Section 2.1.

## 26    **2.4.4    Recovery or persistence of extreme flows**

27    The long term mean flow bands (Section 2.4.2) are used to assess the chance of flows recovering to ‘normal’ by  
28    each month, in each region, for each case study. For example, for the Summer 2022 drought case study, in each  
29    region each ensemble member is assessed to see whether the flows have reached the ‘normal’ flow band (or  
30    higher) by month  $m$  but are still in a lower flow band for each month of the simulation prior to month  $m$ . Similarly,  
31    for the Autumn 2023 flood case study, in each region each ensemble member is assessed to see whether the flows  
32    have reached the ‘normal’ flow band (or lower) by month  $m$  but are still in a higher flow band for each month of  
33    the simulation prior to month  $m$ . The number of ensemble members recovering to ‘normal’ is then expressed as a  
34    percentage of the full ensemble, and plotted for each region and each case study. In a similar way, the chance of  
35    flows remaining at least exceptionally low (high) is assessed for Summer 2022 (Autumn 2023).



1 **3 Results**

2 **3.1 Fidelity tests**

3 The results of the flow fidelity tests are summarised in Table 2 for the each case study, indicating a pass ('1') or  
 4 fail ( $\leq 0$ ) for each month and region, with a negative number indicating which statistic the test failed on if it only  
 5 failed on one of the four. These show overall pass-rates of 86% and 78% for the Summer 2022 drought and  
 6 Autumn 2023 flood events respectively. Note that there are no failures only on the mean ('-1'), because of the  
 7 correction applied to GCM precipitation data for mean monthly rainfall (Section 2.1).

8 Most of the failures occur in February; the pass-rates leaving out February are 96% and 96% respectively. The  
 9 failures in February are only related to the standard deviation ('-2'), which is due to this being too low for rainfall  
 10 in the climate model runs (Kelder et al. 2022). This is partly due to the extremely wet Feb 2020 (Davies et al.  
 11 2021); removing this from the February fidelity testing leads to passes in 6 out of 8 regions for both the Summer  
 12 2022 drought and Autumn 2023 flood (compared to only 2 when Feb 2020 is included). In each case, the test still  
 13 fails in Thames and SE England. There are limited failures in months other than February, but notably for SE  
 14 England in August—the latter fails on both skewness and kurtosis which seems to be due to the WBM UNSEEN  
 15 ensemble providing more extreme high flows than the WBM Obs ensemble, but lower flows are better  
 16 represented.

17 **Table 2 The results of fidelity tests for the Summer 2022 drought (left) and Autumn 2023 flood (right) case studies. '1'**  
 18 **indicates a PASS for all four statistics, while  $\leq 0$  indicates a FAIL for at least one statistic. If there is only a FAIL on**  
 19 **one statistic then the negative number indicates which (-1 mean, -2 sd, -3 skewness, -4 kurtosis).**


|              | Summer 2022 drought |     |     |     |     |     |     | Autumn 2023 flood |     |     |     |
|--------------|---------------------|-----|-----|-----|-----|-----|-----|-------------------|-----|-----|-----|
|              | Aug                 | Sep | Oct | Nov | Dec | Jan | Feb | Nov               | Dec | Jan | Feb |
| NWEngland    | 1                   | 1   | 1   | 1   | 1   | 1   | -2  | 1                 | 1   | 1   | -2  |
| Northumbria  | 1                   | 1   | 1   | 1   | 1   | 1   | -2  | 1                 | -3  | 1   | -2  |
| Severn-Trent | 1                   | 1   | 1   | 1   | 1   | 1   | -2  | 1                 | 1   | 1   | -2  |
| Yorkshire    | 1                   | 1   | 1   | 1   | 1   | 1   | -2  | 1                 | 1   | 1   | -2  |
| Thames       | 1                   | 1   | 1   | 1   | 1   | 1   | -2  | 1                 | 1   | 1   | -2  |
| Anglian      | 1                   | 1   | -4  | 1   | 1   | 1   | 1   | 1                 | 1   | 1   | 1   |
| Wessex       | 1                   | 1   | 1   | 1   | 1   | 1   | 1   | 1                 | 1   | 1   | 1   |
| SEEngland    | 0                   | 1   | 1   | 1   | 1   | 1   | -2  | 1                 | 1   | 1   | -2  |

20


21 **3.2 Extreme flows**

22 Plots of the median and range of regional mean flows are shown in Figure 2 for the Summer 2022 drought case  
 23 study, and Figure 3 for the Autumn 2023 flood case study. These show that using the large ensemble of UNSEEN  
 24 data gives more extreme flows than using all of the historical observed data (red dotted lines vs grey dotted lines),  
 25 although the 5<sup>th</sup>, 50<sup>th</sup> and 95<sup>th</sup> percentiles from WBM UNSEEN and WBM Obs are all similar (orange dashed and  
 26 dot-dashed vs grey dashed and dot-dashed), as expected from the fidelity tests. It should be emphasised that, for  
 27 both WBM UNSEEN and WBM Obs, the min and max (dotted lines) represent the overall envelope of the  
 28 ensemble of simulations for each month, so they do not necessarily represent a plausible monthly evolution of the  
 29 flows (see Section 3.3). On average across the 8 regions, for the 7 months of the Summer 2022 drought case study

1 the low envelope of the WBM UNSEEN flows is 28% lower than that of the WBM Obs flows, and for the 4  
 2 months of the Autumn 2023 flood case study the high envelope of the WBM UNSEEN flows is 42% higher than  
 3 that of the WBM Obs flows.



4  
 5 **Figure 2 Regional mean flows for the Summer 2022 drought case study. The dashed/dot-dashed/dotted lines show the**  
 6 **median/5th-95<sup>th</sup>/min-max across the ensemble using all the observed driving data (grey) and all the UNSEEN driving**  
 7 **data (orange/red) for the WBM initialised from end-July 2022. The WBM flows driven by observed data for Jan 2021-**  
 8 **Feb 2023 and initialised at the start of every month are also shown (WBM Obs 1m; solid black line), as are the WBM**  
 9 **flows driven by observed data for Aug 2022-Feb 2023 but only initialised at the start (WBM Obs n-m; dashed black**  
 10 **line). For historical context, the coloured areas show the ranges from WBM Obs 1m for 1963-2020: min, 5<sup>th</sup>, 13<sup>th</sup>, 28<sup>th</sup>,**  
 11 **72<sup>th</sup>, 87<sup>th</sup>, 95<sup>th</sup>, max percentiles. Note the log-scale on the y-axis, to emphasise lower flows.**



1

2 **Figure 3 As Figure 2 but for the Autumn 2023 flood case study. For historical context, the coloured areas show the**  
 3 **ranges from WBM Obs 1m for 1963-2021: min, 5<sup>th</sup>, 13<sup>th</sup>, 28<sup>th</sup>, 72<sup>th</sup>, 87<sup>th</sup>, 95<sup>th</sup>, max percentiles.**

4 Figure 2 also shows that the simulations from WBM Obs 1m (black solid line; initialised at the start of every  
 5 month) and WBM Obs n-m (black dashed line; only initialised at the start of the case study period) are very  
 6 similar. However, Figure 3 shows that the WBM Obs n-m simulation tends to over-estimate high flows relative  
 7 to the WBM Obs 1m re-initialised simulation; see Section 4.2 of Discussion.

8 The effect of initialisation is clear in all regions, but longer-lasting in some. For example, for the Summer 2022  
 9 drought study the conditions in July 2022 are obviously very dry (below normal in all regions, and exceptionally  
 10 low in some). In Severn-Trent the max from WBM Obs (top grey dotted) has increased from below the long-term

1 max to match it by Feb 2023, whereas in Thames the WBM Obs max is still well below the long-term max by  
2 then. In all regions though, the min from WBM Obs (lower grey dotted) stays below the long-term min.


3 Similarly, for the Autumn 2023 flood study the conditions in October 2023 are very wet (above normal in all  
4 regions, and exceptionally high in some). In Northumbria the max from WBM Obs (top grey dotted) has decreased  
5 from above the long-term max to match it by Dec 2023, whereas in Anglian the WBM Obs max is still well above  
6 the long-term max even by Feb 2024. This is due to the presence of significant groundwater stores in regions to  
7 the south/east of England, which typically respond much more slowly to weather conditions than the shallower  
8 stores typically found to the north/west of England (Svensson et al. 2015). The WBM has information about the  
9 spatial differences in sub-surface stores from the data it takes from long historical runs of G2G (Section 2.2).

### 10 **3.3 Temporal and spatial variation in extreme flows**

11 It is important to note that the WBM UNSEEN extremes for consecutive months are often not given by the same  
12 UNSEEN climate ensemble member (ensemble, initialisation-year and realisation; Section 2.1). Similarly, the  
13 extremes for different regions, even neighbouring ones, are often not given by the same ensemble member. This  
14 is illustrated for four regions for the Summer 2022 drought (Figure 4) and for the Autumn 2023 flood (Figure 5).  
15 Similarly, the WBM Obs extremes for consecutive months will likely not be given by the same ensemble  
16 member.


17 For the Summer 2022 drought, Figure 4 shows that the ensemble members giving the lowest flows in August,  
18 September or October 2022 each have a fast recovery in flows through the autumn. In contrast, the ensemble  
19 members giving the lowest flows in late autumn and early winter 2022 are often low (but not lowest) earlier in  
20 the year too, particularly in the Thames and Anglian regions, which are more influenced by slower-responding  
21 groundwater systems. In no cases is the ensemble member giving the lowest flows for one month also that giving  
22 the lowest flows for the following month. In only three cases does the ensemble member giving the lowest flows  
23 in a given month and region also give the lowest flows for that month in a neighbouring region (Sep and Oct 2022  
24 in Thames and Anglian, and Jan 2023 in Thames and Severn-Trent).

25 For the Autumn 2023 flood, Figure 5 shows that in no cases is the ensemble member giving the highest flows for  
26 one month also that giving the highest flows for the following month, and only rarely is the ensemble member  
27 giving the highest flows in a given month also very extreme in earlier/later months. The only real exception is the  
28 ensemble member which gives the highest flows in Anglian in November 2023, which also gives flows higher  
29 than previous observed records (although not the highest from the UNSEEN ensemble) for Dec 2023 and Jan  
30 2024, falling back into the ‘exceptionally high’ range in Feb 2024. In only one case does the ensemble member  
31 giving the highest flows in a given month and region also give the highest flows for that month in a neighbouring  
32 region (Dec 2023 in Anglian and Severn-Trent).



1

2 **Figure 4 Regional mean flows for the Summer 2022 drought case study, for the ensemble member giving the lowest**  
 3 **flow in each month (coloured solid lines in order from Aug 2022 to Feb 2023), with labels identifying the ensemble**  
 4 **member by ensemble (Nov or May), initialisation-year and realisation number). As in Figure 2, the dashed/dotted lines**  
 5 **show the median/min-max across the ensemble using all the UNSEEN driving data (orange) for the WBM initialised**  
 6 **from end-July 2022, and the WBM flows driven by observed data for Jul 2022-Feb 2023 and initialised at the start of**  
 7 **every month are also shown (WBM Obs 1m; solid black line). For historical context, the coloured areas show the ranges**  
 8 **from WBM Obs 1m for 1963-2020: min, 5<sup>th</sup>, 13<sup>th</sup>, 28<sup>th</sup>, 72<sup>th</sup>, 87<sup>th</sup>, 95<sup>th</sup>, max percentiles.**



1


2 **Figure 5** As Figure 4 but **showing regional mean flows for the Autumn 2023 flood case study, for the ensemble member**  
 3 **giving the highest flow in each month (coloured solid lines in order from Nov 2023 to Feb 2024).** For historical context,  
 4 **the coloured areas show the ranges from WBM Obs 1m for 1963-2021: min, 5<sup>th</sup>, 13<sup>th</sup>, 28<sup>th</sup>, 72<sup>th</sup>, 87<sup>th</sup>, 95<sup>th</sup>, max**  
 5 **percentiles.**

6 These results also need to be interpreted in the context of the fidelity test results. In particular, in most regions the  
 7 fidelity tests were failed in February (Table 2), although this was mostly related to the extreme February observed  
 8 in 2020 so it may be that the results can otherwise be seen as representative. The fidelity tests were also failed in  
 9 the first month (August) of the Summer 2022 drought study in SE England, so those results may be less reliable.



1 **3.4 Recovery or persistence of extreme flows**

2 The chance of flows recovering to ‘normal’, or remaining exceptionally low (or lower), by each month in each  
 3 region for the Summer 2022 drought is shown in Figure 6. Flows in more northerly regions (NW England,  
 4 Northumbria, Yorkshire) were very likely to have recovered to ‘normal’ by early 2023, whereas other regions  
 5 show a slower recovery (from a lower starting point) — Anglian shows only around an 80% chance of recovery  
 6 by early 2023 (Figure 6a). Conversely, the analysis of persistence of extremes shows around a 3% chance of flows  
 7 remaining exceptionally low in early 2023 in the Anglian region, with a zero chance of persistent low flows in  
 8 northerly regions (Figure 6b).




10 **Figure 6 The percentage of ensemble members showing a) recovery of flows to (at least) normal by each month, or b)**  
 11 **remaining at least exceptionally extreme by each month, for the Summer 2022 drought case study.**

12 For the Autumn 2023 flood, flows in regions to the north are very likely to have recovered to ‘normal’ by early  
 13 2024, whereas flows in regions to the south/east show only around a 70-85% chance of recovery by then, with  
 14 Anglian and Severn Trent worst (Figure 7a). Conversely, the analysis of persistence of extremes shows around a



1 3% chance of flows remaining exceptionally high (or higher) in early 2023 in the Wessex region, with a zero  
 2 chance of persistent high flows in the northerly regions (Figure 7b).



3  
 4 **Figure 7 As Figure 6 but for the Autumn 2023 flood case study.**

5 For both case studies, there are differences between the monthly percentages of recovery and persistence estimated  
 6 from the WBM UNSEEN ensemble and the WBM Obs ensemble (coloured bars vs outlined bars). The percentages  
 7 derived from the WBM UNSEEN ensemble appear to vary more smoothly from month to month than those  
 8 derived from the much smaller WBM Obs ensemble, and the larger ensemble size means that the rarer persistent  
 9 extremes should be estimated more robustly. The general patterns for recovery to normal flows are relatively  
 10 similar, but the patterns for persistence of exceptionally extreme flows are more different. In some regions there  
 11 are persistent extremes in the WBM UNSEEN ensemble that do not exist at all in the WBM Obs ensemble (e.g.  
 12 Wessex for the Summer 2022 drought lasting into 2023, and Thames and Severn-Trent for the Autumn 2023 flood  
 13 lasting into 2024), but in other regions there are persistent extremes in the WBM Obs ensemble which do not exist  
 14 in the WBM UNSEEN ensemble (e.g. SE England for the Summer 2022 drought lasting into 2023, and Yorkshire  
 15 for the Autumn 2023 flood lasting into 2024).

## 4 Discussion

### 4.1 Case study outcomes

Both the flood and drought case studies demonstrate the potential of using the large ensemble of UNSEEN climate data in combination with a simple grid-based national-scale hydrological model. The modelling chain provides a large set of plausible events including extremes outside the range from use of observed data, with the lowest flows around 28% lower on average for the Summer 2022 drought study and the highest flows around 42% higher on average for the Autumn 2023 flood study. It enables the investigation of the temporal evolution and spatial dependence of extremes, including the potential time-scale and speed of recovery of flows to a normal range and possible persistence of extremes across a number of months.

For the Summer 2022 drought case study, Figure 4 showed the existence of ensemble members giving very low flows through to February 2023, particularly in the Thames and Anglian regions which are more influenced by slower-responding groundwater systems. In reality, recovery of regional mean flows to normal happened relatively quickly in autumn 2022 (black line), although there were localised drought concerns in parts of eastern England into summer 2023, and concerns about a further drought developing more widely in early 2023 following a dry winter (Sefton et al. 2023b); some ensemble members show this possibility (e.g. realisation 19 from Nov 1993 in Thames – blue line in Figure 4).

Chan et al. (2023a) also used the summer 2022 drought as a case study to explore storylines of development of extremes into 2023, but using a different seasonal hindcast dataset and only looking at selected catchments and boreholes in the Anglian region of England. They split their large ensemble into four clusters (based on atmospheric circulation indices), and showed that the clusters characterising drier-than-average winters resulted in continuation of the drought into 2023. This highlights a further advantage of the use of large ensembles of climate model data – the ability to characterise the large-scale drivers of extreme events.

For the Autumn 2023 flood case study, Figure 5 showed the ensemble member giving the highest flows in the Anglian region in Nov 2023 also gave flows higher than previous records for Dec 2023 and Jan 2024, before falling back into the ‘exceptionally high’ range in Feb 2024. Also, the ensemble member giving the highest flows in Dec 2023 in Anglian gave the highest flows in the neighbouring Severn-Trent region. In reality, after a wet December 2023 (Turner et al. 2024), severe flooding occurred over much of England very early in January 2024, with over 250 flood warnings issued by the Environment Agency and over 1000 properties flooded ([floodlist.com/europe/united-kingdom/storm-henk-floods-january-2024](https://floodlist.com/europe/united-kingdom/storm-henk-floods-january-2024)). Flows then dropped back to some extent later in January 2024, following drier weather across much of England (Sefton et al. 2024), before rising again in February 2024, which was exceptionally wet across much of England leading to record high flows in some catchments ([nrfa.ceh.ac.uk/sites/default/files/HS\\_202402.pdf](https://nrfa.ceh.ac.uk/sites/default/files/HS_202402.pdf)); some ensemble members show this possibility (e.g. realisation 11 from May 2020 in Anglian – dark red line in Figure 5).

For each case study, the selection of ensemble members illustrating temporal and spatial variation (Section 3.3) focussed on those that gave the most extreme flows for any given region and month. However, Section 3.4 included a summary of the percentage of ensemble members where flows remained at least exceptionally extreme by each month. This illustrated a clear possibility of persistent extremes in some regions to the south/east, with the chance of at least exceptionally low flows persisting from Summer 2022 into 2023 being ~3% in Anglian, and

1 the chance of at least exceptionally high flows persisting from Autumn 2023 into 2024 being ~3% for Wessex.  
2 The selection of ensemble members could also focus on any giving extreme low/high flows across multiple  
3 regions, to enable study of possible spatially extensive extreme events.

#### 4 **4.2 Limitations of the models and data**

5 The WBM applied here is a very simple monthly hydrological model, which has both advantages and  
6 disadvantages. The monthly time-step of the model means that it runs very quickly, so can easily be used for large  
7 climate ensembles such as those applied here. However, it also assumes climatological actual evaporation (AE,  
8 derived from long historical runs of G2G). The effect of this will likely be less in so-called water-limited areas to  
9 the south/east of England (where AE is generally limited by soil water availability, especially in summer) and in  
10 so-called energy-limited areas to the north/west of England (where AE is generally limited by potential  
11 evaporation), but it could have a larger effect in the more energy-water balanced areas in between (Kay et al.  
12 2013), and also during more extreme wet/dry periods when AE should probably be higher/lower. This could be  
13 the reason for the possible over-estimation of high flows from the WBM Obs n-m run, relative to the WBM Obs  
14 1m run which is re-initialised from G2G at the start of each month (Figure 3). Future work will investigate this  
15 possibility, and assess whether simple adjustments can be made to the WBM to improve simulation of extreme  
16 events.

17 The WBM also does not account for snow at all; although G2G has an optional snow module (Bell et al. 2016) it  
18 is not applied for the long historical runs used to provide data for the setup of the WBM, nor for the runs used for  
19 WBM initialisation. The lack of snow accounting will not have a significant effect for most regions of England  
20 most of the time, particularly given the monthly time-step of the WBM, as there are very few large and/or long-  
21 lasting accumulations of snow in most of England, although it is more important in parts of Scotland (Kay 2016).

22 The monthly time-step of the WBM, and the resolution of the UNSEEN data, is likely sufficient for investigating  
23 extreme low flows/droughts, which typically evolve relatively slowly due to rainfall deficits over extended periods  
24 of time. For extreme high flows/floods, a finer time-step would really be required as they can develop and recede  
25 much more quickly, particularly for smaller or flashier catchments. Despite this, the WBM can give a good  
26 indication of flood potential for most of England, because the main driver of floods here is soil moisture excess  
27 rather than extreme precipitation or snowmelt (Berghuijs et al. 2019). One potential approach could be to use a  
28 simple and fast-running model, like the WBM, to run a large ensemble, then select individual members of interest  
29 based on the outcomes from those runs. A much smaller set of runs of a more detailed model, like G2G, could  
30 then be performed using the particular members of interest to gain the extra temporal (and spatial) detail, provided  
31 (at least) daily precipitation data were available. Only monthly rainfall totals are available for the UNSEEN  
32 ensembles applied here, but future options will be investigated.

33 A simple bias correction was applied to the UNSEEN climate data before use to drive the hydrological model  
34 (Section 2.1). Applying this correction improved the results of the fidelity testing (not shown), although February  
35 in particular still has issues (related to the standard deviation of February rainfall being too low; Section 3.1).  
36 Bias-correction was also applied to climate data prior to use for hydrological modelling by Chan et al. (2023a,b).  
37 Brunner and Slater (2022) apply a bias correction to simulated river flows, but highlight that, compared to

1 observation-based estimates of extreme floods, their method can “introduce biases arising from the simulated  
2 meteorology and hydrological model”.

### 3 **5 Conclusions**

4 The UNSEEN climate datasets provide a large ensemble of alternative historical climate realisations, allowing the  
5 direct sampling of more extreme meteorological events than the available observations, and a better assessment  
6 of the likelihood of events (Thompson et al. 2017; Kelder et al. 2020). When combined with a simple hydrological  
7 model, this similarly allows the direct sampling of more extreme hydrological events and better assessment of  
8 likelihood. Both are conditional on a demonstration of fidelity for the event of interest.

9 An important issue for the hydrological modelling component is antecedent conditions. Here, two recent periods,  
10 one very dry and one very wet, were selected as case studies to initialise and run the simple hydrological model  
11 with the large ensemble of UNSEEN climate data. These case studies illustrate the potential of the approach to  
12 assess the temporal evolution and spatial dependence of unprecedented but plausible hydrological extremes.  
13 Clearly other periods could be similarly simulated and investigated, for example the summer 1976 drought, which  
14 was one of the most extreme and extensive meteorological and hydrological droughts in recent history (Rudd et  
15 al. 2017), and the widespread flooding of winter 2013/14, which was the wettest winter in Britain since records  
16 began (Kay et al. 2018). The method could also be applied to other countries/regions, with an appropriate  
17 hydrological model and using the same or similar global climate ensemble data.

18 Future work could include analysing the large-scale atmospheric drivers of selected hydrological extremes,  
19 whether a record extreme for an individual month or persistently extreme for a number of months, which could  
20 improve understanding of extreme events and their evolution. More detailed hydrological modelling, with (at  
21 least) daily precipitation data, would ideally be used for flood case studies, when results could also be investigated  
22 at a finer spatial scale, and additional hydrodynamic modelling (or pre-modelled design floods) could then provide  
23 information on flood extents and impacts (e.g. Kay et al. 2018). Similarly, additional water resource system  
24 modelling could provide information on drought impacts (e.g. Borgomeo et al. 2014). Soil moisture extremes  
25 could also be investigated, with consequent impacts for agriculture as well as a range of natural hazards (e.g. Kay  
26 et al. 2022), and other variables like groundwater levels could be investigated in a similar framework.

27 Being able to plan for unprecedented but plausible hydrological extremes is important in terms of improving the  
28 resilience of water supply systems to drought (Chan et al. 2023a), and improving flood risk management and  
29 incident response (Brunner and Slater 2022; Ganapathy et al. 2024). The UK water industry now has a statutory  
30 obligation to demonstrate resilience to droughts that are more extreme than those recorded in the past, including  
31 very rare events (e.g. 1:200 and 1:500 year) (Counsell and Durant, 2023). Similarly, for fluvial flood risk  
32 management there is statutory requirement to plan for very extreme (rare) events that may lie outside observational  
33 envelopes. In practice this is achieved through stochastic methods (e.g. using weather generators to produce long  
34 precipitation series that are then run through hydrological and supply system models) or statistical methods  
35 (pooling flood events from many catchments, or Probable Maximum Precipitation/Flood analyses). The UNSEEN  
36 modelling chain described here provides a physically-informed alternative to complement these primarily  
37 statistical approaches, with potential for use in both long-term water resource/flood risk planning and emergency  
38 drought/flood response contexts.

1 The use of methods such as those presented here, deriving unprecedented events from historical case studies, can  
2 aid preparedness by enabling planning for events similar to, but more extreme than, known events (with known  
3 responses and impacts). Increasing resilience to potential extremes in the current climate will also provide some  
4 resilience to the effects of climate change, which is expected to increase both floods and droughts in future in the  
5 UK (e.g. Lane and Kay 2021; Kay et al. 2021; Rudd et al. 2019, 2023).

6 **Data availability.** The data are available from the authors upon reasonable request.

7 **Author contribution.** Conceptualisation and methodology: all authors. Formal analysis and visualisation: AK.  
8 Funding acquisition: JH. Writing – original draft: AK. Writing – review and editing: all authors.

9 **Competing interests.** The authors declare that they have no conflict of interest.

10 **Acknowledgements.** This work was supported by the Natural Environment Research Council award number  
11 NE/X019063/1 (Hydro-JULES). Thanks to UKCEH colleague Wilson Chan for comments on the draft  
12 manuscript.

### 13 **References**

- 14 Barker, L.J., Hannaford, J., Magee, E., Turner, S., Sefton, C., Parry, S., Evans, J., Szczykulska, M., Haxton, T.  
15 (2024). An appraisal of the severity of the 2022 drought and its impacts. *Weather*, doi:10.1002/wea.4531.
- 16 Beever, L., Popescu, I., Pregolato, M., Liu, Y., Wright, N. (2022). Identifying hotspots of hydro-hazards under  
17 global change: A worldwide review. *Front. Water*, 4, 879536, doi: 10.3389/frwa.2022.879536.
- 18 Bell, V.A., Kay, A.L., Jones, R.G., Moore, R.J. (2007). Development of a high resolution grid-based river flow  
19 model for use with regional climate model output. *Hydrology and Earth System Sciences*, 11(1), 532–549.
- 20 Bell, V.A., Kay, A.L., Jones, R.G., Moore, R.J., Reynard, N.S. (2009). Use of soil data in a grid-based  
21 hydrological model to estimate spatial variation in changing flood risk across the UK. *Journal of Hydrology*,  
22 377(3–4), 335–350, doi:10.1016/j.jhydrol.2009.08.031.
- 23 Bell, V.A., Davies, H.N., Kay, A.L., Marsh, T.J., Brookshaw, A., Jenkins A. (2013). Developing a large-scale  
24 water-balance approach to seasonal forecasting: application to the 2012 drought in Britain. *Hydrological  
25 Processes*, 27(20), 3003–3012. doi:10.1002/hyp.9863.
- 26 Bell, V.A., Kay, A.L., Davies, H.N., Jones, R.G. (2016). An assessment of the possible impacts of climate change  
27 on snow and peak river flows across Britain. *Climatic Change*, 136(3), 539–553, doi:10.1007/s10584-016-1637-  
28 x.
- 29 Bell, V.A., Davies, H.N., Kay, A.L., Brookshaw, A., Scaife, A.A. (2017). A national-scale seasonal hydrological  
30 forecast system: development and evaluation over Britain. *Hydrol. Earth Syst. Sci.*, 21, 4681–4691.
- 31 Berghuijs, W.R., Harrigan, S., Molnar, P., Slater, L.J., Kirchner, J.W. (2019). The relative importance of different  
32 flood-generating mechanisms across Europe. *Water Resources Research*, 55, 4582–4593,  
33 doi:10.1029/2019WR024841.
- 34 Bertola, M., Blöschl, G., Bohac, M. et al. (2023). Megafloods in Europe can be anticipated from observations in  
35 hydrologically similar catchments. *Nat. Geosci.* 16, 982–988.
- 36 Borgomeo, E., Hall, J.W., Fung, F., Watts, G., Colquhoun, K., Lambert, C. (2014). Risk-based water resources  
37 planning: Incorporating probabilistic nonstationary climate uncertainties. *Water Resour. Res.*, 50(8), 6850–6873.
- 38 Brunner, M.I., Slater, L.J. (2022). Extreme floods in Europe: going beyond observations using reforecast ensemble  
39 pooling. *Hydrol. Earth Syst. Sci.*, 26, 469–482.

- 1 Chan, W.C.H., Shepherd, T.G., Facer-Childs, K., Darch, G., Arnell, N.W. (2022). Storylines of UK drought based  
2 on the 2010–2012 event. *Hydrol Earth Syst Sci* 26, 1755–1777. doi:10.5194/hess-26-1755-2022.
- 3 Chan, W.C.H., Arnell, N.W., Darch, G., Facer-Childs, K., Shepherd, T.G., Tanguy, M. (2023a). Added value of  
4 seasonal hindcasts for UK hydrological drought outlook. *Nat. Hazards Earth Syst. Sci. Discuss.*,  
5 doi:10.5194/nhess-2023-74.
- 6 Chan, W.C.H., Arnell, N.W., Darch, G., Facer-Childs, K., Shepherd, T.G., Tanguy, M., van der Wiel, K. (2023b).  
7 Current and future risk of unprecedented hydrological droughts in Great Britain, *Journal of Hydrology*,  
8 doi:10.1016/j.jhydrol.2023.130074.
- 9 Counsell, C., Durant, M. (2023). Water supply: observed and projected. In: Environment Agency, 2023. Annex  
10 to the review of the research and scientific understanding of drought. Environment Agency, Bristol. 669pp.  
11 <https://www.gov.uk/government/publications/review-of-the-research-and-scientific-understanding-of-drought>
- 12 Cranston, M., Maxey, R., Tavendale, A., Buchanan, P., Motion, A., Cole, S., Robson, A., Moore, R.J., Minett, A.  
13 (2012). Countrywide flood forecasting in Scotland: challenges for hydrometeorological model uncertainty and  
14 prediction. In: *Weather Radar and Hydrology* (ed. by R.J Moore, S.J. Cole & A.J. Illingworth) (Proc. Exeter  
15 Symp., April 2011), IAHS Publ. no. 351, 538-543.
- 16 Davies, P.A., McCarthy, M., Christidis, N., Dunstone, N., Fereday, D., Kendon, M., Knight, J.R., Scaife, A.A.,  
17 Sexton, D. (2021). The wet and stormy UK winter of 2019/2020. *Weather*, 76(12), 396-402.
- 18 Dawkins, L.C., Osborne, J.M., Economou, T., Darch, G.J.C., Stoner, O.R. (2022). The Advanced Meteorology  
19 Explorer: a novel stochastic, gridded daily rainfall generator. *Journal of Hydrology*, 607, 127478.
- 20 Dunstone, N., Smith, D., Scaife, A. et al. Skilful predictions of the winter North Atlantic Oscillation one year  
21 ahead. *Nature Geosci* 9, 809–814 (2016), doi:10.1038/ngeo2824.
- 22 Ganapathy, A., Hannah, D.M., Agarwal, A. (2024). Improved estimation of extreme floods with data pooling and  
23 mixed probability distribution. *Journal of Hydrology*, 629, 130633. Hannaford, J., Suman, G., Howson, T.,  
24 Bloomfield, J., Clemas, S. (2023). Hydrological summary for the United Kingdom: October 2023. Wallingford,  
25 UK, UK Centre for Ecology & Hydrology, 12pp.
- 26 Hannaford, J., Turner, S., Alton, J., Lewis, M., Clemas, S. (2022). Hydrological summary for the United  
27 Kingdom: July 2022. Wallingford, UK, UK Centre for Ecology & Hydrology, 12pp.
- 28 Harrigan, S., Prudhomme, C., Parry, S., Smith, K., Tanguy, M. (2018). Benchmarking ensemble streamflow  
29 prediction skill in the UK. *Hydrol. Earth Syst. Sci.*, 22, 2023–2039.
- 30 Hough, M.N., Jones, R.J.A. (1997) The United Kingdom Meteorological Office rainfall and evaporation  
31 calculation system: MORECS version 2.0-an overview, *Hydrol. Earth Syst. Sci.*, 1, 227–239, doi:10.5194/hess-  
32 1-227-1997.
- 33 Kay, A.L. (2016). A review of snow in Britain: the historical picture and future projections. *Progress in Physical*  
34 *Geography*, 40(5) 676–698, doi:10.1177/0309133316650617.
- 35 Kay, A.L. (2021). Simulation of river flow in Britain under climate change: baseline performance and future  
36 seasonal changes. *Hydrological Processes*, 35(4), e14137, doi:10.1002/hyp.14137.
- 37 Kay, A.L., Bell, V.A., Blyth, E.M., Crooks, S.M., Davies, H.N., Reynard, N.S. (2013). A hydrological perspective  
38 on evaporation: historical trends and future projections in Britain. *Journal of Water and Climate Change*, 4(3),  
39 193–208, doi:10.2166/wcc.2013.014.
- 40 Kay, A.L., Booth, N., Lamb R., Raven, E. Schaller, N., Sparrow, S. (2018). Flood event attribution and damage  
41 estimation using national-scale grid-based modelling: Winter 2013/14 in Great Britain. *International Journal of*  
42 *Climatology*, 38(14), 5205-5219, doi:10.1002/joc5721.
- 43 Kay, A.L., Davies, H.N., Lane, R.A., Rudd, A.C., Bell, V.A. (2021). Grid-based simulation of river flows in  
44 Northern Ireland: model performance and future flow changes. *Journal of Hydrology: Regional Studies*, 38,  
45 100967, doi:10.1016/j.ejrh.2021.100967.

- 1 Kay, A.L., Lane, R.A., Bell, V.A. (2022). Grid-based simulation of soil moisture in the UK: future changes in  
2 extremes and wetting and drying dates. *Environmental Research Letters*, 17(7), 074029, doi:10.1088/1748-  
3 9326/ac7a4e.
- 4 Kay, A.L., Bell, V.A., Davies, H.N., Lane, R.A., Rudd, A.C. (2023a). The UKSCAPE-G2G river flow and soil  
5 moisture datasets: Grid-to-Grid model estimates for the UK for historical and potential future climates. *Earth Syst.*  
6 *Sci. Data*, 15, 2533–2546, doi:10.5194/essd-15-2533-2023.
- 7 Kay, A.L., Rudd, A.C. and Coulson, J. (2023b). Spatial downscaling of precipitation for hydrological modelling:  
8 assessing a simple method and its application under climate change in Britain. *Hydrological Processes*, 37(2),  
9 e14823, doi:10.1002/hyp.14823.
- 10 Kelder, T., Müller, M., Slater, L.J., Marjoribanks, T.I., Wilby, R.L., Prudhomme, C., Bohlinger, P., Ferranti, L.,  
11 Nipen, T. (2020). Using UNSEEN trends to detect decadal changes in 100-year precipitation extremes. *Climate*  
12 *and Atmospheric Science*, 3, 47.
- 13 Kelder, T., Marjoribanks, T.I., Slater, L.J., Prudhomme, C., Wilby, R.L., Wagemann, J., Dunstone, N. (2022). An  
14 open workflow to gain insights about low-likelihood high-impact weather events from initialized predictions.  
15 *Meteorological Applications*, 29(3), e2065. doi:10.1002/met.2065.
- 16 Kent, C., Dunstone, N., Tucker, S., Scaife, A.A., Brown, S., Kendon, E.J., Smith, D., McLean, L., Greenwood, S.  
17 (2022). Estimating unprecedented extremes in UK summer daily rainfall. *Environ. Res. Lett.*, 17, 014041.
- 18 Kjeldsen, T.R. and Prosdocimi, I. (2016). Assessing the element of surprise of record-breaking flood events.  
19 *Journal of Flood Risk Management*, 11(51), S541-S553.
- 20 Lane, R.A., Kay, A.L. (2021). Climate change impact on the magnitude and timing of hydrological extremes  
21 across Great Britain. *Frontiers in Water*, 3, 684982, doi:10.3389/frwa.2021.684982.
- 22 Met Office, Hollis, D., McCarthy, M., Kendon, M., Legg, T., Simpson, I. (2021). HadUK-Grid gridded and  
23 regional average climate observations for the UK.  
24 <http://catalogue.ceda.ac.uk/uuid/4dc8450d889a491ebb20e724debe2dfb>.
- 25 Mokhtari, M., Faridi, P., Masoodi, M., Mehran Ahmadi, S. (2023). Perspective Chapter: A Global View of Natural  
26 Hazards Related Disasters [Internet]. *Natural Hazards - New Insights*. IntechOpen.  
27 doi:10.5772/intechopen.111582.
- 28 Price, D., Pilling, C., Robbins, G., Lane, A., Boyce, G., Fenwick, K., Moore, R.J., Coles, J., Harrison, T., Van  
29 Dijk, M. (2012). Representing the spatial variability of rainfall for input to the G2G distributed flood forecasting  
30 model: operational experience from the Flood Forecasting Centre. In: Moore, R.J.; Cole, S.J.; Illingworth, A.J.,  
31 (eds.) *Weather Radar and Hydrology*, Proc. Exeter Symp., April 2011. International Association of Hydrological  
32 Sciences, 532-537. (IAHS Publ., 351).
- 33 Prudhomme, C., Hannaford, J., Harrigan, S., Boorman, D., Knight, J., Bell, V., Jackson, C., Svensson, C., Parry,  
34 S., Bachiller-Jareno, N., Davies, H., Davis, R., Mackay, J., McKenzie, A., Rudd, A., Smith, K., Bloomfield, J.,  
35 Ward, R., Jenkins, A. (2017). Hydrological Outlook UK: an operational streamflow and groundwater level  
36 forecasting system at monthly to seasonal time scales, *Hydrological Sciences Journal*, 62:16, 2753-2768, doi:  
37 10.1080/02626667.2017.1395032
- 38 Reynard, N.S., Kay, A.L., Anderson, M., Donovan, B., Duckworth, C. (2017). The evolution of climate change  
39 guidance for fluvial flood risk management in England. *Progress in Physical Geography*, 41(2), 222–237,  
40 doi:10.1177/0309133317702566.
- 41 Rudd, A.C., Bell, V.A., Kay, A.L. (2017). National-scale analysis of simulated hydrological droughts (1891-  
42 2015). *Journal of Hydrology*, 550, 368-385.
- 43 Rudd, A.C., Kay, A.L., Bell, V.A. (2019). National-scale analysis of future river flow and soil moisture droughts:  
44 potential changes in drought characteristics. *Climatic Change*, 156(3), 323–340, doi:10.1007/s10584-019-02528-  
45 0.



- 1 Rudd, A.C., Kay, A.L., Sayers, P.B. (2023). Climate change impacts on flood peaks in Britain for a range of global  
2 mean surface temperature changes. *Journal of Flood Risk Management*, 16(1), e12863, doi:10.1111/jfr3.12863.
- 3 [Sefton, C., Barker, L., Lewis, M., Clemas, S. \(2022\) Hydrological summary for the United Kingdom: June 2022.](#)  
4 [Wallingford, UK, UK Centre for Ecology & Hydrology, 12pp.](#)
- 5 [Sefton, C., Magee, E., Howson, T., Crane, E., Clemas, S. \(2023a\). Hydrological summary for the United](#)  
6 [Kingdom: September 2023. Wallingford, UK, UK Centre for Ecology & Hydrology, 12pp.](#)
- 7 Sefton, C., Suman, G., Smith, R., Crane, E., Clemas, S. (2023**b**). Hydrological summary for the United Kingdom:  
8 February 2023. Wallingford, UK, UK Centre for Ecology & Hydrology, 12pp.
- 9 Sefton, C., Kumar, A., Smith, R., Bloomfield, J., Clemas, S. (2024) Hydrological summary for the United  
10 Kingdom: January 2024. Wallingford, UK, UK Centre for Ecology & Hydrology, 12pp.
- 11 Shepherd, T.G., Boyd, E., Calel, R.A. et al. (2018). Storylines: an alternative approach to representing uncertainty  
12 in physical aspects of climate change. *Climatic Change* 151, 555–571, doi:10.1007/s10584-018-2317-9
- 13 Sillmann, J., Shepherd, T.G., van den Hurk, B., Hazeleger, W., Martius, O., Slingo, J., Zscheischler, J. (2021).  
14 Event-based storylines to address climate risk. *Earth's Future*, 9, e2020EF001783.
- 15 Svensson, C., Brookshaw, A., Scaife, A.A., Bell, V.A., Mackay, J.D., Jackson, C.R., Hannaford, J., Davies, H.N.,  
16 Arribas, A., Stanley, S. (2015). Long-range forecasts of UK winter hydrology. *Environ. Res. Lett.*, 10(6), 064006.
- 17 Thompson, V., Dunstone, N.J., Scaife, A.A., Smith, D.M., Slingo, J.M., Brown, S., Belcher, S.E. (2017). High  
18 risk of unprecedented UK rainfall in the current climate. *Nature Communications*, 8,107, doi:10.1038/s41467-  
19 017-00275-3.
- 20 Turner, S., Magee, E., Howson, T., Crane, E., Clemas, S. (2024). Hydrological summary for the United Kingdom:  
21 December 2023. Wallingford, UK, UK Centre for Ecology & Hydrology, 12pp.
- 22 Williams, K. D., Harris, C. M., Bodas-Salcedo, A., Camp, J., Comer, R. E., Copsey, D., Fereday, D., Graham, T.,  
23 Hill, R., Hinton, T., Hyder, P., Ineson, S., Masato, G., Milton, S. F., Roberts, M. J., Rowell, D. P., Sanchez, C.,  
24 Shelly, A., Sinha, B., Walters, D. N., West, A., Woollings, T., Xavier, P. K.: The Met Office Global Coupled  
25 model 2.0 (GC2) configuration, *Geosci. Model Dev.*, 8, 1509–1524, doi:10.5194/gmd-8-1509-2015, 2015.